# On the Conductivity of a Magnetoactive Turbulent Plasma.

Otto G.Chkhetiani<sup>∗</sup>

Space Research Institute, Russian Academy of Sciences, Profsoyuznaya ul. 84/32, Moscow 117997, Russia

#### Abstract

The problem of determining the effective conductivity tensor of a magnetoactive turbulent plasma is considered in the approximation of isolated particles. Additional gyrotropicterms are shown to appear in the conductivity tensor in the presence of mean, nonzero magnetic helicity. The dispersion of propagating electro- magnetic waves changes, additional modes and additional rotation of the polarization plane appear, and the waves can be amplified. The properties acquired by plasma with helicity are similar those observed in chiral and bianisotropic electrodynamic media.

## 1 INTRODUCTION

An effective description of the propagation of waves and particles in fluctuational magnetic fields in a turbulent conductive medium is of great importance in solving the various problems of plasma physics and astrophysics. The phenomena associated with the presence of smallscale magnetic helicity  $\langle AB \rangle$  ( $B = \text{rot}A$ ), which manifest themselves virtually on all scales of plasma systems, play a special role here. Whereas large-scale helicity contributes to the stability of electromagnetic structures [1], its presence at the level of fluctuations is a nonequilibrium phenomenon that is accompanied by various large-scale instabilities [2]. Other effects produced by small-scale helicity, such as an asymmetry in the particle distribution and acceleration, are well known in the diffusion theory of cosmicray propagation [3–6]. The gyrotropic acceleration effects are also known in a laboratory plasma as helicity input effects [7, 8]. The appearance of additional helicity-related transport was also shown to be possible in [9]. Changes in the transport properties are also directly reflected in the dielectric (conductive) properties of a plasma medium. Thus, for example, it was shown in [10] that in the presence of fluctuational magnetic helicity in the low-conductivity limit, the effective current in an isotropic plasma proves to be dependent on the curl of the electric field (  $\mathbf{j} = \sigma \mathbf{E} + \sigma_{\kappa} \text{rot} \mathbf{E}$ ), which causes the mean magnetic field to grow under certain conditions. In [10], the external magnetic field was disregarded. In natural and laboratory conditions, the plasma is always under the influence of large-scale magnetic fields that affect significantly its properties [11]. Since magnetic helicity also emerges in plasma systems in the presence of a large-scale magnetic field, a study of its influence should take into account this factor. In [12], it was shown for an exactly solvable

<sup>∗</sup>ochkheti@mx.iki.rssi.ru

model of nonlinear dynamo that the diffusion and generation rate are strongly suppressed even in a relatively weak magnetic field, and the regime of fast dynamo transforms into the regime of slow dynamo with a linear growth with time. The goal of this work is to study the effective conductivity of a turbulent magnetoactive plasma with nonzero magnetic helicity. The kinetic approach is commonly used for a thorough theoretical description of plasma problems. However, an allowance for the fluctuational effects of gyrotropy is rather difficult to make and is possible in finished form only with an appreciable number of assumptions and simplifications (see, e.g., [6, 9]). At the same time, many basic plasma properties can be determined in the approximation of isolated particles [11, 13], which will be used below. The statistical parameters of the electromagnetic fluctuations are assumed to be stationary and uniform. In Section 2, we consider the equations of motion for particles and calculate the effective Lorentz force by the functional method with an allowance made for the nonuniformity of the electromagnetic perturbations to within the first order of the perturbation theory. In Section 3, we determine the effective conductivity tensor. Fluctuational magnetic helicity gives rise to new gyrotropic terms. Our analysis of the dispersion relation both in the approximation of  $\delta$ -correlated (in time) fluctuations (Section 4) and in the opposite case of long correlation times and high frequencies (Section 6) for electromagnetic waves and the evolution of the magnetic field (in the low-frequency limit) (Section 5) reveals changes in the dispersion of propagating waves and the presence of instabilities. The characteristic scales and growth rates of the instabilities are determined by the relationship between the fluctuational helicity and the energy and the external magnetic field. A magnetoactive turbulent plasma with helicity acquires properties similar to those of chiral and bianisotropic electrodynamic media, which have been extensively studied in recent years [14, 15]. In Conclusions, we discuss our results and implications of the detected effect.

## 2 BASIC EQUATIONS

Let us consider the motion of a one-component, singly charged plasma in a fluctuational electromagnetic field with given correlation properties. We will consider a cold plasma where the approximation of isolated particles [11, 13] can be used. A regular large-scale nonuniform perturbation of the electromagnetic field that is too weak to change significantly the correlation properties of the electromagnetic fluctuations, which are supposed to be given, stationary, and uniform, is assumed to arise in the system. The expression for the electron velocity  $\bf{v}$  can be written as

$$
\frac{d\mathbf{v}}{dt} = \frac{e}{m} \left( \mathbf{E} + \frac{1}{c} \left[ \mathbf{v} \times \mathbf{B} \right] \right). \tag{1}
$$

where  $e$  and  $m$  are the electron charge and mass, respectively. The electromagnetic field and the velocity can be represented as a sum of the large-scale slow component and the small-scale (with a zero mean) fast component:

$$
\mathbf{E} = \langle \mathbf{E} \rangle + \widetilde{\mathbf{E}}, \quad \mathbf{B} = \mathbf{B}_0 + \langle \mathbf{B} \rangle + \widetilde{\mathbf{B}}, \ \mathbf{v} = \langle \mathbf{v} \rangle + \widetilde{\mathbf{v}}.
$$

As was said above, the mean electric and magnetic fields are assumed to be weak compared to the fluctuational fields, i.e.,  $\langle E \rangle \ll \left\langle \widetilde{E}^2 \right\rangle^{1/2}$ ,  $\langle B \rangle \ll \left\langle \widetilde{B}^2 \right\rangle^{1/2} < B_0$ . Passing to the Fourier representation,  $(F(\mathbf{x}, t) = \int \int \widehat{\mathbf{F}}(\mathbf{k}, w) \exp[i(\mathbf{kx} - wt)] d\mathbf{k} dw)$ , we write

$$
-iw\widehat{\mathbf{v}}(\mathbf{k},w) - \frac{e}{mc}[\widehat{\mathbf{v}}(\mathbf{k},w) \times \mathbf{B}_0] = \frac{e}{m}\widehat{\mathbf{E}}(\mathbf{k},w) + \frac{e}{mc}\int \left[\widehat{\mathbf{v}}(\mathbf{q},s) \times \widehat{\mathbf{B}}(\mathbf{k}-\mathbf{q},w-s)\right]d\mathbf{q}ds. (2)
$$

The equation of motion averaged over the electromagnetic fluctuations takes the form

$$
-iw\langle \widehat{\mathbf{v}}(\mathbf{k},w)\rangle - \frac{e}{mc}[\langle \widehat{\mathbf{v}}(\mathbf{k},w)\rangle \times \mathbf{B}_0] = \frac{e}{m}\langle \widehat{\mathbf{E}}(\mathbf{k},w)\rangle + \frac{e}{mc}\int \left[\langle \widehat{\mathbf{v}}(\mathbf{q},s)\rangle \times \langle \widehat{\mathbf{B}}(\mathbf{k}-\mathbf{q},w-s)\rangle\right]d\mathbf{q}ds + \frac{e}{mc}\int \left[\langle \widehat{\mathbf{v}}(\mathbf{q},s)\times \widehat{\mathbf{B}}(\mathbf{k}-\mathbf{q},w-s)\rangle\right]d\mathbf{q}ds.
$$
\n(3)

<span id="page-2-0"></span>In view of the linear formulation of the problem, below we disregard the term e  $\frac{e}{mc} \int \left[ \langle \hat{\mathbf{v}}(\mathbf{q},s) \rangle \times \langle \hat{\mathbf{B}}(\mathbf{k}-\mathbf{q},w-s) \rangle \right] d\mathbf{q} ds$ . The correlation  $\langle \hat{\vec{\mathbf{v}}}(\mathbf{q},s) \times \hat{\vec{\mathbf{B}}}(\mathbf{k}-\mathbf{q},w-s) \rangle$  $\setminus$ can be expressed in terms of the cumulants of the fluctuational magnetic field using the Furutsu– Novikov [16]:

$$
\left\langle \hat{\tilde{\mathbf{v}}}(\mathbf{q},s) \times \hat{\tilde{\mathbf{B}}}(\mathbf{k}-\mathbf{q},w-s) \right\rangle_{i} =
$$
\n
$$
\varepsilon_{ijk} \int \left\langle \frac{\delta \hat{\tilde{v}}_{j}(\mathbf{q},s)}{\delta \hat{\tilde{B}}_{m}(\mathbf{k}',w')} \right\rangle \left\langle \hat{\tilde{B}}_{m}(\mathbf{k}',w') \hat{\tilde{B}}_{k}(\mathbf{k}-\mathbf{q},w-s) \right\rangle d\mathbf{k}' dw' +
$$
\n
$$
+\varepsilon_{ijk} \int \left\langle \frac{\delta^{2} \hat{\tilde{v}}_{j}(\mathbf{q},s)}{\delta \hat{\tilde{B}}_{m}(\mathbf{k}',w') \delta \hat{\tilde{B}}_{n}(\mathbf{k}'',w'')} \right\rangle \times
$$
\n
$$
\times \left\langle \hat{\tilde{B}}_{m}(\mathbf{k}',w') \hat{\tilde{B}}_{n}(\mathbf{k}'',w'') \hat{\tilde{B}}_{k}(\mathbf{k}-\mathbf{q},w-s) \right\rangle d\mathbf{k}' d\mathbf{k}'' dw' dw'' + \dots \tag{4}
$$

where the variational derivative  $\left\langle \frac{\delta \widehat{v}_{j}(\mathbf{q},s)}{\widehat{\infty}} \right\rangle$  $\delta B_m({\bf k}^\prime,w^\prime)$  $\setminus$ satisfies the equation

$$
\hat{L}_{js}(s) \left\langle \frac{\delta \hat{\tilde{v}}_{s}(\mathbf{q}, s)}{\delta \hat{\tilde{B}}_{m}(\mathbf{k}', w')} \right\rangle = -\frac{es}{mc} \varepsilon_{jlm} \frac{q_{l}}{q^{2}} \delta\left(s - w'\right) \delta\left(\mathbf{q} - \mathbf{k}'\right) + \frac{e}{mc} \varepsilon_{jlm} \langle \hat{v}_{l}(\mathbf{q} - \mathbf{k}', s - w') \rangle \n+ \frac{e}{mc} \varepsilon_{jlr} \int \left\langle \frac{\delta \hat{\tilde{v}}_{l}(\mathbf{q}', s')}{\delta \hat{\tilde{B}}_{m}(\mathbf{k}', w')} \right\rangle \langle \hat{B}_{r}(\mathbf{q} - \mathbf{q}', s - s') \rangle d\mathbf{q}' ds' \n+ \frac{e}{mc} \varepsilon_{jlr} \int \left\langle \frac{\delta^{2} \hat{\tilde{v}}_{l}(\mathbf{q}', s')}{\delta \hat{\tilde{B}}_{m}(\mathbf{k}', w') \delta \hat{\tilde{B}}_{n}(\mathbf{q}' - \mathbf{q}, s' - s)} \right\rangle \hat{Q}_{nr}(\mathbf{q} - \mathbf{q}', s - s') d\mathbf{q}' ds'.
$$
\n(5)

Here,

<span id="page-2-1"></span>
$$
\hat{L}_{js}(s) = -is\delta_{js} + \frac{e}{mc}\varepsilon_{jrs}B_{0r}
$$
\n
$$
\left\langle \hat{B}_n(\mathbf{q} - \mathbf{q}', s - s')\hat{B}_r(\mathbf{k}, w) \right\rangle = \hat{Q}_{nr}(\mathbf{q} - \mathbf{q}', s - s')\delta(\mathbf{k} + \mathbf{q} - \mathbf{q}', w + s - s')
$$
\n(6)

The second variational derivative depends on the third derivative etc. In general, the problem is not closed. In the case of  $\delta$ -correlated (in time) fluctuations, the first term is retained in Eq. [\(4\)](#page-2-0), which corresponds to the Gaussian approximation. This is also a good approximation for short correlation times.

To take into account long correlation times, we can use, in particular, a consistent procedure of allowance for the memory effects similar to that suggested in [17]. Having obtained the equation for the n−th variational derivative, let us substitute the emerging term with the( $n+1$ )−th derivative with an effective relaxation term, which reflects the mixing role of the higher-order moments. This, in turn, gives rise to an effective collision frequency determined by the pulsation amplitude of the magnetic field in the equation for the  $(n-1)-$ th variational derivative, so the frequency s in an operator of the type  $\hat{L}_{js}(s)$  changes to s' s + iw\*. Here, we restrict our analysis to a simpler approach and set the last term in Eq. ([\(5\)](#page-2-1), as for a  $\delta$ correlated (in time) process, equal to zero. We can verify by direct analysis that this is possible when the characteristic frequencies of the electromagnetic fluctuations are much higher than the stochastic Larmor frequency determined from the mean amplitude of the magnetic fluctuations,  $w_{fluct} \gg e \left\langle \tilde{\mathbf{B}}^2 \right\rangle^{1/2} / mc$ . This approximation is similar to the "first post-Markovian" approximation used in the statistical theory of wave propagation in a turbulent medium [17]. Thus, for the first variational derivative, we write

$$
\hat{L}_{js}(s) \left\langle \frac{\delta \hat{\tilde{v}}_{s}(\mathbf{q}, s)}{\delta \hat{\tilde{B}}_{m}(\mathbf{k}', w')} \right\rangle = -\frac{es}{mc} \varepsilon_{jlm} \frac{q_{l}}{q^{2}} \delta \left( s - w' \right) \delta \left( \mathbf{q} - \mathbf{k}' \right) + \frac{e}{mc} \varepsilon_{jlm} \left\langle \hat{v}_{l}(\mathbf{q} - \mathbf{k}', s - w') \right\rangle \n+ \frac{e}{mc} \varepsilon_{jlr} \int \left\langle \frac{\delta \hat{\tilde{v}}_{l}(\mathbf{q}', s')}{\delta \hat{\tilde{B}}_{m}(\mathbf{k}', w')} \right\rangle \left\langle \hat{B}_{r}(\mathbf{q} - \mathbf{q}', s - s') \right\rangle d\mathbf{q}' ds'. \tag{7}
$$

We take into account the nonuniformity of the mean field by successive approximations:

$$
\hat{L}_{js}(s)'\left\langle \frac{\delta \hat{\tilde{v}}_s(\mathbf{q},s)}{\delta \hat{\tilde{B}}_m(\mathbf{k}',w')} \right\rangle^{(0)} = -\frac{es}{mc}\varepsilon_{jlm}\frac{q_l}{q^2}\delta\left(s-w'\right)\delta\left(\mathbf{q}-\mathbf{k}'\right) + \frac{e}{mc}\varepsilon_{jlm}\left\langle \hat{v}_l(\mathbf{q}-\mathbf{k}',s-w')\right\rangle
$$
\n(8)

$$
\hat{L}_{js}(s')\left\langle \frac{\delta\hat{\tilde{v}}_s(\mathbf{q},s)}{\delta\hat{\tilde{B}}_m(\mathbf{k}',w')}\right\rangle^{(1)} = \frac{e}{mc}\varepsilon_{jlr}\int \left\langle \frac{\delta\hat{\tilde{v}}_l(\mathbf{q}',s')}{\delta\hat{\tilde{B}}_m(\mathbf{k}',w')}\right\rangle^{(0)} \left\langle \hat{B}_r(\mathbf{q}-\mathbf{q}',s-s') \right\rangle d\mathbf{q}'ds' \qquad (9)
$$

Retaining only the linear terms, we write

<span id="page-3-0"></span>
$$
\left\langle \frac{\delta \hat{\tilde{v}}_j(\mathbf{q},s)}{\delta \hat{\tilde{B}}_m(\mathbf{k}',w')} \right\rangle = \hat{L}_{js}^{-1}(s) \left( -\frac{es}{mc} \varepsilon_{spm} \frac{q_p}{q^2} \delta\left(s - w'\right) \delta\left(\mathbf{q} - \mathbf{k}'\right) + \frac{e}{mc} \varepsilon_{spm} \left\langle \hat{v}_p(\mathbf{q} - \mathbf{k}',s - w') \right\rangle \right) - \left( \frac{e}{mc} \right)^2 \varepsilon_{jlr} \varepsilon_{tpm} \hat{L}_{js}^{-1}(s) \hat{L}_{lt}^{-1}(w') w' \frac{k'_p}{k'^2} \left\langle \hat{B}_r(\mathbf{q} - \mathbf{k}',s - w') \right\rangle \tag{10}
$$

Here, is the operator  $\hat{L}_{ij}^{-1}(s)$  that is the inverse of  $\hat{L}_{ij}(s)$ :

$$
\hat{L}_{ij}^{-1}(s) = \frac{1}{is\left(s^2 - \Omega_e^2\right)} \left(-\delta_{ij}s^2 + \Omega_{ei}\Omega_{ej} - is\varepsilon_{ijk}\Omega_{ek}\right), \quad \Omega_e = \frac{e\mathbf{B}_0}{mc}
$$
\n(11)

In what follows, we use the relationship between the fields  $\bf{B}$  and  $\bf{E}$  via Maxwell's equation written in the Fourier representation as

$$
\widehat{\mathbf{B}} = \frac{c}{w} \left[ \mathbf{k} \times \widehat{\mathbf{E}} \right]
$$
 (12)

For uniform gyrotropic fluctuations with the anisotropy introduced by a uniform magnetic field, the correlation tensor  $\widehat{Q}_{mk}(\mathbf{q}, s)$  is [18–20]

$$
\widehat{Q}_{mk}(\mathbf{q},s) = \left(\delta_{mk} - \frac{q_m q_k}{q^2}\right) \frac{E_M(q, (\mathbf{l}\mathbf{q}), s)}{4\pi q^2} + i \frac{H_M(q, (\mathbf{l}\mathbf{q}), s)}{8\pi q^4} \varepsilon_{mkt} q_t + \left[\left(l_m q_k + l_k q_m\right) (\mathbf{l}\mathbf{q}) - l_m l_k q^2 - \frac{q_m q_k}{q^2} (\mathbf{l}\mathbf{q})^2\right] \frac{F\left(q, (\mathbf{l}\mathbf{q}), s\right)}{4\pi q^4} - i \left(\delta_{ml} \varepsilon_{kij} + \delta_{kl} \varepsilon_{mij}\right) l_i l_j \left(l_l q^2 - q_l (\mathbf{l}\mathbf{q})\right) \frac{C\left(q, (\mathbf{l}\mathbf{q}), s\right)}{4\pi q^4} \tag{13}
$$

Here, I is a unit vector parallel to the uniform magnetic field,  $I \uparrow B_0$ . All of the correlation functions, except for  $C(q, (\mathbf{Iq}), s)$ , are even in  $(\mathbf{Iq})$ . The symmetry properties also admit the combinations linear in components of the vector l considered in [6, 9]. However, it was shown in [18, 20] that when the anisotropy is attributable to a magnetic field, the only possible combinations are quadratic ones<sup>1</sup>. This is also confirmed by direct calculations of the magnetic field effect on the correlation properties of turbulence [21]. For a weak anisotropy (and for obtaining analytical results), we can use the representation

$$
E_M(q, (\mathbf{lq}), s) = E(q, s) - \frac{(\mathbf{lq})^2}{q^2} E_1(q, s),
$$
  
\n
$$
H_M(q, (\mathbf{lq}), s) = H(q, s) - \frac{(\mathbf{lq})^2}{q^2} H_1(q, s),
$$
  
\n
$$
F(q, (\mathbf{lq}), s) = F(q, s), C(q, (\mathbf{lq}), s) = C_1(q, s) (\mathbf{lq})
$$
\n(14)

Assuming the decay of the correlations with time to be exponential,  $\sim \frac{\tau_*}{\tau_*}$  $\frac{\tau_*}{\tau} \exp\left(-\left|t-t'\right|/\tau\right)$ , we write for the Fourier transform

$$
f(q,s) = f(q) \frac{\tau_*}{\pi (1 + s^2 \tau^2)}.
$$
\n(15)

Here,  $\tau_*$  is the time constant determined by the characteristic frequencies and scales. Thus, for example, for interplanetary plasma turbulence [22],  $\tau_*$  is assumed to be  $\tau_* \sim \frac{\lambda}{n}$  $\frac{\lambda}{v_A} = \frac{\lambda \omega_i}{c \Omega_i}$ where  $\lambda$  is the characteristic fluctuational scale of the magnetic nonuniformities. Clearly, this estimate is also valid for ionospheric plasma. Let us expand the tensor  $\hat{Q}_{mk}(\mathbf{k} - \mathbf{q}, w - s) =$  $Q_{km}(\mathbf{q} - \mathbf{k}, s - w)$  as a series in  $k \ll q$ ,

$$
\widehat{Q}_{km}(\mathbf{q}-\mathbf{k},s-w)=\widehat{Q}_{km}(\mathbf{q},s)-k_r\frac{\partial\widehat{Q}_{km}(\mathbf{q},s)}{\partial q_r}+\frac{k_rk_t}{2}\frac{\partial\widehat{Q}_{km}(\mathbf{q},s)}{\partial q_r\partial q_t}+\ldots,\qquad(16)
$$

and substitute this representation in [\(10\)](#page-3-0), performing the integration over the solid angles, the frequencies s. We then find that, to within the first degree of the expansion in terms of the

$$
h_i(\mathbf{x}) = l_k \frac{\partial^2 P}{\partial x_k \partial x_i} - l_i \Delta P + \varepsilon_{ikj} l_k \frac{\partial T}{\partial x_j}.
$$

<sup>&</sup>lt;sup>1</sup>Indeed, an arbitrary vortex field can be represented as a sum of its toroidal and poloidal components with the basis defined for an arbitrary direction of l:

Choosing the direction of the external stationary uniform magnetic field as this direction, we find that the dependence on the components of this direction appears in the tensor of the pair correlations between the magnetic fluctuations only quadratically.

correlation time  $\tau$  and neglecting the effects quadratic in wave vector ( $\sim k^2$ ), the Lorentz force averaged over the uniform electromagnetic background fluctuations is

$$
\frac{e}{mc} \int \left\langle \widehat{\widetilde{\mathbf{v}}}(\mathbf{q},s) \times \widehat{\widetilde{\mathbf{B}}}(\mathbf{k}-\mathbf{q},w-s) \right\rangle d\mathbf{q}ds = -\left(\frac{e}{mc}\right)^2 \widehat{\mathcal{E}}\tau_* \left\langle \widehat{\mathbf{v}}(\mathbf{k},w) \right\rangle \n+ \frac{e}{m} \left(\frac{e}{mc}\right)^2 \widehat{\mathcal{H}}\tau_* i \left[\mathbf{k} \times \left\langle \widehat{\mathbf{E}}(\mathbf{k},w) \right\rangle \right] + \frac{2}{3} \left(\frac{e}{mc}\right)^2 E_0 \tau_* \tau \left(1 - \frac{1}{5}t_1 + \frac{4}{5}t_2\right) \left[\Omega_e \times \left\langle \widehat{\mathbf{v}}(\mathbf{k},w) \right\rangle\right] \n- \frac{2}{3} \frac{e}{m} \left(\frac{e}{mc}\right)^2 \frac{H_0 \tau \tau_*}{w} \left(1 - \frac{3}{10}g_1\right) \left[\Omega_e \times \left[\mathbf{k} \times \left\langle \widehat{\mathbf{E}}(\mathbf{k},w) \right\rangle\right]\right] - \frac{2}{3} \left(\frac{e}{mc}\right)^2 H_0 \tau_* (1 + i\tau w) \Omega_e \delta(\mathbf{k}) \delta(w),
$$

where

$$
\left[\widehat{\mathcal{H}}\right]_{ij} = \mathcal{H}_{\perp}\delta_{ij} + \left(\mathcal{H}_{\parallel} - \mathcal{H}_{\perp}\right)l_{i}l_{j},\tag{17}
$$

$$
\mathcal{H}_{\perp} = H_0 \frac{2i}{3w} \left( 1 + iw\tau \right) \left( 1 - \frac{3}{10} g_1 \right), \ \mathcal{H}_{\parallel} = H_0 \frac{2i}{3w} \left( 1 + iw\tau \right) \left( 1 - \frac{2}{5} g_1 \right), \tag{18}
$$

$$
\left[\hat{\mathcal{E}}\right]_{ij} = \mathcal{E}_{\perp}\delta_{ij} + \left(\mathcal{E}_{\parallel} - \mathcal{E}_{\perp}\right)l_i l_j \mathcal{E}_{\perp} = \frac{4}{3}E_0\left(1 + i\tau w\right)\left(1 - \frac{3}{10}t_1 + \frac{9}{20}t_2\right),\tag{19}
$$

$$
\mathcal{E}_{\parallel} = \frac{4}{3} E_0 \left( 1 + i \tau w \right) \left( 1 - \frac{2}{5} t_1 + \frac{1}{10} t_2 \right). \tag{20}
$$

Here,

$$
H_0 = \int \frac{H(q)}{q^2} dq = \left\langle \widetilde{\mathbf{A}} \widetilde{\mathbf{B}} \right\rangle_0; H_1 = \int \frac{H_1(q)}{q^2} dq; \tag{21}
$$

$$
E_0 = \int E(q) dq = \left\langle \widetilde{\mathbf{B}}^2 \right\rangle_0; E_1 = \int E_1(q) dq; E_2 = \int F(q) dq. \tag{22}
$$

$$
g_1 = \frac{H_1}{H_0}, \ t_1 = \frac{E_1}{E_0}, \ t_2 = \frac{E_2}{E_0}, q = |\mathbf{q}|.
$$
\n(23)

The subscript 0 corresponds to the isotropic case. As we see, the effective transport coefficients are directly related to the mean energy and helicity of the fluctuational magnetic field. For the time being, let us restrict our analysis to the approximation of a d-correlated process,  $\tau \to 0$ . The effects of finite correlation times will be considered below. For the average Lorentz force, we then obtain

<span id="page-5-0"></span>
$$
\frac{e}{mc} \int \left\langle \widehat{\widetilde{\mathbf{v}}}(\mathbf{q},s) \times \widehat{\widetilde{\mathbf{B}}}(\mathbf{k}-\mathbf{q},w-s) \right\rangle d\mathbf{q}ds =
$$
\n
$$
\frac{e}{mc} \left( \frac{e}{mc} \right)^2 \widehat{\mathcal{H}}_{\tau \to 0} \tau_1 i \left[ \mathbf{k} \times \left\langle \widehat{\mathbf{E}}(\mathbf{k},w) \right\rangle \right] - \left( \frac{e}{mc} \right)^2 \widehat{\mathcal{E}}_{\tau \to 0} \tau_1 \left\langle \widehat{\mathbf{v}}(\mathbf{k},w) \right\rangle - \left( \frac{e}{mc} \right)^2 \frac{2}{3} H_0 \tau_1 \Omega_e
$$
\n(24)

The last term on the right-hand side of Eq. [\(24\)](#page-5-0) has the meaning of constant acceleration along the external magnetic field. To all appearances, the possibility of such acceleration was first pointed out in  $[5]$  (see also  $[6]$ ) and was also considered in detail in  $[7, 8]$  when the helicity input was discussed. It was suggested, as an explanation, that the acceleration is produced by the electric field generated by a fluctuational dynamo effect. Attention to the relationship between the acceleration effect and the transfer of electromagnetic field momentum to particles of the medium was drawn in [23]. Assuming that  $\langle E \rangle, \langle B \rangle = 0$  are equal to zero, we find that in the nonrelativistic collisionless limit, a charged particle reaches a velocity  $\mathbf{v}_{max} \cong -\frac{1}{2}$  $\frac{1}{2} \langle AB \rangle / \langle B^2 \rangle \Omega_e$  i.e., does not depend on the correlation time and is determined by the Larmor frequency in external magnetic field and by the scale specified by the relationship between magnetic helicity and energy. In what follows, we disregard this effect. This is possible for  $|\mathbf{kv}_{max}|/\omega_e \ll 1$ , where  $\omega_e^2 = \frac{4\pi n e^2}{m}$  $\frac{me^2}{m}$  and *n* is the electron density.

## 3 THE CONDUCTIVITY TENSOR

Given the fluctuational friction specified by the term  $-\left(\frac{e}{e}\right)$ mc  $\int^2 \widehat{\mathcal{E}}_0 \tau_* \langle \widehat{\mathbf{v}}(\mathbf{k}, w) \rangle$  the inverse operator is

$$
\hat{L}_{ij}^{-1}(w) = \frac{-\delta_{ij}\left(w + i\overline{\Omega_{\perp e}^2}\tau_*\right)^2 + \Omega_{ei}\Omega_{ej} - i\left(w + i\overline{\Omega_{\parallel e}^2}\tau_*\right)\varepsilon_{ijk}\Omega_{ek}}{i\left(w + i\overline{\Omega_{\parallel e}^2}\tau_*\right)\left(\left(w + i\overline{\Omega_{\perp e}^2}\tau_*\right)^2 - \Omega_e^2\right)}
$$
(25)

Here  $\overline{\Omega_{\perp e}^2} = \left(\frac{e}{mc}\right)^2 \mathcal{E}_{0\perp}, \overline{\Omega_{\parallel e}^2} = \left(\frac{e}{mc}\right)^2 \mathcal{E}_{0\parallel}.$  Taking into account the explicit form of the tensor  $\widehat{\mathcal{H}}_0$ , let us write the electron velocity as

$$
\langle \widehat{\mathbf{v}}(\mathbf{k},w) \rangle = -\frac{e}{m} \frac{\langle \widehat{\mathbf{E}}(\mathbf{k},w) \rangle}{iw'_{e\parallel}} + \frac{e}{m} \frac{\Omega_e^2 \left( \mathbf{1} \left( \mathbf{1} \langle \widehat{\mathbf{E}}(\mathbf{k},w) \rangle \right) - \langle \widehat{\mathbf{E}}(\mathbf{k},w) \rangle \right)}{iw'_{e\parallel} (w'^2_{e\perp} - \Omega_e^2)} + \frac{e}{m} \frac{\Omega_e \left[ \mathbf{1} \times \langle \widehat{\mathbf{E}}(\mathbf{k},w) \rangle \right]}{(w'^2_{e\perp} - \Omega_e^2)} - \frac{e}{m} \left( \frac{e}{mc} \right)^2 \mathcal{H}_{0\perp} \tau_* \frac{i}{w} \frac{w'^2_{e\perp} \left( [\mathbf{k} \times \mathbf{E}(\mathbf{k})] - \mathbf{1} \left( \mathbf{1} \left[ \mathbf{k} \times \langle \widehat{\mathbf{E}}(\mathbf{k},w) \rangle \right] \right) \right)}{w'_{e\parallel} (w'^2_{e\perp} - \Omega_e^2)} - \frac{e}{m} \left( \frac{e}{mc} \right)^2 \mathcal{H}_{0\perp} \tau_* \frac{i}{w} \frac{\mathbf{1} \left( \mathbf{1} \left[ \mathbf{k} \times \langle \widehat{\mathbf{E}}(\mathbf{k},w) \rangle \right] \right)}{w'_{e\parallel}} - \frac{e}{m} \left( \frac{e}{mc} \right)^2 \mathcal{H}_{0\perp} \tau_* \Omega_e \frac{\left[ \mathbf{1} \times \left[ \mathbf{k} \times \langle \widehat{\mathbf{E}}(\mathbf{k},w) \rangle \right] \right]}{w \left( w'^2_{e\perp} - \Omega_e^2 \right)}, \quad (26)
$$

where  $\Omega_e = \frac{e|\mathbf{B}|_0}{mc}$  $\frac{|\mathbf{B}|_0}{mc}, w'_{e\perp(\parallel)} = w + i\Omega^2_{\perp(\parallel)}\tau_*$ . The calculations for ions are similar, and the ion velocity can be written as

$$
\langle \widehat{\mathbf{v}}(\mathbf{k},w) \rangle_{i} = \frac{e}{M} \frac{\langle \widehat{\mathbf{E}}(\mathbf{k},w) \rangle}{iw'_{i\parallel}} - \frac{e}{M} \frac{\Omega_{i}^{2} \left( \mathbf{1} \left( \mathbf{1} \langle \widehat{\mathbf{E}}(\mathbf{k},w) \rangle \right) - \langle \widehat{\mathbf{E}}(\mathbf{k},w) \rangle \right)}{iw'_{i\parallel} (w_{i\perp}^{\prime 2} - \Omega_{i}^{2})} + \frac{e}{M} \frac{\Omega_{i} \left[ \mathbf{1} \times \langle \widehat{\mathbf{E}}(\mathbf{k},w) \rangle \right]}{(w_{i\perp}^{\prime 2} - \Omega_{i}^{2})} + \frac{e}{M} \left( \frac{e}{Mc} \right)^{2} \mathcal{H}_{0\perp} \tau_{*} \frac{i}{w} \frac{w_{i\perp}^{\prime 2} \left( \left[ \mathbf{k} \times \langle \widehat{\mathbf{E}}(\mathbf{k},w) \rangle \right] - \mathbf{1} \left( \mathbf{1} \left[ \mathbf{k} \times \langle \widehat{\mathbf{E}}(\mathbf{k},w) \rangle \right] \right) \right)}{w'_{i\parallel} (w_{i\perp}^{\prime 2} - \Omega_{i}^{2})} + \frac{e}{M} \left( \frac{e}{Mc} \right)^{2} \mathcal{H}_{0\parallel} \tau_{*} \frac{i}{w} \frac{\mathbf{1} \left( \mathbf{1} \left[ \mathbf{k} \times \langle \widehat{\mathbf{E}}(\mathbf{k},w) \rangle \right] \right)}{w'_{i\parallel}} - \frac{e}{M} \left( \frac{e}{Mc} \right)^{2} \mathcal{H}_{0\perp} \tau_{*} \Omega_{i} \frac{\left[ \mathbf{l} \times \left[ \mathbf{k} \times \langle \widehat{\mathbf{E}}(\mathbf{k},w) \rangle \right] \right]}{w \left( w_{i\perp}^{\prime 2} - \Omega_{i}^{2} \right)} \tag{27}
$$

The subscript *i* refers to the ion analogues of the parameters introduced for electrons:  $\Omega_i = \frac{e|\mathbf{B}|_0}{Mc}, w'_{i\perp(\parallel)} = w + i\overline{\Omega^2_{\perp(\parallel)}}_i\tau_*$ . As we see, averaging over the electromagnetic fluctuations is equivalent, in particular, to an effective collision with frequencies proportional to  $\sim \Omega_{\perp e(i)}^2 \tau_*$ and  $\Omega^2_{\parallel_{e(i)}}\tau_*$ . For the conductivity tensor  $j_k = \hat{\sigma}_{kl}(\mathbf{k}, w) E_l(\mathbf{k}, w)$   $(\mathbf{j} = ne(\langle \mathbf{v} \rangle - \langle \mathbf{v} \rangle_i))$ , we obtain

$$
4\pi \hat{\sigma}_{kl}(\mathbf{k}, w) = -\left(\frac{\omega_e^2 w_{e\perp}^{'2}}{iw'_{e\parallel} (w_{e\perp}^{'2} - \Omega_e^2)} + \frac{\omega_i^2 w_{i\perp}^{'2}}{iw'_{i\parallel} (w_{i\perp}^{'2} - \Omega_i^2)}\right) \delta_{kl}
$$

$$
+ \left(\frac{\omega_e^2 \Omega_e^2}{iw'_{e\parallel} (w_{e\perp}^{'2} - \Omega_e^2)} + \frac{\omega_i^2 \Omega_i^2}{iw'_{i\parallel} (w_{i\perp}^{'2} - \Omega_i^2)}\right) l_k l_l + \left(\frac{\omega_e^2 \Omega_e}{(w_{e\perp}^{'2} - \Omega_e^2)} - \frac{\omega_i^2 \Omega_i}{(w_{i\perp}^{'2} - \Omega_i^2)}\right) \varepsilon_{kml} l_m
$$

$$
+ \left(\frac{\omega_e^2 h_{\perp e} \Omega_e}{w (w_{e\perp}^{'2} - \Omega_e^2)} - \frac{\omega_i^2 h_{\perp i} \Omega_i}{w (w_{i\perp}^{'2} - \Omega_i^2)}\right) (l_m k_m \delta_{kl} - l_l k_k) - i \left(\frac{\omega_e^2 h_{\parallel e}}{w w'_{e\parallel}} + \frac{\omega_i^2 h_{\parallel ei}}{w w'_{i\parallel}}\right) l_k l_m \varepsilon_{mnl} k_n
$$

$$
- i \left(\frac{\omega_e^2 h_{\parallel e} w'_{e\perp}}{w (w_{e\perp}^{'2} - \Omega_e^2)} + \frac{\omega_i^2 h_{\parallel ei} w'_{\perp}}{w (w_{i\perp}^{'2} - \Omega_i^2)}\right) (\varepsilon_{kml} k_m - l_k l_m \varepsilon_{lmn} k_n) \tag{28}
$$

where  $\omega_{e(i)}^2 = \frac{4\pi n e^2}{m(M)}$  $\frac{4\pi ne^2}{m(M)}$  .

The coefficients $h_{\perp e(i)}$  and  $h_{\parallel e(i)}$  have the dimensions of velocity, and it is convenient to represent them as

<span id="page-7-1"></span>
$$
h_{\perp e(i)} = \overline{\Omega_{\perp e(i)}^2} \tau_* \lambda_{\kappa \perp} = \alpha_{\perp e(i)} \frac{\Omega_{e(i)}}{\Omega_{\kappa \perp}} c, \alpha_{\perp e(i)} = \frac{\Omega_{\perp e(i)}^2 \tau_*}{\Omega_{e(i)}},
$$
  
\n
$$
\Omega_{\kappa \perp} = \frac{c}{\lambda_{\kappa \perp}}
$$
  
\n
$$
h_{\parallel e(i)} = \overline{\Omega_{\parallel e(i)}^2} \tau_* \lambda_{\kappa \parallel} = \alpha_{\parallel e(i)} \frac{\Omega_{e(i)}}{\Omega_{\kappa \parallel}} c, \alpha_{\parallel e(i)} = \frac{\overline{\Omega_{\perp e(i)}^2} \tau_*}{\Omega_{e(i)} \tau_*},
$$
  
\n
$$
\Omega_{\kappa \parallel} = \frac{c}{\lambda_{\kappa \parallel}}
$$
\n(29)

where the scale  $\lambda_{\kappa\perp(||)}$  is defined by the ratio of the helicity and energy of the fluctuations:<sup>2</sup>

$$
\lambda_{\kappa\perp(\parallel)} = \frac{\mathcal{H}_{0\perp(\parallel)}}{\mathcal{E}_{0\perp(\parallel)}} \approx \frac{1}{2} \frac{\langle \mathbf{AB} \rangle}{\langle \mathbf{B}^2 \rangle}.
$$
\n(30)

Neglecting the fluctuational damping  $\Omega_{\|\mathcal{T}_{*}}^{2}(\Omega_{\bot}^{2}\tau_{*})$ , we obtain

$$
4\pi \hat{\sigma}_{kl}(\mathbf{k}, w) = iw \left(\frac{\omega_e^2}{w^2 - \Omega_e^2} + \frac{\omega_i^2}{w^2 - \Omega_i^2}\right) \delta_{kl} + \left(\frac{\omega_e^2 \Omega_e^2}{w^2 - \Omega_e^2} + \frac{\omega_i^2 \Omega_i^2}{w^2 - \Omega_i^2}\right) \frac{l_k l_l}{iw} + \left(\frac{\omega_e^2 \Omega_e}{w^2 - \Omega_e^2} - \frac{\omega_i^2 \Omega_i}{w^2 - \Omega_i^2}\right) \varepsilon_{kml} l_m + \left(\frac{\omega_e^2 \Omega_e h_{\perp e}}{w (w^2 - \Omega_e^2)} - \frac{\omega_i^2 \Omega_i h_{\perp i}}{w (w^2 - \Omega_i^2)}\right) (l_m k_m \delta_{kl} - l_l k_k) - i \left(\frac{\omega_e^2 h_{\parallel e}}{w^2} + \frac{\omega_i^2 h_{\parallel i}}{w^2}\right) l_k l_m \varepsilon_{mnl} k_n - i \left(\frac{\omega_e^2 h_{\perp e}}{w^2 - \Omega_e^2} + \frac{\omega_i^2 h_{\perp i}}{w^2 - \Omega_i^2}\right) (\varepsilon_{kml} k_m - l_k l_m \varepsilon_{lmn} k_n).
$$
\n(31)

<span id="page-7-0"></span>Hence, the permittivity tensor is

$$
\varepsilon_{ij} = \delta_{ij} + \frac{4\pi i}{w} \hat{\sigma}_{ij}
$$
\n
$$
\hat{\varepsilon} = \begin{pmatrix}\n\varepsilon_{\perp} + i\chi_0 k_z & ig + \chi_{\perp} k_z & -i\chi_0 k_x - \chi_{\perp} k_y \\
-ig - \chi_{\perp} k_z & \varepsilon_{\perp} + i\chi_0 k_z & -i\chi_0 k_y + \chi_{\perp} k_x \\
\chi_{\parallel} k_y & -\chi_{\parallel} k_x & \varepsilon_{\parallel}\n\end{pmatrix} \tag{32}
$$

<sup>&</sup>lt;sup>2</sup>The characteristic scale of the fluctuational magnetic helicity is known for the solar-wind turbulence [24], where it lies within the range  $0.004 \div 0.02$  AU ( $\sim 6 \cdot 10^8 \div 3 \cdot 10^9$  m).

where

$$
\varepsilon_{\perp} = 1 - \frac{\omega_e^2}{w^2 - \Omega_e^2} - \frac{\omega_i^2}{w^2 - \Omega_i^2}, \varepsilon_{\parallel} = 1 - \frac{\omega_e^2}{w^2} - \frac{\omega_i^2}{w^2}, g = \frac{\omega_e^2 \Omega_e}{w \left(\Omega_e^2 - w^2\right)} - \frac{\omega_i^2 \Omega_i}{w \left(\Omega_i^2 - w^2\right)},
$$
(33)

$$
\chi_0 = \frac{\omega_e^2 \Omega_e}{w^2 (w^2 - \Omega_e^2)} h_{\perp e} - \frac{\omega_i^2 \Omega_i}{w^2 (w^2 - \Omega_i^2)} h_{\perp i}, \tag{34}
$$

$$
\chi_{\perp} = \frac{h_{\perp e}}{w} \frac{\omega_e^2}{w^2 - \Omega_e^2} + \frac{h_{\perp i}}{w} \frac{\omega_i^2}{w^2 - \Omega_i^2}, \chi_{\parallel} = \frac{h_{\parallel e}}{w} \frac{\omega_e^2}{w^2 - \Omega_e^2} + \frac{h_{\parallel i}}{w} \frac{\omega_i^2}{w^2 - \Omega_i^2}
$$
(35)

As we see from  $(32)$ , fluctuational helicity gives rise to additional gyrotropic terms in the permittivity tensor. To elucidate their role, let us analyze the dispersion relation for electromagnetic waves.

## 4 THE DISPERSION RELATION

Denote the angle between the vectors **n** and  $B_0$  by  $\theta$ . The dispersion relation for the complex refractive index  $\mathbf{n} = c\mathbf{k}/w$  is defined as [11]

$$
\det \|n^2 \delta_{ij} - n_i n_j - \widehat{\varepsilon}_{ij}\| = 0
$$
\n(36)

We also assume that

$$
\mathbf{n} = (n\sin(\theta), 0, n\cos(\theta)).
$$

The dispersion relation is then

<span id="page-8-0"></span>
$$
(g^{2} + (n^{2} - \varepsilon_{\perp}) \varepsilon_{\perp}) \varepsilon_{\parallel} - n^{2} \varepsilon_{\parallel} (n^{2} - \varepsilon_{\perp}) \cos^{2}(\theta) - n^{2} (g^{2} + (n^{2} - \varepsilon_{\perp}) \varepsilon_{\perp}) \sin^{2}(\theta) - n^{2} \varepsilon_{\parallel} w^{2} (-\chi_{0}^{2} + \chi_{\perp}^{2}) / c^{2} \cos^{2}(\theta) - n^{2} w^{2} (g \chi_{0} + \varepsilon_{\perp} \chi_{\perp}) \chi_{\parallel} / c^{2} \sin^{2}(\theta) + i n w \cos(\theta) \varepsilon_{\parallel} (n^{2} \chi_{0} - 2 \varepsilon_{\perp} \chi_{0} - 2 g \chi_{\perp}) / c + i n^{3} w \cos(\theta) (\varepsilon_{\parallel} \chi_{0} / c \cos^{2}(\theta) + (\varepsilon_{\perp} \chi_{0} + g \chi_{\perp} - g \chi_{\parallel}) / c \sin^{2}(\theta)) = 0
$$
\n(37)

Let us consider the waves that propagate along the magnetic field,  $\theta = 0$ . In this case, the dispersion relation has the solutions

$$
n_{1,2} = \frac{1}{2} \left( iw \left( \chi_0 + \chi_+ \right) / c \pm \left( 4 \left( \varepsilon_+ - g \right) - w^2 \left( \chi_0 + \chi_+ \right)^2 / c^2 \right)^{1/2} \right) n_{3,4} = \frac{1}{2} \left( iw \left( \chi_0 - \chi_+ \right) / c \pm \left( 4 \left( \varepsilon_+ + g \right) - w^2 \left( \chi_0 - \chi_+ \right)^2 / c^2 \right)^{1/2} \right)
$$
(38)

Given that

$$
\varepsilon_{\perp} \mp g = 1 - \frac{\omega_e^2}{w(w \pm \Omega_e)} - \frac{\omega_i^2}{w(w \mp \Omega_i)},
$$
  

$$
\chi_0 \pm \chi_{\perp} = \pm \alpha_{\perp e} \frac{\omega_e^2 \Omega_e}{w^2 (w \mp \Omega_e) \Omega_{\kappa \perp}} c \pm \alpha_{\perp i} \frac{\omega_i^2 \Omega_i}{w^2 (w \pm \Omega_i) \Omega_{\kappa \perp}} c,
$$
 (39)

the wave vector is

$$
ck = i \left( \pm \frac{\alpha_{\perp e}}{2\Omega_{\kappa\perp}} \frac{\omega_e^2 \Omega_e}{(w \mp \Omega_e)} \pm \frac{\alpha_{\perp i}}{2\Omega_{\kappa\perp}} \frac{\omega_i^2 \Omega_i}{(w \pm \Omega_i)} \right) \pm \left( w^2 \left( 1 - \frac{\omega_e^2}{w (w \pm \Omega_e)} - \frac{\omega_i^2}{w (w \mp \Omega_i)} \right) - \left( \pm \frac{\alpha_{\perp e}}{2\Omega_{\kappa\perp}} \frac{\omega_e^2 \Omega_e}{(w \mp \Omega_e)} \pm \frac{\alpha_{\perp i}}{2\Omega_{\kappa\perp}} \frac{\omega_i^2 \Omega_i}{(w \pm \Omega_i)} \right)^2 \right)^{1/2},
$$

whence the equation for the frequency is

$$
w^2 - w\left(\frac{\omega_e^2}{(w \pm \Omega_e)} + \frac{\omega_i^2}{(w \mp \Omega_i)}\right) \pm i\frac{ck}{\Omega_{\kappa\perp}} \left(\pm \alpha_{\perp e} \frac{\omega_e^2 \Omega_e}{(w \mp \Omega_e)} \pm \alpha_{\perp i} \frac{\omega_i^2 \Omega_i}{(w \pm \Omega_i)}\right) = c^2 k^2 \tag{40}
$$

At low frequencies with  $\frac{\omega_e^2}{\Omega_e^2} \ll \frac{\omega_i^2}{\Omega_i^2}$  and  $\omega_e^2 \gg \omega_i^2$ , the square of the frequency is

$$
w^2 = \frac{v_A^2 k^2}{\left(1 + \left(1 + \alpha_{\perp i}\right) \frac{v_A^2}{c^2}\right)} \left(1 \pm i\alpha_{\perp e} k \lambda_{\kappa \perp} \frac{\omega_e^2}{c^2 k^2}\right),\tag{41}
$$

where  $v_A^2 = \frac{B_0^2}{4\pi n M}$ . At low values of  $\alpha_{\perp e} k \lambda_{\kappa \perp} \frac{\omega_e^2}{c^2 k^2}$  (for small scales),

$$
w = \left(\frac{v_A k}{\left(1 + \left(1 + \alpha_{\perp i}\right) v_A^2/c^2\right)^{1/2}} \pm i \frac{v_A/c}{\left(1 + \left(1 + \alpha_{\perp i}\right) v_A^2/c^2\right)^{1/2}} \frac{\alpha_{\perp e} \lambda_{\kappa \perp} \omega_e^2}{2c}\right). \tag{42}
$$

In this case, the coefficient of the complex refractive index does not depend on the wave vector. In contrast, at high values of  $\alpha_{\perp e} k \lambda_{\kappa \perp} \frac{\omega_e^2}{c^2 k^2}$  (for large scales),

$$
w = \frac{v_A/c}{\left(1 + \left(1 + \alpha_{\perp i}\right)v_A^2/c^2\right)^{1/2}} \left(\frac{\alpha_{\perp e}k\lambda_{\kappa\perp}}{2}\right)^{1/2} \omega_e \left(1 \pm i\right). \tag{43}
$$

As we see, in the presence of magnetic fluctuation helicity, there is an instability and the amplitude of the electromagnetic waves propagating in a plasma increases. This demonstrates the nonequilibrium existence of reflectional symmetry breaking at the level of fluctuations. Thus, for example, helicity also leads to an instability, an inverse energy cascade, in magnetohydrodynamics [2]. Unstable waves have nonzero helicity, i.e., a vortex component of the electric field. The motion of charged particles in a magnetic field with fluctuational helicity is equivalent to the motion in random helical magnetic fields with preferred helix orientation.

The resonance condition during the motion of particles in a helical magnetic field is satisfied for the particles that move in a direction opposite to the field  $(\mathbf{B}\mathbf{v} < 0)$  [25]. After averaging, this resonance condition will correspond to the following: when the helicities of the perturbations and fluctuations have opposite signs, the perturbations will give up energy to particles of the medium; in contrast, when the helicities of the perturbations and fluctuations have the same signs, the field will be amplified — take away energy from particles of the medium. Indeed, the helicity of growing waves coincides in sign with the small-scale fluctuational helicity. In the opposite case, the perturbation is damped. Note also that the dispersion of the propagating waves changes as well. For large scales,  $w \sim k^{1/2}$ , the dispersion law is similar to that of gravity waves in deep water whose phase velocity increases with scale. Such long waves can be revealed in the spectrum of geoelectromagnetic perturbations. Note that the fast large–scale electric perturbations in the E region of the ionosphere that accompany such catastrophic events as magnetic storms and substorms, earthquakes, and man-made explosions are, to all appearances, of a vortex nature [26]. Let us consider the range of helicon frequencies:  $\Omega_i \ll w \ll \Omega_e, \omega_e^2 \gg w\Omega_e$ . In this case, the frequency can be expressed as

$$
w = \pm \Omega_e \frac{c^2 k^2}{\omega_e^2} \mp i \alpha_{\perp e} k \lambda_{\kappa \perp} \Omega_e \tag{44}
$$

The wave propagation is also accompanied by an instability with the growth rate  $\alpha_{\perp e}k\lambda_{\kappa\perp}\Omega_e$ . Retaining the quadratic terms in the expansion of the Lorentz force  $\overline{1}$  $\widetilde{\mathbf{v}}(\mathbf{q}, s) \times \mathbf{B}(\mathbf{k} - \mathbf{q}, w - s)$ Ň. in terms of large scales  $(k \ll q)$  yields a lower limit for such instability [10], and the perturbations are damped at  $k > k_{crit}$ . Let us consider the waves that propagate perpendicular to the magnetic field,  $\theta = \frac{\pi}{2}$  $\frac{\pi}{2}$ . In this case, the square of the complex refractive index is

<span id="page-10-0"></span>
$$
n_1^2 = \frac{\varepsilon_\perp (\varepsilon_\perp + \varepsilon_\parallel) - g^2 - \kappa + \left( \left( \varepsilon_\perp (\varepsilon_\perp - \varepsilon_\parallel) - g^2 \right)^2 - 2 \left( \varepsilon_\perp (\varepsilon_\perp + \varepsilon_\parallel) - g^2 \right) \kappa + \kappa^2 \right)^{1/2}}{2\varepsilon_\perp}
$$
  

$$
n_2^2 = \frac{\varepsilon_\perp (\varepsilon_\perp + \varepsilon_\parallel) - g^2 - \kappa - \left( \left( \varepsilon_\perp (\varepsilon_\perp - \varepsilon_\parallel) - g^2 \right)^2 - 2 \left( \varepsilon_\perp (\varepsilon_\perp + \varepsilon_\parallel) - g^2 \right) \kappa + \kappa^2 \right)^{1/2}}{2\varepsilon_\perp}
$$

$$
\kappa = w^2 \left( g \chi_0 + \varepsilon_\perp \chi_\perp \right) \chi_\parallel/c^2
$$
(45)

In the absence of helicity, the first and second expressions in [\(45\)](#page-10-0) would correspond to the extraordinary and ordinary waves, respectively. As we see, their propagation conditions change, and elliptical polarization attributable to helicity appears in both types of waves.

## 5 THE OHM LAW FOR LOW FREQUENCIES

Let us consider the case of low frequencies where  $w \ll \Omega_{\parallel e(i)}^2 \tau_* (\Omega_{\perp e(i)}^2 \tau_*)$  and take into account the collision frequency  $\nu = 1/\tau_c \gg w (1/\tau_c'$  for ions). To simplify our calculations, we also assume a weak anisotropy of  $\Omega^2_{\perp e(i)} \approx \Omega^2_{\parallel e(i)}$  and  $\alpha_{\perp e(i)}\lambda_{\kappa\perp} \approx \alpha_{\parallel e(i)}\lambda_{\kappa\parallel} = \alpha_{e(i)}\lambda_{\kappa}$ .

The conductivity tensor [\(28\)](#page-7-1) in this limit will then appear as

$$
4\pi\hat{\sigma}_{kl}(\mathbf{k},w) = \left(\frac{\omega_e^2 \tau_e}{1 + \Omega_e^2 \tau_e^2} + \frac{\omega_i^2 \tau_i}{1 + \Omega_i^2 \tau_i^2}\right)\delta_{kl} + \left(\frac{\omega_e^2 \Omega_e^2 \tau_e^3}{1 + \Omega_e^2 \tau_e^2} + \frac{\omega_i^2 \Omega_i^2 \tau_i^3}{1 + \Omega_i^2 \tau_i^2}\right)l_kl_l
$$

$$
- \left(\frac{\omega_e^2 \Omega_e \tau_e^2}{1 + \Omega_e^2 \tau_e^2} - \frac{\omega_i^2 \Omega_i \tau_i^2}{1 + \Omega_i^2 \tau_e^2}\right)\varepsilon_{kml}l_m
$$

$$
+ i\left(\alpha_e \frac{\omega_e^2 \Omega_e \tau_e^2}{1 + \Omega_e^2 \tau_e^2} \left(1 + \frac{\tau_e}{\tau_c}\right) - \alpha_i \frac{\omega_i^2 \Omega_i \tau_i^2}{1 + \Omega_i^2 \tau_i^2} \left(1 + \frac{\tau_i}{\tau_c}\right)\right)\lambda_{\kappa} (l_m k_m \delta_{kl} - l_l k_k)
$$

$$
+ i\left(\alpha_e \omega_e^2 \left(\tau_c + \tau_e\right) + \alpha_i \omega_i^2 \left(\tau_c' + \tau_i\right)\right)\lambda_k l_k l_m \varepsilon_{mnl} k_n
$$

$$
+ i\left(\alpha_e \frac{\omega_e^2 \left(\tau_c + \tau_e\right)}{1 + \Omega_e^2 \tau_e^2} + \alpha_i \frac{\omega_i^2 \left(\tau_c' + \tau_i\right)}{1 + \Omega_i^2 \tau_i^2}\right)\lambda_{\kappa} \left(\varepsilon_{kml} k_m - l_k l_m \varepsilon_{lmn} k_n\right) \tag{46}
$$

Here, we introduced the following characteristic time scales:

<span id="page-10-1"></span>
$$
\tau_e = \frac{\tau_c}{1 + \overline{\Omega_{\parallel e}^2} \tau_* \tau_c}, \tau_i = \frac{\tau_c'}{1 + \overline{\Omega_{\parallel_i}^2} \tau_*' \tau_c}
$$
(47)

Having defined the conductivities

$$
\sigma_{0e(i)} = \frac{\omega_{e(i)}^2 \tau_{e(i)}}{4\pi}, \sigma_{e(i)\perp} = \frac{\sigma_{0e(i)}}{1 + \Omega_{e(i)}^2 \tau_{e(i)}^2}; \sigma_{e(i)\parallel} = \frac{\sigma_{0e(i)} \Omega_{e(i)}^2 \tau_{e(i)}^2}{1 + \Omega_{e(i)}^2 \tau_{e(i)}^2},
$$

the current after applying the inverse Fourier transform can be written as

$$
\mathbf{j} = (\sigma_{e\perp} + \sigma_{i\perp}) \langle \mathbf{E} \rangle + (\sigma_{e\parallel} + \sigma_{i\parallel}) \mathbf{1} (\mathbf{1} \langle \mathbf{E} \rangle) - (\sigma_{e\perp} \Omega_e \tau_e - \sigma_{i\perp} \Omega_i \tau_i) [\mathbf{1} \langle \mathbf{E} \rangle] \n+ (\alpha_e \sigma_{e\perp} \Omega_e \tau_e \left( 1 + \frac{\tau_e}{\tau_c} \right) - \alpha_i \sigma_{i\neq\perp} \Omega_i \tau_i \left( 1 + \frac{\tau_i}{\tau_c'} \right)) \lambda_\kappa ((\mathbf{I} \nabla) \langle \mathbf{E} \rangle - \nabla (\mathbf{1} \langle \mathbf{E} \rangle)) \n+ (\alpha_e (\sigma_{0e} - \sigma_{e\perp}) \left( 1 + \frac{\tau_c}{\tau_e} \right) + \alpha_i (\sigma_{0i} - \sigma_{i\perp}) \left( 1 + \frac{\tau_c'}{\tau_i} \right)) \lambda_\kappa \mathbf{1} (\text{Irot} \langle \mathbf{E} \rangle) \n+ (\alpha_e \sigma_{e\perp} \left( 1 + \frac{\tau_e}{\tau_e} \right) + \alpha_i \sigma_{i\perp} \left( 1 + \frac{\tau_c'}{\tau_i} \right)) \lambda_\kappa \text{rot} \langle \mathbf{E} \rangle
$$
\n(48)

The influence of fluctuations and external magnetic field primarily causes the conductivity to decrease, while the presence of helicity leads to an additional dependence of the current on the vortex component of the electric field. Let us consider the mean magnetic field in a medium with the Ohm law [\(48\)](#page-10-1). We will disregard the ion component (the EMHD approximation). Neglecting the displacement current, we obtain the following expression for the growth rate  $\gamma$ of wave field perturbations of the form

$$
\langle \mathbf{E} \rangle = \exp(\gamma t) (E_x(z), E_y(z), 0),
$$
  

$$
\langle \mathbf{B} \rangle = \exp(\gamma t) (B_x(z), B_y(z), 0)
$$

$$
\gamma = -\frac{c^2 k^2}{4 \pi \sigma_{e\perp}} \frac{(\tau_c + i \alpha_e k \lambda \zeta \Omega_e \tau_e)}{((1 - i \tau_e \Omega_e) - \alpha_e k \lambda \zeta (\tau_c / \tau_e - i \Omega_e \tau_e)) ((1 + i \Omega_e \tau_e) + \alpha_e k \lambda \zeta (\tau_c / \tau_e + i \Omega_e \tau_e))}
$$
\n(49)

where  $\zeta = 1 + \tau_e/\tau_c$ . For the wave vectors

$$
\alpha_e |k \lambda| > \frac{\left(1 + \Omega_e^2 \tau_e^2\right)^{1/2}}{\left(1 + \tau_e/\tau_c\right) \left(\tau_c^2/\tau_e^2 + \left(2\tau_c/\tau_e - 1\right) \Omega_e^2 \tau_e^2\right)^{1/2}}
$$

the perturbations grow. For intense magnetic fluctuations,  $\tau_e \approx 1/\left(\overline{\Omega_{\parallel}^2} \tau_*\right) \ll \tau_c$ , and we obtain for the threshold wave number

$$
\alpha_e |k\,\lambda| \gtrsim \frac{1}{\Omega_\|^2 \tau_* \tau_c} \left(1 + \frac{\Omega_e^2}{2\left(\overline{\Omega_\|^2} \tau_*\right)^2}\right).
$$

In the collisionless limit  $\tau_c \to 0$ , the threshold wave number is

$$
\alpha_e |k \lambda| > \frac{1}{2}
$$

i.e., the threshold instability scale also increases with fluctuation amplitude (parameter  $\alpha_e$ ). On this threshold scale, the waves with the following frequency propagate at  $\tau_e \ll \tau_c$ :

$$
w = \frac{c^2}{8 \pi \sigma_{0e}} \frac{1}{\alpha_e^2 \lambda^2 \overline{\Omega_{\parallel}^2 \tau_* \tau_c}} \frac{1 + \Omega_e^2 \tau_c^2}{\Omega_e \tau_c} \left( 1 + \frac{\Omega_e^2}{2 \left( \overline{\Omega_{\parallel}^2 \tau_* \tau_c} \right)^2} \right)
$$

Retaining the quadratic terms in the permittivity tensor gives rise to dissipative terms of the form  $-\sigma_*\Delta \langle E \rangle + \sigma'_* \nabla \text{div} \langle E \rangle$  [10] in the Ohm law. Their influence restricts the instability region, and the field perturbations are damped on small scales.

## 6 FINITE CORRELATION TIMES

Consider the effects of finite correlation times for high frequencies,  $w\tau \gg 1$ , with the anisotropy effects disregarded. These also include the case of long correlation times. In this limit, the effective Lorentz force is

$$
\frac{e}{mc} \int \left\langle \hat{\mathbf{v}}(\mathbf{q}, s) \times \hat{\mathbf{B}}(\mathbf{k} - \mathbf{q}, w - s) \right\rangle d\mathbf{q} ds = -\frac{4}{3} i w \tau \left( \frac{e}{mc} \right)^2 \hat{\mathcal{E}} \tau_* \left\langle \hat{\mathbf{v}}(\mathbf{k}, w) \right\rangle \n- \frac{2\tau}{3} \frac{e}{m} \left( \frac{e}{mc} \right)^2 H_{0\perp} \tau_* i \left[ \mathbf{k} \times \left\langle \hat{\mathbf{E}}(\mathbf{k}, w) \right\rangle \right] \n+ \frac{2}{3} \left( \frac{e}{mc} \right)^2 E_0 \tau_* \tau \left[ \mathbf{\Omega}_{\mathbf{e}} \times \left\langle \hat{\mathbf{v}}(\mathbf{k}, w) \right\rangle \right] - \frac{2}{3} i \tau w \left( \frac{e}{mc} \right)^2 H_0 \tau_* \mathbf{\Omega}_{\mathbf{e}} \delta(\mathbf{k}) \delta(w),
$$

The gyrotropic fluctuational acceleration will be replaced with oscillations. The frequencies will acquire a negative shift, and the permittivity tensor will take the form

$$
\widehat{\varepsilon} = \begin{pmatrix}\n\varepsilon_{\perp} - w\tau \chi_0 k_z & ig + iw\tau \chi_{\perp} k_z & w\tau \chi_0 k_x - iw\tau \chi_{\perp} k_y \\
-ig - iw\tau \chi_{\perp} k_z & \varepsilon_{\perp} - w\tau \chi_0 k_z & w\tau \chi_0 k_y + iw\tau \chi_{\perp} k_x \\
iw\tau \chi_{\parallel} k_y & -iw\tau \chi_{\parallel} k_x & \varepsilon_{\parallel}\n\end{pmatrix}.
$$
\n(50)

Consider the waves propagating along the magnetic field,  $\theta = 0$ . In this case, the dispersion relation [\(37\)](#page-8-0) has the solutions

$$
n_{1,2} = (\varepsilon_{\perp} + g) \left( 1 \pm \frac{2w\tau(\chi_0 - \chi_{\perp})}{(4c^2(\varepsilon_{\perp} + g)^2 + (\chi_0 - \chi_{\perp})^2 w^4 \tau^2)^{1/2} \mp (\chi_0 - \chi_{\perp})cw^2 \tau} \right)
$$
  
\n
$$
n_{3,4} = (\varepsilon_{\perp} - g) \left( 1 \pm \frac{2w^2\tau(\chi_0 + \chi_{\perp})}{(4c^2(\varepsilon_{\perp} - g)^2 + (\chi_0 - \chi_{\perp})^2 w^4 \tau^2)^{1/2} \mp (\chi_0 + \chi_{\perp})cw^2 \tau} \right)
$$
\n(51)

Assuming the helical additions to be small, we can write

$$
n_{1,2} = \left(\varepsilon_{\perp} + g \pm \frac{w^2 \tau (\chi_0 - \chi_{\perp})}{c}\right) =
$$
  
\n
$$
1 - \frac{\omega_e^2}{w (w - \Omega_e)} - \frac{\omega_i^2}{w (w + \Omega_i)} \mp \left(\alpha_{\perp e} \frac{\omega_e^2 \Omega_e \tau}{(w + \Omega_e) \Omega_{\kappa \perp}} + \alpha_{\perp i} \frac{\omega_i^2 \Omega_i \tau}{(w - \Omega_i) \Omega_{\kappa \perp}}\right) (52)
$$
  
\n
$$
n_{3,4} = \left(\varepsilon_{\perp} - g \pm \frac{w^2 \tau (\chi_0 + \chi_{\perp})}{c}\right) =
$$
  
\n
$$
1 - \frac{\omega_e^2}{w (w + \Omega_e)} - \frac{\omega_i^2}{w (w - \Omega_i)} \pm \left(\alpha_{\perp e} \frac{\omega_e^2 \Omega_e \tau}{(w - \Omega_e) \Omega_{\kappa \perp}} + \alpha_{\perp i} \frac{\omega_i^2 \Omega_i \tau}{(w + \Omega_i) \Omega_{\kappa \perp}}\right) (53)
$$

It is easy to see that an additional rotation of the polarization plane appears here.

For the waves propagating perpendicular to the magnetic field,  $\theta = \frac{\pi}{2}$  $\frac{\pi}{2}$ , we obtain the following solutions:

$$
n_1^2 = \frac{\varepsilon_\perp (\varepsilon_\perp + \varepsilon_\parallel) - g^2 + \kappa' + \left( \left( \varepsilon_\perp (\varepsilon_\perp - \varepsilon_\parallel) - g^2 \right)^2 - 2 \left( \varepsilon_\perp (\varepsilon_\perp + \varepsilon_\parallel) - g^2 \right) \kappa' + \kappa'^2 \right)^{1/2}}{2\varepsilon_\perp}
$$
  
\n
$$
n_2^2 = \frac{\varepsilon_\perp (\varepsilon_\perp + \varepsilon_\parallel) - g^2 + \kappa' - \left( \left( \varepsilon_\perp (\varepsilon_\perp - \varepsilon_\parallel) - g^2 \right)^2 - 2 \left( \varepsilon_\perp (\varepsilon_\perp + \varepsilon_\parallel) - g^2 \right) \kappa' + \kappa'^2 \right)^{1/2}}{2\varepsilon_\perp}
$$
  
\n
$$
\kappa' = w^4 \tau^2 \left( g \chi_0 + \varepsilon_\perp \chi_\perp \right) \chi_\parallel / c^2
$$
\n(54)

As in the approximation of a  $\delta$ -correlated random process [\(45\)](#page-10-0) considered above, the propagation conditions change, and elliptical polarization attributable to helicity appears in both the ordinary and extraordinary waves. Note that in the case of infinite correlation times or high frequencies (frozen fluctuations), the properties of a plasma medium with magnetic helicity are similar to those of chiral and bianisotropic media [14, 15]. In real systems,  $w\tau$  is finite, and the effects of both instability considered in Section 4 and the appearance of additional wave modes must simultaneously manifest themselves.

### 7 CONCLUSIONS

The influence of magnetic fluctuations on the motion of the particles of a cold magnetoactive plasma primarily reduces to the appearance of an effective fluctuational collision frequency determined by the statistical parameters and to the decrease in conductivity. Reflectional symmetry breaking—nonzero mean magnetic helicity of the fluctuations—leads to a change in the dispersion of the propagating waves and the appearance of additional modes. The waves can be unstable, reflecting both the nonequilibrium nature of the turbulent magnetic helicity and the peculiarities of the particle motion in random helical magnetic fields. The instability growth rate is proportional to the helicity of the fluctuational magnetic field and the amplitude of the large scale uniform magnetic field. An allowance for the finite correlation times and for the additional fluctuational quadratic dispersion effects restricts the action of this instability. In contrast to the turbulent dynamo effects considered in the MHD and EMHD approximations [2], here there is a natural restriction of the instability region on large scales determined by the relationship between the fluctuational helicity and energy and the large-scale magnetic field. The plasma acquires properties similar to those observed in chiral and bianisotropic media [14, 15]. Consequently, it can have properties characteristic of these media, such as anomalous absorption [27, 28] and additional wave conversion effects [29, 30]. In contrast to the artificial external origin of the chirality in chiral media, this property is natural in a turbulent magnetoactive plasma with helicity. The deviations in the rotation of the polarization plane attributable to fluctuational helicity can serve as a tool for diagnosing it. The results were obtained in the approximation of isolated particles whose advantages and disadvantages are well known. It is easy to see that the above effects are preserved when the thermal and collisional effects are taken into account and can be obtained in terms of the kinetic approach.

I am grateful to S.N. Artekha and N.S. Erokhin for helpful discussions. This work was supported by the Russian Science Support Foundation.

#### References

- [1] J.B.Taylor, Phys.Rev.Lett. 33, 1139 (1974).
- [2] S. I. Vainshte n, Ya. B. Zel'dovich, and A.A.Ruzma kin, The Turbulent Dynamo in Astrophysics (Nauka, Moscow, 1980) [in Russian].
- [3] K.Hasselmann, G. Wibberenz, Zs. Geophys. 34, 353 (1968).
- [4] M. L.Goldstein, W. H. Matthaeus, Proc. 17th Internat. Cosmic Ray Conf., 3, 294 (1981).
- [5] L. L. Kichatinov, Pis'ma Zh. Eksp. Teor. Fiz. 37, 43 (1983) [JETP Lett. 37, 51 (1983)].
- [6] Yu.I.Fedorov, V.E.Katz, L.L.Kichatinov & M.Stehlic, Astron. and Astrophys 260, 499 (1992).
- [7] R.R.Mett, J.A.Tataronis, Phys.Rev.Lett. 63, 1380 (1989).
- [8] J.B.Taylor, Phys.Rev.Lett. 63, 1384 (1989).
- [9] A.V.Chechkin, V.V.Yanovsky and A.V.Tur, Phys.Plasmas 1, 2566 (1994).
- [10] L. L. Kichatinov, Pis'ma Zh. Eksp. Teor. Fiz. 37, 43 (1983) [JETP Lett. 37, 51 (1983)].
- [11] V. L. Ginzburg and A. A. Rukhadze, Waves in Magnetized Plasmas (Nauka, Moscow, 1975) [in Russian].
- [12] S.I.Vaihnstein, Phys.Rev.Lett. 80, 4879 (1998).
- [13] A. F. Aleksandrov, and A. A. Rukhadze, Lectures on Electrodynamics of Plasma-like Media (Mosk.Gos. Univ., Moscow, 1999) [in Russian].
- [14] D.L.Jaggard, A.R.Mickelson , C.H.Papas, Applied Physics 18, 211, (1979).
- [15] B. Z. Katsenelenbaum, E. N. Korshunova, A. N. Sizov, and A. D. Shatrov, Usp. Fiz. Nauk 167, 1201 (1997) [Phys. Usp. 40, 1149(1997)].
- [16] E. A. Novikov, Zh. Eksp. Teor. Fiz. 47, 1919 (1964) [Sov. Phys. JETP 20, 1290 (1964).
- [17] V. I. Klyatskin and V. I. Tatarski , Izv. Vyssh. Uchebn. Zaved., Radiofiz. 15, 1433 (1972).
- [18] S. I. Vainshtein, Zh. Eksp. Teor. Fiz. 58, 153 (1970) [Sov. Phys. JETP 31, 87 (1970)].
- [19] W.H.Matthaeus, C.Smith, Phys.Rev. A 24, 2135 (1981).
- [20] S.Oughton, K.–H.R¨adler, W.H.Matthaeus, Phys.Rev. E 56, 2875 (1997).
- [21] F. Krause and K.-H. Radler, Mean-Field Magnetohydrodynamics and Dynamo Theory (Akademie, Berlin, 1980; Mir, Moscow, 1984).
- [22] I. N. Toptygin, Cosmic Rays in Interplanetary Magnetic Fields (Nauka, Moscow, 1983; Reidel, Dordrecht, 1985).
- [23] A.V.Chechkin, D.P.Sorokin, V.V.Yanovsky,<http://xxx.lanl.gov/ps/hep-th/9306159>
- [24] C. W. Smith, J. W. Bieber, in Solar Wind Eight, AIP Conference Proceedings 382, 498, AIP, New York (1996).
- [25] A. I. Morozov and L. S. Solov'ev, Zh. Tekh. Fiz. 30, 271 (1960) [Sov. Phys. Tech. Phys. 5, 250 (1960)].
- [26] A. G.Khantadze, G. D. Aburdzhania, G. V. Dzhandieri, and Kh. Z. Chargaziya, Fiz.Plazmy (Moscow) 30, 88 (2004) [Plasma Phys. Rep. 30, 83 (2004)].
- [27] D.L.Jaggard, N.Engheta, Electronics Lett. 25, 1060 (1989).
- [28] V. A. Neganov and O. V. Osipov, Izv. Vyssh. Uchebn. Zaved., Radiofiz. 42, 870 (1999).
- [29] N.S.Erokhin, S.S.Moiseev, On the mode conversion in weakly inhomogeneous chiral plasma, Preprint No. 1948, IKI AN RAN (Inst. for Space Research, Russian Academy of Sciences, Moscow, 1996).
- [30] H.T.Torres, P.H.Sakanaka, N.Reggiani, J.Phys.Soc. of Japan 67, 850 (1998).