

Approximation of the energy levels of the particle on a segment by energy levels of the particle in monomial potential well

Michal Demetrian*
Comenius University
Mlynská Dolina M, 842 48, Bratislava IV, Slovak Republic

November 21, 2018

Abstract

This simple text considers an application of Bohr-Sommerfeld quantization rule. It might be of interest for the students of physics.

1 Particle on a segment

The quantum stationary states and possible values of energy of the particle on the segment of a line is given by the Schrödinger equation

$$-\frac{\hbar^2}{2m} \frac{d^2\psi(x)}{dx^2} = E\psi(x) \quad (1)$$

and boundary conditions

$$\psi(0) = \psi(a) = 0, \quad (2)$$

where m is the mass of the particle and a is the length of the segment. It is well known that the conditions (1) and (2) lead to the energy eigenfunctions

$$\psi_n(x) = \sqrt{\frac{2}{a}} \sin\left(\frac{n\pi x}{a}\right), \quad \text{where } n \in \{1, 2, 3, \dots\} \quad (3)$$

and energy eigenvalues

$$E_n = \frac{\pi^2 \hbar^2}{2ma^2} n^2 \sim n^2. \quad (4)$$

Particle on the segment $[0, a]$ can be modeled as particle moving in the line $(-\infty, \infty)$ in the "intuitive" potential energy

$$V(x) = \begin{cases} 0, & x \in [0, a], \\ \infty, & x \notin [0, a]. \end{cases} \quad (5)$$

This kind of potential energy can be, also intuitively as plotted in the figure (1), imagined as "the limit" of the following sequence of potential energy functions

$$V_k(x) = A_0^{(k)} x^{2k}, \quad k \in \{1, 2, 3, \dots\} \quad (6)$$




Figure 1: There are plotted the first six functions (6) with $A_0^{(k)} = 1$ on the panel.

with some positive constants $A_0^{(k)}$.

Our task is to show that the energy levels (4) can be approximated by the energy levels of the particle moving in the potentials (6). The tool we are going to use for this purpose is the well-known Bohr-Sommerfeld quantization rule.

2 Approximative expression for the energy levels of the particle in the potential well $A_0^{(k)} x^{2k}$

The Hamiltonian of the particle in the potential well $V = A_0^{(k)} x^{2k}$ is given by

$$H(p, x) = \frac{p^2}{2m} + A_0^{(k)} x^{2k}. \quad (7)$$

Let $\mathcal{C}(E)$ be the classical periodic trajectory of the particle with the energy E in the phase space (there are no trajectories of other kind in our case). The Bohr-Sommerfeld quantization rule chooses from the set of all classical trajectories only the trajectories for which

$$\oint_{\mathcal{C}(E)} p dx = 2\pi\hbar(n + \gamma_n), \quad (8)$$

where n is natural number and γ_n is the quantity of the order of unity. For our purpose, it is sufficient to write down the rule in less accurate form

$$\oint_{\mathcal{C}(E)} p dx = 2\pi\hbar n. \quad (9)$$

In this way, Bohr-Sommerfeld quantization rule gives us the method to determine approximately the energy levels of the particle by solving the equation (9) with the unknown E and parameter n . The accuracy of this formula grows with growing n . In our case, the particle with the energy E moves in the range $[-x_M(E), x_M(E)]$, where the energy conservation law determines $x_M(E)$ as

$$x_M(E) = \left(\frac{E}{A_0^{(k)}} \right)^{\frac{1}{2k}}. \quad (10)$$

*demetrian@fmph.uniba.sk

Therefore, the left-hand side of Eq. (9) reads

$$\begin{aligned} \oint_{\mathcal{C}(E)} p dx &= 4 \int_0^{x_M} p(x) dx = 4\sqrt{2m} \int_0^{x_M} \sqrt{E - A_0^{(k)} x^{2k}} dx = \\ 4x_M \sqrt{2mE} \int_0^1 \sqrt{1 - y^{2k}} dy &= \frac{4x_M \sqrt{2mE}}{2k} \int_0^1 (1-t)^{1/2} t^{\frac{1}{2k}-1} dt = \\ \frac{2}{k} (2m)^{1/2} \left(A_0^{(k)}\right)^{-\frac{1}{2k}} E^{\frac{1}{2} + \frac{1}{2k}} \mathcal{B}\left(\frac{3}{2}, \frac{1}{2k}\right), \end{aligned}$$

where \mathcal{B} is the Euler's beta-function. Above written equalities together with (9) allow for expressing energetic levels of the system in closed form

$$E_n^{(k)} = \left[\frac{\pi \hbar k}{(2m)^{1/2}} \right]^{\frac{2k}{k+1}} \left(A_0^{(k)}\right)^{\frac{1}{k+1}} \left[\frac{1}{\mathcal{B}\left(\frac{3}{2}, \frac{1}{2k}\right)} \right]^{\frac{2k}{k+1}} n^{\frac{2k}{k+1}}. \quad (11)$$

We are interested in asymptotic behavior of $E_n^{(k)}$ at fixed n and k running to infinity. By making use the facts

$$\mathcal{B}(x, y) = \frac{\Gamma(x)\Gamma(y)}{\Gamma(x+y)} \quad \text{and} \quad \Gamma(x) = \frac{1}{x} - \gamma_E + \mathcal{O}(x) \text{ for } x \rightarrow 0^+,$$

where γ_E is the Euler's constant, and expressing the constant $A_0^{(k)}$ in the form appropriate to keep the width of the well equal to a constant $2a$

$$A_0^{(k)} = \frac{V_0}{a^{2k}}, \quad V_0 = \text{const}, \quad V_0 > 0,$$

we can derive that for $k \rightarrow \infty$

$$E_n^{(k)} = \left[\frac{\pi^2 \hbar^2}{2ma^2} k^2 + k + \mathcal{O}(1) \right] \left[\Gamma\left(\frac{1}{2k}\right) \right]^{-\frac{2k}{k+1}} \left[n^2 + \mathcal{O}\left(\frac{1}{k}\right) \right]$$

and therefore

$$\lim_{k \rightarrow \infty} E_n^{(k)} = \frac{\pi^2 \hbar^2}{2m(2a)^2} n^2 \quad (12)$$

as it should be.