Confinement of hydrodynamic modes on a free surface and their quantum analogs

M. Torres^{*},¹ J. P. Adrados,¹ P. Cobo,² A. Fernández,² C. Guerrero,³ G.

Chiappe,⁴ E. Louis,⁴ J.A. Miralles,⁴ J. A. Vergés,⁵ and J.L. Aragón^{†6}

¹Instituto de Física Aplicada, Consejo Superior de Investigaciones Científicas, Serrano 144, 28006 Madrid, Spain.

²Instituto de Acústica, Consejo Superior de Investigaciones Científicas, Serrano 144, 28006 Madrid, Spain

³Laboratorio de Física de Sistemas Pequeños y Nanotecnología,

Consejo Superior de Investigaciones Científicas, Serrano 144, 28006 Madrid, Spain.

⁴Departamento de Física Aplicada and Unidad Asociada del Consejo Superior de Investigaciones Científicas,

Universidad de Alicante, San Vicente del Raspeig, Alicante 03690, Spain.

⁵Departamento de Teoría de la Materia Condensada, Instituto de Ciencia de Materiales de Madrid,

Consejo Superior de Investigaciones Científicas, Cantoblanco, Madrid 28049, Spain.

⁶Centro de Física Aplicada y Tecnología Avanzada,

Universidad Nacional Autónoma de México, Apartado Postal 1-1010, Querétaro 76000, México.

(Dated: November 21, 2018)

A subtle procedure to confine hydrodynamic modes on the free surface of a fluid is presented here. The experiment consists of a square vessel with an immersed square central well vibrating vertically so that the surface waves generated by the meniscus at the vessel boundary interfere with the bound states of the well. This is a classical analogy of a quantum well where some fundamental phenomena, such as bonding of states and interference between free waves and bound states, can be visualized and controlled. The above mentioned interference leads to a novel hydrodynamic transition from quasiperiodic to periodic patterns. Tight binding numerical calculations are performed here to show that our results could be transferred to design quantum confinements exhibiting electronic quasiperiodic surface states and their rational approximants for the first time.

The interest in experiments of classical analogs, which accurately model the salient features of some quantum systems or other fundamental undulatory phenomena, was firstly raised by the acoustic experiments of Maynard [1], where the analogy between both the acoustic and Schrödinger equations is invoked. In a general overview, a hydrodynamic analogy has also been used to describe the flow of electrons in quantum semiconductor devices [2]. Some experiments with liquid surface waves have been reported later and they presented abstract concepts, such as Bloch states, domain walls and band-gaps in periodic systems [3, 4], Bloch-like states in quasiperiodic systems [5] or novel findings as the superlensing effect [6] in an visually clear manner. On the other hand, the correspondence between the shallow water wave equation and the acoustic wave equation has also been demonstrated [4, 7]. Such correspondences could be exploited to investigate and address formally similar quantum effects as those observed in quantum corrals [8], where an optical analogy has already been proposed [9], and in grain boundaries or simple surface steps [10]. Our main goal is to build up the hydrodynamic analogy of the interference between bound states in a finite quantum well and free states. Then we realized an experimental set up consisting of a square vessel with a single square well drilled at its bottom; both squares are concentric and well diagonals are parallel to the box sides. The immersed well

was expected to work as a weak potential binding surface waves upon dependence on the liquid depth. When the vessel vibrated vertically with such amplitudes that the Faraday instability was prevented, such geometry produced three kinds of linear or weakly non-linear patterns on the free surface of the liquid. The first pattern is a sort of bound state restricted to the surface area occupied by the immersed well that works as a weak potential binding standing plane waves. The second pattern is produced by the meniscus at the vessel walls [11] and it can invade the region of the well depending on the liquid depth and the vessel vibration frequency. Finally, the last pattern arises from the interference between the vessel meniscus waves and the bound states of the well. Patterns observed depend on the liquid depth h_1 , which plays the role of an order parameter by controlling the amplitudes of the bound states inside the immersed well.

Our main observation is summarized in Fig. 2(a), which clearly shows the binding of the surface wave produced by the drilled well when the vibration amplitude is 60 μ m. This pattern will be detailed below but the physics of its origin can be discussed as follows. The bound states arise from an inertial hydrodynamic instability, balanced by the liquid surface tension [12], that grows over the square well region [5]. The bound state amplitudes increase on increasing $1/a^2$; where $a^2 = T/\rho g$, a is the capillary length, T is the liquid surface tension, ρ is the liquid density and $g = g_0 \pm \alpha \omega^2$ is the effective gravity; g_0 is the acceleration due to gravity, α is the vessel vibration amplitude and ω is the vibration angular frequency of the vessel. On the contrary, the meniscus wave amplitude depends on the variation of

^{*}Electronic address: manolo@iec.csic.es

[†]Electronic address: aragon@fata.unam.mx

the meniscus volume during each vessel oscillation and, it grows accordingly when a^2 increases [11, 12]. In our experiment α is about 60 μ m and the meniscus wave amplitude reaches a maximum at a vessel vibration frequency of 64 Hz. On the other hand, the vessel vibration frequency represents the wave state energy. The frequency and the wavelength are related through the well known dispersion relation of the gravity-capillary waves [3, 4, 5, 12]. The present experiment is essentially monochromatic as it occurs in the optical analogs [9] of quantum corrals. To show that Fig. 1(a) is not a Faraday pattern, we present a snapshot of the system vibrating at about 70 μ m in Fig. 1(b), when the Faraday instability is really triggered in the square well. Figure 1(b) shows a higher wavelength that matches with the corresponding period doubling related to a Faraday wave pattern.

We chose a square vessel and the orientated square well configuration to verify that the immersed well confined wave states. Then we used a square methacrylate box with side L of 8 cm where a single square well with depth d of 2 mm and side l of 3.5 cm was drilled at its bottom. The bottom of the vessel was covered with a shallow ethanol layer of depth h_1 . The liquid depth over the well was then $h_2 = h_1 + d$.

As it was already mentioned, the vessel vibration amplitude was 60 μ m, below the Faraday instability threshold set at about 70 μ m. The vessel vibrated vertically at a single frequency lying within the range from 35 to 60 Hz. An optimum frequency value was 50 Hz. Depending on the depth h_1 , our experimental results can be separated into three cases.

Case I. For h_1 lower than 1 mm, the experiment shows two square lattices rotated 45° between them, namely the lattice of the immersed square well is separated from the vessel square lattice (Fig. 1a). The external wave amplitudes are lower than the wave amplitudes within the well and, furthermore, the external wave reflection at the well step [13, 14, 15] is strong. For very shallow liquid layers, waves are only present within the well at the vibration amplitudes of the experiment.

Case II. At a vessel vibration frequency of 50 Hz, when h_1 is 1.2 mm, a quasicrystalline standing wave pattern appears inside the immersed square region whereas the outer wave pattern is a square network (Fig. 2(a)). The immersed square well works as a weak potential and binds standing plane waves with eigenvectors $k_2 \mathbf{u}'_r$ and $k_2 \mathbf{u}'_{u}$, parallel to the well sides. Nevertheless, it is transparent for the standing waves of the square box which tunnel the immersed well framework under the experimental conditions. Inside the square region, the vessel eigenstates have eigenvectors $k_2 \mathbf{u}_x$ and $k_2 \mathbf{u}_y$, parallel to the outer box sides. Thus, the standing state is finally described by $\psi_2(k_2) + \psi'_2(k_2) = A_2[\exp(ik_2x) + \exp(ik_2y)] +$ $A'_{2}[\exp(ik_{2}x') + \exp(ik_{2}y')]$ inside the region of the square well. Equal phases along x and y, and x' and y', respectively, were assumed and $|A_2| = |A'_2|$ for the mentioned

liquid depth and vibration frequency. The interference of standing patterns ψ_2 with ψ'_2 increases the symmetry in the well from square crystalline to octagonal quasicrystalline. Such pattern matches well with the outer one described by $\psi_1(k_1) = A_1[\exp(ik_1x) + \exp(ik_1y)]$. According to the dispersion relation for gravity-capillary waves described elsewhere [3, 4, 5, 12], the difference between k_1 and k_2 is about 2% and the refraction bending of about 1.1° at the boundary of the central window is negligible. Furthermore, the external wave reflection at the well step [13, 14, 15] is also negligible with such parameters. On the other hand, slender outgoing evanescent waves are emitted at the boundary of the well and they play an important role in the matching between patterns. It is important to remark that the parameters of the experiment do not allow through the well known dispersion relation [3, 4, 5, 12] that the wave inside the inner well were a subharmonic Faraday wave and the outer wave were a harmonic meniscus wave. If this were the case, then k_1 and k_2 should be very different.

Case III. When h_1 is increased, the potential of the immersed well is seen increasingly weaker by the system, and A'_2 decreases accordingly. Under such conditions, transitional patterns from a quasicrystalline form to a crystalline one appear gradually on the hydrodynamic window. Figure 2(b) shows a transitional pattern corresponding to a liquid depth h_1 of 1.5 mm and an excitation frequency of 50 Hz. According to crystallographic techniques of image processing described elsewhere [5], the Fourier transforms of the experimental patterns are calculated (Figs. 2(a) and 2(b)) and they are used to depict Fig. 3, where the fast decay of $|A'_2/A_2|^2$ is shown on increasing h_1 . Figure 4 shows four frames of the numerical simulation of the quasicrystal-crystal transition on the hydrodynamic central square window. The complete movie is available as Auxiliary Material.

The appearance of a quasiperiodic pattern inside the immersed square region, in Case II, confirms that the square well is binding standing plane waves. Such a pattern can only be produced by the interference of two square patterns rotated by 45° , namely the pattern of bound standing waves of the square well is transparent to the pattern of standing meniscus waves of the vessel if the adequate conditions of the experimental parameters are fulfilled. On the other hand, the square pattern of the vessel is only observed beyond the region of the central square well. Moreover, there are evanescent waves coming out of the boundary of the square well. The observation of such waves also confirms the analogy with a quantum well [16].

The description of the observed standing wave patterns given in Case II is the same as the used to describe the stationary states of a particle moving in a potential given by a square well. Using these functions, we show a numerical simulation of the observed quasicrystalline pattern in Fig. 3(b). A trial and error numerical method was used to fit at the boundaries and a better matching is obtained if slender outgoing evanescent waves emitted at the boundary of the well are considered.

To test the viability of a quantum scenario analogous to our experimental results, we numerically studied the quantum confinement of a double square well by using a tight-binding Hamiltonian in a $L \times L$ cluster of the square lattice with a single atomic orbital per lattice site [17],

$$\hat{H} = \sum_{m,n} \epsilon_{m,n} |m,n\rangle \langle m,n|$$

$$- \sum_{\langle mn;m'n'\rangle} t_{m,n;m'n'} |m,n\rangle \langle m',n'|,$$

where $|m, n\rangle$ represents an atomic orbital at site (m, n), and $\epsilon_{m,n}$ its energy. In order to simulate the inner square we have explored several possibilities. Outside the inner square, the energy of all orbitals are taken equal to zero. Besides, the hopping energies between nearestneighbor sites (the symbol $\langle \rangle$ denotes that the sum is restricted to nearest-neighbors) were all taken equal to 1. Instead, inside the inner square, we either changed the orbital or the hopping energies. Some illustrative results are shown in Fig. 5. Figures 5(a) and (b) correspond to wavefunctions close to the band bottom, *i.e.*, long wavelengths and high linearity. In Fig. 5(a) the octagonal symmetry within the inner square is clearly visible. Figure 5(b) illustrates an effect that is purely quantum, namely, the effect of the outer square is visible even on a wavefunction localized in the inner square. Although one cannot expect a one-to-one correspondence between the experiments discussed here and this simple quantum simulation, the results clearly suggest that similar effects could be observed in a suitable quantum system. Figure 5(c) shows a peculiar null energy state which is not located at the band bottom. In this state the hopping energy outside the square is equal to 1, whereas it is 0.7 inside the square and the general pattern conspicuously resembles our experiment. The Fourier spectrum of the inner square pattern has been performed and it corresponds to a rational approximant of an octagonal quasicrystal generated by the following vectors: $(2/\sqrt{5},0), (0,2/\sqrt{5}), 2(2/\sqrt{5},1/\sqrt{5}), 2(2/\sqrt{5},-1/\sqrt{5}),$ $2(1/\sqrt{5}, 2/\sqrt{5}), 2(-1/\sqrt{5}, 2/\sqrt{5}).$ Fourier spectra of Figs. 5(a) and (b) correspond to quasicrystalline states. This suggests that the hopping energy could be a parameter to induce quasicrystal-crystal transitions in confined quantum states.

In conclusion, we have described here a hydrodynamic experiment that gives rise to the confinement of wave states on a free surface. It constitutes a stirring macroscopic experimental scenario which models some salient features of a quantum well and stimulates the study and visualization of confined quasiperiodic quantum states for the first time.

Technical support from C. Zorrilla and S. Tehuacanero

is gratefully acknowledged. This work has been partially supported by the Spanish MCYT (BFM20010202 and MAT2002-04429), the Mexicans DGAPA-UNAM (IN-108502-3) and CONACyT (D40615-F), the Argentineans UBACYT (x210 and x447) and Fundación Antorchas, and the University of Alicante.

- S. He and J.D. Maynard, Phys. Rev. Lett. 62, 1888 (1989); J.D. Maynard, Rev. Mod. Phys. 73, 401 (2001).
- [2] C.L. Gardner, SIAM J. Appl. Math. 54, 409 (1994).
- [3] M. Torres, J.P. Adrados and F.R. Montero de Espinosa, Nature (London) 398, 114 (1999)
- [4] M. Torres, J.P. Adrados, F.R. Montero de Espinosa, D. García-Pablos and J. Fayos, Phys. Rev. E 63, 11204 (2000).
- [5] M. Torres, J.P. Adrados, J.L. Aragón, P. Cobo and S. Tehuacanero, Phys. Rev. Lett. 90, 114501 (2003). See also Physical Review Focus: http://focus.aps.org/story/v11/st11
- [6] X. Hu, Y. Shen, X. Liu, R. Fu, and J. Zi, Phys. Rev. E 69, 030201(R) (2004); M. Peplow, Nature (London) 428, 713 (2004).
- [7] T. Chou, J. Fluid Mech. **369**, 333 (1998).
- M.F. Crommie, C.P. Lutz and D.M. Eigler, Nature (London) 363, 524 (1993); M.F. Crommie, C.P. Lutz and D.M. Eigler, Science 262, 218 (1993).
- [9] Gérard Colas des Francs, et al., Phys. Rev. Lett. 86, 4950 (2001); C. Chicanne, et al., Phys. Rev. Lett. 88, 097402 (2002).
- [10] Y. Hasegawa and P. Avouris, Phys. Rev. Lett. 71, 1071 (1993).
- [11] S. Douady, J. Fluid Mech. 221, 383 (1990).
- [12] L.D. Landau and E.M. Lifshitz, *Fluid Mechanics* (Pergamon Press, London, 1959).
- [13] H. Lamb, *Hydrodynamics* (Cambridge University Press, Cambridge, 1932).
- [14] E.F. Bartholomeusz, Proc. Cambridge Phil. Soc. 54, 106 (1958).
- [15] J.W. Miles, J. Fluid Mech. 28, 755 (1967).
- [16] A. Messiah, Mècanique Quantique (Dunod, Paris, 1962).
- [17] E. Cuevas, E. Louis, and J. A. Vergés Phys. Rev. Lett. 77, 1970 (1996).

FIG. 1: (a) Snapshot of the system below the Faraday instability onset, vibrating at 35 Hz with $h_1 = 0.35$ mm. Two clearly separated wave patterns are shown. The amplitude of the meniscus wave is much lower than that of the well-bound states. The reflexion coefficient at the well step is about 0.5. (b) Snapshot of the system at the onset of the Faraday instability in the square well.

FIG. 2: (a) Snapshot of the system vibrating at 50 Hz for $h_1 = 1.2$ mm. The well is transparent for the meniscus wave. The interference between such wave and the well-bound states gives rise to a perfect quasicrystalline pattern (Top). Fourier transform of the quasicrystalline wave pattern. Some secondary rings with a lower intensity appear. Such rings are due to weak nonlinearities corresponding to the parametrically driven experiment (Bottom). (b) Snapshot of the system vibrating at 50 Hz for $h_1 = 1.5$ mm. The amplitude of the bound states is lower than that of the meniscus wave. A quasicrystal-crystal intermediate state appears on the well (Top). Fourier transform of the transitional pattern. In the main octagonal ring there are four brighter spots (Bottom).

FIG. 3: (a) Decay of $|A'_2/A_2|^2$ when the depth of the liquid layer increases; the spots (\blacklozenge) on the figure are experimental points. (b) Simulation of the pattern in Fig. 2(a) by means of a numerical trial and error fitting of the standing waves of the box and those waves bound by the square well, as well as the evanescent waves outgoing from the well boundary.

FIG. 4: Four frames of the numerical simulation of the quasicrystal- crystal transition are shown as the liquid depth h_1 increases. The corresponding Fourier transforms are shown as insets. A complete movie is available as Auxiliary Material.

FIG. 5: Quantum wavefunctions of the tight binding Hamiltonian described in the text.

This figure "figure1.jpg" is available in "jpg" format from:

This figure "figure2.jpg" is available in "jpg" format from:

This figure "figure3.jpg" is available in "jpg" format from:

This figure "figure4.jpg" is available in "jpg" format from:

This figure "figure5.jpg" is available in "jpg" format from: