Is E112 a relatively inert element? Benchmark relativistic correlation study of spectroscopic constants in E112H and its cation.

N.S. Mosyagin,* T.A. Isaev, and A.V. Titov

Petersburg Nuclear Physics Institute, Gatchina,

St.-Petersburg district 188300, Russia

(Dated: July 9, 2018)

Abstract

We report the first results of relativistic correlation calculation of the spectroscopic properties for the ground state of E112H and its cation in which spin-orbit interaction is taken into account non-perturbatively. Studying the properties of E112 (eka-Hg) is required for chemical identification of its long-lived isotope, 283 112. It is shown that appropriate accounting for spin-orbit effects leads to dramatic impact on the properties of E112H whereas they are not so important for E112H⁺. The calculated equilibrium distance, $R_e^{\rm calc}$ =1.662 Å, in E112H is notably smaller than $R_e^{\rm expt}$ =(1.738 ± 0.003) Å and $R_e^{\rm calc}$ =1.738 Å in HgH, whereas the dissociation energy, $D_e^{\rm calc}$ =0.42 eV, in E112H is close to $D_e^{\rm expt}$ =0.46 eV and $D_e^{\rm calc}$ =0.41 eV in HgH. These data are quite different from $R_e^{\rm NH}$ =1.829 Å and $D_e^{\rm NH}$ =0.06 eV obtained for E112H within the scalar-relativistic Douglas-Kroll approximation [Nakajima and Hirao, Chem. Phys. Lett., **329**, 511 (2000)]. Our results indicate that E112 should not be expected to be "more inert" than Hg in opposite to the results by other authors.

Introduction. The superheavy element 112 (eka-Hg) was discovered at GSI (Darmstadt) in 1996 within the "cold" fusion reaction¹. The recent observation at FLNR (Dubna) of the " α –SF" chain, attributed to 4 sec α -decay branch of ²⁸³112 followed by a 0.2 sec spontaneous fission of ²⁷⁹110 (Ds)², brought up the question, what species was observed in the previous "hot" fusion FLNR experiment³. Moreover, the production of ²⁸³112 in the reaction of ⁴⁸Ca and ²³⁸U was not confirmed at LBNL (Berkeley)^{4,5}. However, a very specific decay mode of the short " α –SF" chain offered a unique chance to unambiguously identify ²⁸³112 in a chemical experiment.

To our knowledge, starting from the papers of Pitzer⁶ and Fricke⁷ in 1975 it was mainly suggested by other authors that E112 behaves rather like a rare gas than Hg. In Ref. 8, the confusing conclusions about both relative inertness of E112 as compared to Hg and similarity of E112 with Hg were made in the abstract and conclusion, respectively. The first attempt to identify E112 chemically was made at FLNR^{9,10} but no spontaneous fissions were detected. It was interpreted as indication of the Rn-like behavior of E112 as well.

The chemical experiments on studying properties of E112 are currently under way at FLNR¹⁰, similar work is in progress at GSI and PSI (Villigen)¹¹ that involves the attempt to clarify the recent observation of the decay chains and fission products associated with the production of E114 and E116²: being their decay product, E112 should be detectable in gasphase chromatographic experiments. The experimental study of superheavy element (SHE) properties (see Refs. 12,13 and references) is very difficult because of their short half-lives and extremely small quantities, only single atoms are available for research. In this connection, reliable theoretical prediction of their properties based on benchmark *ab initio* calculations is highly desirable. As a first step in the extensive study of chemistry of E112, bonding in simple diatomic molecules such as E112H should be studied. The earlier studies of ekamercury fluorides¹⁴ are not so sensitive because both mercury and xenon are known to burn in fluorine atmosphere. On the other hand, the Hg₂, Xe₂ and E112₂ dimers^{8,15} are Van der Waals systems with a small dissociation energy. By contrast, the ground state RnH and XeH molecules are not observed in the gas phase, whereas HgH can be obtained by radiofrequency discharge in hydrogen and metal vapor (see, e.g., Ref. 16).

It was shown in Ref. 17 on example of E112 and other SHE's that the errors in calculations due to employing the point nucleus (instead of the realistic Fermi nuclear model) reach 0.4 eV for transition energies between low-lying states, whereas neglecting the Breit effects leads to

the errors up to 0.1 eV. The generalized relativistic effective core potentials (GRECP's)¹⁸ were generated for E112 and other SHE's¹⁷ which allow one to simulate Breit interaction and Fermi nuclear model by economic way but with very high precision¹⁹. The accuracy of these GRECP's and of the RECP's of other groups was estimated in atomic finite-difference SCF calculations with Coulomb two-electron interaction and point nucleus as compared to the corresponding all-electron Dirac-Fock-Breit calculations with the Fermi nuclear model. It was justified and checked in Refs. 17,19 that the GRECP method allows one to carry out reliable calculations of SHE's and their compounds within the level of "chemical accuracy" (1 kcal/mol, 0.043 eV, or 350 cm⁻¹ for valence transition energies) when the valence and outer core shells are appropriately treated. Hence, the overall accuracy of calculations in heavy-atom molecules is limited, in practice, by current possibilities of the correlation methods and codes and not by the GRECP approximation.

In this paper, we present GRECP calculations of spectroscopic constants for the ground states of the E112H molecule and its cation exploiting MOLGEP code²⁰. To our knowledge, only three calculations on E112H and E112H⁺ were published^{8,14,21}. In Ref. 21, the third order Douglas-Kroll (DK3) method was applied to calculation of E112H and its ions. The correlations were taken into account by the second-order Møller-Plesset perturbation theory (MP2) and by the coupled cluster method with single, double (and triple) cluster amplitudes, CCSD(T), for 19 external electrons of the E112H molecule. It is not clear from Ref. 21 why the correlations with the 6p shell of E112 are considered but correlations with the 6p and 5p shells are not whereas the small value of calculated dissociation energy, $D_e^{\rm NH}=0.06\,{\rm eV}$, could be strongly influenced by the latters. The 6p and 6p shells are closely localized in space, whereas the 5p and 6p shells have close orbital energies. The effect of the finite nuclear size was taken into account but the Breit effects and even the spin-orbit (SO) interaction were neglected (i.e., only scalar-relativistic calculations were made). It is clear that both effects are increased with the nuclear charge Z, therefore, these approximations can be inappropriate for SHE compounds even if they are justified for their lighter analogues.

In Refs. 8,14, the RECP calculations of E112H⁺ (but not E112H) were carried out by the MP2 and CCSD(T) methods. In Ref. 14, complete active space SCF and multireference configuration interaction (MRCI) calculations of E112H⁺ were also performed⁴¹. The SO, finite nuclear size and Breit effects were taken into account at the generation stage of the pseudopotential (PP) of Seth *et al.*¹⁴. It should be noted that the parameters of this PP

were fitted with the help of the adjustment procedure based on the LS-coupling scheme (which was, in general, inappropriate for SHE's since the errors of these PP's were up to 4 eV for valence energies in our test calculations accounting for the SO interaction non-perturbatively). Recently, a new PP for E112 was generated by Seth et al. using the jj-coupling scheme⁴², but we are not informed about molecular calculations with this PP. The SO interaction was taken into account in the calculations from Ref. 8. However, the author applied the 20-electron RECP of Nash et al.²². The Breit effects were not considered at the generation stage of this RECP. It is not clear from Refs. 8,22 which nuclear model was used there. In our test calculations¹⁷, the errors of this RECP in reproducing the results of the all-electron Dirac-Fock-Breit calculations with the Fermi nuclear model were up to 1 eV for transition energies between low-lying states. Moreover, the basis set superposition errors (BSSE's)^{23,24} were not estimated in the calculations from Ref. 8.

Methods and calculations. The GRECP^{17,18,19,25}, Fock-space relativistic CC-SD (RCC-SD)^{26,27} and spin-orbit direct configuration interaction (SODCI)^{28,29} methods used for the present calculations are well described in literature. The gaussian expansions of our GRECP and (16, 21, 16, 12, 14)/[4, 6, 4, 2, 1] basis set for E112 are available at our website http://www.qchem.pnpi.spb.ru/Basis/. In the SODCI calculation, the relativistic scheme of configuration selection was applied³⁰.

Two series of Fock-space RCC-SD calculations were performed for E112H with the GRECP. The ground state of the cation E112H⁺ served as reference in the first series (denoted by RCC-SD-1), and the Fock-space scheme was

$$E112H^+ \to E112H, \tag{1}$$

with an electron added in the lowest unoccupied σ orbital of E112H⁺. The second series (RCC-SD-2) started from the ground state of the anion E112H⁻ as reference using the Fock-space scheme

$$E112H^{-} \rightarrow E112H \rightarrow E112H^{+}, \tag{2}$$

with electrons removed from the highest occupied σ orbital of E112H⁻. Moreover, the RCC-SD calculations of E112 (to calculate counterpoise corrections and D_e) were carried out where the $6d^{10}7s^2$ ground state of the E112 atom was used as reference and the Fock-space scheme was

$$E112 \to E112^+, \tag{3}$$

with an electron removed from the 6d or 7s shell.

Our test atomic RCC calculations on E112 showed that at least 34 external electrons of the atom (occupying the $5f, 6s, 6p, 6d, 7s, \ldots$ shells) should be correlated and the basis set should include up to *i*-type harmonics (l=6) in order to calculate the excitation and ionization energies with "chemical accuracy". Nevertheless, we expect that the contributions of the core correlations will be less important for the molecule than for the atom as was in our similar calculations^{25,31,32} on Hg and HgH. This is, in particular, supported by a large orbital energy separation between 5p and 5d shells in Hg ($\varepsilon[5p_{3/2};5d_{3/2}]\approx-2.8;-0.65$ a.u.) and by a comparable separation between 6p and 6d shells in E112 ($\varepsilon[6p_{3/2};6d_{3/2}]\approx-2.4;-0.56$ a.u.) Therefore, only 13 external electrons for the E112H molecule (12 electrons for E112H⁺) were correlated in the present calculations. The calculations with the larger number of correlated electrons when SO interaction is explicitly treated are rather time-consuming and suggested in future.

In scalar-relativistic CC-SD calculations, we have also estimated (see Table I) that correlations with the 6p shell of E112 give relatively small contributions to the spectroscopic constants in E112H and E112H⁺ except for D_e in E112H⁺. This cation dissociates to E112⁺($6d_{3/2}^46d_{5/2}^57s_{1/2}^2$) + H(1 s^1)³³ in contrast to HgH⁺, which dissociates to Hg⁺($5d_{3/2}^45d_{5/2}^66s_{1/2}^1$) + H(1 s^1). The 6p shell is closely localized to the 6d shell, therefore, the correlations between these shells have to be important for transitions with an essential change in the occupation number for the 6d shell. In principle, the D_e value for E112H⁺ can be easily corrected using our atomic RCC results for ionization potential of the $6d_{5/2}$ subshell of E112. However, we observed large compensations between contributions accounting for correlations with core shells and for basis functions with high angular momenta. Thus, the above ionization potential from calculation with 12 correlated electrons in the basis including up to g-harmonics (l=4) differs from that with 52 electrons and l up to 8 only on +773 cm⁻¹ (0.10 eV).

The calculations were carried out for 15 internuclear distances from 2.3 a.u. to 3.7 a.u. with interval of 0.1 a.u. The spectroscopic constants were calculated by the Dunham method in the Born-Oppenheimer approximation. All our RCC and SODCI results reported in Table I, were improved using counterpoise corrections $(CPC's)^{23,24}$ calculated for the E112 $6d^{10}7s^2$ state with a ghost H atom. CPC's calculated for the ground state of the H atom are about 1 cm^{-1} , and are, therefore, ignored.

Results and discussion. Our results for the ground states of E112H and E112H⁺ are collected in Table I. The corresponding results^{25,32} for HgH and HgH⁺ and the results of other groups^{8,14,21} for E112H and E112H⁺ are also presented for comparison. In the GRECP/RCC-SD-1 calculations one can observe the bond length contraction for E112H and E112H⁺ on 0.07 and 0.06 Å with respect to HgH and HgH⁺. Detailed comparison of our results for HgH and HgH⁺ with the results of other groups and the experimental data can be found in Ref. 32.

Our RCC-SD values for spectroscopic constants show considerable differences between two Fock-space schemes, GRECP/RCC-SD-1 and GRECP/RCC-SD-2. Such differences are caused by the truncation of the CC operator, they indicate significant contributions of the omitted higher-order (triple, etc.) cluster amplitudes (HOCA). HOCA influence on the total energies in each point of the potential curves were estimated with the help of the configuration interaction corrections on HOCA³⁴ calculated as differences in the total energies of the SODCI and RCC-SD-1 values. In these calculations, the same numbers of electrons were correlated as in the above RCC-SD case, but a reduced basis set, [4, 4, 3, 1] on E112 and [3, 2] on H, was used because approaching the full configuration interaction limit in SODCI calculations becames too time-consuming for larger basis sets.

Except for the dissociation limit, HOCA has small effect on E112H⁺ since the cation is a closed shell system. It is well known that CC approach works particularly well for the closed-shell states which is confirmed by the comparison of the GRECP/12e-RCC-SD-1 results with and without HOCA correction for E112H⁺. The change in D_e is mainly due to different ionization potentials for the $6d_{5/2}$ electron of the E112 atom in the RCC-SD-1 and SODCI calculations. The GRECP/12e-RCC-SD-2 results show considerable dictinctions from the results corrected by HOCA because E112H⁺ is calculated in the high Fock-space sector, (2,0), in which some lost of accuracy takes place. The HOCA contribution for E112H is important. Similar trend was observed in our GRECP/RCC calculations³² on HgH when the effect of the triple cluster amplitudes was taken into account for 13 electrons that essentially improved the agreement with the experimental data.

The differences between DK3/CCSD(T)²¹ and our GRECP/RCC-SD-1 + HOCA results are small for E112H⁺, but are essentially larger for E112H. In particular, unlike our GRECP/13e-RCC-SD-1 + HOCA bond length, the DK3/19e-CCSD(T) value²¹ for E112H is larger than the experimental data^{16,35,36,37} for HgH. It worth to note that the DK3/CCSD(T)

calculations²¹ are scalar-relativistic whereas our RCC and SODCI calculations are performed with the spin-dependent GRECP. To check the effect of SO interaction, we have also carried out scalar-relativistic CC calculations with the spin-averaged GRECP part. The same basis, number of correlated electrons, Fock-space schemes, etc. were taken as in the RCC calculations. One can see from comparison of our CC-SD and RCC-SD results that the SO effect is small for E112H⁺ (except for D_e) but is very essential for E112H.

The R_e and w_e values for E112H⁺ by the PP/CCSD(T) method from Ref. 14 (see the footnote in Table I) differ from the DK3/18e-CCSD(T) results²¹ by -0.017 Å and +45 cm⁻¹. The corresponding PP/CCSD(T)+SO values differ from our GRECP/12e-RCC-SD-1 + HOCA results by -0.023 Å and +126 cm⁻¹. The RECP/RCCSD(T) equilibrium distance for E112H⁺ from Ref. 8 differs from our GRECP/RCC-SD-1 + HOCA result by +0.04 Å and even more from the results of other groups^{14,21}. The difference between the RECP/RCCSD(T) equilibrium distance calculated in Ref. 8 for HgH⁺ and the experimental datum is -0.04 Å. Thus, one can observe the larger bond length for E112H⁺ by 0.03 Å in comparison with HgH⁺ in the RECP/RCCSD(T) calculations⁸ (in contrast to our results).

One, however, can expect rather some increase in R_e for E112H⁺ and decrease for E112H when accounting for correlations with the innermore shells. In our scalar-relativistic calculations, some small increase in D_e was observed with enlarging the basis set.

Conclusions. It is well known that properties of SHE's are somewhat different from those of their lighter analogues due to very strong relativistic effects first of all (see Ref. 27 and references). Therefore, even those approximations which work well for the lighter analogues (neglecting the SO interaction for Σ states, the effective state of a considered atom in a molecule, the preferred valency, etc.) should not be used for SHE's without serious checking and analyzing. The calculated equilibrium distance, R_e , in E112H is notably smaller than that in HgH. Therefore, one can also expect smaller bond lengths for the other E112 compounds in comparison with the Hg ones. There is a long-term discussion in scientific community whether E112 will behave like Hg or Rn. The ground state RnH and XeH molecules are not observed in the gas phase. Our calculations for the E112H molecule do not predict large dissociation energy, D_e , but it is yet close to that of HgH. Therefore, we believe that the singly-valent ground state E112 compounds will rather resemble the Hg compounds than the noble gas patterns that is also supported by calculations of other E112 compounds^{38,39,40}.

ACKNOWLEDGMENTS. The SODCI code of R.J.Buenker and his colleagues H.-P.Liebermann and A.B.Alekseyev together with the RCC-SD code of U.Kaldor, E.Eliav, and A.Landau were used in our calculations. We are grateful to A.Zaitsevskii and Yu.Tchuvil'sky for discussions and corrections in the paper. The present work is supported by the RFBR grant 03–03–32335. N.M. thanks Russian Science Support Foundation.

- *Electronic address: mosyagin@pnpi.spb.ru; URL: http://www.qchem.pnpi.spb.ru
- ¹S. Hofmann, V. Ninov, F. P. Hessberger, et al., Z. Phys. A **354**, 229 (1996).
- ² Y. T. Oganessian, V. K. Utyonkov, Y. V. Lobanov, et al., Phys. Rev. C **70**, 064609 (2004).
- ³ Y. T. Oganessian, A. V. Yeremin, G. G. Gulbekian, et al., Eur. Phys. J. A 5, 63 (1999).
- ⁴ W. Loveland, K. E. Gregorich, J. B. Patin, et al., Phys. Rev. C **66**, 044617 (2002).
- ⁵ K. E. Gregorich, W. Loveland, D. Peterson, et al., Phys. Rev. C **72**, 014605 (2005).
- ⁶ K. S. Pitzer, J. Chem. Phys. **63**, 1032 (1975).
- ⁷B. Fricke, Struct. Bond. **21**, 89 (1975).
- ⁸ C. S. Nash, J. Phys. Chem. A **109**, 3493 (2005).
- ⁹ A. Yakushev, G. V. Bulkanov, M. L. Chelnokov, et al., Radiochim. Acta **89**, 743 (2001).
- ¹⁰ A. Yakushev, I. Zvara, Y. T. Oganessian, et al., Radiochim. Acta **91**, 433 (2003).
- ¹¹ R. Eichler and S. Soverna, Physics of Atomic Nuclei **66**, 1146 (2003).
- ¹² M. Schädel, ed., The Chemistry of Superheavy Elements (Kluwer, Dordrecht, 2003), 318 pp.
- ¹³ M. Schädel, Angew. Chem. Int. Ed. **45**, 368 (2006), [English].
- 14 M. Seth, P. Schwerdtfeger, and M. Dolg, J. Chem. Phys. **106**, 3623 (1997).
- 15 J. Anton, B. Fricke, and P. Schwerdtfeger, Chem. Phys. $\mathbf{311}$, 97 (2005).
- ¹⁶ J. Dufavard, B. Majournat, and O. Nedelec, Chem. Phys. **128**, 537 (1988).
- ¹⁷ N. S. Mosyagin, A. N. Petrov, A. V. Titov, and I. I. Tupitsyn, in *Recent Advances in the Theory of Chemical and Physical Systems* (2006), vol. 15 of *Progr. Theor. Chem. Phys.*, [arXiv: physics/0505207].
- ¹⁸ A. V. Titov and N. S. Mosvagin, Int. J. Quantum Chem. **71**, 359 (1999).
- ¹⁹ A. N. Petrov, N. S. Mosyagin, A. V. Titov, and I. I. Tupitsyn, J. Phys. B **37**, 4621 (2004).
- ²⁰ A. V. Titov, A. N. Petrov, A. I. Panin, and Y. G. Khait, program package for calculation of molecular matrix elements with the Generalized RECP.
- $^{21}\,\mathrm{T}.$ Nakajima and K. Hirao, Chem. Phys. Lett. $\mathbf{329},\,511$ (2000).
- ²² C. S. Nash, B. E. Bursten, and W. C. Ermler, J. Chem. Phys. **106**, 5133 (1997), [Erratum: JCP 111 (1999) 2347].
- ²³ M. Gutowski, J. H. van Lenthe, J. Verbeek, F. B. van Duijneveldt, and G. Chałasiński, Chem. Phys. Lett. 124, 370 (1986).

- ²⁴ B. Liu and A. D. McLean, J. Chem. Phys. **91**, 2348 (1989).
- ²⁵ N. S. Mosyagin and A. V. Titov, J. Chem. Phys. **122**, 234106 (2005).
- 26 E. Eliav and U. Kaldor, Chem. Phys. Lett. **248**, 405 (1996).
- ²⁷ U. Kaldor, E. Eliav, and A. Landau, in *Relativistic Electronic Structure Theory. Part 2. Applications*, edited by P. Schwerdtfeger (Elsevier, Amsterdam, 2004), p. 81.
- ²⁸ R. J. Buenker and S. Krebs, in *Recent Advances in Multireference Methods*, edited by K. Hirao (World Scientific, Singapore, 1999), pp. 1–29.
- ²⁹ A. B. Alekseyev, H.-P. Liebermann, and R. J. Buenker, in *Recent Advances in Relativistic Molecular Theory*, edited by K. Hirao and Y. Ishikawa (World Scientific, Singapore, 2004), pp. 65–105.
- ³⁰ A. V. Titov, N. S. Mosyagin, A. B. Alekseyev, and R. J. Buenker, Int. J. Quantum Chem. 81, 409 (2001).
- ³¹ N. S. Mosyagin, E. Eliav, A. V. Titov, and U. Kaldor, J. Phys. B **33**, 667 (2000).
- ³² N. S. Mosyagin, A. V. Titov, E. Eliav, and U. Kaldor, J. Chem. Phys. **115**, 2007 (2001).
- $^{33}\,\mathrm{E}.$ Eliav, U. Kaldor, and Y. Ishikawa, Phys. Rev. A $\mathbf{52},\,2765$ (1995).
- ³⁴ T. A. Isaev, N. S. Mosyagin, M. G. Kozlov, A. V. Titov, E. Eliav, and U. Kaldor, J. Phys. B 33, 5139 (2000).
- ³⁵G. Herzberg, Spectra of Diatomic Molecules, Molecular spectra and Molecular structure (Van Nostrand-Reinhold, New York, 1950).
- ³⁶ W. C. Stwalley, J. Chem. Phys. **63**, 3062 (1975).
- ³⁷ K. P. Huber and G. Herzberg, Constants of Diatomic Molecules (Van Nostrand-Reinhold, New York, 1979).
- ³⁸ V. Pershina, T. Bastug, T. Jacob, B. Fricke, and S. Varga, Chem. Phys. Lett. **365**, 176 (2002).
- ³⁹ A. V. Zaitsevskii et al. (2006), to be published.
- ⁴⁰ A. N. Petrov et al. (2006), to be published.
- ⁴¹ 12 electrons were correlated in MRCI whereas that number is not explicitly declared for CCSD(T) calculations with 20-electron PP; most likely, all they are correlated.
- ⁴² P. Schwerdtfeger, private communication, 2003.

TABLE I: Spectroscopic constants of the ground states of the E112H molecule and the E112H⁺ ion from two-component RCC-SD and scalar-relativistic CC-SD calculations with GRECP in the H (8,4,3)/[4,2,1] ANO and E112 (16,21,16,12,14)/[4,6,4,2,1] basis set. Our corresponding results for HgH and HgH⁺ and the results of other groups for E112H and E112H⁺ are also presented for comparison. R_e is in Å, D_e in eV, Y_{02} in 10^{-6} cm⁻¹, other values in cm⁻¹.

Molecula	Method	R_e	w_e	D_e	B_e	$w_e x_e$	α_e	$-Y_{02}$
Our calculations:								
$\mathrm{HgH^{+}}$	GRECP/12e-RCC-SD-1	1.596	2037	2.67	6.60	39	0.200	279
$_{\rm HgH^+}$	GRECP/12e-RCC-SD(T)-1	1.599	2013	2.68	6.58	41	0.208	282
HgH ⁺	Experiment ^{35,37}	1.594 ± 0.000	2031 ± 3	$(2.75\pm0.36)^{a}$	$6.61 {\pm} 0.00$	44 ± 3	$0.206 {\pm} 0.000$	285 ± 0
Our calculations:								
$E112H^{+}$	GRECP/20e-CC-SD-1	1.537	2587	4.60	7.10	46	0.198	215
$E112H^{+}$	GRECP/20e-CC-SD-2	1.531	2681	4.46	7.15	35	0.168	205
$E112H^{+}$	GRECP/18e-CC-SD-1	1.537	2588	4.61	7.10	47	0.198	215
$E112H^{+}$	GRECP/18e-CC-SD-2	1.531	2680	4.46	7.16	35	0.169	205
$E112H^{+}$	GRECP/12e-CC-SD-1	1.535	2590	4.96	7.12	47	0.200	216
$E112H^{+}$	GRECP/12e-CC-SD-2	1.527	2679	4.75	7.19	37	0.175	208
$E112H^{+}$	GRECP/12e-RCC-SD-1	1.537	2569	3.96	7.11	47	0.201	218
$E112H^{+}$	GRECP/12e-RCC-SD-2	1.519	2752	3.80	7.28	45	0.187	204
${f E}112{f H}^+$		1.540	2547	4.35	7.08	45	0.195	220
Other groups' calculations:								
$E112H^{+}$	$\mathrm{DK}3/18\mathrm{e\text{-}CCSD}^{21}$	1.528	2621		7.18			
$E112H^{+}$	$DK3/18e-CCSD(T)^{21}$	1.532	2595		7.15			
$E112H^{+}$	$PP/CCSD(T)^{14b}$	1.515	2640	5.15		51		
$E112H^{+}$	PP/MRCI+SO ^{14b}	1.503	2620	3.86				
$E112H^{+}$	$PP/CCSD(T)+SO^{14b}$	1.517	2673	4.09		52		
$E112H^{+}$	RECP/RCCSD(T) ^{8b}	1.583		3.50				
Our calculations:								
HgH	GRECP/13e-RCC-SD-1	1.709	1575	0.35	5.76	56	0.262	312
HgH	GRECP/13e-RCC-SD(T)-1	1.738	1395	0.41	5.56	83	0.348	363
HgH	Experiment 16,35,36	1.738 ± 0.003	1403 ± 18	0.46 ± 0.00	5.57 ± 0.02	98±23	0.337 ± 0.067	345 ± 1
HgH	Experiment ³⁷	[1.766] ^c	$[1203]^{c}$	0.46	$[5.39]^{c}$			$[395]^{c}$
Our calculations:								
E112H	GRECP/21e-CC-SD-1	1.742	1438	-0.03	5.53	113	0.409	340
E112H	GRECP/21e-CC-SD-2	1.801	1104	-0.02	5.15			
E112H	GRECP/19e-CC-SD-1	1.741	1439	-0.03	5.53	113	0.409	340
E112H	GRECP/19e-CC-SD-2	1.801	1102	-0.02	5.15			
E112H	GRECP/13e-CC-SD-1	1.746	1402	-0.03	5.50	119	0.429	354
E112H	GRECP/13e-CC-SD-2	1.808	1038	-0.05	5.10			
E112H	GRECP/13e-RCC-SD-1	1.638	1859	0.36	6.25	95	0.338	288
E112H	GRECP/13e-RCC-SD-2	1.663	1649	0.32	6.06	123	0.425	340
E112H	GRECP/13e-RCC-SD-1 + HOCA	1.662	1800	0.42	6.07	152	0.385	287
Other groups' calculations:								
E112H	$\mathrm{DK}3/19\mathrm{e\text{-}CCSD}^{21}$	1.823	991	0.04	5.05			
E112H	$DK3/19e\text{-}CCSD(T)^{21}$	1.829	1007	0.06	5.02			

 $^{^{\}rm a}$ Cited in Refs. 35,37 as uncertain.

^b Note that the RCCSD(T), RMRCI+SO, RCCSD(T)+SO values from Ref. 14 and the CCSD(T) values from Ref. 8 are listed. The acronyms for these calculations (the last one is in the jj-coupling scheme, the other ones are scalar-relativistic where the second and third ones are corrected for the SO effects) are redefined in accordance with the other notations of the present paper.

^c Cited in Ref. 37 as corresponding to the zero vibrational level.