

Molecular orbitals and strong-field approximation

Thomas Kim Kjeldsen and Lars Bojer Madsen

Department of Physics and Astronomy, University of Aarhus, 8000 Århus C, Denmark

V.I. Usachenko and S.-I. Chu [Phys. Rev. A **71**, 063410 (2005)] discuss the molecular strong-field approximation in the velocity gauge formulation and indicate that some of our earlier velocity gauge calculations are inaccurate. Here we comment on the results of Usachenko and Chu. First, we show that the molecular orbitals used by Usachenko and Chu do not have the correct symmetry, and second, that it is an oversimplification to describe the molecular orbitals in terms of just a single linear combination of two atomic orbitals. Finally, some values for the generalized Bessel function are given for comparison.

PACS numbers: 33.80.Rv,32.80.Rm

I. INTRODUCTION

The molecular strong-field approximation (MO-SFA) was formulated in the velocity gauge in Ref. [1], and more recently in the length gauge [2]. The latter work shows that the velocity gauge MO-SFA with the initial state of the highest occupied molecular orbital (HOMO) obtained in a self-consistent Hartree-Fock (HF) calculations does not account for the observed orientation-dependent ionization of N_2 [3]. Only the corresponding length gauge MO-SFA [2] and the molecular tunneling theory [4] give the correct behavior.

The orientation dependence of the strong-field ionization of N_2 was revisited in Ref. [5]. There a simple model is made for the $3\sigma_g$ HOMO in terms of a *single* linear combination of atomic orbitals (LCAO) using two atomic p orbitals. Then the MO-SFA is applied in the velocity gauge to calculate the ionization rate, and it is concluded that the velocity MO-SFA, with such an approximation for the HOMO, is capable of predicting the observed minimum [3] in the rate for perpendicular orientation of the internuclear axis with respect to the linear polarization of the laser. Based on these findings it is stated [5] that our earlier velocity gauge calculations [2] are inaccurate because of inaccuracies in our (i) molecular orbital and (ii) generalized Bessel function. Here we refute both these claims by giving details of our orbitals and generalized Bessel functions. Additionally, we show that apart from having the wrong symmetry, the model used for the $3\sigma_g$ HOMO of N_2 in Ref. [5] is too simple and we therefore believe that the agreement in [5] with experiment is accidental. To account for the molecular structure more LCAOs have to be included as, e.g., in HF calculations. With such a self-consistent wave function, the velocity gauge MO-SFA, indeed gives the wrong prediction for the orientation-dependent ionization [2]. Finally, we present accurate values from [2] of the generalized Bessel function since also in this case some discrepancies exist with Ref. [5] – a discrepancy which can be attributed to the difference in wavelength (we use 800 nm light [2], in [5] 795 nm light is used [6]).

II. MOLECULAR ORBITALS

The idea of the LCAO approach to molecular orbital theory is to identify the molecular orbitals in a basis of atomic orbitals. For a homonuclear diatomic molecule, we write a molecular wave function $\Psi(\mathbf{r})$ as a superposition of atomic orbitals which are centered on each of the atomic cores. The nuclei are located at the positions $\pm\mathbf{R}/2$ with respect to the center of the molecule. The molecular orbitals are the eigenfunctions of the Fock operator, i.e., of the kinetic energy operator and the full HF potential. These eigenfunctions can be obtained by a diagonalization of the matrix representation of the Fock operator in the atomic centered basis, $\phi_{nlm}(\mathbf{r} \mp \mathbf{R}/2)$. A great simplification of the problem is achieved if the basis functions are symmetrized according to the molecular point group ($D_{\infty h}$). Consider, e.g., the atomic $1s$ [$\phi_{100}(\mathbf{r} \mp \mathbf{R}/2)$] orbitals. A change of basis leads to a new set of basis functions

$$\psi_{\sigma_g 1s}(\mathbf{r}) = \mathcal{N}_{\sigma_g 1s}[\phi_{100}(\mathbf{r} - \frac{\mathbf{R}}{2}) + \phi_{100}(\mathbf{r} + \frac{\mathbf{R}}{2})] \quad (1)$$

$$\psi_{\sigma_u 1s}(\mathbf{r}) = \mathcal{N}_{\sigma_u 1s}[\phi_{100}(\mathbf{r} - \frac{\mathbf{R}}{2}) - \phi_{100}(\mathbf{r} + \frac{\mathbf{R}}{2})]. \quad (2)$$

These new basis functions belong to definite symmetry classifications in $D_{\infty h}$, namely Σ_g^+ ('+' combination) and Σ_u^+ ('-' combination). The $2s$, $3s$, etc. orbitals are symmetrized similarly. If we extend the basis to contain atomic $2p$ orbitals, the properly symmetrized basis functions in the Σ blocks are

$$\psi_{\sigma_g 2p}(\mathbf{r}) = \mathcal{N}_{\sigma_g 2p}[\phi_{210}(\mathbf{r} - \frac{\mathbf{R}}{2}) - \phi_{210}(\mathbf{r} + \frac{\mathbf{R}}{2})] \quad (3)$$

$$\psi_{\sigma_u 2p}(\mathbf{r}) = \mathcal{N}_{\sigma_u 2p}[\phi_{210}(\mathbf{r} - \frac{\mathbf{R}}{2}) + \phi_{210}(\mathbf{r} + \frac{\mathbf{R}}{2})], \quad (4)$$

for the $2p_0$ ($2p_z$) orbitals, while the '+' and '-' combination of the $2p_{\pm 1}$ are of π_u and π_g symmetry, respectively, and hence not of interest in our present discussion focusing on the $3\sigma_g$ HOMO of N_2 .

It is important to note that for a quantitative HF calculation, the basis is further extended until convergence is obtained. We may easily check the correct parity (g or u) of the basis function by letting $\mathbf{r} \rightarrow -\mathbf{r}$ and using the fact




FIG. 1: (Color online). Planar cuts through the $3\sigma_g$ HOMO of N_2 obtained in a HF calculation with a $(12s7p)/[6s4p]$ basis (see text). Left (right) column: coordinate (momentum) space wave function. Upper row: contribution from s type basis functions, middle row: contribution from p type basis functions, lower row: total HF result in the $(12s7p)/[6s4p]$ basis.

that the parity of ϕ_{nlm} is $(-1)^l$. We emphasize this point since in Ref. [5], the ‘+’ combination of the $2p_z$ orbitals is erroneously referred to as the gerade state. Although not of relevance for our fully numerical HF calculations, we note that in the hydrogenic basis used in Ref. [5] the Fourier transform of the antisymmetric combination of atomic $2p_z$ orbitals is

$$\tilde{\psi}_{\sigma_g 2p}(\mathbf{p}) = \mathcal{N}_{\sigma_g 2p} \sin(\mathbf{p} \cdot \mathbf{R}/2) \frac{p}{(1 + 4p^2)^3} Y_{10}(\hat{\mathbf{p}}), \quad (5)$$

with $\mathcal{N}_{\sigma_g 2p}$ the normalization constant. We may verify the gerade parity symmetry by letting $\mathbf{p} \rightarrow -\mathbf{p}$ and using $Y_{lm}(-\hat{\mathbf{p}}) = (-1)^l Y_{lm}(\hat{\mathbf{p}})$. Equation (5) is the correctly symmetrized version of Eq. (20) of Ref. [5]. The difference lies in the sine-factor of Eq. (5) compared with a cosine-factor in Ref. [5]. A calculation with the p -type basis functions alone would, because of this sine-factor, predict suppressed ionization of N_2 compared with the companion Ar atom – in contrast to experimental and theoretical findings [1, 7, 8]. Thus having pointed out the importance of working with properly symmetrized basis function, we now confront the *ad hoc* single LCAO approach [5] with our HF calculations. Clearly, but contrary to the approach in Ref. [5], one cannot expect the

basis functions themselves to be eigenfunctions, instead the eigenfunctions are linear combinations of basis functions.

III. HARTREE-FOCK CALCULATIONS

In Ref. [2] we used the GAMESS [9] program for our quantitative HF calculations of N_2 with a $(11s6p)/[5s3p]$ Cartesian Gaussian basis [10]. The way we specify our basis is standard in the quantum chemistry literature [11], but is recalled here for completeness. The wave function is expanded in 11 s and 6 p orbitals centered on each atom. These basis orbitals all have different exponentials in the radial parts. The orbitals of our HF calculation are obtained by a variation of the expansion coefficients until the energy is minimized. This procedure is equivalent to an iterative diagonalization of matrix representations of the Fock operator. With little loss of accuracy but significant gain in computational speed, the optimization procedure can be simplified with a contracted basis set. This means that all the expansion coefficients are not varied independently, contrary a fixed relationship is kept between some of the coefficients. Such a simplification is used in most quantum chemistry software. Five coefficients corresponding to s orbitals and three coefficients corresponding to the p orbitals are optimized independently. In order to describe the asymptotic behavior we augment the basis by adding an extra diffuse s and p orbital. In total, the wave function is effectively expanded in a basis of six s and four p orbitals on each atom denoted by $(12s7p)/[6s4p]$ basis.

In Fig. 1 we show results of our calculations for the $3\sigma_g$ HOMO of N_2 [Fig. 1 (c)] and in its decomposition of s [Fig. 1 (a)] and p [Fig. 1 (b)] type basis functions. We see that the orbital is composed of both types of basis functions. At first sight the p contribution alone seems to resemble the orbital quite well. However, in the proximity of the nuclei, it is the s contribution that dominates, since the p functions have nodes on the nuclei. It is the s type contribution that dominates the momentum space wave function at low momenta. This fact becomes clear in Figs. 1 (d)-(f), where we show the Fourier transforms of the functions in Figs. 1 (a)-(c). It is clear that both the s and p type of basis functions contribute significantly to the momentum wave function. Therefore, as discussed in Sec. II, it is insufficient to consider each type exclusively: The s and p_z basis orbitals are not separately eigenfunctions of the Fock operator, contrary, the mixture turns out to be essential. At this point the authors of Ref. [5] introduce an oversimplification by considering the HOMO to be constructed only from a single linear combination of two p -orbitals.

IV. IMPLICATIONS TO THE MOLECULAR STRONG-FIELD APPROXIMATION

In the MO-SFA one calculates the transition rate from the HOMO to a continuum Volkov state through absorption of n photons. For a linearly polarized, monochromatic field of frequency ω and field strength F_0 , the velocity gauge result for the angular differential ionization rate in atomic units is [1]

$$\frac{dW}{d\mathbf{p}} = 2\pi N_e p_n \left| \tilde{\Psi}(\mathbf{p}_n) \right|^2 (U_p - n\omega)^2 \times J_n^2 \left(\frac{|\mathbf{F}_0 \cdot \mathbf{p}_n|}{\omega^2}, -\frac{U_p}{2\omega} \right), \quad (6)$$

with N_e the number of electrons occupying the HOMO, $p_n = \sqrt{2(n\omega - U_p - I_p)}$ the momentum, $U_p = F_0^2/(4\omega^2)$ the ponderomotive energy, I_p the ionization potential of the HOMO (0.5725 in the case of N_2 [12]), $\tilde{\Psi}(\mathbf{p})$ the momentum space wave function and $J_n(u, v)$ a generalized Bessel function [13].

According to the discussion in Sec. III, it is important to apply the correct momentum space wave function in the evaluation of Eq. (6). To illustrate this fact, we consider the total ionization rate integrated over all angles of the outgoing electron and summed over all numbers of accessible photon absorptions. We present this total ionization rate for different angles θ between the internuclear axis and the polarization axis for the HOMO $3\sigma_g$ orbital of N_2 . We consider three descriptions of the molecular orbital: (i) using s -type basis functions only, (ii) using p -type basis functions only and (iii) using the self-consistent HF solution. Clearly the approaches (i) and (ii) only give a very poor description of the HOMO and therefore these results deviate from the result obtained by using the true HOMO. In Fig. 2 we present the results for the three cases at (a) 800 nm [2] and (b) 795 nm [6]. Additionally, we show in Fig. 2 the results obtained with an initial HOMO derived from a grid-based HF calculation [14]. The excellent agreement with the $(12s7p)/[6s4p]$ basis-state calculation proves the convergence of the latter approach. We note that panels (a) and (b) are very similar and hence the differences between the results in [2] and [5] are not due to the slight difference in wavelength.

From Fig. 1 (d) we see that the momentum wave function in case (i) is nearly spherically symmetric, and hence, in Fig. 2 the ionization rate in this case is nearly independent of the molecular orientation. In case (ii) we use the momentum function from Fig. 1 (e). Although exceptions occur, as evident from Fig. 3(a) below, the generalized Bessel functions are typically maximized when \mathbf{p} is nearly parallel to the polarization direction [Fig. 3(b)]. In a qualitative analysis, one may therefore expect that the maximum rate is obtained, when the polarization axis coincides with a direction where $|\tilde{\Psi}(\mathbf{p})|^2$ is large. Correspondingly, in case (ii) we see from Fig. 2 that the rate is maximized when $\theta = 0$, i.e. when the polarization is




FIG. 2: Ionization rate at $2 \times 10^{14} \text{ W/cm}^2$ of the HOMO of N_2 ($3\sigma_g$) as a function of the angle θ between the internuclear axis and the polarization axis at (a) 800 nm [2] and (b) 795 nm [6]. The short-dashed and solid curve are the calculations with the HF wave function derived from a grid calculation and with the $(12s7p)/[6s4p]$ basis set in Fig. 1 (f), respectively. The long-dashed curve with a maximum and the chained curve are obtained with the s [Fig. 1(d)] and p [Fig. 1(e)] contributions only, respectively. All rates are normalized to parallel ($\theta = 0$) geometry.

horizontal in Fig. 1 (e) and overlaps with a large value of the momentum wave function. Finally, we turn to the full wave function in case (iii) which is the coherent sum of s and p contributions, Fig. 1 (f). It is the low momenta that contribute mostly to the total rate. At these small momenta, we see from Fig. 1 (f) that the absolute square of the momentum wave function is maximized along the direction perpendicular to the molecular axis and therefore the rate is maximized with the molecule aligned perpendicularly to the laser polarization, $\theta = \pi/2$. This result is also shown in Fig. 2 verifying the result of Ref. [2]. Note that the rate using the total wave function is not the sum of rates of the s and p contributions since their contributions to the wave function should be added coherently and with the correct self-consistent amplitudes before entering $\tilde{\Psi}(\mathbf{p})$ in Eq. (6). We note that the admixture of atomic s and p orbitals for the description of the HOMO of N_2 was also recently pointed out in related work on high-harmonic generation [15].

In Refs. [2, 5], the values of the generalized Bessel function in Eq. (6) do also not agree. Figure 3(a) shows our previous result for the square of the generalized Bessel function corresponding to 18 photon absorption at 800 nm and at an intensity of $2 \times 10^{14} \text{ W/cm}^2$. Panel (b) shows the corresponding result at 795 nm [6]. Hence, Panel (b) corresponds to Fig. 2(a) of Ref. [5] (since




FIG. 3: Polar plot of the generalized Bessel function of order $n = 18$ squared for 2×10^{14} W/cm², and (a) 800 nm [2], (b) 795 nm [6]. The polar angle θ_p is measured from the horizontal line.

$J_{-n}(u, v) = (-1)^n J_n(u, -v)$ [13]) and by comparison we see that the results agree. We may therefore attribute the difference between the generalized Bessel functions in [2] and [5] to stem from the difference in the wavelengths used. It is clear from Fig. 3 that the values of the generalized Bessel function are very sensitive to the exact value of the arguments. We reproduce the values for the generalized Bessel function given in Ref. [16] with the input parameters used therein, and we therefore believe that our previous and present calculations of the

generalized Bessel function are correct. Due to the excellent agreement at 795 nm with the Bessel functions of Ref. [5] we likewise believe that the latter are evaluated correctly.

V. CONCLUSION

In conclusion, the velocity gauge MO-SFA when based on a transition to the continuum from an initial HOMO, determined by a HF calculation using standard quantum chemistry software, gives the wrong prediction for the orientation-dependent rate. In this approximation only the length gauge MO-SFA (and the molecular tunneling theory) [2] give the observed minimum [3] in the rate for the perpendicular geometry. This conclusion contrasts the findings in a recent work [5]. In that work, however, only a single LCAO of two p -orbitals (of wrong symmetry) was considered for the description of the HOMO and no self-consistent wave function was applied. From Fig. 2 we clearly see, that an application of a too simple wave function, e.g., one with only p basis functions, leads to a prediction accidentally in qualitative agreement with experiment [3]. If one nevertheless would be encouraged by this agreement to believe that the single LCAO of p orbitals describes the physics well, one should note that such a wave function due to the presence of the sine-factor in (5) would predict suppressed ionization in N₂ compared with Ar which is in contrast with experiments and theory [1, 7, 8]. In closing we note that also in the atomic case, the length gauge version of the SFA was recently shown to be superior to the velocity gauge version when compared with results obtained by integrating the time-dependent Schrödinger equation [17].

Acknowledgments

LBM is supported by the Danish Natural Science Research Council (Grant No. 21-03-0163).

-
- [1] J. Muth-Böhm, A. Becker, and F. H. M. Faisal, *Phys. Rev. Lett.* **85**, 2280 (2000).
 [2] T. K. Kjeldsen and L. B. Madsen, *J. Phys. B: At. Mol. Opt. Phys.* **37**, 2033 (2004).
 [3] I. V. Litvinyuk, K. F. Lee, P. W. Dooley, D. M. Rayner, D. M. Villeneuve, and P. B. Corkum, *Phys. Rev. Lett.* **90**, 233003 (2003).
 [4] Z. X. Zhao, X. M. Tong, and C. D. Lin, *Phys. Rev. A* **67**, 043404 (2003).
 [5] V. I. Usachenko and S.-I. Chu, *Phys. Rev. A* **71**, 063410 (2005).
 [6] The first author of Ref. [5] has informed us that he used 795 nm light.
 [7] A. Talebpour, C. Y. Chien, and S. L. Chin, *J. Phys. B: At. Mol. Opt. Phys.* **29**, L677 (1996).
 [8] C. Guo, M. Li, J. P. Nibarger, and G. N. Gibson, *Phys. Rev. A* **58**, R4271 (1998).
 [9] M. W. Schmidt, K. K. Baldridge, J. A. Boatz, S. T. Elbert, M. S. Gordon, J. H. Jensen, S. Koseki, N. Matsunaga, K. A. Nguyen, S. J. Su, et al., *J. Comput. Chem.* **14**, 1347 (1993).
 [10] T. H. Dunning, *J. Chem. Phys.* **55**, 716 (1971).
 [11] P. W. Atkins and R. S. Friedman, *Molecular Quantum Mechanics* (Oxford University Press, Oxford, 1997), 3rd ed.
 [12] K. P. Huber and G. H. Herzberg, *Molecular Spectra and Molecular Structure, IV Constants of Diatomic Molecules* (Van Nostrand Reinhold Company, New York, 1979).

- [13] H. R. Reiss, Phys. Rev. A **22**, 1786 (1980).
- [14] J. Kobus, L. Laaksonen, and D. Sundholm, Comput. Phys. Commun. **98**, 346 (1996).
- [15] B. Zimmermann, M. Lein, and J. M. Rost, Phys. Rev. A **71**, 033401 (2005).
- [16] H. R. Reiss and V. P. Krainov, J. Phys. A: Math. Gen. **36**, 5575 (2003).
- [17] D. Bauer, D. B. Milosević, and W. Becker, Phys. Rev. A **72**, 023415 (2005).