

Corrections to the energy levels of a spin-zero particle bound in a strong field

R.N. Lee ^{(a)*}, A.I. Milstein^{(a)†}, S.G. Karshenboim^{(b,c)‡}

^(a)*Budker Institute of Nuclear Physics,*

630090 Novosibirsk, Russia

^(b)*D. I. Mendeleev Institute for Metrology (VNIIM),*

St. Petersburg 198005, Russia

^(c)*Max-Planck-Institut für Quantenoptik,*

85748 Garching, Germany

(Dated: October 11, 2018)

Abstract

Formulas for the corrections to the energy levels and wave functions of a spin-zero particle bound in a strong field are derived. General case of the sum of a Lorentz-scalar potential and zero component of a Lorentz-vector potential is considered. The forms of the corrections differ essentially from those for spin-1/2 particles. As an example of application of our results, we evaluated the electric polarizability of a ground state of a spin-zero particle bound in a strong Coulomb field.

PACS numbers: 31.30.Jv

* Electronic address: R.N.Lee@inp.nsk.su

† Electronic address: A.I.Milstein@inp.nsk.su

‡ Electronic address: sek@mpq.mpg.de

I. INTRODUCTION

As known, in many cases the perturbation theory (PT) is a very fruitful method to obtain analytic results for various corrections. The formulas of PT for the Dirac equation are similar to that for the Schrödinger equation and have simple forms (see, e.g. [1]). However, it is essentially more complicated to derive the formulas of PT for the Klein-Gordon-Fock equation, because this equation contains second derivative over time. In the present paper, we derive the first and the second order corrections to the energy levels, and the first order correction to the wave function of a spin-zero particle bound in a strong field. We consider a general case of a sum of the Lorentz-scalar and the Lorentz-vector potentials. This may be useful at the consideration of the effects of the strong interaction in pionic atoms. Using our formula for the correction to the energy level, we evaluate the electric polarizability of a ground state of a spin-zero particle bound in a strong Coulomb field.

II. PERTURBATION THEORY

The relativistic equation for the wave function of a spin-zero particle (Klein-Gordon-Fock equation) bound in the external time-independent field has the form

$$\left[\left(i \frac{\partial}{\partial x^\mu} - e A_\mu(\mathbf{r}) \right)^2 - m^2 - 2mV(\mathbf{r}) \right] \Psi(\mathbf{r}, \mathbf{t}) = 0, \quad (1)$$

where $V(\mathbf{r})$ is the Lorentz-scalar potential, $A_\mu(\mathbf{r})$ is the Lorentz-vector potential, e and m are the charge and the mass of the particle, respectively; we set $\hbar = c = 1$. The corresponding current, which obeys the continuity equation, reads

$$J_\mu = \Psi^*(\mathbf{r}, \mathbf{t}) i \overset{\leftrightarrow}{\partial} x^\mu \Psi(\mathbf{r}, \mathbf{t}) - 2e A_\mu(\mathbf{r}) |\Psi(\mathbf{r}, \mathbf{t})|^2. \quad (2)$$

For $\mathbf{A} = 0$, the solution with the fixed energy E_n , $\Psi(\mathbf{r}, \mathbf{t}) = \exp(-iE_n t) \Phi_n(\mathbf{r})$, obeys the equation

$$[(E_n - U(\mathbf{r}))^2 - \mathbf{p}^2 - m^2 - 2mV(\mathbf{r})] \Phi_n(\mathbf{r}) = 0, \quad (3)$$

where $U(\mathbf{r}) = eA_0(\mathbf{r})$. The non-relativistic approximation of this equation is the Schrödinger equation with the potential $U_{nr}(\mathbf{r}) = U(\mathbf{r}) + V(\mathbf{r})$. From Eq.(2) we find the normalization of the wave function

$$2 \int d\mathbf{r} [E_n - U(\mathbf{r})] |\Phi_n(\mathbf{r})|^2 = 1. \quad (4)$$

Multiplying both sides of Eq.(3) by $\Phi_k(\mathbf{r})$ with $E_k \neq E_n$ and taking the integral over \mathbf{r} , we obtain

$$\int d\mathbf{r} [E_k + E_n - 2U(\mathbf{r})] \Phi_k^*(\mathbf{r}) \Phi_n(\mathbf{r}) = 0. \quad (5)$$

Then we represent the potentials in the form $V(\mathbf{r}) = V_0(\mathbf{r}) + \delta V(\mathbf{r})$, and $U(\mathbf{r}) = U_0(\mathbf{r}) + \delta U(\mathbf{r})$. Let $\phi_n(\mathbf{r})$ be the solution of the Klein-Gordon-Fock equation in the potentials $V_0(\mathbf{r})$, $U_0(\mathbf{r})$ with the corresponding energy ε_n . Let us write

$$\Phi_n(\mathbf{r}) = \phi_n(\mathbf{r}) + \delta\phi_n(\mathbf{r}) \quad , \quad E_n = \varepsilon_n + \delta\varepsilon_n.$$

The first order correction to the wave function, $\delta\phi_n^{(1)}(\mathbf{r})$, with respect to perturbations $\delta V(\mathbf{r})$ and $\delta U(\mathbf{r})$ obeys the equation

$$\begin{aligned} & [(\varepsilon_n - U_0(\mathbf{r}))^2 - \mathbf{p}^2 - m^2 - 2mV_0(\mathbf{r})] \delta\phi_n^{(1)}(\mathbf{r}) \\ & = \{2m\delta V(\mathbf{r}) - 2[\varepsilon_n - U_0(\mathbf{r})][\delta\varepsilon_n^{(1)} - \delta U(\mathbf{r})]\} \phi_n(\mathbf{r}). \end{aligned} \quad (6)$$

Multiplying both sides of this equation by $\phi_n^*(\mathbf{r})$, taking the integral over \mathbf{r} , and using normalization condition (5), we obtain

$$\delta\varepsilon_n^{(1)} = \int d\mathbf{r} \{2m\delta V(\mathbf{r}) + 2\delta U(\mathbf{r})[\varepsilon_n - U_0(\mathbf{r})]\} |\phi_n(\mathbf{r})|^2. \quad (7)$$

Then we multiply Eq.(3) by $\phi_n^*(\mathbf{r})$, take the integral over \mathbf{r} , and collect the terms of the second order in $\delta V(\mathbf{r})$ and $\delta U(\mathbf{r})$. We have

$$\begin{aligned} \delta\varepsilon_n^{(2)} & = \int d\mathbf{r} \{2m\delta V(\mathbf{r}) - 2[\delta\varepsilon_n^{(1)} - \delta U(\mathbf{r})][\varepsilon_n - U_0(\mathbf{r})]\} \phi_n^*(\mathbf{r}) \delta\phi_n^{(1)}(\mathbf{r}) \\ & \quad - \int d\mathbf{r} [\delta\varepsilon_n^{(1)} - \delta U(\mathbf{r})]^2 |\phi_n(\mathbf{r})|^2. \end{aligned} \quad (8)$$

Let us introduce the Green's function $\mathcal{D}_n(\mathbf{r}, \mathbf{r}')$, which obeys the equation

$$\begin{aligned} & [(\varepsilon_n - U_0(\mathbf{r}))^2 - \mathbf{p}^2 - m^2 - 2mV_0(\mathbf{r})] \mathcal{D}_n(\mathbf{r}, \mathbf{r}') \\ & = \delta(\mathbf{r} - \mathbf{r}') - \frac{1}{N} \phi_n(\mathbf{r}) \phi_n^*(\mathbf{r}'), \\ & N_n = \int d\mathbf{r} |\phi_n(\mathbf{r})|^2, \end{aligned} \quad (9)$$

and the conditions

$$\int d\mathbf{r}' \mathcal{D}_n(\mathbf{r}, \mathbf{r}') \phi_n(\mathbf{r}') = 0, \quad \int d\mathbf{r} \phi_n^*(\mathbf{r}) \mathcal{D}_n(\mathbf{r}, \mathbf{r}') = 0. \quad (10)$$

It is obvious that $\mathcal{D}_n(\mathbf{r}, \mathbf{r}')$ is nothing but the reduced Green function of the Schrödinger equation, divided by $2m$, with the effective energy $\tilde{E}_n = (\varepsilon_n^2 - m^2)/2m$, and the effective potential $\tilde{V}(\mathbf{r}) = V_0(\mathbf{r}) + (\varepsilon_n/m)U_0(\mathbf{r}) - U_0^2(\mathbf{r})/2m$. We assume that Eq.(9) can be solved. For instance, the solution is well known for the pure Coulomb field, $U_0(\mathbf{r}) = -Ze^2/r$, $V_0(\mathbf{r}) = 0$, where Z is an atomic charge number and e is the electron charge.

It is easy to check that the solution of Eq.(6) has the form

$$\delta\phi_n^{(1)}(\mathbf{r}) = \int d\mathbf{r}' \mathcal{D}_n(\mathbf{r}, \mathbf{r}') \{2m\delta V(\mathbf{r}') - 2[\delta\varepsilon_n^{(1)} - \delta U(\mathbf{r}')] [\varepsilon_n - U_0(\mathbf{r}')]\} \phi_n(\mathbf{r}') + \beta\phi_n(\mathbf{r}), \quad (11)$$

where the constant β is fixed by the normalization condition (4). We have

$$\begin{aligned} \beta &= -2 \iint d\mathbf{r}d\mathbf{r}' \phi_n^*(\mathbf{r}) [\varepsilon_n - U_0(\mathbf{r})] \mathcal{D}_n(\mathbf{r}, \mathbf{r}') \\ &\quad \times \{2m\delta V(\mathbf{r}') - 2[\delta\varepsilon_n^{(1)} - \delta U(\mathbf{r}')] [\varepsilon_n - U_0(\mathbf{r}')]\} \phi_n(\mathbf{r}') \\ &\quad - \int d\mathbf{r} [\delta\varepsilon_n^{(1)} - \delta U(\mathbf{r})] |\phi_n(\mathbf{r})|^2. \end{aligned} \quad (12)$$

Substituting Eq.(11) in Eq.(8), we finally obtain

$$\begin{aligned} \delta\varepsilon_n^{(2)} &= \iint d\mathbf{r}d\mathbf{r}' \phi_n^*(\mathbf{r}) \{2m\delta V(\mathbf{r}) - 2[\delta\varepsilon_n^{(1)} - \delta U(\mathbf{r})] [\varepsilon_n - U_0(\mathbf{r})]\} \mathcal{D}_n(\mathbf{r}, \mathbf{r}') \\ &\quad \times \{2m\delta V(\mathbf{r}') - 2[\delta\varepsilon_n^{(1)} - \delta U(\mathbf{r}')] [\varepsilon_n - U_0(\mathbf{r}')]\} \phi_n(\mathbf{r}') \\ &\quad - \int d\mathbf{r} [\delta\varepsilon_n^{(1)} - \delta U(\mathbf{r})]^2 |\phi_n(\mathbf{r})|^2. \end{aligned} \quad (13)$$

Note that the term $\beta\phi_n(\mathbf{r})$ in Eq. (11) does not contribute to Eq.(13) due to Eq.(7).

III. ELECTRIC POLARIZABILITY

As an application of our results, let us consider the electric polarizability α_{1s} of a ground state of a spin-zero particle bound in a strong Coulomb field. For spin-1/2 particle the polarizability was obtained in Ref.[3, 4].

For the perturbation $\delta U(\mathbf{r}) = -e\mathcal{E} \cdot \mathbf{r}$, we have $\delta\varepsilon^{(1)} = 0$, and $\delta\varepsilon^{(2)}$ can be written in the form

$$\delta\varepsilon^{(2)} = -\frac{1}{2}\alpha_{1s}\mathcal{E}^2. \quad (14)$$

From Eq.(13), we find

$$\begin{aligned} \alpha_{1s} &= \frac{2}{3}\alpha \left\{ \int d\mathbf{r} r^2 \phi_{1s}^2(r) - 4 \iint d\mathbf{r}d\mathbf{r}' \phi_{1s}(r) \left(\varepsilon_{1s} + \frac{Ze^2}{r} \right) \mathcal{D}_{1s}(\mathbf{r}, \mathbf{r}') \right. \\ &\quad \left. \times \left(\varepsilon_{1s} + \frac{Ze^2}{r'} \right) \phi_{1s}(r') (\mathbf{r} \cdot \mathbf{r}') \right\}, \end{aligned} \quad (15)$$

where $\alpha = e^2 = 1/137$ is the fine structure constant. The wave function and the energy of the ground state has the form (see, e.g.,[5])

$$\begin{aligned}\phi_{1s}(r) &= \sqrt{A} (2\kappa r)^{\gamma-1/2} \exp(-\kappa r), \quad \varepsilon_{1s} = m\sqrt{1/2 + \gamma}, \\ \gamma &= \sqrt{1/4 - (Z\alpha)^2}, \quad \kappa = m\sqrt{1/2 - \gamma}, \quad A = \frac{Z\alpha\kappa^2}{\pi\Gamma(2\gamma + 2)},\end{aligned}\quad (16)$$

Let us introduce the function $\mathbf{F}(\mathbf{r})$

$$\mathbf{F}(\mathbf{r}) = \int d\mathbf{r}' \mathcal{D}_{1s}(\mathbf{r}, \mathbf{r}') \left(\varepsilon_{1s} + \frac{Ze^2}{r'} \right) \mathbf{r}' \phi_{1s}(r'). \quad (17)$$

It is convenient to represent it in the form

$$\mathbf{F}(\mathbf{r}) = \frac{2\sqrt{A}\varepsilon_{1s}}{\kappa^2} (2\kappa r)^{\gamma-3/2} \exp(-\kappa r) g(2\kappa r) \mathbf{r}, \quad (18)$$

where the function $g(x)$ satisfies the equation, following from Eq.(9):

$$g''(x) + \left(\frac{2\gamma + 1}{x} - 1 \right) g'(x) - \frac{2}{x^2} g(x) = \frac{x + 1 - 2\gamma}{8}. \quad (19)$$

Substituting Eqs.(17) and (18) in Eq.(15), we obtain

$$\begin{aligned}\alpha_{1s} &= \frac{\alpha\varepsilon_{1s}}{3\kappa^4} \left\{ (1/2 - \gamma)(\gamma + 3/2)(\gamma + 1) \right. \\ &\quad \left. - \frac{1}{\Gamma(2\gamma + 1)} \int_0^\infty dx (x + 1 - 2\gamma) x^{2\gamma+1} \exp(-x) g(x) \right\}.\end{aligned}\quad (20)$$

The general solution of Eq.(19) has the form

$$\begin{aligned}g(x) &= -\frac{x}{4} - \frac{x^2}{16} + \frac{x}{4} {}_2F_2 \left[\begin{matrix} 1, 1 \\ \gamma + 2 + \nu, \gamma + 2 - \nu \end{matrix} \middle| x \right] \\ &\quad + a x^{-\gamma+\nu} {}_1F_1 \left[\begin{matrix} -\gamma + \nu \\ 1 + 2\nu \end{matrix} \middle| x \right] + b x^{-\gamma-\nu} {}_1F_1 \left[\begin{matrix} -\gamma - \nu \\ 1 - 2\nu \end{matrix} \middle| x \right],\end{aligned}\quad (21)$$

where ${}_pF_q$ is the hypergeometric function, $\nu = \sqrt{\gamma^2 + 2}$, and a and b are some constants to be determined from the boundary conditions at $x = 0$ and $x = \infty$. The condition at $x = 0$ gives $b = 0$. The constant a should be chosen to provide the cancellation of the exponentially large terms of $g(x)$ at $x \rightarrow \infty$. The large- x asymptotics of the hypergeometric functions in Eq. (21) can be calculated from their integral representations, see [6]. The asymptotics have the form

$$\begin{aligned}{}_2F_2 \left[\begin{matrix} 1, 1 \\ \gamma + 2 + \nu, \gamma + 2 - \nu \end{matrix} \middle| x \right] &\sim \Gamma[2 + \gamma + \nu, 2 + \gamma - \nu] \frac{\exp(x)}{x^{2\gamma+2}} \sum_n \frac{(1 + \gamma + \nu)_n (1 + \gamma - \nu)_n}{x^n n!} \\ x^{-\gamma\pm\nu} {}_1F_1 \left[\begin{matrix} -\gamma \pm \nu \\ 1 \pm 2\nu \end{matrix} \middle| x \right] &\sim \frac{\Gamma[1 \pm 2\nu]}{\Gamma[\pm\nu - \gamma]} \frac{\exp(x)}{x^{2\gamma+1}} \sum_n \frac{(1 + \gamma + \nu)_n (1 + \gamma - \nu)_n}{x^n n!}.\end{aligned}\quad (22)$$




FIG. 1: Polarizability in units $(m\alpha)^{-3} Z^{-4}$ as a function of $Z\alpha$.

Thus, we obtain

$$g(x) = -\frac{x}{4} - \frac{x^2}{16} + \frac{x}{4} {}_2F_2 \left[\begin{matrix} 1, 1 \\ \gamma + 2 + \nu, \gamma + 2 - \nu \end{matrix} \middle| x \right] - \frac{\Gamma[2 + \gamma + \nu, 2 + \gamma - \nu, \nu - \gamma]}{4\Gamma[1 + 2\nu]} x^{-\gamma+\nu} {}_1F_1 \left[\begin{matrix} -\gamma + \nu \\ 1 + 2\nu \end{matrix} \middle| x \right], \quad (23)$$

Using the integral representation of ${}_pF_q$, see [6], we obtain the following identity

$$\begin{aligned} & {}_2F_2 \left[\begin{matrix} 1, 1 \\ \gamma + 2 + \nu, \gamma + 2 - \nu \end{matrix} \middle| x \right] - \frac{\Gamma[2 + \gamma + \nu, 2 + \gamma - \nu, \nu - \gamma]}{\Gamma[1 + 2\nu]x^{1+\gamma-\nu}} {}_1F_1 \left[\begin{matrix} -\gamma + \nu \\ 1 + 2\nu \end{matrix} \middle| x \right] \\ &= \int_0^1 dz (1-z)^{\gamma+\nu} \int_0^1 dt t^{\nu-\gamma-2} \exp[zx(1-1/t)] \end{aligned} \quad (24)$$

Substituting Eqs. (24) and (23) in Eq.(20) and integrating over x , z , and t , we come to the final result for the polarizability α_{1s}

$$\begin{aligned} \alpha_{1s} = & \frac{\alpha(\gamma+1)\sqrt{\gamma+1/2}}{3m^3(1-2\gamma)} \left\{ \frac{12\gamma^2 + 64\gamma + 37}{2(1-2\gamma)} \right. \\ & - \frac{2(2\gamma+1)}{\gamma+\nu+1} \left(\frac{1-2\gamma}{\gamma+\nu+2} {}_3F_2 \left[\begin{matrix} 1, 2\gamma+3, \gamma+\nu+1 \\ \gamma+\nu+2, \gamma+\nu+3 \end{matrix} \middle| 1 \right] \right. \\ & \left. \left. + \frac{(2\gamma+3)}{\gamma+\nu+3} {}_3F_2 \left[\begin{matrix} 1, 2\gamma+4, \gamma+\nu+1 \\ \gamma+\nu+2, \gamma+\nu+4 \end{matrix} \middle| 1 \right] \right) \right\}. \end{aligned} \quad (25)$$

The polarizability for arbitrary state can be obtained similarly. Figure 1 shows the polarizability α_{1s} in units $(m\alpha)^{-3} Z^{-4}$ as a function of $Z\alpha$.

At $Z\alpha \ll 1$, we have

$$\alpha_{1s} = \frac{9}{2(m\alpha)^3 Z^4} \left[1 - \frac{121}{27}(Z\alpha)^2 + \left(\frac{229}{72} + \frac{4\pi^2}{81} \right) (Z\alpha)^4 + \dots \right]. \quad (26)$$

At $Z\alpha \rightarrow 1/2$ the polarizability has finite value $\sim 3.84\alpha/m^3$. Our formula for polarizability can not be applied for $Z\alpha > 1/2$ because we did not take into account the effect of finite

nuclear size. Besides, in a pionic atom it is necessary to account for the effects of the strong interaction that become dominant for the ground state at large Z . It is interesting to compare the polarizabilities of a ground state, calculated for spin-0 particle and spin-1/2 particle. It turns out that the functions $f(Z) = Z^4\alpha_{1s}$ obey the relation $f(Z)_{S=0} \approx f(2Z)_{S=1/2}$ with accuracy of a few percent.

In summary, we have derived the formulas for the corrections to the energy levels and wave functions of a spin-zero particle bound in a strong field. These formulas may be useful at the consideration of various effects in pionic atoms. As an example, we evaluated the electric polarizability of a ground state of a spin-zero particle bound in a strong Coulomb field.

ACKNOWLEDGMENTS

A.I.M. gratefully acknowledges the School of Physics at the University of New South Wales, and the Max-Planck-Institute for Quantum Optics, Garching, for warm hospitality and financial support during the visit. The work was supported in parts by RFBR Grants 03-02-16510, 03-02-04029, by DFG Grant GZ 436 Rus 113/769/0-1 and by Russian Science Support Foundation.

-
- [1] V.B. Berestetski, E.M. Lifshits, and L.P. Pitaevsky, *Quantum Electrodynamics*, Pergamon, Oxford (1982).
 - [2] L.D. Landau, E.M. Lifshits, *Quantum Mechanics, Nonrelativistic Theory*, Pergamon, Oxford (1965).
 - [3] B.A. Zon, N.L. Manakov, and L.P. Rapoport, *Sov. J. Nucl. Phys.* **15**, 282 (1972).
 - [4] V. Yakhontov, *Phys. Rev. Lett.*, **91**, 093001 (2003).
 - [5] A. S. Davidov. *Quantum Mechanics*, Pergamon Press, Oxford. 1976. See Sect. 58.
 - [6] L.J. Slater, *Generalized hypergeometric functions*, Cambridge University Press, 1966.