

Harmonium in intense laser fields: excitation, absorption, and transparency

O. Kidun and D. Bauer

Max-Planck-Institut für Kernphysik, Postfach 10 39 80, 69029 Heidelberg, Germany

Abstract. It is known that the dynamics of two (Coulomb-interacting) nonrelativistic electrons confined by a parabolic potential and driven by a classical, intense laser field (in dipole approximation) is exactly soluble. We calculate the time-dependent population of the harmonic oscillator states and the energy absorbed from the laser. It turns out that the key entity on which all observables sensitively depends is the modulus square of the Fourier-transformed vector potential of the laser field, evaluated at the harmonic oscillator frequency. The system is transparent to laser field configurations for which this entity vanishes. We discuss the Poisson statistics behavior of the transition probabilities and analyze the conditions for the complete survival and full depletion of the initial state.

1. Introduction

The parabolic well is a model of great practical importance, as it approximates any arbitrary potential close to equilibrium. In nanotechnology, potentials of simple shape such as quantum dots are often well approximated by parabolic potentials. Superpositions of harmonic oscillators are also used to describe continua, environments, and field modes.

Correlations, for instance, those introduced by the Coulomb interaction between electrons, attract growing attention. Correlated electron dynamics can nowadays be analyzed experimentally with great precision using “reaction microscopes” [1]. Two electrons confined in a parabolic potential, sometimes called “harmonium” or the “Hooke’s atom” version of helium [2], is of great interest since the problem represents one of the few exactly soluble problems involving correlation [3, 4]. As such harmonium serves as a testing ground for, e.g. new exchange-correlation potentials in density functional theory.

In the field of strong laser atom interaction, exactly soluble models are rare as well, in particular as it comes to electron correlation. Fortunately, the two-electron wavefunction of the laser-driven harmonium can still be expressed in analytical form as long as the laser field is treated classically and in dipole approximation [5].

The outline of the paper is as follows. In Sec. 2 we derive the exact solution of the time-dependent Schrödinger equation of two interacting electrons confined in a parabolic potential and driven by a classical (laser) field (in dipole approximation). In Sec. 3 the

probabilities for two-electron excitations from an arbitrary state m to a state n are calculated. The latter are the basis for the discussion of optimal energy absorption and transparency in Sec. 4. Finally, we conclude in Sec. 5.

2. Exact states of two-electrons in a parabolic well driven by a laser field

In the absence of the laser field the Hamiltonian of harmonium reads[‡]

$$H(\mathbf{r}_1, \mathbf{r}_2) = -\frac{\hbar^2}{2m}\nabla_{\mathbf{r}_1}^2 - \frac{\hbar^2}{2m}\nabla_{\mathbf{r}_2}^2 + \frac{m}{2}\omega^2\mathbf{r}_1^2 + \frac{m}{2}\omega^2\mathbf{r}_2^2 + \frac{e^2}{|\mathbf{r}_1 - \mathbf{r}_2|} \quad (1)$$

with ω the harmonic oscillator frequency and m the mass of the electron. The Hamiltonian (1) is separable when written in the center-of-mass (CM) and relative coordinates $\mathbf{R} = \frac{1}{2}(\mathbf{r}_1 + \mathbf{r}_2)$ and $\mathbf{r} = \mathbf{r}_1 - \mathbf{r}_2$, respectively:

$$H = H_R + H_r = \left(-\frac{\hbar^2}{4m}\nabla_{\mathbf{R}}^2 + m\omega^2\mathbf{R}^2\right) + \left(-\frac{\hbar^2}{m}\nabla_{\mathbf{r}}^2 + \frac{m}{4}\omega^2\mathbf{r}^2 + \frac{e^2}{|\mathbf{r}|}\right). \quad (2)$$

As a consequence, the eigenfunctions $\Phi(\mathbf{r}_1, \mathbf{r}_2)$ are products of the eigenfunctions of the CM and relative Hamiltonians H_R and H_r , i.e. with $H_R|\phi\rangle = E_R|\phi\rangle$ and $H_r|\xi\rangle = E_r|\xi\rangle$ follows

$$\Phi(\mathbf{r}_1, \mathbf{r}_2)e^{-\frac{i}{\hbar}Et} = \Phi(\mathbf{R}, \mathbf{r})e^{-\frac{i}{\hbar}(E_R+E_r)t} = \phi(\mathbf{R})e^{-\frac{i}{\hbar}E_R t} \cdot \xi(\mathbf{r})e^{-\frac{i}{\hbar}E_r t}. \quad (3)$$

Due to the spherical symmetry of both H_R and H_r the eigenfunctions are conveniently expressed in the form $\phi_{NLM}(\mathbf{R}) = \frac{U_{NL}(R)}{R}Y_{LM}(\Theta, \chi)$ and $\xi_{nlm}(\mathbf{r}) = \frac{u_{nl}(r)}{r}Y_{lm}(\theta, \varphi)$.

The CM radial wavefunction and energy are readily obtained from the standard problem of the 3D harmonic oscillator to give $U_{NL}(R) = C_{NL}R^{L+1}H_N(R)e^{-m\omega R^2/\hbar}$ and $E_R = \hbar\omega(2N + L + \frac{3}{2})$. $H_N = {}_1F_1(-N, L + \frac{3}{2}, m\omega R^2/\hbar)$ is the N -th order Hermite polynomial, C_{NL} is the normalization constant. The radial wavefunction $u_{nl}(r)$ has the closed analytical form (unnormalized) $u_{nl}(r) = r^{\ell+1}e^{-m\omega r^2/4\hbar} \sum_{k=0}^{n-1} A^k r^k$ only for certain oscillator frequencies [3]. For other frequencies, it has to be evaluated numerically. Once it is found, the eigenenergy has the simple form $E_r = \hbar\omega(n + \ell + \frac{1}{2})$.

Adding the interaction with an electromagnetic field affects only the CM Hamiltonian so that the problem still separates. Having in mind the interaction with near infrared or visible laser pulses, we adopt the dipole approximation and obtain

$$H = \left(-\frac{\hbar^2}{m}\nabla_{\mathbf{r}}^2 + \frac{m}{4}\omega^2\mathbf{r}^2 + \frac{e^2}{|\mathbf{r}|}\right) + \left(\frac{1}{2 \cdot 2m} \left[-i\hbar\nabla_{\mathbf{R}} + \frac{2e}{c}\mathbf{A}(t)\right]^2 + m\omega^2\mathbf{R}^2\right) \quad (4)$$

with $\mathbf{A}(t)$ the vector potential and c the light velocity. The electric field of the laser is given by $\mathbf{E} = -\partial_t\mathbf{A}$ while the magnetic field is neglected[§] in dipole approximation. The total wave function reads

$$\Psi(\mathbf{R}, \mathbf{r}, t) = \psi(\mathbf{R}, t) \cdot \xi(\mathbf{r})e^{-\frac{i}{\hbar}E_r t} \quad (5)$$

with $\xi(\mathbf{r})$ the same as before.

[‡] Spin is omitted for brevity.

[§] Two parabolically confined electrons in constant magnetic fields were studied in [6].

The solution of the time-dependent Schrödinger equation governing the CM motion with twice the electron mass $\mu = 2m$ and charge $\epsilon = 2e$

$$\left(-i\hbar\partial_t + \frac{1}{2\mu} \left[-i\hbar\nabla_{\mathbf{R}} + \frac{\epsilon}{c}\mathbf{A}(t)\right]^2 + \frac{1}{2}\mu\omega^2\mathbf{R}^2\right)\psi(\mathbf{R}, t) = 0 \quad (6)$$

is known [7, 8]. In the case of a linearly polarized laser field $\mathbf{A}(t) = A(t)\mathbf{e}_z$ one has

$$\psi(\mathbf{R}, t) = U(X)e^{-\frac{i}{\hbar}E_X t}U(Y)e^{-\frac{i}{\hbar}E_Y t}U(Z, t), \quad (7)$$

with the two unaffected harmonic oscillator eigenfunctions $U(X) = C_{N_X}H_{N_X}(X)e^{-\mu\omega X^2/2\hbar}$, $U(Y) = C_{N_Y}H_{N_Y}(Y)e^{-\mu\omega Y^2/2\hbar}$ and the respective eigenenergies $E_X = \hbar\omega(N_X + \frac{1}{2})$, $E_Y = \hbar\omega(N_Y + \frac{1}{2})$. Instead, the eigenfunction $U_{N_Z}(Z)$ becomes dressed by the laser field and reads

$$\begin{aligned} U_{N_Z}(Z, t) &= C_{N_Z}H_{N_Z}\left(\sqrt{\frac{\mu\omega}{\hbar}}\left[Z - \frac{\epsilon g(t)}{\mu c}\right]\right) \times \\ &\times \exp\left\{-i\omega\left(N_Z + \frac{1}{2}\right)t + \frac{i\epsilon^2\omega^2}{2\mu\hbar c^2}\int_{t_0}^t\left(f^2(\tau) - g^2(\tau) - \frac{A^2(\tau)}{\omega^2}\right)d\tau + \right. \\ &\left. + \frac{i\epsilon\omega}{\hbar c}f(t)\left[Z - \frac{\epsilon g(t)}{\mu c}\right] - \frac{\mu\omega}{2\hbar}\left[Z - \frac{\epsilon g(t)}{\mu c}\right]^2\right\}. \end{aligned} \quad (8)$$

The normalization constants are given by $C_k = \left(\sqrt{\pi\hbar/\mu\omega} 2^k k!\right)^{-1/2}$. The laser field is turned on at time $t = t_0$. Before, the system is assumed to be in an eigenstate determined by the quantum numbers N_X , N_Y , and N_Z . The functions $f(t)$ and $g(t)$ are given by

$$f(t) = \cos\omega t \int_{t_0}^t A(\tau) \sin\omega\tau d\tau - \sin\omega t \int_{t_0}^t A(\tau) \cos\omega\tau d\tau, \quad (9)$$

$$g(t) = \sin\omega t \int_{t_0}^t A(\tau) \sin\omega\tau d\tau + \cos\omega t \int_{t_0}^t A(\tau) \cos\omega\tau d\tau. \quad (10)$$

From the structure of (8) one can infer (i) that the center of the CM wave packet describes motion along a trajectory $\mathbf{R}(t) = [0, 0, \frac{\epsilon}{\mu c}g(t)]$ where, in fact, $g(t)$ is proportional to the excursion of a driven, classical harmonic oscillator, and (ii) that the time-dependent solution $U_0(Z, t)$ represents a so-called *coherent* wave packet, i.e. a state of minimum uncertainty $\Delta p \Delta Z = \hbar/2$, equally distributed over the momentum p and the spatial coordinate Z (see, e.g. [9]).

3. Photoexcitation of the electron pair

Let us consider the photoexcitation of an electron pair confined in a parabolic well (e.g. two electrons occupying the low-lying states of a quantum dot). The transition amplitude is given by the overlap of the exact time-dependent two-electron wave function $\Psi(\mathbf{r}_1, \mathbf{r}_2, t)$ with the asymptotic stationary solution $\Phi(\mathbf{r}_1, \mathbf{r}_2)$ when the field is switched off:

$$T(t) = \langle\Phi|\Psi(t)\rangle = \langle\phi_{N_L N_M}|\psi_{N_x N_y N_z}(t)\rangle = \langle U_{K_z}|U_{N_z}(t)\rangle. \quad (11)$$

Here we made use of the fact that neither the sub-problem of relative motion of the two electrons nor the X and Y components of the CM motion are affected by the laser field. What is left is the probability to find the CM quasiparticle occupying a stationary state $U_{N_Z^{\text{fin}}}(Z)$ of the free oscillator after the action of the laser.

Let us first discuss the case where we start from the ground state, i.e. $N_X = N_Y = N_Z^{\text{ini}} = 0$, $\langle \mathbf{R} | \psi(t) \rangle = \psi_{000}(\mathbf{R}, t)$. The squared modulus of the corresponding transition amplitude to some final state with the quantum number $N_Z^{\text{fin}} \equiv n$ reads

$$\begin{aligned} |T^{(0 \rightarrow n)}(t)|^2 &= C_0^2 C_n^2 \left| \exp \left\{ in\omega t + i \frac{e^2 \omega^2}{2\mu \hbar c^2} \int_{t_0}^t \left(f^2(\tau) - g^2(\tau) - \frac{A^2(\tau)}{\omega^2} \right) d\tau \right\} \right|^2 \times \\ &\times \left| \int_{-\infty}^{\infty} dZ H_n \left(\sqrt{\frac{\mu\omega}{\hbar}} Z \right) \exp \left\{ -\frac{\mu\omega}{2\hbar} Z^2 + i \frac{e\omega}{\hbar c} f(t) \left[Z - \frac{e g(t)}{\mu c} \right] - \frac{\mu\omega}{2\hbar} \left[Z - \frac{e g(t)}{\mu c} \right]^2 \right\} \right|^2. \end{aligned} \quad (12)$$

One can omit all purely time-dependent imaginary exponents, producing unity, and reduce the integral to the table form [10] by change of variables

$$|T(t)|^2 = C_0^2 C_n^2 e^{-\frac{e^2 \omega}{2\mu \hbar c^2} (g^2(t) + f^2(t))} \left| \int_{-\infty}^{\infty} dx H_n(x) e^{-(x-y)^2} \right|^2 \quad (13)$$

with $x = \sqrt{\frac{\mu\omega}{\hbar}} Z$ and $y = \frac{e}{2c} \sqrt{\frac{\omega}{\mu\hbar}} (g(t) + i f(t))$.

As a result, the probabilities to find the CM of the electron pair in the ground (that is the survival probability) or in the n -th excited state are

$$P^{(0 \rightarrow 0)}(t) = |T^{(0 \rightarrow 0)}(t)|^2 = e^{-\sigma}, \quad (14)$$

$$P^{(0 \rightarrow n)}(t) = |T^{(0 \rightarrow n)}(t)|^2 = \frac{\sigma^n}{n!} e^{-\sigma}, \quad (15)$$

$$\sigma(t) = \frac{e^2 \omega}{2\mu \hbar c^2} [g^2(t) + f^2(t)]. \quad (16)$$

where we introduce the important parameter $\sigma(t)$, which depends on the laser field and the oscillator frequency [cf. eqs. (9) and (10)].

The formula for the photoexcitation probability (15) describes the distribution of the electronic wave packet among the oscillator states as a function of time and the EM field parameter. It has the form of a Poisson distribution, which is characteristic for stochastic processes where the system absorbs the first amount of energy from the field independently from the possibility of the absorption of the second portion and so on.

In Fig. 1 we illustrate the above expression for $P^{(0 \rightarrow n)}$ for a laser pulse of frequency $\omega_l = 2$ atomic units (au) with a Gaussian envelope and halfwidth $t_p = 1$ au, $A(t) = \sin \omega_l t \exp\left(-\frac{1}{2}(t/t_p)^2\right)$ (dashed curve). The survival probability $P^{0 \rightarrow 0}(t)$, changing from unity to some final value, has one maximum during the pulse due to the oscillation of the CM wave packet inside the potential and the associated maximum overlap with the ground state as the center of the wave packet passes through $Z = 0$. The transition probabilities of the excited states behave in the opposite way, i.e. they are minimum at these times.

In the same way as for the ground state excitations [cf. eqs. (11 – 15)], the general expression for the transition probability from an arbitrary initial m^{th} oscillator state

($t \leq t_0$) to the final n^{th} state can be found; it reads

$$P^{(m \rightarrow n)}(t) = {}_2F_0^2\left(-m, -n; -\frac{1}{\sigma}\right) \frac{\sigma^{n+m}}{n! m!} e^{-\sigma}, \quad (17)$$

with the hypergeometric function

$${}_2F_0\left(-m, -n; -\frac{1}{\sigma}\right) = \sum_{k=0}^{\min(n,m)} \frac{n!}{(n-k)!} \frac{m!}{(m-k)!} \frac{1}{k!} \left(-\frac{1}{\sigma}\right)^k. \quad (18)$$

Equation (17) circumscribes the time dependent population of the two-electron excited states following the laser pulse. As the Poisson distribution (15) characterizes the stochastic absorption of n photons, the distribution (17) is responsible for two processes, namely the independent emission of $m - k$ photons and the (re)absorption of $n - k$ photons. The polynomial sum (18) in the expression accounts for the transitions through all the allowed intermediate states k .

Of particular interest are the asymptotic excitation probabilities $\lim_{t \rightarrow +\infty} P^{(m \rightarrow n)}(t)$ since they are experimentally accessible observables. In Fig. 2 the distribution $P^{(m \rightarrow n)}$ is portrayed as a 2D intensity plot vs the (analytically continued) excitation quantum number n and the field-strength parameter σ . The latter is proportional to the laser intensity (see the discussion in the next section). Plots (a) and (b) correspond to the different initial states with $m = 0$ (ground state) and $m = 3$ accordingly. For $m = 0$ the quantum number of the maximally populated final state n^* is governed by the equation

$$\frac{d}{dn} P^{(0 \rightarrow n)} = \frac{d}{dn} \left(\frac{1}{n!} \sigma^n e^{-\sigma} \right) = 0, \quad (19)$$

which leads to the simple relation

$$\sigma = e^{\Lambda(n^*+1)} = n^* + \frac{1}{2} + o\left(10^{-1-\log n^*}\right). \quad (20)$$

$\Lambda(n) = \left. \frac{d^n}{dx^n} \Gamma(x) \right|_{x=0}$ is the Euler polygamma function.

4. Energy absorption

Let us now calculate the energy absorbed from the laser field by the two-electron system initially prepared in the ground state $E_{\text{abs}} = E(t \rightarrow \infty) - E(t = t_0)$, where $E(t) = \langle \psi_0(t) | \hat{H}(t) | \psi_0(t) \rangle$ with $\hat{H}(t)$ the total Hamiltonian^{||}

$$\langle \psi_0(t) | \hat{H}(t) | \psi_0(t) \rangle = \frac{\epsilon^2 \omega^2}{2\mu c^2} \left[g^2(t) + f^2(t) \right] + 2f(t) \frac{A(t)}{\omega} + \frac{A^2(t)}{\omega^2} + \frac{\hbar \omega}{2}, \quad (21)$$

$$E_{\text{abs}} = \lim_{t \rightarrow \infty} \frac{\epsilon^2 \omega^2}{2\mu c^2} \left[g^2(t) + f^2(t) \right] = \hbar \omega \lim_{t \rightarrow \infty} \sigma(t). \quad (22)$$

This expression for the absorbed energy is obtained independently of the initial state, i.e. using in eq. (21) the wave function $|\psi_m(t)\rangle$ (starting from an initial state m) also leads to the result (22). The absorbed energy depends on $\lim_{t \rightarrow \infty} \sigma(t)$ only but not on

^{||} We disregard the possible spontaneous decay of the excited system.

the initial state or the population over the final states of the system. The parameter of central importance thus clearly is $\sigma(t \rightarrow \infty) \equiv \sigma_o$, which can be recast into

$$\begin{aligned} \sigma_o &= \lim_{t \rightarrow \infty} \frac{\epsilon^2 \omega}{2\mu \hbar c^2} [g^2(t) + f^2(t)] = \frac{\epsilon^2 \omega}{2\mu \hbar c^2} \lim_{t \rightarrow \infty} |g(t) + if(t)|^2 = \\ &= \frac{\epsilon^2 \omega}{2\mu \hbar c^2} \left| \int_{-\infty}^{+\infty} dt' A(t') e^{i\omega t'} \right|^2 = \frac{\epsilon^2 \omega}{2\mu \hbar c^2} |A(\omega)|^2, \end{aligned} \quad (23)$$

showing that the spectral power of the incoming laser field *evaluated at the harmonic oscillator frequency* determines the energy absorption E_{abs} .

The expression for the absorbed energy (22) can be alternatively derived via the average number $\langle n \rangle$ of absorbed photons calculated using the photon statistics:

$$E_{\text{abs}} = \hbar \omega \langle n \rangle = \hbar \omega \sum_{n=0}^{\infty} n \frac{\sigma_o^n}{n!} e^{-\sigma_o} = \hbar \omega \sigma_o. \quad (24)$$

The essential fact is that the field strength parameter contains the cumulative mass and the charge of the CM subsystem with different powers $\sigma \sim \frac{\epsilon^2}{\mu}$. Therefore, the observables in the single and in the double-electron excitation differ from each other. In particular, the energy absorption from the same laser pulse by a (2e)-system is twice as large as in the one electron case.

In the generalized many-particle problem the parabolic potential contains an arbitrary number of electrons N . As in the (2e)-case the N -electron CM subhamiltonian accounting for the EM field can be separated from the field-independent relative motion of the electrons. Then, all the formulae describing two-electron transitions are valid for the many-electron case up to the effective charge and mass. Hence the energy absorption changes linearly with the number of electrons in the well, $E_{\text{abs}}^{(Ne)} / E_{\text{abs}}^{(1e)} = N$. Given this property, the number of active electrons in a harmonic system may be experimentally accessible.

Let us now discuss the two extreme cases of complete survival and full depletion of the initial state, that is its maximum and minimum survival probability. If the survival probability at the detection time equals unity, $P^{(m \rightarrow m)}(t \rightarrow \infty) \rightarrow 1$, the quantum system does not absorb energy from the laser field and becomes transparent for the corresponding pulse. This is strictly realized at the zero value of the field-strength parameter $\sigma_o = 0$ in the exponent in (17). Hence, according (23), the absence of the oscillator frequency in the spectrum of the vector potential, $A(\omega) = 0$, precludes absorption, [cf. eq. (22)]. Note, that this statement is correct for any field intensity and not just in a first order perturbative treatment as for nonparabolic confinements.

As an example, consider a laser field consisting of two consecutive Gaussians shifted in time by the interval $2a$: $A(t) = \sin[\omega_l(t+a)]e^{-\frac{1}{2}[(t+a)/t_p]^2} + \sin[\omega_l(t-a)]e^{-\frac{1}{2}[(t-a)/t_p]^2}$. Setting e.g., the laser frequency $\omega_l = 1$ and the halfwidth of pulses $t_p = \sqrt{2}/2$, we aim at finding a relation between the time delay $2a$ and the oscillator frequency ω , for which $0 = A(\omega; a) = -i\sqrt{\pi/\sqrt{e}} \cos \omega a \sinh \frac{\omega}{2} e^{-\omega^2/4}$. The result is $\omega a = \pi(k + 1/2)$ with k an integer. This is, in fact, a condition for the destructive interference (in time) of the two waves within the characteristic time-domain $2\pi/\omega$ of the system. In terms

of the excitation picture the explanation is the following: the electronic wave packet, distributed over the excited levels by the first pulse, may be assembled back to the initial state by the second pulse, if the phase difference between the pulses is chosen properly.

The opposite case amounts to the minimum survival probability, i.e. the maximum depletion of the initial state. On account of $e^{-\sigma_o} \rightarrow 0$ this is the case if the spectral power at the transition energy (coinciding with the oscillator frequency ω) is infinitely large. It is equivalent to the presence of an infinite, monochromatic laser pulse $A(t) = A_0 e^{-i\omega_l t}$ with $A(\omega) \sim A_0 \delta(\omega - \omega_l)$. Such a pulse resonantly depletes, for instance, the ground state. For a finite pulse, e.g., $A(t) = \sin \omega_l t e^{-\frac{1}{2}(t/t_p)^2}$ with a power spectrum $E_{\text{abs}} \sim |A(\omega)|^2 = 2\pi t_p^2 \sinh^2(\omega \omega_l t_p^2) e^{-(\omega^2 + \omega_l^2)t_p^2} \neq 0$, there is always nonvanishing absorption. However, only for the resonance $\omega_l = \omega$ the absorption exists in the limit of large pulse durations, $t_p \rightarrow \infty$.

In addition, if the initial state differs from the ground state, its full depletion occurs at such values of σ_o satisfying $\sigma_o^m \cdot {}_2F_0\left(-m, -n; -\frac{1}{\sigma_o}\right) = 0$. For example, the well prepared state with $m = 1$ will be exhausted and distributed over other levels by the laser field of the strength $\sigma_o = 1$ [cf. eq. (17)]: $P^{(1 \rightarrow 1)} = (\sigma_o - 1)^2 e^{-\sigma_o}$. Similar relations can be obtained for other higher order initial states, e.g. $\sigma_o = 2 \pm \sqrt{2}$ for $m = 2$, $\sigma_o - 3 - \alpha - \frac{3}{\alpha} = 2\sigma_o - 6 + \alpha - \frac{3}{\alpha} \pm i\sqrt{3} \left(\alpha - \frac{3}{\alpha}\right) = 0$, $\alpha = \left(3 + 3i\sqrt{2}\right)^{1/3}$ for $m = 3$ and so on.

5. Conclusion

In this paper we considered the exactly solvable problem of the excitation of two interacting electrons, confined in a parabolic well in the presence of a laser field. The expression for the time-dependent population of the excited two-electron states was derived. The physical features of the process, such as the Poisson statistics of the transition probabilities and the linear dependence of the energy absorption on the laser power, were analyzed, in particular, the conditions for the complete survival and full depletion of the initial state. Our approach can be used for the testing of approximate theories dealing with few-body multiphoton excitations.

6. Acknowledgments

One of us (O.K.) would like to thank A. Voitkiv and N. Fominykh for stimulating discussion. This work was supported by the Deutsche Forschungsgemeinschaft.

References

- [1] J. Ullrich *et al*, Rep. Prog. Phys. **66**, 1463 (2003).
- [2] N. R. Kestner and O. Sinanoglu, Phys. Rev. **128**, 2687 (1962).
- [3] M. Taut, Phys. Rev. A **48**, 3561 (1993).
- [4] S. Kais *et al*, J. Chem. Phys **9**, 417 (1993).
- [5] U. Schwengelbeck, Phys. Lett. A **253**, 168 (1999).

- [6] M. Taut, J. Phys. A **27**, 1045 (1994).
 [7] K. Husimi, Progr. Theor. Phys. **9**, 381 (1953).
 [8] P. Mulser *et al* Phys. Rev. A **48**, 4547 (1993).
 [9] M. O. Scully and M. S. Zubairy, *Quantum optics* (Cambridge UP, Cambridge, 1997).
 [10] I. S. Gradshteyn and I. M. Ryzhik, *Table of Integrals*, 5th ed., AP (1994) eq. 7.374-6, p. 843




Figure 1. The time evolution of the survival probability in the ground state, $n = 0$ (black curve), and the excitation probabilities to the states with the quantum numbers $n = 1, 2, 3, 4$ (green, blue, orange and olive curves, respectively) is shown together with the laser pulse (red dashed line) $A(t) = \sin \omega_l t e^{-(t/t_p)^2/2}$. The laser frequency, the half-width of the pulse and the oscillator frequency were chosen $\omega_l = 2$ au, $t_p = 1$ au, and $\omega = 1$ au respectively. For better visibility, the fast decrease of the probabilities with the quantum number $P^{(0 \rightarrow n)} \sim \frac{1}{n!}$ is compensated by multiplication with the factors given in brackets.




Figure 2. The asymptotic distribution $P^{(m \rightarrow n)}(t \rightarrow \infty)$ of the two-electron wave packet over the excited states, i.e. their population vs. the (analytically continued) quantum number and the field intensity is illustrated as a 2D intensity plot. Figures (a) and (b) correspond to the different initial states with $m = 0$ (ground state) and $m = 3$, respectively. The color scale of the population probability is common for both figures. The red line in (a) [determined by eq. (20)] denotes the position (the quantum number) of the maximally populated state for the given field-intensity parameter σ_0 (c.f. eq. (23)).