

A survey of energy loss calculations for heavy ions between 1 and 100 keV

A. Mangiarotti^{a,1}, M. I. Lopes^{a,b}, M. L. Benabderrahmane^c, V. Chepel^{a,b}, A. Lindote^{a,b},
J. Pinto da Cunha^{a,b}, P. Sona^d

^a*Laboratório de Instrumentação e Física Experimental de Partículas (LIP), 3004-516 Coimbra, Portugal*

^b*Departamento de Física, Universidade de Coimbra, 3004-516 Coimbra, Portugal*

^c*Physikalisches Institut der Universität Heidelberg, D-69120 Heidelberg, Germany*

^d*Dipartimento di Fisica, Università di Firenze and INFN Sez. Firenze, Italy*

Abstract

The original Lindhard-Scharff-Schiøtt (LSS) theory and the more recent Tilinin theory for calculating the nuclear and electronic stopping powers of slow heavy ions are compared with predictions from the SRIM code by Ziegler. While little discrepancies are present for the nuclear contribution to the energy loss, large differences are found in the electronic one. When full ion recoil cascade simulations are tested against the elastic neutron scattering data available in the literature, it can be concluded that the LSS theory is the more accurate.

Key words: Energy loss, Stopping power, Nuclear recoils, Dark matter.

PACS: 34.50.Bw, 78.70.-g, 95.35.+d.

1. Introduction

It is well known that an ion moving inside a medium can lose energy by collisions with both electrons and nuclei. Theoretically, it is necessary to account for both processes to reach an accurate description of the energy loss below a few keV/amu. From the experimental point of view, most detectors are just sensitive to electronic energy loss. Knowledge of the energy loss sharing between the two processes is mandatory for a detailed understanding of the response to particles

interacting with the detecting medium through nuclear recoils. This is the case for Weakly Interacting Massive Particles (WIMPs), which are possible constituents of the galactic dark matter.

The purpose of the present work is to explore the different available descriptions for the two parts of the energy loss between 1 and 100 keV. Range measurements are difficult at such low energies and almost all published data have been obtained employing elastic neutron scattering for transferring small and known amounts of energy to atoms of the detecting material. To reduce the theoretical difficulties, the attention is focused on pure substances: hence only symmetric projectile/target atom combinations will be investigated. From all the avail-

¹ Corresponding author. LIP, Dept. de Física da Universidade de Coimbra, Portugal. Tel.: +351-239-410657; fax: +351-239-822358. *E-mail address:* alessio@lipc.fis.uc.pt

able measurements, known to the authors, the following are then selected: Si [3,4], Ge [5,6], and liquid Xe [7,8,9,10]. Liquid Ar will also be considered on account of its interest for dark matter searches.

Among the few calculations from first principles without free parameters, there are still the original theory of Lindhard [1] and its reevaluation by Tiliin [2]; they will be examined here.

From all the available codes, only SRIM [11] will be discussed, because: i) it gives separately nuclear and electronic energy losses, ii) it covers the low energy range of interest, iii) from an independent survey [12] it was found the most accurate.

2. The nuclear stopping power

According to the Lindhard-Scharff-Schiøtt (LSS) theory [1], the nuclear stopping power S_n of a heavy ion is best described by rescaling its energy E and range R to the non-dimensional variables ϵ and ρ , respectively, defined as

$$\begin{cases} \epsilon = C_{\text{TF}} \frac{A_T}{A_{\text{tot}}} \frac{E/(2 E_B)}{Z_P Z_T Z^{1/2}} \\ \rho = 4\pi (a_B C_{\text{TF}})^2 \frac{A_P A_T}{A_{\text{tot}}^2} \frac{R N}{Z} \end{cases}, \quad (1)$$

with $Z = Z_P^{2/3} + Z_T^{2/3}$ and $A_{\text{tot}} = A_P + A_T$. In Eq. (1) N is the number density of the target material, Z_P and A_P are the atomic and atomic mass numbers of the projectile atom, respectively, Z_T and A_T are the correspondents for the target atom, a_B is the Bohr radius, E_B the Bohr energy and C_{TF} the Thomas-Fermi constant $(9\pi^2/27)^{1/3}$. In fact, while the nuclear part of the energy loss $(dE/dx)_n$ depends on the projectile and target ions, $(d\epsilon/d\rho)_n$ depends on the Thomas-Fermi interaction potential alone and is a universal function $f(\epsilon)$, which can be calculated numerically [13]. The values of S_n are reproduced with the physical units in Fig. 1.

A similar rescaling is employed in SRIM (Z in all appearances in Eq. (1) is replaced by an empirical $Z^{1/2} = Z_P^{0.23} + Z_T^{0.23}$), but now the universal function $f(\epsilon)$ is determined not from first principles but from a fit to experimental data. The results are also shown in Fig. 1.




Fig. 1. Nuclear stopping power S_n as a function of the ion energy for the symmetric projectile/target combinations considered in the present study.

While differences are small, it is known from sputtering data [14] that the LSS theory overestimates S_n . This problem is reduced in SRIM, as it is based on data themselves. Lindhard et al. [13] had also warned that the Thomas-Fermi treatment might be inadequate for very low energies (i.e. $\epsilon < 10^{-2}$), where mostly the tails of the ion-ion potential are probed. In the present study, this is of concern only for Xe nuclear recoils below 15 keV.

3. The electronic stopping power

The electronic energy loss $(dE/dx)_e$ of a unitary charge particle with a velocity β was described as an interaction with an electron plasma in the original work of Fermi and Teller. They explicitly distinguished two cases, for β above and below the Fermi velocity β_F . For $\beta < \beta_F$, $(dE/dx)_e$ was found to be proportional to β , with a proportionality coefficient being a unique function of the electron plasma density n_0 , usually expressed in terms of the Wigner-Seitz radius $r_s = (3n_0/(4\pi))^{1/3}$ [15]. Their result is plotted in Fig. 2. Typically, r_s needs to be corrected because tightly bound electrons contribute only marginally to $(dE/dx)_e$ much below the Bragg peak. In crystals, like Si and Ge, this effective density of the free electron plasma can be deduced from optical properties [16]. In liquids, like Ar and Xe, the problem is much more difficult and no correction was attempted in Fig. 2. Successively, Lindhard calculated in a self consistent way the local increase of the electron plasma density

around the intruder particle due to its Coulomb field. This leads to a higher $(dE/dx)_e$ as can be seen in Fig. 2. Finally, Ritchie considered the case where the Coulomb field is exponentially screened, slightly decreasing $(dE/dx)_e$ (see Fig. 2). The last effect is of particular relevance for ions, which can accommodate bound states while sweeping through the electron plasma. For a bare ion, a scaling with Z_P^2 to the elementary particle case is expected, as assumed in Fig. 2. In reality, this is not correct and the theory was extended to a partially ionized intruder by Ferrell and Ritchie [15], but the determination of the equilibrium charge of a given ion remains a difficult task. Lindhard also independently investigated this problem [1] and, using the Thomas-Fermi theory, arrived at a closed form for the proportionality coefficient. In terms of the non-dimensional variables introduced in Eq. (1), his result can be expressed as $(d\epsilon/d\rho)_e = \kappa \sqrt{\epsilon}$ where

$$\kappa = \frac{32}{3\pi} \sqrt{\frac{m_e c^2}{m_{\text{amu}} c^2}} \frac{Z_P^{1/2} Z_T^{1/2}}{Z^{3/4}} \frac{A_{\text{tot}}^{3/2}}{A_P^{3/2} A_T^{1/2}} \xi_e \quad (2)$$

with $\xi_e \approx Z_P^{1/6}$ (which is regarded only as an approximation by Lindhard) [1]. The points corresponding to the projectile/target combinations of interest for the present study are also reported in Fig. 2. The suppression of the electronic energy loss occurs mostly due to the partial ionization of the intruder. It strongly increases with Z_P ($Z_P = Z_T$).

The proportionality of $(dE/dx)_e$ with β is also a feature of the SRIM code, allowing a value of $(dE/dx)_e/(\beta Z_P^2)$ to be extracted (see Fig. 2). While SRIM exceeds the LSS theory for Si, it then decreases consistently below it, up to a factor of 4 for Xe. The reason for this discrepancy is unclear, because the details on the implementation of $(dE/dx)_e$ in SRIM for low velocities are not public, but it probably resides in the estimate of the intruder charge state. It has been verified that data for protons of comparable energies per nucleon on Ar and Xe are well reproduced by SRIM.

The big drawback of the described theoretical approaches is to assume that the electronic and nuclear collisions are uncorrelated [1]. In reality, the screened Coulomb repulsion between the two interacting nuclei makes part of the impact pa-




Fig. 2. Proportionality coefficient of the electronic energy loss to the particle velocity as a function of the Wigner-Seitz radius r_s . For ions a Z_P^2 scaling is applied.

rameter range unavailable for the scattering of the electrons belonging to the target atom in the screened Coulomb field of the projectile and vice versa. Tiliin [2] has shown that the final net effect is a great decrease of S_e for $\epsilon \ll 1$, with a corresponding lack of proportionality to $\sqrt{\epsilon}$. His results can also be recast in the form of Eq. (2) where ξ_e is replaced by a function $\tau(\epsilon, Z_P/Z_T)$ that can be tabulated [2]. In the present case, his theory predicts roughly half the value of S_e expected from LSS and SRIM for Si and approximately agrees with SRIM for Xe.

4. The full ion recoil cascade

As mentioned, resort must be made to response measurements employing elastic neutron scattering, where recombination or quenching may influence the final fraction of the total energy transferred to electrons that is detectable as excitation or ionization, particularly in the case of scintillation yield for LXe. Theoretically, however, an even bigger disadvantage is present; especially for high Z elements: S_n dominates over S_e and most of the primary ion energy is transferred to nuclear recoils. The knocked ion undergoes the same process, resulting in a full cascade of recoils, whose total electronic energy loss must be evaluated. In the case of the SRIM code, a second program, called TRIM [11], reads S_n and S_e from the first and performs the computation. Lindhard and his group solved numerically the transport equations corre-




Fig. 3. Total fraction of the initial ion energy transferred to electrons integrated over the full cascade as a function of the ion energy itself.

sponding to the LSS theory and arrived at a parameterization in terms of ϵ and κ (see Eq. (9) of Ref. [17]). Both results are compared with data in Fig. 3 (the two series of points for Ge are inconsistent). The LSS theory is on average better, while SRIM both overpredicts and strongly underpredicts the data for Si and Xe, respectively. While the first discrepancy could be ascribed to recombination, the second appears even more surprising due to the probable presence of quenching.

5. Conclusions

For slow heavy ions, the nuclear stopping power predicted by the original LSS theory and the current SRIM code differ at most by $\approx 15\%$. On the contrary, for the electronic stopping power, big discrepancies are present between the LSS theory, the theory of Tilinin and SRIM (up to a factor of ≈ 4 for Xe). Judging from the elastic neutron scattering data, the LSS theory seems the best of all. More detailed full cascade simulations will be performed in the future to assess the robustness of this conclusion. New experimental data for Germanium would be highly needed for clarifying the situation.

Acknowledgments This work was supported by FCT/FEDER/POCI-2010 fund (project POCI/FP/63446/2005).

References

- [1] J. Lindhard et al., *Mat. Fis. Medd. Dan. Vid. Selsk.* 33 (1963) No.14.
- [2] I.S. Tilinin, *Phys. Rev. A* 51 (1995) 3058.
- [3] A.R. Sattler, *Phys. Rev.* 138 (1965) A1815.
- [4] G. Gerbier et al., *Phys. Rev. D* 42 (1990) 3211.
- [5] C. Chasman et al., *Phys. Rev. Lett.* 15 (1965) 245.
- [6] A.R. Sattler et al., *Phys. Rev.* 143 (1966) 588.
- [7] F. Arneodo et al., *Nucl. Instr. Meth. A* 449 (2000) 147.
- [8] D. Akimov et al., *Phys. Lett. B* 524 (2002) 245.
- [9] E. Aprile et al., *Phys. Rev. D* 72 (2005) 072006.
- [10] V. Chepel et al., *Astrop. Phys.* 26, (2006) 58.
- [11] J.F. Zigler, Program SRIM/TRIM, version 2003.26, obtained from <http://www.srim.org>.
- [12] H. Paul et al., *Nucl. Instr. Meth. B* 209 (2003) 252.
- [13] J. Lindhard et al., *Mat. Fis. Medd. Dan. Vid. Selsk.* 36 (1968) No.10.
- [14] H. Oetzmann et al., *Phys. Lett. A* 55 (1975) 170.
- [15] T.L. Ferrell et al., *Phys. Rev. B* 16 (1977) 115.
- [16] A. Mann et al., *Phys. Rev. B* 9 (1981) 4999.

[17] J. Lindhard et al., Mat. Fis. Medd. Dan. Vid. Selsk.
33 (1963) No.10.