

Computational Consequences of Neglected First-Order van der Waals Forces

Kevin Cahill^{1,2,*} and V. Adrian Parsegian^{3,†}

¹*Department of Physics and Astronomy, University of New Mexico, Albuquerque, NM 87131*

²*Center for Molecular Modeling, Center for Information Technology,*

National Institutes of Health, Bethesda, Maryland 20892-5624

³*National Institute of Child Health and Human Development,*

National Institutes of Health, Bethesda, Maryland 20892-0924

(Dated: October 28, 2018)

We have computed the widely neglected first-order interaction between neutral atoms. At interatomic separations typical of condensed media, it is nearly equal to the $-1/r^6$ second-order London energy inferred from interactions in gases. Our results, without the exchange forces that lead to covalent bonding, suggest that the quality of non-bonding attraction between neutral atoms of molecules in condensed media differs from the $-1/r^6$ form usually ascribed to it. If we add first-order and all second-order terms, we obtain energies nearly five times the $-1/r^6$ London energies which dominate only at the atomic separations found in gases. For computation, we propose a practical, accurate form of energy to replace the qualitatively inaccurate Lennard-Jones and harmonic forms casually assumed to hold at the interatomic separations found in condensed media.

PACS numbers: 82.35.Pq, 34.20.Cf, 77.84.Jd

Does the van der Waals potential energy between two neutral atoms fall off as $-1/r^6$ at separations of a few Å? Is the widely neglected first-order van der Waals potential zero at the interaction distances within condensed media? Should one use a Lennard-Jones 6-12 potential in computer simulations? Because it is no longer necessary to sacrifice accuracy for computational convenience, we numerically re-examine these old questions, find that common practice leads to qualitative error, and propose an accurate but practical alternative to the mathematically convenient but potentially misleading Lennard-Jones (L-J) and harmonic forms.

Motivated by an intriguing footnote in Landau and Lifshitz [1], we compute the first-order van der Waals potential for two hydrogen atoms, without the exchange interaction that creates covalent bonds. It is nearly equal to the $1/r^6$ second-order London energy found from the same wave functions when evaluated over a range of separations where both exceed kT at room temperature. We then compute the potential energy to second-order, obtaining an expression that reassuringly reproduces the expected dominance of the $1/r^6$ London term as $r \rightarrow \infty$. The net result at separations $3a_0 < r < 5a_0$ (Bohr radius a_0) is an energy 4 to 5 times that expected from a London interaction, which emerges to dominate only in the $r > 10a_0$ limit.

We next examine the additivity of the interactions of several particles, usually taken to be additive in first and second order; the Axilrod-Teller-Muto force [2, 3] is a third-order effect. In fact, energies add in first-order perturbation theory but not in second order unless one neglects first-order terms.

Constructed to fit the measured equilibrium separation r_s and the measured condensation energy $V(r_s)$ as well as to have the large- r asymptotic form $1/r^6$, the

Lennard-Jones potential is too hard for $r < r_s$ and varies incorrectly for $r > r_s$. Harmonic potentials work only near energy minima. We propose a simple, phenomenological potential that can fit spectroscopic data and that behaves appropriately at all distances.

For our calculations, we take the hamiltonian for two interacting hydrogen atoms to be $H = H_0 + W$ in which H_0 is a sum of two isolated-hydrogen-atom hamiltonians and the perturbation W is a function of the distance r between the two protons

$$W = \frac{e^2}{r} + \frac{e^2}{|\vec{r} + \vec{r}_2 - \vec{r}_1|} - \frac{e^2}{|\vec{r} + \vec{r}_2|} - \frac{e^2}{|\vec{r} - \vec{r}_1|}. \quad (1)$$

Here $r = |\vec{r}|$ and e is the charge of the electron in units with $\alpha = e^2/(\hbar c) \approx 1/137$. The first-order correction to the energy is $\Delta E_1(r) = \langle 0|W|0 \rangle$ in which the unperturbed ground state $|0\rangle = |100, 100\rangle$ is two $1s$ states separated by a distance r . Its wave function is $\langle \vec{r}_1, \vec{r}_2|0 \rangle = (1/\pi a_0^3) \exp(-r_1/a_0 - r_2/a_0)$ where $a_0 = \hbar^2/(me^2)$ is the Bohr radius, $r_1 = |\vec{r}_1|$, and $r_2 = |\vec{r}_2|$. Integration of $\langle 0|W|0 \rangle$ yields

$$\Delta E_1(r) = mc^2 \alpha^2 e^{-2r/a_0} \left(\frac{a_0}{r} + \frac{5}{8} - \frac{3r}{4a_0} - \frac{r^2}{6a_0^2} \right). \quad (2)$$

London and others [4, 5, 6] applied second-order perturbation theory to the potential (1), expanded for large r , and got $\Delta E_L(r) = -6.50 mc^2 \alpha^2 a_0^6/r^6$.

Fig. 1 shows that the two energies $\Delta E_1(r)$ and $\Delta E_L(r)$ are nearly equal for $3a_0 < r < 5a_0$, where both are reliable and greater than $kT = 1/40$ eV. Because $\Delta E_1(r)$ is the first-order result, and $\Delta E_L(r)$ is an approximation to second-order perturbation theory, the two can be added. That is, to (approximate) second order in the perturbation W and within the London approximation, the van der Waals energy is $\Delta E_1(r) + \Delta E_L(r)$, different by about

100% from the London-only result for $3a_0 < r < 5a_0$. Because $\Delta E_1(r)$ decreases with r as $\exp(-2r/a_0)$, the London potential $\Delta E_L(r)$ dominates at large r . But it is not until $r = 5.78a_0$ that $|\Delta E_L(r)| > 2|\Delta E_1(r)|$. By then $|\Delta E_L(r)| < kT/5$.

What about second order beyond $\Delta E_L(r)$? The correction to the ground-state energy due to the perturbation W is

$$\Delta E_2(r) = - \sum_{n \neq 0} \frac{|\langle 0|W|n \rangle|^2}{E_n - E_0} \quad (3)$$

in which the sum is over all eigenstates $|n\rangle$ of the hamiltonian H_0 except $|0\rangle$. If the first state to contribute to the sum has energy E_1 , then $0 < E_1 - E_0 \leq E_n - E_0$ or $-1/(E_1 - E_0) \leq -1/(E_n - E_0)$. So the second-order change in the energy is bounded by

$$- \frac{1}{E_1 - E_0} \sum_{n \neq 0} |\langle 0|W|n \rangle|^2 \leq \Delta E_2(r) \leq 0. \quad (4)$$

The first states to contribute to the sum are those in which one of the electrons is in the $n = 2$ level, so $E_1 = -(1/2)\alpha^2 mc^2(1 + 1/4)$. But because the hydrogen wave functions fall off as $\exp(-r/(na_0))$, at long distances the first states to contribute are those with both electrons in the $n = 2$ level; they have $E_1 = -(1/2)\alpha^2 mc^2(1/4 + 1/4)$. Without the energy denominators, the sum over intermediate states follows from completeness:

$$\sum_{n \neq 0} |\langle 0|W|n \rangle|^2 = \langle 0|W^2|0 \rangle - \langle 0|W|0 \rangle^2, \quad (5)$$

where $\langle 0|W|0 \rangle$ is the first-order result $\Delta E_1(r)$, Eq.(2).

Evaluating the mean value of W^2 involves the integral $G(\vec{x}) = (2/\pi) \int d^3y (1/\bar{y}^2) \exp(-2|\vec{y}-\vec{x}|)$, a function only of the length $x = |\vec{x}| = |\vec{r}|/a_0$. We find

$$G(x) = \frac{1}{x} [(1 + 2x) e^{-2x} \text{Ei}(2x) + (1 - 2x) e^{2x} \text{E}_1(2x)] \quad (6)$$

in which Ei and E_1 are exponential integrals. $G(x)$ contributes to the integral $J(x)$

$$J(x) = \frac{1}{4x} \int_0^\infty y G(y) [(1 + 2|x-y|) e^{-2|x-y|} - (1 + 2(x+y)) e^{-2(x+y)}] dy \quad (7)$$

that represents the square of the interaction of the two electrons. The product of electron-electron and electron-proton energies is described by the function $K(x)$

$$K(x) = \left\{ 4 \ln 2 - 3 + \frac{2 \ln 2}{x} + 2x - \left(4 + \frac{2}{x} \right) [\text{Ei}(2x) - \gamma - \ln(2x)] \right\} e^{-2x} + \left(4 - \frac{2}{x} \right) e^{2x} [\text{E}_1(2x) - \text{E}_1(4x)] \quad (8)$$




FIG. 1: The first-order van der Waals energy $\Delta E_1(r)$, Eq.(2), (solid, red) is nearly equal to the London result $\Delta E_L(r)$ (long dashes, green) when both are reliable and greater than kT (dotted, plum). The change in the sum $\Delta E(r) = \Delta E_1(r) + \Delta E_2(r)$ (dashes, blue) as given by second-order perturbation theory, Eq.(9), is 4 to 5 times greater than the London term for $3a_0 < r < 5a_0$. The potential of Eq.(15) with $a = 336$ eV, $b = 1.97a_0^{-1}$, $c = 0.47a_0^{-1}$, $d = 397a_0^6$ eV, and $e = 14093a_0^{12}$ (dashes, light blue) overlaps $\Delta E(r)$.

in which $\gamma = 0.5772\dots$ is the Euler-Mascheroni constant. In terms of ΔE_1 , J , and K , the second-order correction $\Delta E_2(r)$ to the energy at $r = a_0 x$ becomes $-(f a_0/e^2) (\langle 0|W^2|0 \rangle - \langle 0|W|0 \rangle^2)$ or

$$\Delta E_2(r) = -\frac{f a_0}{e^2} \left\{ - \left(\Delta E_1(r) - \frac{e^2}{r} \right)^2 + \frac{2e^4}{r^2} [1 - (1+x) e^{-2x}]^2 + \frac{e^4}{a_0^2} [G(x) + J(x) + K(x)] \right\}. \quad (9)$$

The fudge factor $f = 1.083 = 6.499/6$ is the one used by Pauling and Beach [7] to estimate the effect of the energy denominators in the sum Eq.(3).

To second order in the perturbation W , the change in the energy of two hydrogen atoms at proton separation r is the sum of the first-order term Eq.(2) and the second-order term Eq.(9), $\Delta E(r) = \Delta E_1(r) + \Delta E_2(r)$. Fig. 1 shows $\Delta E(r)$ for $3 \leq r/a_0 \leq 5$ where ΔE is never even close to ΔE_L . (To model non-reactive atoms, we have neglected the exchange terms that cause covalent bonding and only increase $|\Delta E(r)|$.)

What about additivity? Consider three hydrogen atoms at \vec{R}_1 , \vec{R}_2 , and \vec{R}_3 . Again neglect exchange effects and take the unperturbed ground state to be

$$|1, 2, 3\rangle = |01, 02, 03\rangle = |1s, \vec{R}_1; 1s, \vec{R}_2; 1s, \vec{R}_3\rangle. \quad (10)$$

Now the perturbation W is $W = W_{12} + W_{23} + W_{13}$ in which the pair potential W_{ij} is

$$W_{ij} = \frac{e^2}{|\vec{R}_j - \vec{R}_i|} + \frac{e^2}{|\vec{R}_j + \vec{r}_j - \vec{R}_i - \vec{r}_i|} - \frac{e^2}{|\vec{R}_j + \vec{r}_j - \vec{R}_i|} - \frac{e^2}{|\vec{R}_j - \vec{R}_i - \vec{r}_i|}. \quad (11)$$

To first order in the perturbation W , the change in the energy $\Delta E_1 = \langle 1, 2, 3 | W_{12} + W_{23} + W_{31} | 1, 2, 3 \rangle$ is additive: $\Delta E_1 = \langle 1, 2 | W_{12} | 1, 2 \rangle + \langle 2, 3 | W_{23} | 2, 3 \rangle + \langle 1, 3 | W_{13} | 1, 3 \rangle$.

But at smaller separations the cross-terms do not vanish. To second order in W , ΔE_2 is not quite additive. The mean value of $W_{12}W_{23}$, *e.g.*, involves a non-zero sum over the intermediate states $|1, n2, 3\rangle = |01, n2, 03\rangle$,

$$\begin{aligned} \langle 1, 2, 3 | W_{12} W_{23} | 1, 2, 3 \rangle &= \\ &= \sum_n \langle 1, 2, 3 | W_{12} | 1, n2, 3 \rangle \langle 1, n2, 3 | W_{23} | 1, 2, 3 \rangle \\ &= \sum_n \langle 1, 2 | W_{12} | 1, n2 \rangle \langle n2, 3 | W_{23} | 2, 3 \rangle \neq 0. \end{aligned} \quad (12)$$

Failure of additivity already occurs at second order.

In the usual treatment, one expands the perturbation W in powers of $1/r$ keeping only terms as big as $1/r^6$ at large r and neglects terms like Eq.(12). If $\vec{R}_2 - \vec{R}_1$ points in the 3-direction, then the leading term in the matrix element $\langle 1, 2 | W_{12} | 1, n2 \rangle$ involves $(x_1x_2 + y_1y_2 - 2z_1z_2)/|\vec{R}_2 - \vec{R}_1|^3$ which vanishes because of the spherical symmetry of the ground state of atom 1. This neglect of the first-order van der Waals force defers non-additivity to the third order of perturbation theory in the Axilrod-Teller-Muto formulation [2, 3]. This deferral, however, is probably inconsequential compared with the prior omission of purely additive first-order interactions.

Clearly there is room for improvement on the Lennard-Jones [8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23] and harmonic forms

$$V_{LJ}(r) = |V(r_s)| \left[\left(\frac{r_s}{r} \right)^{12} - 2 \left(\frac{r_s}{r} \right)^6 \right] \quad (13)$$

$$V_H(r) = V(r_s) + \frac{(r - r_s)^2}{2} \frac{d^2 V(r_s)}{dr^2}. \quad (14)$$

What to do? We suggest using the form

$$V(r) = ae^{-br} (1 - cr) - \frac{d}{r^6 + er^{-6}}. \quad (15)$$

The terms involving a , b , and c appear in Eq.(2) for $\Delta E_1(r)$, were proposed by Rydberg [24] to incorporate spectroscopic data, but were largely ignored until recently [25, 26]. The constant $d = C_6$ is the coefficient of the London tail. The new term er^{-6} cures the London singularity; as $r \rightarrow 0$, $d/(r^6 + er^{-6}) \rightarrow dr^6/e$ and $V(r) \rightarrow a$, finite. But as $r \rightarrow \infty$, $V(r)$ approaches the London term, $V(r) \rightarrow -d/r^6 = -C_6/r^6$.




FIG. 2: The phenomenological potential (15) with $a = 53.8$ eV, $b = 2.99 \text{ \AA}^{-1}$, $c = 2.453 \text{ \AA}^{-1}$, $d = C_6 = 3.884 \text{ eV \AA}^6$, and $e = 47.6 \text{ \AA}^{12}$ (solid, red) is finite, fits the RKR spectral points for molecular hydrogen (crosses, blue), and gives the correct London tail for $r > 3 \text{ \AA}$. The harmonic potential V_H (dashes, green) is accurate only near its minimum.

To test the ability of the potential $V(r)$ to represent covalent bonds, we used Gnuplot [27] to choose a , b , c , d , and e so as to fit Eq.(15) to the experimental potentials of molecular H_2 , N_2 , and O_2 obtained from spectroscopic data [28, 29, 30] by the Rydberg [24]-Klein [31]-Rees[32] (RKR) method. For H_2 , we set d equal to the London value, $d = C_6 = 6.5 mc^2 \alpha^2 a_0^6$. Fig.(2) shows that the potential of Eq.(15) (solid, red) goes through the RKR points for H_2 (crosses, blue), while the harmonic potential (14) (dashed, green) is accurate only near its minimum and is dreadful for $r \gg r_s$. The fits of $V(r)$ to the N_2 and O_2 RKR points were equally good. The potential $V(r)$ of Eq.(15) represents covalent bonds better than the harmonic potential V_H of Eq. (14) (or V_{LJ}).

Can $V(r)$ represent weak noncovalent bonds? Using Gnuplot, we fitted a , b , c , and e in Eq.(15) to calculated He_2 [33] or empirical Ar_2 and Kr_2 [34] potentials, setting d equal to the respective London coefficient. Fig.(3) shows that $V(r)$ of Eq.(15) (solid, red) fits the empirical points [34] for Kr_2 (crosses, blue), but that the Lennard-Jones potential V_{LJ} of Eq.(13) (dashes, green) is too low for $r > 4 \text{ \AA}$. Fig. (4) shows that $V(r)$ fits the empirical Kr_2 points, but that V_{LJ} is much too hard for $r < 4 \text{ \AA}$. The fits for He_2 and Ar_2 were equally good. Figs.(2-4) would look even better with the London d treated as a free parameter. The potential $V(r)$ of Eq.(15) represents weak noncovalent bonds better than V_{LJ} or V_H . V_{LJ} is unnaturally hard for $r < r_s$; it can wrongly reject Monte Carlo moves that put atoms slightly too close, frustrating M-C searches for the native states of macromolecules.




FIG. 3: The potential V of Kr-Kr attraction (Eq.(15) with $a = 2296$ eV, $b = 2.5136 \text{ \AA}^{-1}$, $c = 0.2467 \text{ \AA}^{-1}$, $d = 78.2146$ eV \AA^6 , and $e = 407923 \text{ \AA}^{12}$) (solid, red) is finite, fits the Kr_2 Aziz points (crosses, blue), and gives the correct London tail. When matched at the minimum, the Lennard-Jones form V_{LJ} (Eq.(13) with $r_s = 4.008 \text{ \AA}$ and $V(r_s) = -0.017338$ eV) (dashes, green) is too low for $r > 4 \text{ \AA}$.




FIG. 4: Positive potential V (Eq.(15) as in Fig. (3)), Kr_2 Aziz points (crosses, blue), L-J form V_{LJ} (Eq.(13) as in Fig. (3)).

The van der Waals potential is often as large in first order as in second. The potential (15) may be useful in macromolecular simulations that involve not only bonded but also non-bonded atoms. Given the differences between Eq.(15) and the popular Lennard-Jones form, it would be worthwhile to examine the consequences of these differences in numerical simulations.

Thanks to E. Arriola and A. Calle for pointing out a

typo in Eq.(7); to S. Valone for RKR data; to S. Atlas, C. Beckel, B. Brooks, J. Cohen, K. Dill, D. Harries, G. Herling, M. Hodoscek, R. Pastor, R. Podgornik, W. Saslow, and C. Schwieters for advice. P. J. Steinbach kindly hosted KC at NIH, where we used the Biowulf computers.

* Electronic address: cahill@unm.edu

† Electronic address: aparsegi@helix.nih.gov

- [1] L. D. Landau and E. M. Lifshitz, *Quantum Mechanics* (Pergamon, 1976), p. 341, 3rd ed.
- [2] B. M. Axilrod and E. Teller, *J. Chem. Phys.* **11**, 299 (1943).
- [3] Y. Muto, *Phys.-Math. Soc. Jap.* **17**, 629 (1943).
- [4] R. Eisenschitz and F. London, *Z. Physik* **60**, 491 (1930).
- [5] F. London, *Z. Physik* **63**, 245 (1930).
- [6] L. Pauling and E. B. Wilson, Jr., *Introduction to Quantum Mechanics* (Dover, Mineola, NY, 1935), s. XIV-47a.
- [7] L. Pauling and J. Y. Beach, *Phys. Rev.* **47**, 686 (1935).
- [8] H. Margenau, *Rev. Mod. Phys.* **11**, 1 (1939).
- [9] R. Pastor, Ph.D. thesis, Harvard University (1984).
- [10] S. R. Atlas, Ph.D. thesis, Harvard University (1988).
- [11] T. A. Halgren, *J. Am. Chem. Soc.* **114**, 7827 (1992).
- [12] W. J. Meath and D. J. Margoliash, *J. Chem. Phys.* **68**, 1426 (1978).
- [13] A. D. Buckingham, P. W. Fowler, and J. M. Hutson, *Chem. Rev.* **88**, 963 (1988).
- [14] D. Ben-Amotz and D. R. Herschbach, *J. Phys. Chem.* **94**, 1038 (1990).
- [15] D. Ben-Amotz and K. G. Willis, *J. Phys. Chem.* **97**, 7736 (1993).
- [16] G. Chałasiński and M. Szcześniak, Małgorzata, *Chem. Rev.* **94**, 1723 (1994).
- [17] B. Jeziorski, R. Moszynski, and K. Szalewicz, *Chem. Rev.* **94**, 1887 (1994).
- [18] F. B. van Duijneveldt, J. G. C. M. van Duijneveldt-van de Rijdt, and J. H. van Lenthe, *Chem. Rev.* **94**, 1873 (1994).
- [19] D. Yin and A. D. MacKerell, Jr., *J. Comp. Chem.* **19**, 334 (1998).
- [20] T. C. Choy, *Phys. Rev. A* **62**, 012506 (2000).
- [21] I. J. Chen, D. Yin, and A. D. MacKerell, Jr., *J. Comp. Chem.* **23**, 199 (2001).
- [22] D. Ben-Amotz and K. G. Willis, *J. Chem. Phys.* **117**, 6590 (2002).
- [23] A. I. Volokitin and B. N. J. Persson, *Phys. Rev. B* **65**, 115419 (2002).
- [24] R. Rydberg, *Z. Phys.* **73**, 376 (1931).
- [25] J. Ferrante, H. Schlosser, and J. R. Smith, *Phys. Rev. A* **43**, 3487 (1991).
- [26] J. N. Murrell, S. Carter, S. C. Farantos, P. Huxley, and A. J. C. Varandas, *Molecular Potential Energy Functions* (John Wiley & Sons, 1984), p. 9.
- [27] <http://www.gnuplot.info/>.
- [28] S. Weissman, J. T. Vanderslice, and R. Battino, *J. Chem. Phys.* **39**, 2226 (1963).
- [29] A. Lofthus and P. H. Krupenie, *J. Phys. Chem. Ref. Data* **6**, 113 (1977).
- [30] P. H. Krupenie, *J. Phys. Chem. Ref. Data* **1**, 423 (1972).
- [31] O. Klein, *Z. Phys.* **76**, 226 (1932).
- [32] A. L. G. Rees, *Proc. Phys. Soc. (London)* **59**, 998 (1947).

- [33] J. B. Anderson, *J. Chem. Phys.* **115**, 4546 (2001).
- [34] R. A. Aziz and M. J. Slaman, *Mol. Phys.* **58**, 679 (1986).