Correction algorithm for finite sample statistics

Thorsten Pöschel¹, Werner Ebeling¹, Cornelius Frömmel¹, and Rosa Ramírez²

¹ Humboldt Universität zu Berlin, Charité, Institut für Biochemie, Monbijoustraße 2, D-10117 Berlin, Germany

 2 Centre Européen de Calcul Atomique et Moléculaire (CECAM), Ecole Normale Supérieure, 46, Allée d'Italie, Fr-69007 Lyon, France

Fr-09007 Lyon, Fra

Received: November 21, 2018/ Revised version:

Abstract. Assume in a sample of size M one finds M_i representatives of species i with $i = 1 \dots N^*$. The normalized frequency $p_i^* \equiv M_i/M$, based on the finite sample, may deviate considerably from the true probabilities p_i . We propose a method to infer rank-ordered true probabilities r_i from measured frequencies M_i . We show that the rank-ordered probabilities provide important informations on the system, e.g., the true number of species, the Shannon- and the Renyi-entropies.

PACS. 02.50.-r Probability theory, stochastic processes, and statistics -02.60-x Numerical approximation and analysis -07.05.Kf Data analysis: algorithms and implementation; data management

Key words. Probability distribution, finite sample statistics, biometrics

1 Introduction

In experimental work one frequently faces the problem to determine the probabilities of occurrence (or concentrations) p_1, p_2, \ldots, p_N of species $1, 2, \ldots, N$. The probability of species *i* is defined by

$$p_i = \lim_{M \to \infty} \frac{M_i}{M}, \qquad i = 1 \dots N$$
 (1)

$$M = \sum_{j=1}^{N} M_j \,, \tag{2}$$

with M_i being the number of representatives of the species i found in a sample of size M. N is the number of different species which will appear in a sample of *infinite* size. Of course M will never be infinite in reality, but a number which is determined mainly by the experimental effort, i.e., usually costs and time. In this article the term "species" is not used in its strict phylogenetic sense, but it stands as a synonym for "distinguishable events which are members of a statistical ensemble".

A prominent example where we cannot reliably infer the probabilities from counted frequencies is the distribution of words of length n in nucleotide sequences such as DNA. Since we have an alphabet of four letters, there are 4^n words which in principle could be constructed. Even for moderate values of n, the number of words exceeds the size of *any* available data base. If we want to compute, the entropy of the word distribution in biosequences we have to apply, therefore, correction methods, e.g. [1,2,3,4,5,6,7].

Virtually each experimental measurement of concentrations (or probabilities) is affected by finite size effects due to the feasible number M of samples which can be investigated. In a real measurement one cannot even expect to find the correct number N of species. Instead, in general, a smaller number N^* is found, depending on the sample size M. We will show that, even if M is a rather large number, the deviations of the observed relative frequencies

$$p_i^* = \frac{M_i}{M}, \qquad M \text{ finite}$$
(3)

from the true probabilities p_i as defined by Eq. (1) may be significant. A method to deduce true probabilities from measured relative frequencies is, therefore, highly desirable.

The aim of this article is to propose a method to correct relative frequencies $p_i^* = M_i/M$ in a finite sample of size M in a way to approximate the true probabilities p_i which would be obtained if a sample of infinite size was investigated. Our method is based on the idea, that the estimation of rank-ordered probabilities is by orders of magnitude more easy than the estimation of the speciesordered probabilities. This is due to the fact that in the rank-ordering procedure the exact relation between the species number i and the probability p_i is ignored. This way it remains to estimate the shape of a function r_i which is monotonously decreasing with the rank i (see Sec. 2 for the definition of r_i). The large interest in rank-ordered distributions is based on the fact that several characteristic quantities as the Renyi-entropies [8]

$$H^{(q)} = \frac{1}{1-q} \log\left(\sum p_i^q\right) = \frac{1}{1-q} \log\left(\sum r_i^q\right)$$
(4)

are invariant with respect to the ordering. Therefore, the rank-ordered distribution suffices to compute the Renyi entropies. We notice the important relations $M = \exp H^{(0)}$, $H = H^{(1)}$. In other words, the true number of species M as well as the Shannon entropies H are exactly calculable from the rank-ordered distributions r_i . In the last section we will show, how our method can be extended also to estimate any mean value of statistical variables.

2 Species ordered distributions and rank-ordered distributions

Assume we draw a sample from a system of N different species which are equally distributed $p_1 = p_2 = \cdots =$ $p_N = 1/N$. Symbols with upper index * such as p_i^* denote observed quantities in a finite sample of size M. Obviously, if M is large enough the relative frequencies approach the probabilities, $p_1^* \to p_1, p_2^* \to p_2 \dots, p_N^* \to p_N$ due to Eq. (1). Figure 1 shows the observed relative frequencies for three different sample sizes M for 1000 equidistributed species with $p_1 = p_2 = \cdots = p_{1000} = 1/1000$. For this figure we produced uniformly distributed random integers from the interval [0, 999] and counted the occurrences of each number. As expected, with increasing sample size Mthe distribution resembles more and more the equidistribution in agreement with the true probabilities p_i . Nevertheless, the deviations of the relative frequencies from the probabilities are significant: even in the case of rather large relative sample size M/N = 1000 (Fig. 1, bottom) the deviation of the relative frequencies from the probabilities can be as large as $p_j^*/p_j \approx 1.11$. For the case M/N = 1 (top of the figure) we can see that many species have not been found at least once.

To quantify the deviations and for practical purposes that will be motivated below, we will use the data representation given in Fig. 2. Here the same data as in Fig. 1 are displayed, however, the abscissa does not show the species label *i* but the species are ordered according to the frequency of their occurrence in the sample. This means the species which occurs with the largest number of representatives appears at the first position (1) of the abscissa, the species found with the second largest frequency is labeled 2, etc. We call this representation *rank-ordered distribution of frequencies* where r_i^* is the observed relative frequency of the species at rank *i*.

Figure 2 clearly reveals that even for large relative sample size M/N the observed rank-ordered frequencies r_i^* may deviate considerably from the probabilities p_i . For smaller sample size M/N = 1 (top of the figure) about 1/3 of the species are not even found once, i.e., the observed number of species may be smaller than the true number, $N^* \leq N$.

The rank-ordered relative frequencies r_i^* form, by definition, always a decaying function. In the limit $M \to \infty$ this function approaches the rank-ordered probabilities r_i which coincide with p_i for the case of the equidistribution as well as if the probabilities p_i are decaying with increasing label number *i* of the species (see examples in the following sections). As mentioned, this limit is difficult to achieve when N is large. For $M = 10^4$ (Fig. 2, middle) we find a distribution that is far from being uniform.



Fig. 1. Observed relative frequencies of N = 1000 equidistributed species found in samples of size $M = 10^3$ (top), $M = 10^4$ (middle), and $M = 10^6$ (bottom).

Even for $M = 10^6$ the inset shows a rank-ordered distribution which deviates considerably from the equidistribution. Hence, from an observation one might erroneously conclude that the events are non-equally distributed.

The rank-ordered probabilities r_i (*i* is the rank index) contain less information than the species-ordered distribution p_i since the co-ordination species \leftrightarrow probability is lost. The problem to infer the rank-ordered probabilities r_i from a sample of size M is, therefore, a much simpler problem than to infer the species related probabilities p_i . In general, the rank-ordered distribution r_i contains about N! times less information than the species-number ordered distribution p_i , since about N! species-ordered distribu-



Fig. 2. Rank-ordered observed relative frequencies N = 1000 equidistributed species in a sample of size $M = 10^3$ (top), $M = 10^4$ (middle), and $M = 10^6$ (bottom). The inserts show the same data with higher resolution.

tions correspond the same rank-ordered distribution:

$$\{r_i, i = 1, \dots, N\} \leftarrow \begin{cases} p_1, p_2, p_3 \dots p_N \\ p_2, p_1, p_3 \dots p_N \\ p_3, p_2, p_1 \dots p_N \\ \dots \\ p_j, j = 1, \dots, N, \{j\} = \operatorname{perm}\{i\} \end{cases}$$
(5)

More precisely, the number of species-number ordered distributions is slightly smaller than the number of permutations of the species numbers N! since there might be species which occur at the same probability so that their permutation does not affect the distribution.

From these arguments we conclude that it is about N! times simpler to infer the rank-ordered probabilities r_i from the investigation of a sample of size M than the species-ordered probabilities p_i . Or, in other words, a sample of size M allows to determine the rank-ordered probabilities up to a much higher accuracy than the species-ordered probabilities.

Before coming to the main point, the estimate of probabilities from finite sample observations, it is helpful first to consider the inverse problem.

3 Predicting observed relative frequencies from a probability distribution

3.1 Equidistributed species

For the description of the species-ordered observed relative frequencies $\{p_i^*, i = 1, ..., N\}$, in general N - 1 numbers are required, whereas for the corresponding rank-ordered relative frequencies, $\{r_i^*\}$, it is sufficient to specify, how many species did not appear in our sample (this quantity will be denoted by k_0), how many species occurred with one representative (k_1) , how many with two representatives (k_2) , etc. An observed rank-ordered distribution of relative frequencies is, hence, determined by a set of occupation numbers $\{k_i, i = 0, 1, 2, ..., M\}$. In this section we describe a method to predict the observed rank-ordered relative frequencies r_i^* from a probability distribution, p_i , for finite sample size M.

The observed distribution $\{r_i^*\}$ is characterized by the cluster distribution $\{k_j\}$: the number of species that appear with j representatives each in a sample of size M. We define the probability distribution $p_c(k_i, i)$ to find *exactly* k_i species each occurring with precisely i representatives in a sample of size M. With the normalization conditions

$$\sum_{i=0}^{M} k_i = N \quad \text{(total number of species)} \tag{6}$$

$$\sum_{i=0}^{M} i k_i = M \quad \text{(total number of individuals)}, \quad (7)$$

for the case of equidistributed species this distribution reads [9] (see also [10,11,12])

$$p_{c}(k_{i},i) = \frac{M!}{N^{M}} \sum_{j=k_{i}}^{\lfloor M/i \rfloor} (-1)^{(j-k_{i})} {j \choose k_{i}} \frac{(N-j)^{(M-ji)}}{(i!)^{j} (M-ij)!},$$
(8)

where |x| denotes the integer of x.

The observable $\langle k_i \rangle$, i.e., the average number of species that occur with *i* representatives when a sample of size *M* is drawn, is the first moment of this probability distribution [10,11] $\langle k_i \rangle = \sum_{k_i} k_i p_c(k_i, i)$, where the summation is to be performed over all cluster distributions which are in agreement with Eqs. (6) and (7):

$$\langle k_i \rangle = \binom{M}{i} N^{(1-i)} \left(1 - \frac{1}{N} \right)^{(M-i)}.$$
 (9)

The occupation numbers $i = 0, 1, 2, \cdots$ are called the *i*clusters; $\langle k_0 \rangle$ is then the average size of the cluster of species which do not appear in our sample, $\langle k_1 \rangle$ is the size of the cluster of species which appear with one representative, etc.

Obviously, for small $M \ll N$, almost all of the N species which could be found in principle, belong to the 0-cluster, i.e., they do not appear in a sample of size M. As M increases the number of single occupations $\langle k_1 \rangle$ increases as well, consequently $\langle k_0 \rangle$ decays. For still growing M the number of multiple occupations becomes larger and, therefore, the sizes of the 0-cluster and 1-cluster decrease. Figure 3 shows the sizes of the first clusters, $\langle k_0 \rangle$ to $\langle k_5 \rangle$, as a function of the sample size M. The lines show the theoretical result Eq. (9) and the symbols in the top of Fig. 3 show the clusters as they have been found in numerical simulations using equally distributed random numbers.



Fig. 3. Expectation values $\langle k_i \rangle$ for cluster sizes i = 0, ..., 5 over the sample size M taken from a set of N = 1000 equidistributed species. The lines show the theoretical result Eq. (9). The symbols show the cluster sizes found by numerical simulations. The lower figure shows the same data for a larger range of the sample size M.

As a special case $\langle k_0 \rangle$ allows to determine the number of different species N^* , which are expected to be found



Fig. 4. Number of species N^* found in a sample of size M if N = 1000 species occur all with the same probability $c_i = 1/1000$. The dashed line shows the analytical result Eq. (11), the impulses show the results of a computer simulation.

in a sample of size M. This number is given by the total number of species N minus the number of species which we expect to find with zero representatives:

$$N^* = N - \langle k_0 \rangle , \qquad (10)$$

i.e.,

$$\frac{N^*}{N} = 1 - \left(1 - \frac{1}{N}\right)^M$$
 (11)

Figure 4 shows the corresponding simulation results for N = 1000. For sample size M = 5000 we notice that the average number of found species is $N^* \approx 993$, i.e., on average about 7 species are not found. For M = 8000 the average number of found species is $N^* \approx 999.67$, here we can be optimistic to have found at least one representative of all species. For practical purposes it may be useful to note that even for rather small values of N, Eq. (11) can be approximated with very good accuracy in the entire range of M by

$$\frac{N^{*\text{approx}}}{N} \approx 1 - \exp\left(-\frac{M}{N}\right) \,. \tag{12}$$

The maximal *absolute* deviation is $N^* - N^{*approx} = 1/e \approx 0.37$ which falls rapidly to $1/(2e) \approx 0.18$ as N goes to infinity.

Using Eq. (9) for the expectation values $\langle k_i \rangle$ we obtain directly the observed rank-ordered distribution of relative frequencies [12,13]:

$$r_i^* = \begin{cases} 0 & \text{for } N \ge i > N - \langle k_0 \rangle \\ 1/M & \text{for } N - \langle k_0 \rangle \ge i > N - \langle k_0 \rangle - \langle k_1 \rangle \\ & \cdots \\ i/M & \text{for } N - \sum_{s=0}^{i-1} \langle k_s \rangle \ge i > N - \sum_{s=0}^{i} \langle k_s \rangle \end{cases}$$
(13)

Using Stirling's formula to expand the expressions in Eq. (9) the analytical result Eq. (13) can be written easily in elementary functions.

Figure 5 shows rank-ordered relative frequencies calculated from a sample of random numbers (dashed lines) together with the theoretical distributions due to Eq. (13) (solid lines). (To plot more than one curve in the same figure we show the absolute frequencies Mr_i^* , i.e., what is shown are the absolute numbers of occurrence of species i in a pool of size M.) The combinatorial theory sketched above predicts the rank-ordered relative frequencies which results from an equi-probability distribution with good accuracy.



Fig. 5. Numerically determined rank-ordered relative frequencies r_i^* for samples of size M (dashed lines). The N = 1,000 species occur with equal probability, $p_i = 1/1000$. The theoretical curves due to Eq. (13) are drawn with solid lines.

The figure demonstrates that indeed we are able to predict analytically the rank-ordered observed frequencies which appear when a sample of size M is drawn from N equally distributed species. We may turn the question around and answer the question: Which sample size is required to make sure that that at least N^* out of Nspecies are observed. The required sample size is in good approximation

$$M = N \log \frac{N}{N - N^*} = -N \log \zeta . \tag{14}$$

Here ζ is the percentage of species which is admittedly is not to be represented in the sample. If we admit that about 5% are not represented we find, e.g., $M \simeq 2N$, i.e. the size of the sample should be about double the estimated species number. This estimate may be important for the planning of observations of nearly equally probable species.

3.2 General distributions

3.2.1 An alternative motivation of Eq. (9)

The derivation of Eq. (8) for the case of equidistributed species requires some algebra and in this work we will not derive a corresponding equation for general distributions. For the equidistribution the order of the rank-ordered empirical frequencies always corresponds the order of the true probabilities since all true probabilities are identical, i.e., reordering the true probabilities leads always to the equidistribution. This is different in the general case: Due to fluctuations it may happen that $p_i^* < p_j^*$ although $p_i > p_j$. The probability for the species to change ranks in the empirical distribution depends on the difference $p_i - p_i$ (the larger the difference the less probable they change ranks due to fluctuations) and on the sample size (the larger the sample size the less are the fluctuations, hence, the smaller the probability to change ranks). A comprehensive calculation must take these exchange probabilities into account.

Nevertheless, we wish to present an hypothesis which can be checked by numerical simulations. We will demonstrate that although the theoretical derivation of $\langle k_i \rangle$ for the case of a non-uniform probability distribution is somewhat simplified, the predicted results agree well with numerics.

Let us discuss an alternative motivation of Eq. (9): Assume there are N species occurring with the same probability $p_1 = p_2 = \cdots = p_N = 1/N$. The probability to find exactly *i* representatives of species *j* in a community of *M* individuals, is given by the binomial distribution

$$P_{j}(i) = \binom{M}{i} p_{j}^{i} (1 - p_{j})^{M-i} .$$
 (15)

The probability to find any species exactly i times in a community of size M is the union of species 1 appearing i times, species 2 appearing i times, etc. Since these events do not exclude each other for i < N/2 one cannot sum directly the probabilities. Instead, one has to apply the inclusion-exclusion principle [9] to subtract the intersection probabilities which in fact has been done to derive Eq. (9), see [10,11]. Let us see what happens if we ignore the intersection probabilities: The expectation value for the number of species which appear in a sample of size M exactly with i representatives reads then

$$\langle k_i \rangle = \sum_{j=1}^{N} P_j(i) = N \binom{M}{i} \left(\frac{1}{N}\right)^i \left(1 - \frac{1}{N}\right)^{M-i}$$
$$= \binom{M}{i} N^{(1-i)} \left(1 - \frac{1}{N}\right)^{M-i}, \tag{16}$$

which is identical with Eq. (9). We want to point out again that the derivation of the first moments is incomplete but it yields the correct result. In contrast to the exact derivation this simple motivation for the equidistribution has the great advantage that it can be generalized to the case of an arbitrary distribution. In general, according to Eqs. (15) and (16), the expectation value for the number of species which appear in a sample of size M exactly i times is

$$\langle k_i \rangle = \sum_{j=1}^{N} {\binom{M}{i}} p_j^i (1-p_j)^{M-i} .$$
 (17)

It can be easily checked that this distribution has the correct normalization imposed by Eqs. (6) and (7).

Having always in mind that we have no rigorous proof for the correctness of this result yet, we want to check its validity by numerical simulations.

3.2.2 Example: step-wise equidistribution

We wish to demonstrate the application of Eq. (17) using a step-wise equidistribution of N = 102 species. Let us assume for the probabilities:

$$r_{i} = p_{i} = \begin{cases} p_{\alpha} = 15/(8N) & \text{for} \quad 1 \le i \le N/3\\ p_{\beta} = 6/(8N) & \text{for} \quad N/3 < i \le 2N/3\\ p_{\gamma} = 3/(8N) & \text{for} \quad 2N/3 < i \le N \,. \end{cases}$$
(18)

The normalization can be checked easily: $\sum_{i=1}^{N} p_i = 1$. For these probabilities of the species we obtain from Eq. (17)

The expected rank-ordered empirical relative frequencies, r_i^* , can be found from Eq. (13) in the same way as previously: $\langle k_0 \rangle$ is the number of species which on average will not be found in a sample of size M, $\langle k_1 \rangle$ is the number of species which will appear with one representative, $\langle k_2 \rangle$ species are with two representatives each, etc. and finally $\langle k_M \rangle$ is the number of species which are expected to be found with M representatives. Obviously, no species can appear with more than M representatives since our sample is of size M. To generate the rank-ordered observed relative frequencies we notice that on average these values jump from (i+1)/N to i/N at rank-positions $N - \sum_{s=0}^{i-1} \langle k_i \rangle$. Hence, the expected empirical relative frequencies are

$$r_{i}^{*} = \begin{cases} 0 & \text{for} & N \ge i > N - \langle k_{0} \rangle \\ 1/M & \text{for} & N - \langle k_{0} \rangle \ge i > N - \langle k_{0} \rangle - \langle k_{1} \rangle \\ 2/M & \text{for} & N - \langle k_{0} \rangle - \langle k_{1} \rangle \ge i > \\ & > N - \langle k_{0} \rangle - \langle k_{1} \rangle - \langle k_{2} \rangle \\ & \dots \\ 1 & \text{for} & N - \sum_{s=0}^{M} \langle k_{s} \rangle \ge i > 0 \end{cases}$$

$$(20)$$

Note that, in general, the average cluster sizes $\langle k_s \rangle$ and, therefore, i are not integers. To check this formula, in Fig. 6 we show the true probability distribution due to Eq. (18) (dashed lines), the prediction of the observed relative frequencies due to Eq. (20) (solid lines) and the results of a Monte Carlo simulation (circles), where the data have been averaged over 100 independent drawings of random numbers. The analytical and numerical results agree with good accuracy.

3.2.3 Example: exponential distribution

As a second example we wish to check the validity of Eq. (17) by means of a (shifted) exponential probability distribution

$$r_i = p_i = \frac{\alpha}{1 - \exp(\alpha N)} \exp(-\alpha i) \tag{21}$$

with $0 \le i \le N$, i.e., $\int_{i=0}^{N} p_i = 1$. From Eq. (17) we obtain

$$\langle k_i \rangle = \frac{M!}{i!(M-i)!} \left(\frac{1-r_i}{1-r_i^N}\right)^i \times \\ \times \sum_{j=1}^N r_i^{i(j-1)} \left(1 - \frac{1-r_i}{1-r_i^N} r_i^{j-1}\right)^{M-i} .$$
(22)

Figure 7 shows the theoretical predictions together with results of numerical simulations. Again, theory agrees well with the numerical results. Due to the excellent agreement of the Eq. (22) with numerics we conclude that Eq. (17) allows to predict the observed relative frequencies, provided the true probability distribution is given.

4 Inferring probabilities from experiments

4.1 How to determine probabilities?

In strict sense, probabilities cannot be determined by experiments for an obvious reason: even for a fair die it is mathematically possible, although not very probable, to cast the die 1000 times and to find 1000 times the six. From such a measurement, of course, one would hardly conclude that the die is fair, i.e., that the sides one to six appear with equal probability. Hence, we have to require that the measurement is *representative*. The strict definition of this term is not easy since for M = 10 both measured sequences, 5-5-3-4-2-6-5-6-1-3 and 6-6-6-6-6-6-6-6-6-6-6 occur with equal probability. The great advantage of working with rank-ordered measurements is that the order of the measured sequence 6-5-6-1-3 would lead to the same measured rankordered frequencies as 2 - 6 - 6 - 3 - 4 - 5 - 5 - 3 - 1 or 5-5-5-3-3-6-6-1-2-4. In this sense, a rank-ordered measurement from a sequence 5-5-3-4-2-6-5-6-1-3represents many more possible measured configurations than 6 - 6 - 6 - 6 - 6 - 6 - 6 - 6 - 6 - 6. Similar as in statistical mechanics for the derivation of the canonical distribution we will call a measurement *representative* if



Fig. 6. Rank-ordered relative frequencies of step-wise equidistributed species due to Eq. (18) for sample sizes M = 50,000, M = 5,000, M = 500, and M = 100 (top to bottom). The dashed lines show the probabilities due to Eq. (18), the full lines show the predicted relative frequencies due to Eq. (20) and the circles show the results of a Monte Carlo simulation.



Fig. 7. Rank-ordered relative frequencies for exponentially distributed species as defined by Eq. (21) with $\alpha = 0.05$, N = 100and M = 200 (top), $\alpha = 0.05$, N = 200, M = 100 (middle), $\alpha = 1.0$, N = 100, M = 500 (bottom). The theoretical results (solid lines) are due to Eqs. (20) and (22), the circles show the distribution of a single set of M random numbers from the interval $[1 \dots N]$ drawn due to the distribution Eq. (21), and the dashed lines show averages over 100 such experiments.

there are many equivalent measurements (permutations) which all belong to the same rank-ordered sequence.

In strict mathematical sense we have to repeat the experiment of drawing a sample of size M an infinite number of times in order to get an averaged and representative set of rank-ordered frequencies. If we, however, had all these measurements the method presented in this article would turn out to be meaningless since for an infinite set of measurements the observed probabilities approximate the true ones, see Eq. (1). Following the same argumenta-

tion, in order to measure the pressure of air in a room we would also need an infinite set of measurements since there is a non-zero probability (although never observed under common conditions) that all air molecules are located in one half of the room and our manometer would show the double pressure or zero, depending on which half of the room is populated. Therefore, there is not much difference between measuring the pressure of air and inferring probabilities from measurements: in both cases one relies on the fact that a *representative* measurement is, by definition, a very probable one.

4.2 Optimization of cluster distributions

Equation (17) allows to predict in a systematic way the expectation values of the clusters sizes $\langle k_i \rangle$, $i = 0 \dots M$, provided the probabilities p_i , $i = 1 \dots N$, are known. Note that the average cluster sizes $\langle k_i \rangle$ based on the species-ordered probabilities p_i are identical with those based on the rank-ordered probabilities r_i . We can say then that Eq. (17) states the relation between the observed rank-ordered frequencies r_i^* and the rank-ordered probabilities r_i . This relation permits to infer the rank-ordered probabilities from a set of observed frequencies r_i^* .

In this section we propose a variational method to estimate the distribution $\{r_i, i = 1, ..., N\}$ from data of a measurement.

Consider a set of experimentally determined cluster sizes k_i^{exp} , i = 1...M. This set can be determined by counting, how many species in a sample of size M appeared with one individual in the sample (k_1^{exp}) , with two individuals (k_2^{exp}) , etc. We assume further that the set of experimentally determined relative frequencies (and, therefore, cluster sizes) is *representative* in the sense as discussed in Section 4.1, i.e., we assume that there exists a (unknown) probability distribution which leads to the *averaged* cluster sizes $\langle k_1 \rangle \approx k_1^{\text{exp}}$, $\langle k_2 \rangle \approx k_2^{\text{exp}}$, etc.

Equation (17) establishing the relation between the probabilities r_i and the averaged cluster sizes $\langle k_i \rangle$ allows to construct a variational scheme. This is done by constructing the dimensionless objective function

$$\psi_{(k)} \equiv \sum_{i=0}^{M} \left(\langle k_i \rangle - k_i^{\exp} \right)^2.$$
(23)

and requiring that it is minimal for the (unknown) set of probabilities r_i . The index (k) of $\psi_{(k)}$ indicates that the objective function refers to the cluster distribution.

Starting from a trial initial set of rank-ordered probabilities r_i , e.g. the equidistribution, and an initial trial number of species N, e.g. the observed number of species N^* (implying that $k_0^{exp} = 0$), the probabilities can be approximated numerically by a gradient method

$$r_i := r_i - \epsilon \frac{\partial \psi_{(k)}}{\partial r_i}, \qquad i = 1, \dots, N, \qquad (24)$$

with ϵ being a small number. Using Eqs. (17) and (15) we obtain

$$\frac{\partial \psi_{(k)}}{\partial r_i} = \sum_{j=0}^M \frac{\partial \psi_{(k)}}{\partial \langle k_j \rangle} \frac{\partial \langle k_j \rangle}{\partial r_i} \qquad (25)$$

$$= 2 \sum_{j=0}^M \left[\left(\langle k_j \rangle - k_j^{\exp} \right) \left(\frac{j}{r_i} - \frac{M-j}{1-r_i} \right) P_i(j) \right].$$

The so modified rank-ordered probabilities \boldsymbol{r}_i have to be normalized

$$\sum_{i=1}^{N} r_i = \sum_{i=1}^{N} p_i = 1.$$
(26)

Equation (24) and subsequent normalization has to be applied till convergence of $\psi_{(k)}(N)$ is achieved. Of course, the initial value N might not be the true number of different species, i.e., on top of the r_i for fixed N we have to optimize the value of N itself. This can be done by performing a sequence of minimizations for different values of N ranging from the observed value N^* till some N_{\max} . The result of this set of minimizations is the function $\psi_{(k)}(N)$, which has to be minimum for the most probable value of N.

For several examples we have been able to determine the probabilities r_i up to good accuracy. This method has, however, a drawback: for the case of rather large sample size M, when the observed relative frequencies approximate the probabilities, we expect that it is simpler to infer probabilities from observed frequencies. Instead, for increasing M it becomes more and more difficult since the cluster sizes $\langle k_i \rangle$ become small. This can be seen, e.g., from Fig. 2: in the upper figure for $M = 10^3$ the typical size of the clusters is $k_i \sim 100$, whereas in the lower figure drawn for $M = 10^6$ the typical size is $k_i \sim 1$. Therefore, the larger the sample size the larger become the fluctuations of the measured cluster sizes k_i^{exp} and the expression in Eq. (24) becomes ill defined. Considering that a typical cluster size is given by the number of observed different species N^* divided by the sample size M, the described method is useful when $N^*/M > 1$. In this case it yields reliable results.

4.3 Direct optimization of the probabilities

To overcome the mentioned problem, the second proposed algorithm deals directly with the rank-ordered distribution r_i instead of the cluster sizes k_i . This method is very similar to a Monte Carlo simulation in which the function to minimize is the deviation between the predicted rank-ordered frequencies and the experimentally observed frequencies.

Given an experimentally determined set of frequencies M_i^{\exp} $(i = 1..N^{\exp})$, e.g., $M_1^{\exp} = 25$, $M_2^{\exp} = 15$, $M_3^{\exp} = 110$, etc., with N^{\exp} being the observed number of different species in the sample, the following algorithm determines approximately the probabilities:

1. Determine the total number of individuals in the sample $M = \sum_{i=1}^{N^{exp}} M_i^{exp}$.

- 2. Order the frequencies according to their rank, i.e., $r_1^{exp} = accepted$ if $\psi_r(N + \Delta N) \leq \psi_{(r)}(N)$, otherwise it is re-110, $r_2^{exp} = 25$, $r_3^{exp} = 15$, etc. 3. Determine a trial initial value of the total number of
- species N, for example by means of Eq. (11), i.e., determine the initial value of N with the assumption that the (unknown) probabilities are identical.
- 4. Initialize the trial rank-ordered probabilities which are to be determined, for example, with $r_i = 1/N$.
- 5. Predict the rank-ordered observed relative frequencies $r_i^*, i = 1, \ldots, N$ which are expected to be found when drawing a sample of M individuals according to the trial probabilities. This can be done by two different methods, either by
 - (i) calculating the expected cluster distribution due to Eq. (17) and then the rank-ordered frequencies via Eq. (20),
 - (ii) or by the following procedure
 - (a) draw M random numbers from the interval [1, N] with probabilities r_i using a Metropolis algorithm,
 - (b) count the occurrences of the numbers $1, 2, \ldots, N$ and sort these frequencies due to their rank,
 - (c) repeat steps (a) and (b) a number of times, e.g. 10, and average the rank-ordered distributions. (Note that it is essential, first to rank-order and then to average.)
- 6. Determine the deviation of the experimentally observed rank-ordered frequencies $\{r_i^{exp}\}$ and the predicted rankordered frequencies $\{r_i^*\}$

$$\psi_{(r)} = \sum_{i=1}^{N} |r_i^{\exp} - r_i^*| . \qquad (27)$$

The index (r) indicates that $\psi_{(r)}$ is computed based on the frequencies r_i .¹

- 7. Modify the probabilities r_i in order to minimize $\psi_{(r)}$.
- 8. proceed with item 5 until the deviation $\psi_{(r)}$ is sufficiently small or until no further progress can be achieved.

The critical step is item 7 when the probabilities are modified. This has been done either in a deterministic way similar to Eq. (24), or by proposing a Monte Carlo trial movement in the rank-ordered frequencies $r_i \to r_i + \Delta r_i$, with Δr_i being some random number and subsequent normalization. This change is accepted if $\psi_{(r)}(r_i + \Delta r_i) \leq$ $\psi_{(r)}(r_i)$, otherwise it is rejected. Both methods (i) and (ii) yield very similar results. In this step the value of the total number of species N has to be also modified: after ttrial movements of the frequencies we propose a trial number of species $N \to N + \Delta N$, with ΔN being some random number such that N keeps smaller than the initial N corresponding to a uniform distribution. This movement is

jected. Alternatively the optimization could be performed for several values of N, as proposed in Sec. 4.2, in order to find the minimum of the function $\psi_{(r)}(N)$.

We wish to demonstrate the performance of this algorithm by an example. Using the step-wise probability distribution given by Eq. (18) we have drawn two samples of sample sizes M = 2000 and M = 5000, respectively. The according rank-ordered frequencies $r_i^{\exp}M$ are shown in Fig. 8 (upper plot, solid lines). These values serve as input (experimental data) to our algorithm, i.e., we apply the algorithm to re-infer the true step-wise probability distribution from these samples. Applying the algo-



Fig. 8. Top, solid lines: rank-ordered normalized frequencies r_i^{exp} generated from the step-wise probability distribution Eq. (18) (data scaled by M). The upper curve corresponds to the sample size M = 5000, the lower one to M = 2000. These curves serve as input to our algorithm. Top, dashed lines: corresponding expected observed probabilities r_i^* for sample sizes M = 5000 and M = 2000, respectively, as generated from the optimized set of probabilities. Bottom: solid line: original probability distribution given by Eq. (18). Dashed curve: rankordered probabilities as inferred by the described optimization algorithm based on a sample of size M = 5000. Dot-dashed curve: same but for M = 2000.

rithm to these data we obtain the results shown in the lower part of Fig. 8. Using the larger data set M = 5000(dashed line) we reproduced the original function (solid line) up to a good accuracy. Given the significant defor-

Formally Eq. (27) is not perfectly correct. Since the indices *i* in the distribution $\{r_i^*\}$ are no integers (see above), $\chi_{(r)}$ has to be computed as the integral difference between two functions with i being the integration variable. Since the precise mathematical notation appears to be cumbersome without contributing to deeper understanding we leave Eq. (27) in its present form.

mation of the measured frequencies shown in the upper part of the figure, the quality of the result surprises. Even for M = 2000 where in the upper part of the figure (lower dashed line) the three-step function can hardly be recognized, the agreement of the numerical result of the optimization procedure (lower plot, dot-dashed line) and the original set of probabilities (solid line) is agreeable.

Figure 9 shows the deviation of the input frequency distributions from the frequency distributions which have been generated from the optimized probability distribution, according to Eq. (27). After about 100,000 optimiza-



Fig. 9. Deviation of the input frequency distributions from the frequency distributions which have been generated from the optimized probability distribution as defined by Eq. (27) over the number of iteration cycles. The upper line shows the deviation for M = 5000, the lower for M = 2000.

tion loops the result does not improve anymore. We expect that at this level the accuracy of the approximation $k_i^{\text{exp}} \approx \langle k_i \rangle$ is reached.

5 Discussion and perspectives

In this article we propose a method to reconstruct the true probability distribution of species from a set of frequency distributions obtained from a small sample. This method relies on the fact that the observed rank-ordered distribution of probabilities r_i^* for a finite sample of size M can be predicted from the true rank-ordered distribution r_i . Although we have not given any mathematical strict proof of the theoretical expression of r_i^* , we have shown that this method gives quantitatively good results for the cases studied.

As mentioned, the rank-ordered distribution contains less information than the species-ordered distribution: the identity of the species is lost. Nevertheless, many statistical quantities are invariant with respect to the species labeling. Any quantity defined as the sum over all species of a function of their probability is insensitive to the order of the species. The Shannon entropy, for example, can be written as

$$H = -\sum_{i,\text{allspecies}} p_i \log p_i = -\sum_{i,\text{allranks}} r_i \log r_i \,.$$
(28)

Equation (20) allows for the prediction of the value of the *observed entropy*, defined by

$$H^* = -\sum_{i=1}^{N^*} r_i^* \log r_i^* = -\sum_{i=1}^{M} k_i^{\exp} \frac{i}{M} \log \frac{i}{M}.$$
 (29)

This quantity is experimentally accessible and serves in practical applications as a measure for deviations from the equidistribution. Since even for a perfect equidistribution of the species H^* deviates from the entropy $H = \log N$ due to finite size effects as shown in Sec. 2 empirical entropies which are based on different sample size M cannot be compared directly with each other. The method proposed here enables us to subdivide the deviations $H^* - H$ into a part due to the finite sample size M,

$$-\sum_{i=1}^{M} \langle k_i \rangle \, \frac{i}{M} \log \frac{i}{M} \,, \tag{30}$$

where the $\langle k_i \rangle$ are given by Eq. (9), and a part which is related to the true deviations from the equiprobability distribution. This way we can compare also distributions which are based on different sample sizes. In the same way we can also quantify sample-size independent deviations from any other distribution if we compute the expected cluster sizes in the expression (30) due to Eq. (17).

Another question which can be solved using the methods developed here is the evaluation of mean values of fluctuating quantities

$$\langle A \rangle = \sum_{i=1}^{N} A_i p_i \,. \tag{31}$$

This quantity may be determined by the following procedure: We introduce first the set $a_i = A_i p_i$ and the corresponding observed set

$$a_i^* = \frac{a_i M_i}{M} \,. \tag{32}$$

The set of the rank ordered numbers a_i and, therefore, $\langle A \rangle$ may be determined from the observed quantities a_i^* in precisely the same way as shown for the probabilities in this paper. We believe that this new method to estimate mean values may have many interesting applications.

The described correction algorithms have been applied to some biological relevant examples such as the spatial distribution of point mutations in genes [14].

Acknowledgment

The authors are grateful to Jan Freund for discussion.

References

 P. Grassberger. Finite sample corrections to entropy and dimension estimates. *Physics Letters A*, 128:369–373, 1988.

- A. O. Schmitt and H. Herzel. Estimating the entropy of DNA sequences. J. Theor. Bio., 188:369–377, 1997.
- A. O. Schmitt, H. Herzel, and W. Ebeling. A new method to calculate higher order entropies from finite samples. *Europhys. Lett.*, 28:303–309, 1993.
- H. Herzel, A. O. Schmitt, and W. Ebeling. Finite sample effects in sequence analysis. *Chaos, Solitons & Fractals*, 4:97–113, 1994.
- T. Pöschel, W. Ebeling, and H. Rosé. Guessing probability distributions from small samples. J. Stat. Phys., 80:1443– 1452, 1995.
- W. Ebeling and T. Pöschel. Entropy and long range correlations in literary English. *Europhys. Lett.*, 26:241–246, 1993.
- W. Ebeling, T. Pöschel, and K.-F. Albrecht. Entropy, transinformation and word distribution of informationcarrying sequences. *Int. J. Bifurcation and Chaos*, 5:51–61, 1995.
- A. Renyi and L. Vekerdi. *Probability Theory*. North-Holland, Amsterdam, 1970.
- J. N. Johnson and S. Kotz. Urn Models and Their Application. Wiley & Sons, New York, 1977.
- J. Freund and T. Pöschel. A statistical approach to vehicular traffic. *Physica A*, 219:95–114, 1995.
- T. Pöschel and J. Freund. Cluster statistics and traffic on a lattice. In L. Schimansky-Geier and T. Pöschel, editors, *Stochastic Dynamics*, volume 484 of *Lecture Notes in Physics*, pages 220–231, Berlin, Heidelberg, New York, 1998. Springer.
- T. Pöschel and J. Freund. Finite-sample frequency distributions originating from an equiprobability distribution. *Phys. Rev. E*, 66:026103, 2002.
- T. Pöschel and J. Freund. How to decide whether small samples comply with an equidistribution. *Biosystems*, 69:63–72, 2003.
- 14. T. Pöschel, C. Frömmel, and C. Gille. Online tool for the discrimination of equi-distributions. *BMC Bioinformatics* (in press), 2003. http://bioinf.charite.de/equifreq/.