

# How does a protein find its site on DNA?

Michael Slutsky<sup>1</sup> and Leonid A. Mirny<sup>1,2</sup>

<sup>1</sup>*Department of Physics, Massachusetts Institute of Technology,  
77 Massachusetts Avenue, Cambridge, MA 02139, USA*

<sup>2</sup>*Harvard-MIT Division of Health Sciences and Technology,  
Massachusetts Institute of Technology, 77 Massachusetts Avenue, Cambridge, MA 02139, USA*

Recognition and binding of specific sites on DNA by proteins is central for many cellular functions such as transcription, replication, and recombination. In the process of recognition, a protein rapidly searches for its specific site on a long DNA molecule and then strongly binds this site. Here we aim to find a mechanism that can provide both a fast search (1-10 sec) and high stability of the specific protein-DNA complex ( $K_d = 10^{-15} - 10^{-8}$  M).

Earlier studies have suggested that rapid search involves the sliding of a protein along the DNA. We consider sliding as a one-dimensional diffusion in rough (random) energy landscape formed by the energy of binding to consecutive sites on the DNA. Our analytical results and computer simulations demonstrate that, in spite of the landscape's roughness, rapid search can be achieved by a combination of 1D and 3D diffusion. We estimate the range of the specific and non-specific DNA-binding energy required for rapid search. We also establish that, paradoxically, realistic energy functions cannot provide both rapid search and strong binding. To reconcile these two fundamental requirements we propose a search-and-recognition mechanism that involves the coupling of protein binding and protein folding.

Our mechanism is supported by experimental evidence and known structures of protein-DNA complexes.

## I. INTRODUCTION

The complex transcription machinery of cells is primarily regulated by a set of proteins, *transcription factors* (TFs), that bind DNA at specific sites. Every TF can have from one to several dozens of such specific sites on the DNA. Upon binding to the site, TF forms a stable protein-DNA complex that can either activate or repress transcription of nearby genes, depending on the actual control mechanism. Fast and reliable regulation of gene expression requires (1) fast ( $\sim 1-10$  sec) search and recognition of the specific site (referred to as the *target* or *cognate* site below) out of  $10^6 - 10^9$  possible sites on the DNA, and (2) stability of the protein-DNA complex ( $K_d = 10^{-15} - 10^{-8}$  M). In spite of its apparent simplicity, such a mechanism is not understood in depth, either qualitatively or quantitatively. Here we focus on a simpler case of bacterial TFs recognizing their cognate (target) sites on the naked DNA. Needless to say that eukaryotic protein-DNA recognition is significantly complicated by chromatin packing of the DNA and multi-subunit structure of TFs.

Vast amounts of experimental data available these days provide the structures of protein-DNA complexes at atomic resolution in crystals and in solution [1, 2, 3, 4, 5], binding constants for dozens of native and hundreds of mutated proteins [6, 7], calorimetry measurements [8], and novel single-molecule experiments [9]. These experimental data contributed most significantly to our present understanding of protein-DNA interaction since the early work of von Hippel, Berg *et. al.* In a series of pioneering articles [10, 11, 12, 13], they have created a conceptual basis for describing of both the kinetics and thermodynamics of protein-DNA interaction, which became

a starting point for practically every subsequent theoretical work on the subject.

We start by reviewing the history of the problem and describing the paradox of the "faster than diffusion" association rate. Next, we present the classical model of protein-DNA "sliding" and explain how this model can resolve the paradox. We outline the problem that the sliding mechanism faces if the energetics of protein-DNA interactions are taken into account. Next we introduce our novel quantitative formalism and undertake in-depth exploration of possible mechanisms of protein-DNA interaction.

### A. "Faster than diffusion" search

The problem of how a protein finds its target site on DNA has a long history. In 1970, Riggs *et. al.* [14, 15] measured the association rate of LacI repressor and its operator on DNA as  $\sim 10^{10} \text{ M}^{-1}\text{s}^{-1}$ . This astonishingly high rate (as compared to other biological binding rates) was shown to be much higher than the maximal rate achievable by 3D diffusion. In fact, if a protein binds its site by 3D diffusion, it has to hit the right site on the DNA within  $b = 0.34$  nm. (A shift by 0.34 nm would result in binding a site that is different from the native one by 1bp. Such a site can be very different, e.g. GCGCAATT vs CGCAATTC). Using the Debye-Smoluchowski equation for the *maximal* rate of a bimolecular reaction (see e. g. [16, 17, 18]), with a protein diffusion coefficient of  $D_{3d} \sim 10^{-7} \text{ cm}^2/\text{s}$  [19] we get

$$k_{DS} = 4\pi D_{3d} b \sim 10^8 \text{ M}^{-1}\text{s}^{-1} \quad (1)$$

This value for the association rate, relevant for *in vitro* measurements, corresponds to target location *in vivo* on a time scale of a few seconds, when each cell contains up to several tens of TF molecules.

To resolve the discrepancy between the experimentally measured rate of  $10^{10} \text{ M}^{-1}\text{s}^{-1}$  and the maximal rate of  $10^8 \text{ M}^{-1}\text{s}^{-1}$  allowed by diffusion, Riggs et. al., Richter et. al. [16] and later Winter, Berg and von Hippel [10, 12] suggested that the dimensionality of the problem changes during the search process. They concluded that while searching for its target site, the protein periodically scans the DNA by “sliding” along it.

## B. Sliding along the DNA

If a protein performs both 3D and 1D diffusion, then the total search process can be considered as a 3D search followed by binding DNA and a round of 1D diffusion. Upon dissociation from the DNA, the protein continues 3D diffusion until it binds DNA in a different place, and so on. Some experimental evidence supports this search mechanism. These include affinity of the DNA-binding proteins for any fragment of DNA (non-specific binding), single molecule experiments where 1D diffusion has been observed and visualized, and numerous other experiments where the rate of specific binding to the target site has been significantly increased by lengthening non-specific DNA surrounding the site [20]. What are the benefits and the mechanism of 1D diffusion and what limits the search rate?

Here we address this question and consider possible search mechanisms that involve both 1D and 3D diffusion, where 1D diffusion along the DNA proceeds along the rough energy landscape. Quantitative analysis of the search process brought us to the following four main results:

- When the roughness of the binding energy landscape is greater than  $\sim 2k_B T$ , the diffusion along the DNA becomes extremely slow with the protein unable to diffuse more than a few base-pairs. The total search process is prohibitively slow.
- If the search proceeds by a combination of 1D and 3D diffusion, non-specific binding to the DNA plays a very important role in controlling the balance between these two processes. The optimal energy of non-specific binding can provide the maximal search rate. Although faster than either 3D or 1D search alone, optimal combination of 3D and 1D diffusion cannot expedite the search if the roughness of the landscape is greater than  $\sim 2k_B T$ .
- Experimentally observed and biologically relevant rates of search can be reached only when 1D sliding proceeds through a fairly smooth landscape with a roughness of the order of  $k_B T$ .

- Paradoxically, the stability of the protein-DNA complex at the target site requires a roughness of the binding energy landscape considerably larger than  $k_B T$ . Rapid search, however, by 1D/3D diffusion is impossible at such roughness.

Finally, we formulate this “search speed–stability” paradox and suggest a search-and-recognition mechanism that can resolve it. The paradox can be resolved if the DNA-binding protein has two distinct (conformational) states in which it exhibits two modes of binding. In the first, weaker binding mode, it searches for its site, while in the second recognition mode, the protein tightly binds DNA sites. Correlation between the energy landscapes in the two modes controls the frequency of transition between the two modes and provides effective pre-selection of low-energy sites. We also quantify the requirements of this two-mode mechanism to provide *both* rapid search and stability. Structures of known DNA-binding protein are known to be flexible and have been reported to exhibit two or more distinct binding modes. This two-state mechanism also agrees well with the results of calorimetric experiments.

## II. THE MODEL

### A. Search time

In our model, the search process consists of  $N$  rounds of 1D search (each takes time of  $\tau_{1d,i}$ ,  $i = 1..N$ ) separated by rounds of 3D diffusion ( $\tau_{3d,i}$ ). The total search time  $t_s$  is the sum of the times of individual search rounds:

$$t_s = \sum_{i=1}^N (\tau_{1d,i} + \tau_{3d,i}). \quad (2)$$

The total number  $N$  of such rounds occurring before the target site is eventually found is very large, so it is natural to introduce probability distributions for the essentially random entities in the problem. The first obvious simplification that can be made without any loss of rigor is to replace  $\tau_{3d,i}$  by its average  $\tau_{3d}$ . Each round of 1D diffusion scans a region of  $n$  sites (where  $n$  is drawn from some distribution  $p(n)$ ). The time  $\tau_{1d}(n)$  it takes to scan  $n$  sites can be obtained from the exact form of the 1D diffusion law (see Appendix A). If, on average,  $\bar{n}$  sites are scanned in each round, then the average number of such rounds required to find the site on DNA of length  $M$  is  $N = M/\bar{n}$ . Using average values, we get a total search time of

$$t_s(\bar{n}, M) = \frac{M}{\bar{n}} [\tau_{1d}(\bar{n}) + \bar{\tau}_{3d}], \quad (3)$$

From (3) it is clear that in general,  $t_s(\bar{n}, M)$  is large for both very small and very large values of  $\bar{n}$ . In fact if  $\bar{n}$  is small, very few sites are scanned in each round of 1D search and a large number of such rounds (alternating with rounds of 3D diffusion) are required to find the

site. On the contrary, if  $\bar{n}$  is large, lots of time is spent scanning a single stretch of DNA, making the search very redundant and inefficient. An optimal value  $\bar{n}_{\text{opt}}$  should exist that provides little redundancy of 1D diffusion and a sufficiently small number of such rounds. For a given diffusion law  $\tau_{1d}(n)$ , function  $t_s(\bar{n}, M)$  can be minimized producing  $\bar{n}_{\text{opt}}$ , the optimal length of DNA to be scanned between the association and the dissociation events [41].

## B. Protein-DNA energetics

While diffusing along DNA, a TF experiences the binding potential  $U(\vec{s})$  of every site  $\vec{s}$  it encounters. The energy of protein-DNA interactions is usually divided into two parts, *specific* and *non-specific* [13, 21]

$$U_i = U(\vec{s} = s_i, \dots, s_{i+l-1}) + E_{\text{ns}}, \quad (4)$$

where  $\vec{s}$  describes a binding DNA sequence of length  $l$ . As its name suggests, the non-specific binding energy  $E_{\text{ns}}$  arises from interactions that do not depend on the DNA sequence that the TF is bound to, e. g. interactions with the phosphate backbone. The specific part of the interaction energy exhibits a very strong dependence on the actual nucleotide sequence. Here and below we use the term "energy" referred to the change in the free energy related to binding  $\Delta G_b$ . This free energy includes the entropic loss of translational and rotational degrees of freedom of the protein and amino acids' side-chains, the entropic cost of water and ion extrusion from the DNA interface, the hydrophobic effect etc.

The energy of specific protein-DNA interactions can be approximated by a weight matrix (also known as "PSSM" or "profile") where each nucleotide contributes independently to the binding energy [13]:

$$U(\vec{s} = s_i, \dots, s_{i+l-1}) = \sum_{j=1}^l \epsilon(j, s_j), \quad (5)$$

where  $s_j$  is a base-pair in position  $j$  of the site and  $\epsilon(j, x)$  is the contribution of base-pair  $x$  in position  $j$ . Most of the known weight matrices of TFs  $\epsilon(j, s_j)$  give rise to uncorrelated energies of overlapping neighboring sites, obtained by one base pair shift [21]. Figure 1 presents distributions of the sequence specific binding energy  $f(U)$  obtained for different bacterial transcription factors and all possible sites in the corresponding genome. The weight matrices for these transcription factors has been derived using a set of known binding sites and standard approximation [13, 22]. Notice that for a sufficiently long site the distribution of the binding energy of random sites (or genomic DNA) can be closely approximated (see Fig.1) by a Gaussian distribution with a certain mean  $\langle U \rangle$  and variance  $\sigma^2$ :

$$f(U_i) = \frac{1}{\sqrt{2\pi\sigma^2}} \exp \left[ -\frac{(U_i - \langle U \rangle)^2}{2\sigma^2} \right]. \quad (6)$$

We also assume independence of the energy of neighboring (though overlapping) sites. Binding energies calculated for bacterial TFs support this assumption. Other physical factors such as local DNA flexibility [23] can create a correlated energy landscape providing a different mode of diffusion that we have described in [24].




FIG. 1: Spectrum of binding energy for three different transcription factors and the Gaussian approximation (solid line).

## C. Diffusion in a random energy landscape

The whole DNA molecule can thus be mapped onto one-dimensional array of sites  $\{\vec{s}_i\}$ , each corresponding to a certain binding sequence comprising bases from the  $i$ -th to the  $(i+l-1)$ -th,  $l$  being the length of the motif (see Fig. 2). At each site, there is a probability  $p_i$  of hopping to site  $i+1$  and a probability  $q_i$  of hopping to site  $i-1$ . These probabilities depend on the specific binding energies  $U_i$  and  $U_{i\pm 1}$  at the  $i$ -th site and at the adjacent sites respectively; it is most natural to assume the regular activated transport form

$$p_i \propto e^{-\beta(U_{i+1}-U_i)}, \quad q_i \propto e^{-\beta(U_{i-1}-U_i)}, \quad (7)$$

where  $\beta \equiv (k_B T)^{-1}$ ,  $k_B$  is the Boltzmann constant and  $T$  is the ambient temperature. Having defined that, we have a one-dimensional random walk with position-dependent hopping probabilities.

As has been shown in numerous papers throughout the last two decades, the properties of 1D random walks can vary dramatically depending on the actual choice of probabilities  $\{p_i\}$  (for review, see e. g. [25]). Here we employ the mean first-passage time formalism [26] to derive the diffusion law  $\tau_{1d}(\bar{n})$  for protein sliding along the DNA given the sequence-dependent binding energy. (7).

## III. RESULTS

Using the model described above, we studied the following problems:




FIG. 2: The Model Potential.

- How fast is the 1D search on DNA as a function of the “roughness”  $\sigma$  of the binding energy landscape?
- How significant is the role of non-specific binding energy  $E_{ns}$  in determining the search time?
- How fast is the search for the native site under conditions that provide stability to the protein-DNA complex at the target site?

### A. Diffusion along the DNA

We state here the main results without a derivation (which can be found in the Appendix A). For a given set of probabilities  $\{p_i\}$ , the mean first-passage time (MFPT) from  $i = 0$  to  $i = L$  (in terms of number of steps) is [26]

$$\bar{t}_{0,L} = L + \sum_{k=0}^{L-1} \alpha_k + \sum_{k=0}^{L-2} \sum_{i=k+1}^{L-1} (1 + \alpha_k) \prod_{j=k+1}^i \alpha_j, \quad (8)$$

where  $\alpha_i \equiv q_i/p_i$ . The relation (8) gives the MFPT for one given realization of probabilities. Assuming that the specific binding energies  $\{U_i\}$  have a normal distribution with variance  $\sigma^2$  (see above), we plug the probabilities in (7) into (8) and after a somewhat lengthy but straightforward calculation, we obtain an expression for the MFPT averaged over genomic sequences for  $L \gg 1$ :

$$\langle \bar{t}_{FP}(L) \rangle \simeq \tau_0 L^2 e^{11\beta^2\sigma^2/4} (1 + \beta^2\sigma^2/2)^{-1/2}, \quad (9)$$

where  $\tau_0$  is the reciprocal of the effective “attempt frequency” for hopping to a neighboring site.

The main result is that the 1D search by hopping to neighboring sites proceeds by normal diffusion with  $t \sim L^2/D_{1d}$ , where the diffusion coefficient

$$D_{1d}(\sigma) \simeq \frac{1}{2\tau_0} \left(1 + \frac{\beta^2\sigma^2}{2}\right)^{1/2} e^{-11\beta^2\sigma^2/4} \quad (10)$$

exhibits an exponential dependence on the “roughness” of the binding energy landscape  $\sigma$ , dropping rapidly as  $\sigma$  becomes greater than few  $k_B T$  [24]. Hence, rapid diffusion of a protein along the DNA is possible only if the roughness of the binding energy landscape is small compared to  $k_B T$ . ( $\beta\sigma < 1$ ). This requirement imposes strong constraints on the allowed energy of specific binding interactions.

### B. Optimal time of 3D/1D search

When 1D scanning is combined with 3D diffusion, what is the optimal time a protein has to spend in each of the two regimes? To answer this question we compute the optimal number of sites the protein has to scan by 1D diffusion in order to get the fastest overall search. Results of this section are rather general and are not limited to the particular scenario of slow 1D diffusion on a rough landscape discussed above.

Each time the protein binds DNA it performs a round of 1D diffusion. If the round lasts  $\tau_{1d}$  then on average the protein scans  $\bar{n} = \sqrt{\frac{16}{\pi} D_{1d} \tau_{1d}}$  bps [27]. By plugging this relation into Eq. (3) for search time  $t_s$ , and minimizing  $t_s$  with respect to  $\bar{n}$ , we get the optimal total search time and the optimal number of sites to be scanned in each round:

$$t_s^{\text{opt}} = t_s(\bar{n}_{\text{opt}}) = \frac{M}{2} \sqrt{\frac{\pi \tau_{3d}}{D_{1d}}} \quad \bar{n}_{\text{opt}} = \sqrt{\frac{16}{\pi} D_{1d} \tau_{3d}} \quad (11)$$

This analysis brings us to the following conclusions.

First, and most importantly, we obtain that in the *optimal* regime of search

$$\tau_{1d}(\bar{n}_{\text{opt}}) = \tau_{3d}, \quad (12)$$

i. e. the protein spends equal amounts of time diffusing along non-specific DNA and diffusing in the solution. This striking result is very general, not limited to any particular regime of 1D or 3D diffusion, and is true irrespective of the values of diffusion coefficients  $D_{1d}$  or  $D_{3d}$ , or size of the genome  $M$ . In fact it follows directly from the diffusion law  $\bar{n} \sim \sqrt{\tau_{1d}}$ . More importantly this central result can be verified experimentally by either single-molecule techniques or by traditional methods.

Also note that the optimal region of the DNA scanned in a single round of 1D diffusion  $\bar{n}_{\text{opt}}$  does not depend on  $M$ , i.e. is the same irrespective of the size of the genomes to be searched for a specific site.

Second, the optimal 1D/3D combination reached at  $\tau_{1d} = \tau_{3d}$  leads to a significant speed up of the search process. In fact, an optimal 1D/3D search is  $\bar{n}_{\text{opt}}$  times faster than a search by 3D diffusion alone, and  $(M/\bar{n}_{\text{opt}})$  times faster than a search by 1D diffusion alone. For example, if the protein operates in the optimal 1D/3D regime and scans  $\bar{n}_{\text{opt}} = 100$ bp during each round of

DNA binding, then the experimentally measured rate of binding to the specific site can be 100 times greater than the rate achievable by 3D diffusion alone.

Third, we can estimate  $\bar{n}_{opt}$ , the maximal number of sites a protein can scan in each round of 1D search. If we set  $D_{1d}$  to its maximum, i. e.  $D_{1d} \sim D_{3d}$  and  $\bar{\tau}_{3d} \sim l_d^2/D_{3d}$ , with  $l_m \sim 0.1\mu\text{m}$ , we get

$$\bar{n}_{opt}^{\max} \sim 500 \text{ bp.} \quad (13)$$

For a smaller 1D diffusion coefficient, e. g.  $D_{1d} \sim D_{3d}/100$ , we get  $\bar{n}_{opt}^{\max} \sim 50\text{bp}$ . Again, single molecule experiments can provide estimates of these quantities for different conditions of diffusion.

Finally, we obtain estimates of the shortest possible total search time. If  $M \approx 10^6$  bp and 1D diffusion is at its fastest rate, i. e.  $D_{1d} \sim D_{3d} = 10^{-7}\text{cm}^2/\text{s}$ , then using Eq. (11) we get

$$t_s^{opt} \sim \frac{M}{2} \sqrt{2\pi\bar{\tau}_{3d}\tau_0} \sim 5 \text{ sec,} \quad (14)$$

where we estimate  $\tau_0 \sim a_0^2/D_{3d} \sim 10^{-8}$  sec.

One can also estimate the search time using *in vitro* experimentally measured binding rates in water  $k_{on}^{\text{water}} \approx 10^{10}\text{M}^{-1}\text{s}^{-1}$  [14, 15]. The diffusion coefficient of a protein in the cytoplasm is 10 – 100 times lower than that in water leading to the estimated binding rate of  $k_{on}^{\text{cytoplasm}} \approx 10^8 - 10^9\text{M}^{-1}\text{s}^{-1}$  (see Appendix D). From this we obtain the time it takes for one protein to bind one site in a cell of  $1\mu\text{m}^3$  volume (i.e.  $[\text{TF}] \approx 10^{-9}\text{M}$ ) as

$$t_s^{exp} = (k_{on}^{\text{cytoplasm}}[\text{TF}])^{-1} \sim 1 - 10 \text{ sec.} \quad (15)$$

One can see perfect agreement between our theoretical estimates and experimentally measured binding rates.

As we mentioned above, there are usually several TF molecules searching in parallel for the target site. Naturally, in this case, the search is sped up proportionally to the number of molecules.

### C. Non-specific binding

While the diffusion of the TF molecules along DNA is controlled by the specific binding energy, the dissociation of the TF from the DNA depends on the total binding energy, i. e. on the non-specific binding as well as on the specific one. Moreover, since the dissociation events are much less frequent than the hopping between neighboring base-pairs (roughly by a factor of  $\bar{\tau}_{3d}/\langle\tau\rangle$ ), the non-specific energy  $E_{ns}$  makes a sensibly larger contribution to the total binding energy.

For a TF at rest bound to some DNA site  $i$ , the dissociation rate  $r_i$  would be given by the Arrhenius - type relation,

$$r_i = \frac{1}{\tau_0} e^{-\beta(E_{ns}-U_i)}. \quad (16)$$

Given the specific  $U_i$  non-specific  $E_{ns}$  energy one can calculate the average time  $\tau_{1d}$  a protein spends before dissociating from the DNA (see Appendix B). We obtain

$$E_{ns} = k_B T \left[ \ln \left( \frac{\tau_{1d}}{\tau_0} \right) - \frac{1}{2} \left( \frac{\sigma}{k_B T} \right)^2 \right], \quad (17)$$

and in the optimal regime where  $\tau_{1d} = \bar{\tau}_{3d}$

$$E_{ns}^{opt} = k_B T \left[ \ln \left( \frac{\bar{\tau}_{3d}}{\tau_0} \right) - \frac{1}{2} \left( \frac{\sigma}{k_B T} \right)^2 \right]. \quad (18)$$

### D. The parameter space

Since for a given value of  $\sigma$ , the non-specific binding controls the dissociation rate, the search time will deviate from the optimum if  $E_{ns}$  moves from this predetermined value. In Fig. 3a we plot the search time as a function of the non-specific binding energy for different values of  $\sigma$ .

We now define the *tolerance factor*  $\zeta$  as the ratio between the acceptable value of the search time  $t_s$  and the optimal search time  $t_s^{opt}$ . Experimental data suggest  $\zeta \leq 5$ , but for the moment we allow for much larger values of  $\zeta \sim 10 - 100$  (this can be done when, for instance, there are many protein molecules searching in parallel). As we can see from Fig. 3a, for each value of  $\sigma$ , there is a range of possible values of  $E_{ns}$  such that the resulting search time is within the region of tolerance (see Appendix B). Notice the dramatic increase in the search time as  $E_{ns}$  deviates from its optimal value.

Specifying  $\zeta$ , we can define our parameter space, i. e. the values of specific and non-specific energy producing a total search time within the region of tolerance. In Fig. 3b, we consider three values of  $\zeta$ . The most relaxed requirement  $\zeta = 100$  provides a search time  $t_s \leq 500$  sec. If 100 proteins are searching for a single site, then the first one will find it after  $\sim 5$  sec, leading however to a fairly low binding rate of  $k_{on} \approx 1/500 \text{ sec} \cdot 10^9 \text{ M}^{-1} = 2 \cdot 10^6 \text{ M}^{-1}\text{s}^{-1}$  (compared to experimentally measured  $10^{10} \text{ M}^{-1}\text{s}^{-1}$ ). Importantly, in order to comply with even this most relaxed search time requirement, the characteristic strength of specific interaction must be smaller than  $\sim 1.8 k_B T$ .

These results bring us to a very important conclusion that a protein cannot find its site in biologically relevant time if the roughness of the specific binding landscape is greater than  $\sim 2 k_B T$ . Although an optimal 1D/3D combination can speed up the search, it cannot overcome the slowdown of 1D diffusion. Only fairly smooth landscapes ( $\sigma < 1k_B T$ ) can be effectively navigated by proteins.

## IV. SPEED VERSUS STABILITY

To check the applicability of the above considerations, we simulated one - dimensional diffusion of *PurR* transcription factor on the *E. coli* chromosome.




FIG. 3: (a) Dependence of the search time on the non-specific binding energy. (b) The parameter space. The dashed line corresponds to optimal parameters  $\sigma$  and  $E_{ns}$  connected by Eq. (18).

The specific energy profile was built using a weight matrix derived from 35 *PurR* binding sites following a standard procedure described elsewhere ([13], [22]). The resulting energy profile is random and uncorrelated and has a standard deviation  $\sigma \simeq 6.5 k_B T$ . This profile was used as an input for calculating mean first passage time at different temperatures [42]. The result of these calculations is presented in Fig. 4a. It is clear that when the temperature becomes lower than  $\sim \sigma$ , the diffusion proceeds extremely slowly so a natural requirement for sufficiently fast diffusion is, as before,  $\sigma \leq k_B T$ .

In Fig. 4b, we present the probability  $P_b$  of binding the strongest target site with energy  $U_{\min} = U_0$  [21]. In equilibrium,  $P_b$  equals the fraction of time the protein spends at the target site:

$$P_b = \frac{\exp[-\beta U_0]}{\sum_{i=0} \exp[-\beta U_i]}. \quad (19)$$

Since the target site is not separated from the rest of the distribution by a significant energy gap,  $P_b$  is comparable to 1 (which is the natural requirement for a good regulatory site) only at  $\sigma$  *much greater than*  $k_B T$ . Recalling that a protein can effectively search the target site *at*  $\sigma$  *smaller than*  $1 - 2k_B T$ , this brings us to the central result that *the ability to translocate rapidly along the DNA clearly cannot comply with the stability requirement.*

Requirement of high stability at the target site  $P_b \sim 1$  (or  $P_b \sim 1/N_p$  if  $N_p$  copies of the protein are present) yields an estimate for the minimal  $\sigma$ ,

$$\sigma \sim k_B T \sqrt{2 \ln M} \simeq 5 k_B T, \quad (20)$$

given a genome size  $M = 10^6$ .

From the above analysis, an obvious conflict arises: *the same energy landscape cannot allow for both rapid translocation and high stability of states formed at sites with the lowest energy.* This conflict is similar to the speed-stability paradox of protein folding formulated by

Gutin *et al.* [28]: rapid search in conformation space requires a smooth energy landscape, but then the native state is unstable. In protein folding, this conflict is resolved by the presence of a large energy gap between the native state and the rest of the conformations [29, 30].

As evident from Figure 1, no such energy gap separates cognate sites from the bulk of other (random) sites. In fact, the energy function in the form of (5) cannot, in principle, provide a significant energy gap. Increasing the number of TFs cannot resolve the paradox either (see Appendix D,E). An alternative solution must be sought.

## V. THE TWO-MODE MODEL

The “search speed - stability” paradox has already been qualitatively anticipated by Winter, Berg and von Hippel [11], who therefore concluded that a conformational change of some sort should exist that would allow fast switching between “specific” and “non-specific” modes of binding. In the non-specific mode, the protein is “sliding” over an essentially equipotential surface (in our terms,  $\sigma_{\text{non-spec}} = 0$ ) whereas site-binding takes place in the “specific” mode ( $\sigma_{\text{spec}} \gg k_B T$ ). A protein in the non-specific binding mode is “unaware” of the DNA sequence it is bound to. Thus, it should permanently alternate between the binding modes, probing the underlying sites for specificity.

This model naturally rises a question about the nature of the conformation change. Originally, it was described as a “microscopic” binding of the protein to the DNA accompanied by water and ion extrusion. However, numerous calorimetry measurements and calculations [8] show that such a transition is usually accompanied by a large heat capacity change  $\Delta C$ . This  $\Delta C$  cannot be accounted for, unless additional degrees of freedom, namely, protein folding, are taken into account. On-site folding of the transcription factor may involve significant structural




FIG. 4: (a) The mean first passage time vs travelling distance for *purR* transcription factor at different temperatures. The horizontal line indicates the optimal regime,  $\tau_{1d} \sim \bar{\tau}_{3d}$ . (b) Binding probability of the strongest site as a function of temperature.

change [17, 18] and take a time of  $\sim 10^{-6}$  sec (compared to a characteristic on-site time of  $\tau_0 \sim 10^{-8}$  sec).

If the TF is to probe every site for specificity in this fashion, it would take hours to locate the native site. We note, however, that if there was a way to probe only a very limited set of sites, i. e. only those having high potential for specificity, the search time would be dramatically reduced. From the previous section it is clear that a relatively weak site-specific interaction (i.e. smooth landscape,  $\sigma \sim k_B T$ ) does not affect significantly the diffusive properties of the DNA and the total search time. If this landscape, however, is correlated with the actual specific binding energy landscape (with  $\sigma \sim 5 - 6 k_B T$ ), the specific sites will be the strongest ones in both modes. The protein conformational changes should occur therefore mainly at these sites, which constitute “traps” in the smooth landscape. Since such sites constitute a very small fraction of the total number of sites, the transitions between the modes are very rare.

We therefore suggest that there are two modes of protein-DNA binding: the *search* mode and the *recognition* mode. In the search mode, the protein conformation is such that it allows only a relatively weak site-specific interaction ( $\sigma_s \sim 1.0..1.5k_B T$ ). In the recognition mode, the protein is in its final conformation and interacts very strongly ( $\sigma_r \geq 5 k_B T$ ) with the DNA. The two energy profiles are very strongly correlated so that the lowest lying energy levels in the search mode ( $\leq -5 k_B T$ ) are likely to correspond to the strongest sites in the recognition mode, putatively, the specific ones. The transitions between the two modes happen mainly when the protein is trapped at a low-energy site of the search landscape. In this fashion, the 1D diffusion coefficient  $D_{1d}$  is about 10–100 times smaller than the ideal limit, but the search time in the optimal regime is reduced only by a factor of  $\sim 3 - 10$  (see Eq. (11)).

The coupling between the conformational change and association at a site with a low-energy trap is likely

to take place through time conditioning. Namely, *the folding occurs only if the protein spends some minimal amount of time bound to a certain site*. This statement is basically equivalent to saying that the free energy barrier that the protein must overcome to transform to the final state must be comparable to the characteristic energy difference that controls hopping to the neighboring sites.

The protein conformation in recognition mode should be stabilized by additional protein-DNA interactions. If these interactions are unfavorable, the folded structure is destabilized, then the search conformation is rapidly restored and the diffusion proceeds as before. If the new interactions are favorable, the folded structure is stable and the protein is trapped at the site for a very long time.

The proposed mechanism of specific site location (see Fig. 5) is akin to kinetic proofreading [31], which is a very general concept for a broad class of high-specificity biochemical reactions. The required specificity is achieved in kinetic proofreading through formation of an intermediate metastable complex that paves the way for irreversible enzymatic reaction. If the reaction is much slower than the life-time of the complex, then substrates that spend enough time in the complex are subject to the enzymatic reaction, while substrates that form short-lived complexes are released back to the solvent before the reaction takes place. In other words, the substrates are selected by kinetic partitioning.

In contrast to kinetic proofreading that increases equilibrium specificity for the price of energy consumption, the search-and-recognition model doesn’t require any additional source of energy. The two mode search-and-recognition model provides faster “on-rate” of binding while keeping the equilibrium binding constant unchanged. Naturally, the “off-rate” is increased as well. This makes our two-mode model thermodynamically “neutral”.

The idea of coupling between local folding and site




FIG. 5: Cartoon demonstrating the two-mode search-and-recognition mechanism. Top: search mode, bottom: recognition mode (a) two conformations of the protein bound to DNA: partially unfolded (top) and fully folded (bottom). (b) The binding energy landscape experienced by the protein in the corresponding conformations. (c) The spectrum of the binding energy determining stability of the protein in the corresponding conformations.

binding has been around for some time and was recently reassessed in the much broader context of intrinsically unstructured proteins [32, 33, 34]. Heat capacity measurements on a vast variety of protein-DNA complexes report a large negative heat capacity change in site-specific recognition, which is a clear indication of a phase transition. These measurements supplemented by X-ray crystallography and NMR structural data were interpreted by Spolar *et. al.* [8] mainly in terms of hydrophobic and conformational contributions to entropy. Thus, folding - binding coupling is now considered a well established effect for a large set of transcription factors.

However, real-time kinetic measurements were not performed until recently, so that the question of the actual mechanism was left open. Serious advances in this direction were made by Kalodimos *et. al.* [35, 36], who observed a two-step site recognition by *Lac* repressor. The deciphered NMR data unambiguously demonstrates site pre-selection by  $\alpha$ -helices bound in the major groove followed by folding of hinge helices that bind to the minor groove elements and complete the specific site recognition. Though the experiments in this field were performed with a single model system, their implications are likely to have a general character.

It should be mentioned, that no transition of this kind is observed when the protein is unbound from DNA. A possible reason for this can be a significant reduction of the free energy barrier for folding, entropic in essence, that accompanies protein-DNA association. Entropy barrier reduction is a natural consequence of relative anchoring of the various parts of the protein on the DNA “scaffold”. Thermal fluctuations that the as-

sociated protein is subject to are generally of the order of  $\sim k_B T$ , and their main effect is protein translocation along the DNA. From the above analysis, it follows that the translocation actually takes place only if the protein encounters barriers of  $\sigma_s \sim k_B T$  on its way. In a large enough collection of sites ( $M \gg 1$ ), however, potential “wells” of depth  $\sim \sigma_s \sqrt{2 \ln M}$  will be present. If the “well” depth is larger than the folding barrier height, the probability of on-site (“in-well”) folding increases, leading eventually to a stable complex formation. More detailed computational analysis of coupling between folding and binding will be published elsewhere.

## VI. CONCLUSIONS

We have developed a quantitative picture of protein-DNA interaction that provides an insight into the mechanism of fast target site location. To reconcile it with the requirement of the specific complex stability, we propose a two-step site recognition mechanism based on specific site pre-selection coupled with protein folding. The mechanism agrees qualitatively with the results of recent structural experiments.

The presence of a corrugated search-mode energy landscape appears to be crucial for the robust functioning of transcription factors. It creates a distribution of waiting times that spans three orders of magnitude and allows conformational changes in the protein at the sites corresponding to the extrema of the distribution.

The natural question therefore arises of the actual form of this landscape. Whereas much is known about the



stability and structure of specific complexes, the experimental data on non-specific complexes, especially their structure, is rather scarce. Proper understanding of the entire mechanism will hardly be possible without further experimental effort in these directions.

## VII. ACKNOWLEDGMENTS

We are thankful to A. Finkelstein, M. Kardar, W. Bialek and A. van Oijen for useful discussions. LM is an Alfred P. Sloan Research Fellow.

## APPENDIX A: DIFFUSIVE PROPERTIES OF THE DNA.

The microscopic model of protein translocation along DNA described in section Model allows for exact analytical treatment. To make the description complete, we formulate the model once again. The whole DNA molecule is mapped onto one-dimensional array of sites  $\{S_i\}$ , each corresponding to a certain binding sequence comprising bases from the  $i$ -th to the  $(i+l-1)$ -th,  $l$  being the length of the sequence. At each site, there is a probability  $p_i$  to hop to site  $i+1$  and a probability  $q_i$  to hop to site  $i-1$ . These probabilities depend on the specific binding energies  $U_i$  and  $U_{i\pm 1}$  at the  $i$ -th site and at the adjacent sites respectively; it is most natural to assume the regular activated transport form

$$p_i \propto e^{-\beta(U_{i+1}-U_i)}, \quad q_i \propto e^{-\beta(U_{i-1}-U_i)}, \quad (\text{A1})$$

where  $\beta \equiv (k_B T)^{-1}$ ,  $k_B$  is the Boltzmann constant and  $T$  is the ambient temperature.

The derivation consists of two steps. First, we describe the random walk along the DNA in terms of number of steps. Next, we calculate the mean time between successive steps in a random energetic landscape which provides the time - scale for the problem. Such a decoupling, strictly speaking, does not hold when the number of steps is small, i. e. when the number of visited sites is small and the random quantities are not averaged properly. However, since we are dealing with large numbers of steps ( $\sim 10^5 - 10^6$ ) this approach is legal, which is also confirmed by numerical simulations.

### 1. The MFPT.

To derive the diffusion law, we calculate the mean first passage time (MFPT) from site #0 to site #L, defined as the mean number of steps the particle is to make in order to reach the site #L for the first time. The derivation here follows the one in [26].

Let  $P_{i,j}(n)$  denote the probability to start at site #i and reach the site #j in exactly  $n$  steps. Then, for example,

$$P_{i,i+1}(n) = p_i T_i(n-1), \quad (\text{A2})$$

where  $T_i(n)$  is defined as the probability of returning to the  $i$ -th site after  $n$  steps *without* stepping to the right of it. Now, all the paths contributing to  $T_i(n-1)$  should start with the step to the left and then reach the site #i in  $n-2$  steps, not necessarily for the first time. Thus, the probability  $T_i(n-1)$  can be written as

$$T_i(n-1) = q_i \sum_{m,l} P_{i-1,i}(m) T_i(l) \delta_{m+l,n-2}. \quad (\text{A3})$$

We now introduce generating functions

$$\tilde{P}_{i,j}(z) = \sum_{n=0}^{\infty} z^n P_{i,j}(n), \quad \tilde{T}_i(z) = \sum_{n=0}^{\infty} z^n T_i(n). \quad (\text{A4})$$

One can easily show (see e. g. [37]) that

$$\tilde{P}_{0,L}(z) = \prod_{i=0}^{L-1} \tilde{P}_{i,i+1}(z). \quad (\text{A5})$$

Knowing  $\tilde{P}_{i,i+1}(z)$ , one calculates the MFPT straightforwardly as

$$\begin{aligned} \bar{t}_{0,L} &= \frac{\sum_n n P_{0,L}(n)}{\sum_n P_{0,L}(n)} = \left[ \frac{d}{dz} \ln \tilde{P}_{0,L}(z) \right]_{z=1} \\ &= \sum_{i=0}^{L-1} \left[ \frac{d}{dz} \ln \tilde{P}_{i,i+1}(z) \right]_{z=1}. \end{aligned} \quad (\text{A6})$$

Using (A2) and (A3), we obtain the following recursion relation for  $\tilde{P}_{i,i+1}(z)$ :

$$\tilde{P}_{i,i+1}(z) = \frac{z p_i}{1 - z q_i \tilde{P}_{i-1,i}(z)}. \quad (\text{A7})$$

To solve for  $\bar{t}_{0,L}$ , we must introduce boundary conditions. Let  $p_0 = 1$ ,  $q_0 = 0$ , which is equivalent to introducing a reflecting wall at  $i = 0$ . This boundary condition clearly influences the solution for short times and distances. However, as numerical simulations and general considerations suggest, its influence relaxes quite fast, so that for longer times, the result is clearly independent of the boundary. The benefit of setting  $p_0 = 1$  becomes clear when we observe that

$$\tilde{P}_{0,1}(1) = 1 \quad \Rightarrow \quad \forall i \quad \tilde{P}_{i,i+1}(1) = 1. \quad (\text{A8})$$

Hence,

$$\bar{t}_{0,L} = \sum_{i=0}^{L-1} \tilde{P}'_{i,i+1}(1). \quad (\text{A9})$$

The recursion relation for  $\tilde{P}'_{i,i+1}(1)$  is readily obtained from (A7) :

$$\tilde{P}'_{i,i+1}(1) = \frac{1}{p_i} + \frac{q_i}{p_i} \tilde{P}'_{i-1,i}(1) = 1 + \alpha_i \left[ 1 + \tilde{P}'_{i-1,i}(1) \right], \quad (\text{A10})$$

with  $\alpha_i \equiv p_i/q_i$ . Thus, the expression for  $\bar{t}_{0,L}$  is obtained in closed form

$$\bar{t}_{0,L} = L + \sum_{k=0}^{L-1} \alpha_k + \sum_{k=0}^{L-2} \sum_{i=k+1}^{L-1} (1 + \alpha_k) \prod_{j=k+1}^i \alpha_j. \quad (\text{A11})$$

This solution expression gives MFPT in terms of a given realization of disorder producing a certain set of probabilities  $\{p_i\}$ , whereas we are interested in the behavior averaged over all realizations of disorder. Since in our case,

$$\alpha_i = \exp[\beta(U_{i+1} - U_{i-1})], \quad (\text{A12})$$

the cumulative products in (A11) reduce to either of the two forms:

$$e^{\beta(U_i - U_j)} \quad \text{or} \quad e^{\beta(U_i - U_j + U_l - U_k)}, \quad (\text{A13})$$

which after being averaged over Gaussian disorder produce factors of  $e^{\beta^2\sigma^2}$  and  $e^{2\beta^2\sigma^2}$  respectively. After the summations are carried out, the expression for MFPT becomes

$$\langle \bar{t}_{0,L} \rangle = (L-1)^2 e^{2\beta^2\sigma^2} + L(2e^{\beta^2\sigma^2} + 1) - e^{\beta^2\sigma^2}, \quad (\text{A14})$$

which for  $L \gg 1$  reduces to

$$\langle \bar{t}_{0,L} \rangle \simeq L^2 e^{2\beta^2\sigma^2}. \quad (\text{A15})$$

Thus, the diffusion law appears to be the classical one, with a renormalized diffusion coefficient.

## 2. The time constant.

Consider a particle at site  $\#i$ . The particle will eventually escape to one of the neighboring sites  $\#(i \pm 1)$ , the escape rate being

$$r_i = \frac{1}{2\tau_0} \left( e^{-\beta(U_{i+1} - U_i)} + e^{-\beta(U_i - U_{i-1})} \right), \quad (\text{A16})$$

where  $\tau_0$  is the natural time scale for the problem. The mean time between the successive steps can be calculated therefore as the average over all possible configurations of  $U_i, U_{i \pm 1}$  of the reciprocal of the escape rate, i. e.

$$\begin{aligned} \langle \tau \rangle &= \left\langle \frac{1}{r_i} \right\rangle = \\ &= \int \frac{dU_i dU_{i+1} dU_{i-1}}{r_i} f(U_i) f(U_{i+1}) f(U_{i-1}) \end{aligned} \quad (\text{A17})$$

Assuming as above Gaussian energy statistics, this integral is evaluated as follows

$$\langle \tau \rangle = \frac{\tau_0}{\pi} \frac{e^{\beta^2\sigma^2/2}}{\pi} \int_{-\infty}^{\infty} dx dy \frac{e^{-(x^2+y^2)/2}}{e^{-\beta\sigma x} + e^{-\beta\sigma y}}. \quad (\text{A18})$$

After the change of variables

$$s = \frac{1}{\sqrt{2}}(x+y), \quad t = \frac{1}{\sqrt{2}}(x-y), \quad (\text{A19})$$

the integral factorizes leading to

$$\begin{aligned} \langle \tau \rangle &= \frac{\tau_0}{2\pi} \frac{e^{\beta^2\sigma^2/2}}{\pi} \int_{-\infty}^{\infty} ds e^{-s^2/2 + \beta\sigma s/\sqrt{2}} \int_{-\infty}^{\infty} dt \frac{e^{-t^2/2}}{\cosh(\beta\sigma t/\sqrt{2})} = \frac{\tau_0}{\sqrt{2\pi}} \frac{e^{3\beta^2\sigma^2/4}}{\pi} \int_{-\infty}^{\infty} dt e^{-t^2/2 - \ln[\cosh(\beta\sigma t/\sqrt{2})]} \\ &\simeq \frac{\tau_0}{\sqrt{2\pi}} \frac{e^{3\beta^2\sigma^2/4}}{\pi} \int_{-\infty}^{\infty} dt e^{-t^2(1+\beta^2\sigma^2/2)/2} = \tau_0 e^{3\beta^2\sigma^2/4} [1 + \beta^2\sigma^2/2]^{-1/2} \end{aligned} \quad (\text{A20})$$

Now, multiplying (A15) by  $\langle \tau \rangle$ , we obtain the diffusion coefficient as

$$D_{1d}(\sigma) \simeq \frac{1}{2\tau_0} \left( 1 + \frac{\beta^2\sigma^2}{2} \right)^{1/2} e^{-11\beta^2\sigma^2/4}. \quad (\text{A21})$$

## APPENDIX B: NON-SPECIFIC ENERGY

To find how the non-specific energy  $E_{ns}$  is related to the average time  $\tau_{1d}$  protein spends scanning a single region of the DNA we use simple observation that

$$\left\langle \sum_i \tau_i r_i \right\rangle = 1 \quad \left\langle \sum_i \tau_i \right\rangle = \tau_{1d} \quad (\text{B1})$$

which states that "on average" protein dissociates ones from the region it scans.

Since some massive hopping from site to site takes place before the particle eventually dissociates, the dissociation rates and, consequently, the non-specific binding energy should satisfy the following equation

$$\begin{aligned} \left\langle \sum_i \tau_i r_i \right\rangle &= \frac{1}{\tau_0} \left\langle \sum_i \tau_i e^{-\beta(E_{ns} - U_i)} \right\rangle = \\ &= \frac{1}{\tau_0} \int_{-\infty}^{\infty} e^{-\beta(E_{ns} - U)} \tau(U) f(U) dU = 1, \end{aligned} \quad (\text{B2})$$

and this subject to a condition

$$\left\langle \sum_i \tau_i \right\rangle = \int_{-\infty}^{\infty} \tau(U) f(U) dU = \tau_{1d}. \quad (\text{B3})$$

Where  $\tau_i$  is the time TF spends at the  $i$ -th site and  $\tau_{1d}$  is the average time of 1D search to dissociation. The average lifetime  $\tau_i = \tau(U_i)$  at that site is proportional to  $\exp(-\beta U_i)$ . In this specific case, the particle usually escapes to one of the neighboring sites, and we should average over their energies. Hence, the explicit form  $\tau(U)$  as calculated from (B3) is

$$\tau(U) = \tau_{1d} e^{-\beta^2 \sigma^2 / 2} e^{-\beta U}. \quad (\text{B4})$$

Substituting this into (B2), we have

$$\frac{\tau_{1d}}{\tau_0} e^{-\frac{1}{2}\beta^2 \sigma^2 - \beta E_{\text{ns}}} = 1, \quad (\text{B5})$$

or

$$E_{\text{ns}} = k_B T \left[ \ln \left( \frac{\tau_{1d}}{\tau_0} \right) - \frac{1}{2} \left( \frac{\sigma}{k_B T} \right)^2 \right]. \quad (\text{B6})$$

Next we recall that, in the optimal regime,  $\tau_{1d} = \bar{\tau}_{3d}$ . Thus, to ensure optimal performance,  $E_{\text{ns}}$  should be equal the expression in (B6) with  $\tau_{1d}$  replaced by  $\bar{\tau}_{3d}$ :

$$E_{\text{ns}} = k_B T \left[ \ln \left( \frac{\bar{\tau}_{3d}}{\tau_0} \right) - \frac{1}{2} \left( \frac{\sigma}{k_B T} \right)^2 \right]. \quad (\text{B7})$$

The meaning of this relation is quite transparent. The logarithm gives  $E_{\text{ns}}$  in a system with zero or constant specific binding energy. The second term introduces suppression of  $E_{\text{ns}}$  due to disorder, so that the dissociation events in a system with disorder are more frequent to compensate partially for the 1D diffusion slowdown. This relation obviously holds as long as  $E_{\text{ns}} > 0$ . Negative values of  $E_{\text{ns}}$  mean simply that the non-specific interaction became overshadowed by the specific one and has no direct physical sense anymore.

Since for a given value of  $\sigma$ , the non-specific binding controls the dissociation rate, the search time will deviate from the optimum if  $E_{\text{ns}}$  moves from this predetermined value. In Fig.3a we plot the search time as a function of the non-specific binding energy for different values of  $\sigma$ .

We now define the *tolerance factor*  $\zeta$  as the ratio between the maximal acceptable value of the search time  $t_s$  and the minimal time  $t_{s0}$ . Experimental data suggest  $\zeta \leq 5$ , but we for the moment allow for much larger values of  $\zeta \sim 10 - 100$  (this can be done when, for instance, there are many protein molecules searching in parallel). As we can see from Fig.3a, for each value of  $\sigma$ , there is a range of possible values of  $E_{\text{ns}}$  such that the resulting search time is within the region of tolerance. This range is easily calculated producing the values of non-specific

energy between

$$E_{\text{ns}}^{\pm}(\sigma, \zeta) = \frac{2}{\beta} \ln \left[ \sqrt{\frac{D_{1d}(\sigma) \bar{\tau}_{3d}}{D_{1d}(0) \tau_0}} \left( \zeta \pm \sqrt{\zeta^2 - \frac{D_{1d}(0)}{D_{1d}(\sigma)}} \right) \right] - \frac{\sigma^2 \beta}{2} \quad (\text{B8})$$

### APPENDIX C: ROLE OF DNA CONFORMATION

Central parameter here is  $\tau_{3d}$ , the interval of time between a dissociation of the protein from DNA till the next binding to DNA. Exact calculation of  $\tau_{3d}$  is a very difficult task, considering the nontrivial packaging of the DNA molecule inside a bacterial cell, electrostatic effects and the inhomogeneity of the cytoplasm.

Considering the microscopic picture one can easily obtain a reasonable estimate of  $\tau_{3d}$  as a characteristic time of 3D diffusion across the nucleoid (the region of a bacterial cell to which the DNA is confined). The corresponding diffusion length depends on the conformation of the DNA molecule. Indeed, if the DNA molecule was a single homogeneous globule, there would be a single relevant length scale, which is the molecule characteristic size  $l_m$  (the gyration radius). On the other hand, as Fig. 6 shows, diffusion of a protein molecule inside a more realistic non-homogeneous multi-domain molecule involves at least one additional length scale  $l_d$ , which is a characteristic size of a domain. These two lengths may differ by a factor of  $\sim 10$  [38], making the ratio of the resulting diffusion times  $\tau_{3d}^m / \tau_{3d}^d \sim 10^2$ . In the original problem (a single protein molecule searching for a single site on the DNA), the search process is dominated by the larger time-scale, since at least few domains must be explored before the target site is located. However, there are usually about  $10^2$  TF molecules present in a cell, so it is reasonable to assume that the domains are scanned in parallel, making the inter-domain transfer processes irrelevant.

### APPENDIX D: STABILITY REQUIREMENT

In fact, it is not hard to estimate analytically the  $(\sigma/k_B T)$  ratio for a genome of length  $M$  such that the probability of binding to the lowest site is comparable to the probability of binding to the rest of the genome. , i. e. their contributions to the partition function are of the same order of magnitude. The partition sum for the Gaussian energy level statistics is.

$$\Omega = \frac{M}{\sqrt{2\pi\sigma^2}} \int_{-\infty}^{\infty} e^{-\beta U - U^2/(2\sigma^2)} dU = M e^{\beta^2 \sigma^2 / 2} \sim \exp[-\beta U_{\text{min}}] \sim \exp\left(\beta \sigma \sqrt{2 \ln M}\right) \quad (\text{D1})$$

so that for  $M = 10^6$

$$\sigma \sim k_B T \sqrt{2 \ln M} \simeq 5 k_B T. \quad (\text{D2})$$




FIG. 6: Effect of DNA conformation on the effective diffusion distance: (a) Single globule; (b) Multi-domain conformation.

Strictly speaking, for a large though finite set of energy levels, the integration limits are cut off at  $\pm\sigma\sqrt{2\ln M}$  so that for  $\beta\sigma \gg \sqrt{\ln M}$  the partition function is dominated by the lower edge of the distribution. The estimate for  $\beta\sigma$  gives therefore the crossover value between the regime of multiple-site contribution to  $\Omega$  and the regime with single-site domination [43].

If  $N_p$  proteins are searching and binding a single target site, then the probability of being occupied is given by

$$P(N_p) = 1 - (1 - P_b)^{N_p} \approx N_p P_b \quad (\text{D3})$$

where  $P_b$  is the probability of the site being occupied by a single protein (eq 19 of the paper) and approximation is for  $P_b \ll 1/N_p$ . As evident from Fig 4b, requirement of the rapid search is satisfied if  $P_b(\sigma/T \approx 1) \sim 10^{-5}$ . An unfeasible amount of  $10^4$  copies of a single TF are required to saturate such weak binding site.

#### APPENDIX E: ENERGY GAP

Large energy gap between the cognate site  $\vec{s}_c$  and the bulk of genomic sites would solve the paradox of rapid search and stability. One may seek parameters  $\epsilon(j, s)$  of the energy function

$$U(\vec{s} = s_i, \dots, s_{i+l-1}) = \sum_{j=1}^l \epsilon(j, s_j), \quad (\text{E1})$$

to maximize the energy gap by minimizing the Z-score

$$Z(\vec{s}_c) = \frac{U(\vec{s}_c) - \langle U \rangle}{\sigma}, \quad (\text{E2})$$

where averaging and variance is taken over all possible sequences of length  $l$  (or over genomic words of length  $l$ ).

It's easy to see that  $Z(\vec{s}_c)$  is minimal if

$$\epsilon^{\text{opt}}(j, s) = -\delta(s, s_{c_j}) \quad (\text{E3})$$

where  $\delta(x, y)$  is Kronecker delta. For  $K$  types of nucleotides assuming their equal frequency in genome we obtain the maximal reachable energy gap of

$$Z^{\text{min}} = -\sqrt{lK}. \quad (\text{E4})$$

For  $K = 4$  and  $l \approx 8$  we get  $Z^{\text{min}} \approx -5$ . For the genome of  $10^6 - 10^7$  bp the energy spectrum of the genomic DNA ends at  $Z \approx -5$ . While sufficient to provide stability of the bound complex (see main text), such energy gap is unable to resolve the search-stability paradox.

#### APPENDIX F: 3D DIFFUSION

The diffusion coefficient of a protein molecule in water can be estimated as [39]

$$D \simeq \frac{k_B T}{3\pi\eta d}, \quad (\text{F1})$$

where  $d$  is the diameter of the molecule and  $\eta$  is the water viscosity. Setting  $\eta \sim 10^{-2}$  g/(sec · cm) and  $d \sim 10$  nm, we obtain at room temperature

$$D \sim 10^2 \mu\text{m}^2/\text{sec}. \quad (\text{F2})$$

Diffusion coefficient measurements for GFP in *E. coli* [19] produce values of about  $1 - 10 \mu\text{m}^2/\text{sec}$ . This difference in diffusion coefficients may account for more than order of magnitude difference in the theoretically calculated and measured target location times.

[1] Luscombe, N. M., Austin, S. E., Berman, H. M., and

Thornton, J. M., *Genome Biol.* **1**, 1 (2000).

- [2] Bell, C. E. and Lewis, M., *Curr. Opin. Struct. Biol.* **11**, 19 (2001).
- [3] Bell, C. E. and Lewis, M., *Nat. Struct. Biol.* **7**, 209 (2000).
- [4] Lewis, M., Chang, G., Horton, N. C., Kercher, M. A., Pace, H. C., Schumacher, M. A., Brennan, R. G., and Lu, P., *Science* **271**, 1247 (1996).
- [5] Schumacher, M. A., Choi, K. Y., Zalkin, H., and Brennan, R. G., *Science* **266**, 763 (1994).
- [6] Takeda, Y., Sarai, A., and Rivera, V. M., *Proc. Natl. Acad. Sci. USA* **86**, 439 (1989).
- [7] Grillo, A. O., Brown, M. P., and Royer, C. A., *J. Mol. Biol.* **287**, 539 (1999).
- [8] Spolar, R. S. and Record, M. T., *Science* **263**, 777 (1994).
- [9] Shimamoto, N., *J. Biol. Chem.* **274**, 15293 (1999).
- [10] Berg, O. G., Winter, R. B., and von Hippel, P. H., *Biochemistry* **20**, 6929 (1981).
- [11] Winter, R. B., Berg, O. G., and von Hippel, P. H., *Biochemistry* **20**, 6961 (1989).
- [12] von Hippel, P. H. and Berg, O. G., *J. Biol. Chem.* **264**, 675 (1989).
- [13] Berg, O. G. and von Hippel, P. H., *J. Mol. Biol.* **193**, 723 (1987).
- [14] Riggs, A. D., Suzuki, H., and Bourgeois, S., *J. Mol. Biol.* **48**, 67 (1970).
- [15] Riggs, A. D., Bourgeois, S., and Cohn, M., *J. Mol. Biol.* **53**, 401 (1970).
- [16] Richter, P. H. and Eigen, M., *Biophys. Chem.* **2**, 255 (1974).
- [17] Flyvbjerg, H., Jülicher, F., Ormos, P., and David, F., eds., *Physics of bio-molecules and cells* (Springer-Verlag Heidelberg, 2002), vol. 75 of *Les Houches*, chap. 1.
- [18] Bruinsma, R. F., *Physica A* **313**, 211 (2002).
- [19] Elowitz, M. B., Surette, M. G., Wolf, P. E., Stock, J. B., and Leibler, S., *J. Bacteriol.* **181**, 197 (1999).
- [20] Kim, J. G., Takeda, Y., Matthews, B. W., and Anderson, W. F., *J. Mol. Biol.* **196**, 149 (1987).
- [21] Gerland, U., J. D. Moroz, and Hwa, T., *Proc. Natl. Acad. Sci. USA* **99**, 12015 (2002).
- [22] Stormo, G. D. and Fields, D. S., *Trends Biochem. Sci.* **23**, 109 (1998).
- [23] Erie, D. A., Yang, G., Schultz, H. C., and Bustamante, C., *Science* **266**, 1562 (1994).
- [24] Slutsky, M., Kardar, M., and Mirny, L.A. (2003), preprint q-bio.BM/0310008.
- [25] Bouchaud, J. P. and Georges, A., *Phys. Rep.* **195**, 127 (1990).
- [26] Murthy, K. P. N. and Kehr, K. W., *Phys. Rev. A* **40**, 2082 (1989), **41**, 1160(E) (1990).
- [27] Hughes, B. D., *Random Walks and Random Environments* (Clarendon Press, 1995).
- [28] Gutin, A., Sali, A., Abkevich, V., Karplus, M., and Shakhnovich, E.I., *J. Chem. Phys.* **108**, 6466 (1998).
- [29] Finkelstein A.V. and Ptitsyn O.B., *Protein Physics* (Academic Press, 2002).
- [30] Pande, V., Grosberg, A., and Tanaka, T., *Rev. Mod. Phys.* **72**, 259314 (2000).
- [31] Hopfield, J. J., *Proc. Natl. Acad. Sci. USA* **71**, 4135 (1974).
- [32] Wright, P. E. and Dyson, H. J., *J. Mol. Biol.* **293**, 321 (1999).
- [33] Dyson, H. J. and Wright, P. E., *Curr. Opin. Struct. Biol.* **12**, 54 (2002).
- [34] Uversky, V. N., *Protein Sci.* **11**, 739 (2002).
- [35] Kalodimos, C. G., Boelens, R., and Kaptein, R., *Nat. Struct. Biol.* **9**, 193 (2002).
- [36] Kalodimos, C. G., Folkers, G. E., Boelens, R., and Kaptein, R., *Proc. Natl. Acad. Sci. USA* **98**, 6039 (2001).
- [37] Goldhirsh, I. and Gefen, Y., *Phys. Rev. A* **33**, 2583 (1986).
- [38] Neidhardt, F. C., Curtiss, R., and Lin, E. C., eds., *Escherichia Coli and Salmonella* (ASM Press, 1996), chap. 12.
- [39] Landau L. D. and Lifshitz E. M., *Fluid Mechanics* (Butterworth-Heinemann, 1987).
- [40] Derrida, B., *Phys. Rev. B* **24**, 2613 (1981).
- [41] Naturally, we assume here that  $\tau_{1d}(\bar{n})$  grows with  $\bar{n}$  at least as  $O(\bar{n}^{1+\alpha})$ , with  $\alpha > 0$ .
- [42] Since the magnitude of the interaction is fixed, in these calculations we vary temperature rather than binding strength
- [43] In the Random Energy Model [40], the analog of this effect would be the thermodynamical freezing.