

Vibrational energy relaxation (VER) of a CD stretching mode in cytochrome c

Hiroshi FUJISAKI, Lintao BU^y and John E. STRAUB^z

Department of Chemistry, Boston University, 590 Commonwealth Ave.,
Boston, Massachusetts, 02215, USA

October 23, 2019

Abstract

We first review how to determine the rate of vibrational energy relaxation (VER) using perturbation theory. We then apply those theoretical results to the problem of VER of a CD stretching mode in the protein cytochrome c. We model cytochrome c in vacuum as a normal mode system with the lowest-order anharmonic coupling elements. We find that, for the lifetime width parameter $\Gamma = 3 \pm 30 \text{ cm}^{-1}$, the VER time is $0.2 \pm 0.3 \text{ ps}$, which agrees rather well with the previous classical calculation using the quantum correction factor method, and is consistent with spectroscopic experiments by Romesberg's group. We decompose the VER rate into separate contributions from two modes, and find that the most significant contribution, which depends on the lifetime width parameter, comes from those modes most resonant with the CD vibrational mode.

1 Introduction

Vibrational energy relaxation (VER) is fundamentally important to our understanding of chemical reaction dynamics as it influences reaction rates significantly [1]. In general, estimating VER rates for selected modes in large molecules is a challenging problem because large molecules involve many degrees of freedom and, furthermore, quantum effects cannot be ignored [2]. If we assume a weak interaction between the "system" and the surrounding "bath", however, we can derive an estimate of the VER rate through Fermi's golden rule [3, 4, 5, 6]: A VER rate is written as a Fourier transformation of a force-force correlation function. Though it is not trivial to define and justify a separation of a system and a bath, such a formulation has been successfully applied to many VER processes in liquids [7] and in proteins [8].

Here we apply such theories of VER to the problem of estimating the vibrational population relaxation time of a CD stretching mode, in short, a CD mode, in the protein cytochrome c [9]. (We will define the CD mode to be the system and the remainder of the protein to be the bath.) Recently Romesberg's group succeeded in selectively deuterating a terminal methyl group of a methionine residue in cytochrome c [10]. The resulting CD mode has a frequency $\nu_s \approx 2100 \text{ cm}^{-1}$, which is located in a transparent region of the density of states of the protein. As such, spectroscopic detection of this mode provides clear evidence of the protein dynamics, including the VER of the CD vibrational mode. Note that at room temperature ($T = 300 \text{ K}$) $h\nu_s \approx 10$ where $\beta = 1/(k_B T)$, hence quantum effects are not negligible for this mode.

^ffujisaki@bu.edu

^ybulit@bu.edu

^zstraub@bu.edu

in mitochondrial inner-membranes, chloroplasts of plants, and bacteria [11]. Its function is related to cell respiration [12], and cyt c, using its heme molecule, "delivers" an electron from cytochrome bc₁ to cytochrome oxidase (two larger proteins both embedded in a membrane). Recently it was also found that cyt c is released when apoptosis occurs [13]. In this sense, cyt c governs the "life and death" of a cell.

The heme molecule in cyt c has a large oscillator strength, and serves as a good optical probe. As a result, many spectroscopic experiments have been designed to clarify VER and the (un)folding properties of cyt c [14]. Cyt c is often employed in numerical simulations [15, 16] because a high resolution structure was obtained [17] and its simulation has become feasible. Attempts have also been made to characterize cyt c through ab initio (DFT) calculations [18, 19].

VER of the selected CD mode in the terminal methyl group of methionine (Met80) was previously addressed by Bu and Straub [9]: They used equilibrium simulations for cyt c in water with the quantum-correction factor (QCF) method [20], and predicted that the VER time for the CD mode is on the order of 0.3 ps. However, their results are approximate: The use of the QCF method is not justified a priori, and their analysis is based on a harmonic model for cyt c. To extend that previous analysis, in this work, we model cyt c in vacuum as a normal mode system and include the lowest anharmonic coupling elements. A similar analysis has been completed for another protein myoglobin by Kidera's group [21] and by Leitner's group [22]. Use of a reduced model Hamiltonian allows us to investigate the VER rate of the CD mode in cyt c more "exactly" and to move beyond the use of quantum correction factors and the harmonic approximation.

This paper is organized as follows: In Sec. 2, we derive the principle VER formula employed in our work, and mention the related Maradudin-Fein formula. In Sec. 3, we apply those theoretical results for the rate of VER to the CD mode in cyt c, and compare our results with the classical simulation by Bu and Straub, and the experiments by Romesberg's group. In Sec. 4, we provide a summary of our results, and discuss further aspects of VER processes in proteins.

2 Vibrational energy relaxation (VER)

2.1 Perturbation expansion for the interaction

We begin with the von Neumann equation for the complete system written as

$$i\hbar \frac{d}{dt} |\psi(t)\rangle = [H; |\psi(t)\rangle]; \quad (1)$$

The interaction representation for the von Neumann equation is

$$i\hbar \frac{d}{dt} |\tilde{\psi}(t)\rangle = [V(t); |\tilde{\psi}(t)\rangle]; \quad (2)$$

where

$$H = H_0 + V = H_S + H_B + V \quad (3)$$

and

$$|\tilde{\psi}(t)\rangle = e^{iH_0 t/\hbar} |\psi(t)\rangle e^{-iH_0 t/\hbar}; \quad V(t) = e^{iH_0 t/\hbar} V e^{-iH_0 t/\hbar}; \quad (4)$$

Here H_S is the system Hamiltonian representing a vibrational mode, H_B the bath Hamiltonian representing solvent or environmental degrees of freedom, and V the interaction Hamiltonian describing the coupling between the system and the bath. If we assume that V is small in some

$$\begin{aligned}
\tilde{\rho}(t) &= \rho(0) + \frac{1}{i\hbar} \int_0^t dt' [\mathcal{V}(t'); \tilde{\rho}(t')] \\
&= \rho(0) + \frac{1}{i\hbar} \int_0^t dt' [\mathcal{V}(t'); \rho(0)] + \frac{1}{(i\hbar)^2} \int_0^t dt' \int_0^{t'} dt'' [\mathcal{V}(t'); [\mathcal{V}(t''); \rho(0)]] + \dots \quad (5)
\end{aligned}$$

Let us calculate the following probability:

$$P_v(t) = \text{Tr}_v \rho_v(t) g = \text{Tr}_v \rho_v(0) e^{iH_0 t/\hbar} \tilde{\rho}(t) e^{-iH_0 t/\hbar} g \quad (6)$$

$$\rho_v(0) = \sum_B |j_0\rangle \langle j_0| e^{-H_B} = Z_B^{-1} \quad (7)$$

$$Z_B = \text{Tr}_B e^{-H_B} g \quad (8)$$

$$Z_B = \text{Tr}_B e^{-H_B} g \quad (9)$$

where the initial state is assumed to be a direct product state of $\rho_s = \sum_j |j_0\rangle \langle j_0|$ and $\rho_B = e^{-H_B} = Z_B^{-1}$. Here $|j_0\rangle$ is the vibrational eigenstate for the system Hamiltonian H_s , i.e., $H_s |j_0\rangle = E_{j_0} |j_0\rangle$. The VER rate $\Gamma_{v_0 \rightarrow v}$ may be defined as follows:

$$\Gamma_{v_0 \rightarrow v} = \lim_{t \rightarrow \infty} \frac{d}{dt} P_v(t) \quad (10)$$

Note that the results derived from this definition are equivalent to those derived from Fermi's golden rule [23]. Hence we refer to them as a Fermi's golden rule formula.

2.2 General formula for VER

First we notice that

$$P_v(t) = \text{Tr}_v \rho_v(0) e^{iH_0 t/\hbar} \tilde{\rho}(t) e^{-iH_0 t/\hbar} g = \text{Tr}_v \rho_v(0) \tilde{\rho}(t) g \quad (11)$$

as ρ_v commutes with H_0 . If we assume that $v \neq v_0$, then $\rho_v(0) = 0$. Using this fact, we obtain the lowest (second) order result

$$\begin{aligned}
P_v(t) &\approx \frac{1}{(i\hbar)^2} \int_0^t dt' \int_0^{t'} dt'' \text{Tr}_v \rho_v(0) [\mathcal{V}(t'); [\mathcal{V}(t''); \rho(0)]] g \\
&= \frac{1}{\hbar^2} \int_0^t dt' \int_0^{t'} dt'' \text{Tr}_v \rho_v(0) \mathcal{V}(t') \mathcal{V}(t'') + \text{Tr}_v \rho_v(0) \mathcal{V}(t'') \mathcal{V}(t') g \\
&= \frac{1}{\hbar^2} \int_0^t dt' \int_0^{t'} dt'' [e^{i!_{v_0 v}(t'' - t')} C(t'' - t') + e^{i!_{v_0 v}(t' - t'')} C(t' - t'')] \quad (12)
\end{aligned}$$

where

$$C(t) = \text{Tr}_B \rho_B \mathcal{V}_{v_0 v}(t) \mathcal{V}_{v v_0}(0) i \quad \text{Tr}_B \rho_B \mathcal{V}_{v_0 v}(t) \mathcal{V}_{v v_0}(0) g \quad (13)$$

$$\mathcal{V}_{v v_0}(t) = \langle v | \mathcal{V}(t) | v_0 \rangle i \quad (14)$$

$$!_{v_0 v} = (E_{v_0} - E_v) = \hbar \omega \quad (15)$$

Hence the lowest order estimate of the VER rate is given by

$$\begin{aligned}
\Gamma_{v_0 \rightarrow v} &= \lim_{t \rightarrow \infty} \frac{1}{t} \frac{d}{dt} \int_0^t dt' \int_0^{t'} dt'' [e^{i!_{v_0 v}(t'' - t')} C(t'' - t') + e^{i!_{v_0 v}(t' - t'')} C(t' - t'')] \\
&= \frac{1}{\hbar^2} \int_0^{\infty} dt e^{i!_{v_0 v} t} C(t) \quad (16)
\end{aligned}$$

If we assume that a system variable q is small in some sense, the interaction Hamiltonian is expressed as

$$V = \sum_k q_k F_k(q; p_k; g) \quad (17)$$

force applied to the system by the bath. Thus we finally obtain the following Fermi's golden rule formula [3, 4, 5, 6]

$$\Gamma_{v_0 \rightarrow v} = \frac{\hbar \omega_{v_0 v}^2}{\hbar^2} \int_0^{\infty} dt e^{i(\omega_{v_0 v} - \omega) t} \langle \mathbf{F}^*(t) \mathbf{F}(0) \rangle \quad (18)$$

where $\omega_{v_0 v} = \hbar \omega_{v_0 v} / \hbar$ and $\langle \mathbf{F}^*(t) \mathbf{F}(0) \rangle = \text{Tr}_{\mathcal{B}} \{ \rho_{\mathcal{B}} \mathbf{F}^*(t) \mathbf{F}(0) \}$. In most situations, the transition from $v_0 = 1$ to $v = 0$ is considered. In such a case, $\Gamma_{1 \rightarrow 0} = \frac{1}{\hbar} \frac{1}{2m_s \omega_s}$, where m_s is the system mass and $\omega_s = \omega_{10}$ is the system frequency in the harmonic approximation. Hence

$$\Gamma_{1 \rightarrow 0} = \frac{1}{2m_s \hbar \omega_s} \int_0^{\infty} dt e^{i\omega_s t} \langle \mathbf{F}^*(t) \mathbf{F}(0) \rangle \quad (19)$$

2.3 Use of a symmetrized auto-correlation function

It is useful to define a symmetrized force-force correlation function as [3, 4, 5, 6]

$$S(t) = \frac{1}{2} \langle \mathbf{F}^*(t) \mathbf{F}(0) + \mathbf{F}^*(0) \mathbf{F}(t) \rangle \quad (20)$$

Since $S(t)$ is real and symmetric with respect to t , $S(t) = S(-t)$, we consider it to be analogous to $S_{cl}(t)$, the classical limit of the correlation function. Hereafter we drop the tilde on F for simplicity. By half-Fourier transforming $S(t)$ with the use of the relation $\langle \mathbf{F}^*(t) \mathbf{F}(0) \rangle = \langle \mathbf{F}^*(t - i\hbar) \mathbf{F}(0) \rangle$, we have

$$\begin{aligned} \int_0^{\infty} dt e^{i\omega t} S(t) &= \frac{1}{2} \int_0^{\infty} dt e^{i\omega t} \langle \mathbf{F}^*(t) \mathbf{F}(0) \rangle + \frac{1}{2} \int_0^{\infty} dt e^{i\omega t} \langle \mathbf{F}^*(0) \mathbf{F}(t) \rangle \\ &= \frac{1}{2} \int_0^{\infty} dt e^{i\omega t} \langle \mathbf{F}^*(t) \mathbf{F}(0) \rangle + \frac{1}{2} \int_0^{\infty} dt e^{i\omega t} \langle \mathbf{F}^*(t - i\hbar) \mathbf{F}(0) \rangle \\ &= \frac{1}{2} \int_0^{\infty} dt e^{i\omega t} \langle \mathbf{F}^*(t) \mathbf{F}(0) \rangle + \frac{1}{2} e^{-\hbar\omega} \int_0^{\infty} dt e^{i\omega t} \langle \mathbf{F}^*(t) \mathbf{F}(0) \rangle \\ &= \frac{1}{2} (1 + e^{-\hbar\omega}) \int_0^{\infty} dt e^{i\omega t} \langle \mathbf{F}^*(t) \mathbf{F}(0) \rangle \end{aligned} \quad (21)$$

Taking the real parts of both sides, we have

$$\int_0^{\infty} dt \cos(\omega t) S(t) = \frac{1}{4} (1 + e^{-\hbar\omega}) \int_0^{\infty} dt e^{i\omega t} \langle \mathbf{F}^*(t) \mathbf{F}(0) \rangle \quad (22)$$

where we have used the fact that $S(-t) = S(t)$ is real and $\langle \mathbf{F}^*(t) \mathbf{F}(0) \rangle = \langle \mathbf{F}^*(0) \mathbf{F}(t) \rangle = \langle \mathbf{F}^*(-t) \mathbf{F}(0) \rangle$. Hence, Eq. (19) can be rewritten as [6]

$$\Gamma_{1 \rightarrow 0} = \frac{1}{m_s \hbar \omega_s} \frac{1}{1 + e^{-\hbar\omega_s}} \int_0^{\infty} dt \cos(\omega_s t) S(t) \quad (23)$$

Note that this expression diverges in the classical limit because $\Gamma_{1 \rightarrow 0} / \omega_s = \hbar$. According to Bader-Beme [4], to make contact with the classical limit, we introduce another VER rate as

$$\begin{aligned} \frac{1}{T_1} &= (1 - e^{-\hbar\omega_s}) \Gamma_{1 \rightarrow 0} \\ &= 2C(\omega_s; \hbar\omega_s) \int_0^{\infty} dt \cos(\omega_s t) S(t) \end{aligned} \quad (24)$$

where

$$C(\omega; \hbar\omega_s) = \frac{1}{m_s \hbar \omega_s} \frac{1 - e^{-\hbar\omega}}{1 + e^{-\hbar\omega_s}} \quad (25)$$

This is a natural quantum expression, which can be interpreted as an energy relaxation rate and be used to estimate the VER rate.

In the limit $\hbar \rightarrow 0$, Eq. 24) reduces to

$$\frac{1}{T_1^{\text{cl}}} = \frac{1}{m_s} \int_0^{\infty} dt \cos(\omega_s t) S_{\text{cl}}(t) \quad (26)$$

This result can also be derived from a classical theory of Brownian motion, and is known as the Landau-Teller-Zwanzig (LTZ) formula. Since the full quantum calculation of $S(t)$ is demanding even for the smallest molecules, the LTZ formula is often invoked to estimate VER rates in the literature [8]. However, it is not certain that such a classical calculation can accurately describe actual VER processes, particularly in the region where $\hbar \omega_s \approx 1$. When a vibrational mode with a high frequency is involved, one anticipates that quantum effects will contribute to the process of VER.

An intuitive and readily applied theoretical approach that addresses this problem is the quantum-correction factor (QCF) method [20]:

$$\frac{1}{T_1^{\text{QCF}}} = \frac{Q(\omega_s)}{\hbar \omega_s} \frac{1}{T_1^{\text{cl}}} \quad (27)$$

where the classical VER rate is "scaled" by an appropriate QCF $Q(\omega_s) = (\hbar \omega_s)$ to obtain an estimate of the VER rate that includes the quantum effects of the initial vibrational state distribution. There exist several QCFs corresponding to different VER processes, and these classic results have been unified and substantially extended by Skinner and coworkers [20]. However, a problem with the QCF method is that we do not know a priori which VER process is dominant for the system considered. Furthermore, it is possible that several VER processes may compete [24]. Hence one must be careful in the application of the QCF method, and Skinner and coworkers have provided a number of examples of how this can be accomplished.

Alternatively, one may calculate $S(t)$ itself systematically using controlled approximations. Calculating a correlation function for large systems has a long history in chemical physics [25], including recent applications to VER processes in liquid [26, 27]. The vibrational self-consistent field (VSCF) method [28] will also be useful in this respect.

On the other hand, if we approximate F as a simple function of $f_{\mathbf{k}g}; p_{\mathbf{k}g}$, we can calculate $S(t)$ rather easily and, in a sense, more "exactly." In the next section, we explore such an approach.

2.5 Approximations for the force-force correlation function

2.5.1 Taylor expansion of the force

We can formally Taylor-expand the force as a function of the bath variables $f_{\mathbf{k}g}; p_{\mathbf{k}g}$:

$$F(f_{\mathbf{k}g}; p_{\mathbf{k}g}) = \sum_{\mathbf{k}} A_{\mathbf{k}}^{(1)} q_{\mathbf{k}} + \sum_{\mathbf{k}} B_{\mathbf{k}}^{(1)} p_{\mathbf{k}} + \sum_{\mathbf{k}; \mathbf{k}^0} A_{\mathbf{k}; \mathbf{k}^0}^{(2)} q_{\mathbf{k}} q_{\mathbf{k}^0} + \sum_{\mathbf{k}; \mathbf{k}^0} B_{\mathbf{k}; \mathbf{k}^0}^{(2)} p_{\mathbf{k}} p_{\mathbf{k}^0} + \quad (28)$$

where the expansion is often truncated in the literature following the first term. Depending on the system-bath interaction considered, higher order coupling including the third and fourth terms can be relevant. For example, the fourth term appears in benzene to represent the interaction between the CH stretch and CCH wagging motion [29] through the Wilson G matrix [30]. In the case of a CD stretching mode in cyt c, as discussed below, or a CN stretching mode in water [24], the third term is relevant for VER.

If the first term $\sum_k A_k^{(1)} q_k$ is dominant in the force, then the force-force correlation function becomes

$$\langle F(t)F(0) \rangle = \sum_k A_k^{(1)} A_k^{(1)} \langle q_k(t)q_k(0) \rangle = \sum_k \frac{\hbar (A_k^{(1)})^2}{2! \omega_k} [(n_k + 1)e^{-i\omega_k t} + n_k e^{i\omega_k t}] \quad (29)$$

where we have used

$$q_k(t) = \frac{\hbar}{2! \omega_k} (a_k e^{-i\omega_k t} + a_k^y e^{i\omega_k t}) \quad (30)$$

and $\langle a_k^y a_k \rangle = n_k$ with $n_k = 1/(e^{\hbar \omega_k} - 1)$ because

$$\langle a_k^y a_k \rangle = e^{-\hbar \omega_k} = e^{-\hbar \omega_k (a_k^y a_k + 1/2)} \quad (31)$$

Here we have assumed that the bath Hamiltonian is an ensemble of harmonic oscillators:

$$H_B = \sum_k \hbar \omega_k (a_k^y a_k + 1/2) = \sum_k \left(\frac{p_k^2}{2} + \frac{1}{2} \omega_k^2 q_k^2 \right) \quad (32)$$

where ω_k is the k -th mode frequency for the bath, and

$$p_k = i \frac{\hbar \omega_k}{2} (a_k - a_k^y) \quad (33)$$

Thus we obtain

$$S(t) = \sum_k \frac{\hbar (A_k^{(1)})^2}{2! \omega_k} (2n_k + 1) \cos \omega_k t \quad (34)$$

and

$$\frac{1}{T_1} = \frac{\hbar C(\hbar \omega_s)}{\hbar C(\hbar \omega_s) D^{(1)}(\omega_s) [2n(\omega_s) + 1]} \sum_k \frac{(A_k^{(1)})^2}{\omega_k} (2n_k + 1) [(\omega_s - \omega_k) + (\omega_s + \omega_k)] \quad (35)$$

where we have assumed that $\omega_s, \omega_k > 0$ and defined $D^{(1)}(\omega) = [A^{(1)}(\omega)]^2$ as a continuum limit of $(A_k^{(1)})^2 = \omega_k$. (The contribution from the second term $\sum_k B_k^{(1)} p_k$ is calculated in the same way.)

2.5.3 Contribution from the third term

If the third term $\sum_k A_k^{(2)} q_k q_k$ is dominant in the force, then

$$\begin{aligned} \langle F(t)F(0) \rangle &= \sum_k A_k^{(2)} A_k^{(2)} \langle q_k(t)q_k(t)q_k(0)q_k(0) \rangle \\ &= R_{--}(t) + R_{++}(t) + R_{+}(t) \end{aligned} \quad (36)$$

where

$$R_{--}(t) = \frac{\hbar^2}{4} \sum_k A_k^{(2)} A_k^{(2)} \langle q_k^y q_k^y q_k^y q_k^y \rangle e^{-i(\omega_k + \omega_k)t}; \quad (37)$$

$$R_{++}(t) = \frac{\hbar^2}{4} \sum_k A_k^{(2)} A_k^{(2)} \langle q_k^y q_k^y q_k^y q_k^y \rangle e^{i(\omega_k + \omega_k)t}; \quad (38)$$

$$\begin{aligned} R_{+}(t) &= \frac{\hbar^2}{4} \sum_k A_k^{(2)} A_k^{(2)} \langle q_k^y q_k^y q_k^y q_k^y \rangle e^{i(\omega_k - \omega_k)t} \\ &+ \langle q_k^y q_k^y (a_k^y a_k + a_k a_k^y) \rangle e^{i(\omega_k - \omega_k)t} \end{aligned} \quad (39)$$

$$D_{k;k^0;k^0;k^0}^{(2)} = \frac{A_{k;k^0}^{(2)} A_{k^0;k^0}^{(2)}}{k! k^0! k^0! k^0!} : \quad (40)$$

Using the following

$$h a_k a_{k^0} a_{k^0}^y a_{k^0}^y i = (1 + n_k) (1 + n_{k^0}) (k k^0 k^0 k^0 + k k^0 k^0 k^0); \quad (41)$$

$$h a_k^y a_{k^0}^y a_{k^0} a_{k^0} i = n_k n_{k^0} (k k^0 k^0 k^0 + k k^0 k^0 k^0); \quad (42)$$

$$h a_k a_{k^0}^y (a_{k^0}^y a_{k^0} + a_{k^0} a_{k^0}^y) i = (1 + n_k) (1 + 2n_{k^0}) k k^0 k^0 k^0 + (1 + n_k) n_{k^0} (k k^0 k^0 k^0 + k k^0 k^0 k^0); \quad (43)$$

$$h a_k^y a_{k^0} (a_{k^0}^y a_{k^0} + a_{k^0} a_{k^0}^y) i = n_k (1 + 2n_{k^0}) k k^0 k^0 k^0 + n_k (1 + n_{k^0}) (k k^0 k^0 k^0 + k k^0 k^0 k^0); \quad (44)$$

we have

$$R_{k;k^0}(t) = \frac{\hbar^2 X}{2} D_{k;k^0;k^0;k^0}^{(2)} (1 + n_k) (1 + n_{k^0}) e^{i(l_k + l_{k^0})t}; \quad (45)$$

$$R_{++}(t) = \frac{\hbar^2 X}{2} D_{k;k^0;k^0;k^0}^{(2)} n_k n_{k^0} e^{i(l_k + l_{k^0})t}; \quad (46)$$

$$R_+(t) = \hbar F(0) i^2 + \hbar^2 X D_{k;k^0;k^0;k^0}^{(2)} (1 + n_k) n_{k^0} e^{i(l_k - l_{k^0})t} \quad (47)$$

where we have used $A_{k;k^0}^{(2)} = A_{k^0;k^0}^{(2)}$, and

$$\hbar F(0) i^2 = \frac{\hbar^2 X}{4} D_{k;k^0;k^0;k^0}^{(2)} (1 + 2n_k) (1 + 2n_{k^0}); \quad (48)$$

Hence we obtain

$$S_{k;k^0}(t) = \sum_{k;k^0} h^{(+)} \cos(l_k + l_{k^0})t + \sum_{k;k^0} h^{(-)} \cos(l_k - l_{k^0})t + \hbar F(0) i^2 \quad (49)$$

and

$$\frac{1}{T_1} = C(\hbar; S) \sum_{k;k^0} n^{(+)} [l_k + l_{k^0} - l_S] + \sum_{k;k^0} n^{(-)} [l_k - l_{k^0} - l_S] \quad (50)$$

where we have assumed $l_S \notin 0$ and defined

$$h^{(+)}_{k;k^0} = \frac{\hbar^2}{2} D_{k;k^0;k^0;k^0}^{(2)} (1 + n_k + n_{k^0} + 2n_k n_{k^0}); \quad (51)$$

$$h^{(-)}_{k;k^0} = \frac{\hbar^2}{2} D_{k;k^0;k^0;k^0}^{(2)} (n_k + n_{k^0} + 2n_k n_{k^0}); \quad (52)$$

Though its appearance is rather different, the formula Eq. 50) is equivalent to that derived by Kenkre, Tokmako, and Fayer [5] as well as by Shiga and Okazaki [24]. There is also a similar result known as the Maradudin-Fein formula [31]

$$W = W_{\text{decay}} + W_{\text{coll}}; \quad (53)$$

$$W_{\text{decay}} = \frac{\hbar}{2m_S l_S} \sum_{k;k^0} \frac{(A_{k;k^0}^{(2)})^2}{k! k^0!} (1 + n_k + n_{k^0}) (l_S - l_k - l_{k^0}); \quad (54)$$

$$W_{\text{coll}} = \frac{\hbar}{m_S l_S} \sum_{k;k^0} \frac{(A_{k;k^0}^{(2)})^2}{k! k^0!} (n_k + n_{k^0}) (l_S + l_k + l_{k^0}); \quad (55)$$

group [22]. As was demonstrated in [5], this formula is also equivalent to Eq. (50); in the following we make use of Eq. (50). A problem with this formula is that we cannot take its continuum limit as easily as Eq. (35). As a remedy, a width parameter related to the vibrational lifetime is usually introduced leading to a definite value for the VER rate. We will discuss this problem in Sec. 3.3.

3 Application to a CD stretching mode in cytochrome c

3.1 Definition of system and bath

We take horse heart cytochrome c (cyt c) as an example of how one may estimate the rate of VER for selected modes in proteins. We use the CHARMM program [33] to describe the force field, to minimize the structure, and to calculate the normal modes for the system. Starting from the 1HRC structure for cyt c in Protein Data Bank (PDB) [34] one hydrogen atom of the terminal methyl group of Met80 was deuterated. The energy of the protein structure was minimized in vacuum using the conjugate gradient algorithm. We diagonalized the hessian matrix (second derivatives of the potential) for that mechanically stable conformation of the protein:

$$K_{ij} = \frac{\partial^2 V_{\text{CHARMM}}}{\partial x_i \partial x_j} = \frac{1}{\sqrt{m_i m_j}} \frac{\partial^2 V_{\text{CHARMM}}}{\partial x_i \partial x_j} \quad (56)$$

where V_{CHARMM} is the CHARMM potential, and $x_i = \sqrt{m_i} x_{i1}$ are mass weighted Cartesian coordinates. The number of atoms in cyt c is 1745 (myoglobin has 2475 atoms), so the hessian matrix is 5235×5235 , and its diagonalization was readily accomplished using the vibrancy facility in CHARMM [33].

The result of this calculation was the density of states (DOS) for the system as shown in Fig. 1. The DOS consists of three regions: (1) below around 1700 cm^{-1} , (2) from around 1700 cm^{-1} to around 2800 cm^{-1} , (3) above around 2800 cm^{-1} . The first region corresponds to rotational and torsional motions of the protein, and the third to bond stretching motions such as CH bonds. The second is rather "transparent" but one can observe one mode localized around the CD bond stretching mode in Met80 with frequency 2129.1 cm^{-1} as shown in Fig. 2 (a). Hence we refer to this as a CD stretching mode, or CD mode; the dynamics of which is the focus of our study.




Figure 1: Density of states for cytochrome c in vacuum.




Figure 2: Normal modes of cytochrome c in vacuum. (a) 4357th mode (CD mode) with $\omega = 2129.1 \text{ cm}^{-1}$, (b) 3330th mode with $\omega = 1330.9 \text{ cm}^{-1}$, (c) 1996th mode with $\omega = 829.9 \text{ cm}^{-1}$. Only vectors on the terminal methyl group of Met80 in cyt c are depicted.

At this level of description, the system is an ensemble of harmonic oscillators, i.e., normal modes. Since we are interested in VER of the CD mode, we represent it as a system

$$H_S = \frac{P_{CD}^2}{2} + \frac{1}{2} \omega_{CD}^2 Q_{CD}^2 \quad (57)$$

while other degrees of freedom are treated as a bath

$$H_B = \sum_k \left(\frac{P_k^2}{2} + \frac{1}{2} \omega_k^2 Q_k^2 \right) \quad (58)$$

The interaction between the system and bath is described by the interaction Hamiltonian

$$V = H_{\text{cytc}} + H_S + H_B \quad (59)$$

where H_{cytc} is the Hamiltonian for the full cytochrome c protein. We will discuss the content of V in the following section.

3.2 Calculation of the coupling constants

As in Eq. (17), we assume that the interaction Hamiltonian is of the form

$$V = \sum_k F_k Q_k \quad (60)$$

and Taylor expand the force as Eq. (28). The first and second terms do not appear because this is a normal mode expansion, and the fourth term does not appear as the original coordinates are Cartesian coordinates. As in the first approximation, we take the force to be

$$F_k = \sum_{k',l} A_{kk'}^{(2)} Q_{k'} Q_{l'} \quad (61)$$

The coupling coefficients $A_{k,l}^{(2)}$ are calculated as

$$A_{k,l}^{(2)} = \frac{1}{2} \frac{\partial^3 V}{\partial Q_{CD} \partial Q_k \partial Q_l} \quad (62)$$

A problem arises: How does one calculate these coupling coefficients? The most direct approach is to use a finite difference method:

$$A_{k,l}^{(2)} = \frac{1}{2} \frac{V_{+++} - V_{++} - V_{+} - V_{+} + V_{+} + V_{+} + V_{+} - V}{(2 \omega_{CD}) (2 \omega_k) (2 \omega_l)} \quad (63)$$

the approximation [22, 24]:

$$A_{k;l}^{(2)}, \quad \frac{1}{2} \sum_{ij} U_{ik} U_{jl} \frac{K_{ij}(q_{CD}) - K_{ij}(q_D)}{2 q_{CD}} \quad (64)$$

where U_{ik} is an orthogonal matrix that diagonalizes the hessian matrix at the mechanically stable structure K_{ij} , and $K_{ij}(q_D)$ is a hessian matrix calculated at a shifted structure along the direction of the CD mode with a shift q_D . This expression is approximate but readily implemented using the CHARMM facility to compute the hessian matrix. A comparison between Eqs. (63) and (64) is made in Table 1. We also examined the convergence of the results by changing q_{CD} , and found that $q_{CD} = 0.02A$ is sufficient for the following calculations.

Table 1: Comparison between the finite difference method Eq. (63) and the formula Eq. (64). We have used $q_{CD} = 0.02A$, and $A_{k;l}^{(2)}$ is given in kcal/mol/ \AA^3 .

$(k;l)$	formula, Eq. (64)	Finite difference
(3330, 1996)	22.3	22.4
(3330, 4399)	-29.6	-29.5
(3327, 1996)	-5.7	-5.8
(1996, 678)	0.64	0.63

The numerical results for the coupling elements are shown in Fig. 3. The histogram for the elements is shown in Fig. 4. As one can see from these figures, most of the elements are small, while the largest coupling elements are rather large. See Table 2. Note that the combination (3330, 1996) is particularly significant for the CD mode because it approximately satisfies the resonant condition [21]:

$$j_{CD} - \omega_k - \omega_l \approx 0 \quad (A_{k;l}^{(2)}) \quad (65)$$

As shown in Fig. 2 (b), (c), these modes are localized near the terminal methyl group of Met80 as well as the CD mode. In such a case, resonant energy transfer (Fermi resonance) is expected as shown by Moritsugu-Miyashita-Kidera [21]. We have observed similar behavior in cytochrome when the CD mode was excited and the energy migration to other normal modes facilitated by resonance was followed.

Table 2: The largest coupling elements. The value of $A_{k;l}^{(2)}$ are given in kcal/mol/ \AA^3 .

$(k;l)$	$A_{k;l}^{(2)}$
(1996, 1996)	42.9
(4399, 3330)	29.6
(4622, 3170)	27.3
(3330, 1996)	22.3

3.3 Assignment of the "lifetime" parameter

We cannot directly evaluate Eq. (50) as it contains delta functions. Evaluation of this expression for a finite system like a protein leads to a null result. To circumvent this problem, we "thaw"




Figure 3: Distribution of the coupling elements. Left: $50.0 > A_{k;l}^{(2)} > 5.0$, Right: $5.0 > A_{k;l}^{(2)} > 0.5$. The value of $A_{k;l}^{(2)}$ are given in units of kcal/mol/Å³.




Figure 4: Histogram for the amplitude of the coupling elements.

$$\langle x \rangle = \frac{1}{2 + x^2} \quad (66)$$

using a width parameter γ . Physically this means that each normal mode should have a lifetime $\tau = 1/\gamma$ due to coupling to other degrees of freedom, i.e., the surrounding environment including water (or we might be able to interpret $1/\gamma$ as a time resolution). It is difficult to derive γ from first principles, so we take it to be a phenomenological parameter as in the literature [22, 32].

As a result, the VER rate, Eq. (50), for the CD mode becomes

$$\frac{1}{T_1} = C(\gamma; h\nu_S) \sum_{k;k^0} \left[\frac{2}{4} \frac{k;k^0^{(+)}}{2 + (\nu_k + \nu_{k^0} + \nu_S)^2} + \frac{k;k^0^{(+)}}{2 + (\nu_k + \nu_{k^0} + \nu_S)^2} \right. \\ \left. + \frac{k;k^0^{(-)}}{2 + (\nu_k + \nu_{k^0} + \nu_S)^2} + \frac{k;k^0^{(-)}}{2 + (\nu_k + \nu_{k^0} + \nu_S)^2} \right] \quad (67)$$

We employ this expression in our subsequent calculations.

3.4 Results

3.4.1 Classical calculation

Classical force-force correlation functions and their (cosine) Fourier transformations are shown in the left and middle column of Fig. 5 for five different trajectories. Here we have defined a function as

$$\langle C(t) \rangle = \frac{1}{m_S} S_{cl}(t); \quad (68)$$

and its (cosine) Fourier transformation as $\tilde{C}(\nu)$, i.e., $1/T_1^{cl} = \tilde{C}(\nu_S)$. Note that these data are obtained from molecular dynamics simulations of cytochrome c in water [9]. As can be seen, the correlation functions oscillate wildly, and the (cosine) Fourier transformations are messy. As such, it is difficult to extract a reliable and stable value for the VER rate.

To address this problem, we introduce the window function

$$w(t) = \exp(-\gamma|t|); \quad (69)$$

The $\langle C(t) \rangle$ functions are multiplied by this function and (cosine) Fourier transformed. This corresponds to broadening each peak of a spectrum with a Lorentzian with width γ . The results for five trajectories are shown in the right column of Fig. 5. (The width parameter is taken as $\gamma = 3 \text{ cm}^{-1}$.) The results in the right column are better behaved than those in the middle column but there still remain some fluctuations.

According to Bu and Straub simulations of cytochrome c in water [9], we take $\nu_S = 2135 \text{ cm}^{-1}$ to investigate the γ dependence of the result as shown in the left of Fig. 6. We see that $\tilde{C}(\nu_S) \approx 1:1 - 1:2 \text{ ps}^{-1}$ for $\gamma = 3 - 30 \text{ cm}^{-1}$. Since $Q(\nu_S) = (h\nu_S) \approx 2:4 - 3:0$ for two-phonon processes [9], this corresponds to a VER time of $0:3 - 0:4 \text{ ps}$ according to Eq. (27).¹

3.4.2 Quantum calculation

We use the formula Eq. (67) as a quantum mechanical estimate of the VER rate. The dependence of the result is shown on the right hand side of Fig. 6. We see that, for $\gamma = 3 - 30 \text{ cm}^{-1}$, the quantum mechanical estimate gives $T_1 \approx 0:2 - 0:3 \text{ ps}$, which is similar to the classical estimate Eq. (27): $T_1 \approx 0:3 - 0:4 \text{ ps}$.

¹In the VER calculation of myoglobin [22], Leitner's group took $\gamma = 0:5 - 10 \text{ cm}^{-1}$ to be the width, and concluded that the result is relatively insensitive to the choice of γ in this range.




Figure 5: Left: Classical data for the force-force correlation function. Middle: Fourier spectra for the correlation function. Right: The corresponding coarse-grained Fourier spectra. The lifetime width parameter $\tau = 3 \text{ cm}^{-1}$.




Figure 6: Left: Classical VER rate for five trajectories as a function of the lifetime width parameter γ . Right: VER rate calculated by Eq. (67) as a function of γ .

In Tables 3, we list the largest contributions to the VER rate for different width parameters. For the case of $\gamma = 3 \text{ cm}^{-1}$, the largest contribution is due to modes (3823,1655). This combination of modes is nearly resonant with the CD mode as $|j_{3823} + j_{1655} - j_{\text{CD}}| = 0.1 \text{ cm}^{-1}$. Though the coupling element for the combination is small ($\mathcal{A}_{3823;1655}^{(2)} < 0.05 \text{ kcal/mol/\AA}^3$), this mode combination contributes significantly to the VER rate. On the other hand, for the case of $\gamma = 30 \text{ cm}^{-1}$, the largest contribution results from the combination of modes (3330,1996). This combination is somewhat off-resonant, i.e., $|j_{3330} + j_{1996} - j_{\text{CD}}| = 32 \text{ cm}^{-1}$, but the coupling element is very large ($\mathcal{A}_{3823;1655}^{(2)} = 22.3 \text{ kcal/mol/\AA}^3$), and the contribution is significant. In both cases, one combination of modes dominates the VER rate ($\sim 20\%$) though there are non-negligible contributions from other combinations of modes.

Table 3: The largest contributions to the VER rate (in units of ps^{-1}) for $\gamma = 3 \text{ cm}^{-1}$ (left) and $\gamma = 30 \text{ cm}^{-1}$ (right).

$(k;l)$	contribution	$(k;l)$	contribution
(3823, 1655)	1.10 (19 %)	(3330, 1996)	0.88 (22 %)
(3823, 1654)	0.43 (8 %)	(3823, 1655)	0.11 (3 %)
(3822, 1655)	0.37 (6 %)	(3170, 2196)	0.07 (2 %)
(3330, 1996)	0.17 (3 %)	(1996, 1996)	0.05 (1 %)
(3822, 1654)	0.15 (3 %)	(3823, 1654)	0.04 (1 %)
(3823, 1661)	0.14 (3 %)	(3170, 2202)	0.04 (1 %)
(3822, 1661)	0.05 (1 %)	(3822, 1655)	0.04 (1 %)
(3822, 1656)	0.05 (1 %)	(3327, 1996)	0.03 (1 %)
(3823, 1658)	0.04 (1 %)	(3330, 1655)	0.02 (1 %)

We have also examined the temperature dependence of the VER rates using Eq. (67). As shown in Fig. 7, for $T < 300 \text{ K}$ there is little temperature dependence as has been addressed in the case of myoglobin [22]. Thus we can say that the relaxation of the CD mode is quantum mechanical rather than thermal because the decay at 300 K is similar to that at 0.3 K .

3.4.3 Discussion

We examine the relationship between the theoretical results described above and the corresponding experiments of Romesberg's group which has studied the spectroscopic properties of




Figure 7: Temperature dependence of the VER rate calculated by Eq. (67). The width parameter is $\gamma = 3 \text{ cm}^{-1}$.

the CD mode in cytochrome c [10]. They measured the shifts and widths of the spectra for different forms of cytochrome c; the widths of the spectra (FWHM) were found to be $\Delta_{\text{FWHM}} \approx 6.0 - 13.0 \text{ cm}^{-1}$. A rough estimate of the VER rate leads to

$$T_1 \approx 5.3 \gamma_{\text{FWHM}} \text{ (ps)} \quad (70)$$

which corresponds to $T_1 \approx 0.4 - 0.9 \text{ ps}$. This estimate is similar to the semi-classical prediction computed using Eq. (27) and appropriate QCFs (0.3 - 0.4 ps) and the perturbative quantum mechanical estimate using the reduced model (0.2 - 0.3 ps). This result appears to justify the use of QCFs and the reduced model in this situation, and suggests that the effects of the protein solvation (by water) are negligible in describing the VER of the CD mode. Of course, we must be careful in comparing the estimate derived from (70) as there may be inhomogeneity in the experimental spectra. As such, it is more desirable to calculate not only VER rates but spectroscopic observables themselves to compare with experiments.

Finally we discuss the relation between this work and previous work on carbon monoxide myoglobin (MbCO). Though there are many experimental studies on Mb [35], we focus on the experiments of Annu's group [36] and Fayer's group [37] on MbCO. The former group found that the VER time for CO in the heme pocket (photolyzed MbCO) is $\approx 600 \text{ ps}$, whereas the latter group found that the VER time for CO bounded to the heme is $\approx 20 \text{ ps}$. This difference is interpreted as follows: CO is covalently bonded to the heme for the latter case, whereas CO is "loosely" in the pocket in the former case, i.e., the force applied to CO for the latter case is stronger than that for the former case. This difference in the magnitude of the force causes the slower VER for the "loosely" CO. In this respect, the CD bond is expected to be stronger than the CO-heme coupling. This may explain a VER time $\approx 0.1 \text{ ps}$, which is similar to the VER times for the CH (CD) stretching modes in benzene (or perdeuterobenzene) [29, 38]. It will be interesting to apply a similar reduced model to the analysis of the VER of CO in MbCO.

4 Summary and further aspects

After reviewing the VER rate formula derived from quantum mechanical perturbation calculations, we applied it to the analysis of VER of a CD stretching mode in cytochrome c. We modeled cytochrome c in vacuum as a normal mode system with the third-order anharmonic coupling elements, which were calculated from the CHARMM potential. We found that, for the width parameter

classical calculation using the quantum correction factor (QCF) method, and is consistent with the experiments by Romesberg's group. This result indicates that the use of QCFs or a reduced model Hamiltonian can be justified a posteriori to describe the VER problem. We decomposed the VER rate into contributions from two modes, and found that the most significant contribution, which depends on the lifetime width parameter, results from modes most resonant with the CD mode.

Finally we note several future directions which should be studied: (a) Our results for the VER rate depend on a width parameter. Unfortunately we do not know which value is the most appropriate for. Non-equilibrium simulations (with some quantum corrections [39]) might help this situation, and are useful to investigate energy pathways or sequential IVR (intramolecular vibrational energy redistribution) [40] in a protein. (b) This work is motivated by pioneering spectroscopic experiments by Romesberg's group. The calculation of the VER rate and the linear or nonlinear response functions, related to absorption or 2D-IR (or 2D-Raman) spectra [41, 42, 43, 44], is desirable. (c) Romesberg's group investigated a spectroscopic change due to the oxidation or reduction of Fe in the heme; such an electron transfer process [45] is fundamental for the functionality of cyt c. To survey this process dynamically, it will be necessary to combine some quantum chemistry (ab initio) calculations with MD simulations [18, 46, 47].

One of the authors (H.F.) thanks Dr. J. Gong for noting Ref. [5], Prof. A. Kidera, Prof. S. Okazaki, Prof. D.M. Leitner, Prof. Y. Mizutani, Dr. T. Takami, Dr. T. Miyadera, Dr. Y. Kawashima, Dr. S. Fuchigami, Mr. H. Teramoto for useful discussions, Prof. A. Stuchebrukhov for sending reprints, and Dr. M. Shigemori for providing perl scripts used in this work.

References

- [1] B.J. Beme, M. Borkovec, and J.E. Straub, "Classical and Modern Methods in Reaction Rate Theory," *J. Phys. Chem.* 92, 3711 (1988); J.I. Steinfeld, J.E. Francisco, and W.L. Hase, *Chemical Kinetics and Dynamics*, Prentice-Hall (1989); A. Stuchebrukhov, S. Ionov, and V. Letokhov, "IR spectra of highly vibrationally excited large polyatomic molecules and intramolecular relaxation," *J. Phys. Chem.* 93, 5357 (1989); T. Uzer, "Theories of intramolecular vibrational energy transfer," *Phys. Rep.* 199, 73 (1991).
- [2] D.E. Logan and P.G. Wolynes, "Quantum localization and energy flow in many-dimensional Fermi resonant systems," *J. Chem. Phys.* 93, 4994 (1990); S.A. Schoeld and P.G. Wolynes, "A scaling perspective on quantum energy flow in molecules," *J. Chem. Phys.* 98, 1123 (1993); S.A. Schoeld, P.G. Wolynes, and R.E. Wyatt, "Computational study of many-dimensional quantum energy flow: From action diffusion to localization," *Phys. Rev. Lett.* 74, 3720 (1995); S.A. Schoeld and P.G. Wolynes, "Rate theory and quantum energy flow in molecules: Modeling the effects of anisotropic diffusion and dephasing," *J. Phys. Chem.* 99, 2753 (1995); D.M. Leitner and P.G. Wolynes, "Vibrational mixing and energy flow in polyatomics: Quantitative prediction using local random matrix theory," *J. Phys. Chem. A* 101, 541 (1997).
- [3] D.W. Oxtoby, "Vibrational population relaxation in liquids," *Adv. Chem. Phys.* 47, 487 (1981).
- [4] J.S. Bader and B.J. Beme, "Quantum and classical relaxation rates from classical simulations," *J. Chem. Phys.* 100, 8359 (1994).
- [5] V.M. Kenkre, A. Tokmako, and M.D. Fayer, "Theory of vibrational relaxation of polyatomic molecules in liquids," *J. Chem. Phys.* 101, 10618 (1994).

- J. Chem. Phys. 105, 7047 (1996).
- [7] R. Rey and J.T. Hynes, \Vibrational energy relaxation of HOD in liquid D₂O," J. Chem. Phys. 104, 2356 (1996); \Vibrational phase and energy relaxation of CN in water," *ibid.* 108, 142 (1998).
- [8] D.E. Sagnella and J.E. Straub, \A study of vibrational relaxation of B-state carbon monoxide in the heme pocket of photolyzed carboxymyoglobin," Biophys. J. 77, 70 (1999).
- [9] L. Bu and J.E. Straub, \Vibrational frequency shifts and relaxation rates for a selected vibrational mode in cytochrome c," Biophys. J. 85, 1429 (2003); Erratum, to be published.
- [10] J.K. Chin, R. Jimenez, and F. Romesberg, \Direct observation of protein vibrations by selective incorporation of spectroscopically observable carbon-deuterium bonds in cytochrome c," J. Am. Chem. Soc. 123, 2426 (2001); \Protein dynamics and cytochrome c: Correlation between ligand vibrations and redox activity," *ibid.* 124, 1846 (2002).
- [11] David Keilin, *The History of Cell Respiration and Cytochrome*, Cambridge University Press, Cambridge (1966); Richard E. Dickerson, \Cytochrome c and the evolution of energy metabolism," *Sci. Am.* 242, 136 (1980); G.W. Pettigrew and G.R. Moore, *Cytochromes c: Evolutionary, Structural, and Physicochemical Aspects*, Springer-Verlag, Berlin (1990).
- [12] B. Alberts, D. Bray, J. Lewis, M. Raff, and K. Roberts, *Molecular Biology of the Cell*, 3rd edition, Garland Pub. (1994); G. Karp, *Cell and Molecular Biology: Concepts and Experiments*, 3rd edition, Wiley Text Books (2002).
- [13] J. Yang, X. Liu, K. Bhalla, C.N. Kim, A.M. Ibrado, J. Cai, T.-I. Peng, D.P. Jones, and X. Wang, \Prevention of Apoptosis by Bcl-2: Release of Cytochrome c from Mitochondria Blocked," *Science*, 275, 1129 (1997); R.M. Kluck, E. Bossy-Wetzel, D.R. Green, D.D. Newmeyer, \The Release of Cytochrome c from Mitochondria: A Primary Site for Bcl-2 Regulation of Apoptosis," *ibid.* 275, 1132 (1997).
- [14] S.-R. Yeh, S. Han, and D.L. Rousseau, \Cytochrome c folding and unfolding: A biphasic mechanism," *Acc. Chem. Res.* 31, 727 (1998); S.W. Englander, T.R. Sosnick, L.C. Mayne, M. Shtilerman, P.X. Qi, and Y. Bai, \Fast and slow folding in cytochrome c," *ibid.* 31, 737 (1998); W.W. Wang, X. Ye, A.A. Denkov, F. Rosca, T. Sjodin, W. Cao, M. Sheeran, and P.M. Champion, \Femtosecond multicolor pump-probe spectroscopy of ferrous cytochrome c," *J. Phys. Chem. B* 104, 10789 (2000).
- [15] S.H. Northrup, M.P. Pear, J.A. McCammon, and M. Karplus, \Molecular dynamics of ferrocyclochrome c," *Nature* 286, 304 (1980); C.F. Wong, C. Zheng, J. Shen, J.A. McCammon, and P.G. Wolynes, \Cytochrome c: A molecular proving ground for computer simulations," *J. Phys. Chem.* 97, 3100 (1993); A.E. Garcia and G. Hummer, \Conformational dynamics of cytochrome c: Correlation to hydrogen exchange," *Proteins: Struct. Funct. Genet.* 36, 175 (1999); A.E. Cardenas and R. Elber, \Kinetics of cytochrome c folding: Atomically detailed simulations," *ibid.* 51, 245 (2003); X. Yu and D.M. Leitner, \A nonabundant diusion of vibrational energy in proteins," *J. Chem. Phys.* 119, 12673 (2003).
- [16] L. Bu and J.E. Straub, \Simulating vibrational energy flow in proteins: Relaxation rate and mechanism for heme cooling in cytochrome c," *J. Phys. Chem. B* 107, 12339 (2003).
- [17] G.W. Bushnell, G.V. Louie, and G.D. Breyer, \High-resolution three-dimensional structure of horse heart cytochrome c," *J. Mol. Biol.* 214, 585 (1990).

- for computational biology: First applications and perspective," *IBM J. Res. & Dev.* 45, 397 (2001).
- [19] F. Sato, T. Yoshihiro, M. Era, and H. Kashiwagi, "Calculation of all-electron wavefunction of hemoprotein cytochrome c by density functional theory," *Chem. Phys. Lett.* 341, 645 (2001).
- [20] J.L. Skinner and K. Park, "Calculating vibrational energy relaxation rates from classical molecular dynamics simulations: quantum correction factors for processes involving vibration-vibration energy transfer," *J. Phys. Chem. B* 105, 6716 (2001).
- [21] K. Moritsugu, O. Miyashita, and A. Kidera, "Vibrational Energy Transfer in a Protein Molecule," *Phys. Rev. Lett.* 85, 3970 (2000); "Temperature Dependence of Vibrational Energy Transfer in a Protein Molecule," *J. Phys. Chem. B* 107, 3309 (2003).
- [22] D.M. Leitner, "Vibrational energy transfer in helices," *Phys. Rev. Lett.* 87, 188102 (2001); X. Yu and D.M. Leitner, "Vibrational energy transfer and heat conduction in a protein," *J. Phys. Chem. B* 107, 1698 (2003).
- [23] J.J. Sakurai, *Modern Quantum Mechanics*, 2nd edition, Addison-Wesley (1994); G.C. Schatz and M.A. Ratner, *Quantum Mechanics in Chemistry*, Dover (2002).
- [24] M. Shiga and S. Okazaki, "An influence functional theory of multiphonon processes in molecular vibrational energy relaxation," *J. Chem. Phys.* 109, 3542 (1998); "Molecular dynamics study of vibrational energy relaxation of CN in H₂O and D₂O solutions: An application of path integral influence functional theory to multiphonon processes," *ibid.* 111, 5390 (1999); T. Mikami, M. Shiga and S. Okazaki, "Quantum effect of solvent on molecular vibrational energy relaxation of solute based upon path integral influence functional theory," *ibid.* 115, 9797 (2001).
- [25] W.H. Miller, "The semiclassical initial value representation: A potentially practical way for adding quantum effects to classical molecular dynamics simulations," *J. Phys. Chem. A* 105, 2942 (2001).
- [26] Q. Shi and E. Geva, "Semiclassical theory of vibrational energy relaxation in the condensed phase," *J. Phys. Chem. A* 107, 9059 (2003); "Vibrational energy relaxation in liquid oxygen from a semiclassical molecular dynamics simulation," *ibid.* 107, 9070 (2003); "On the calculation of vibrational energy relaxation rate constants from centroid molecular dynamics simulations," *J. Chem. Phys.* 119, 9030 (2003).
- [27] H. Kim and P.J. Rossky, "Evaluation of quantum correlation functions from classical data," *J. Phys. Chem. B* 106, 8240 (2002); J.A. Poulsen, G. Nymann, and P.J. Rossky, "Practical evaluation of condensed phase quantum correlation functions: A Feynman-Kleinert variational linearized path integral method," *J. Chem. Phys.* 119, 12179 (2003).
- [28] R.B. Gerber, V. Buch, and M.A. Ratner, "Time-dependent self-consistent field approximation for intramolecular energy transfer. I. Formulation and application to dissociation of van der Waals molecules," *J. Chem. Phys.* 77, 3022 (1982); A. Roitberg, R.B. Gerber, R. Elber, and M.A. Ratner, "Anharmonic Wave Functions of Proteins: Quantum Self-Consistent Field Calculations of BPTI," *Science*, 268, 1319 (1995); S.K. Gregurick, E. Fredj, R. Elber, and R.B. Gerber, "Vibrational spectroscopy of peptides and peptide-water complexes: Anharmonic coupled-mode calculations," *J. Phys. Chem. B* 101, 8595 (1997); Z. Bihary, R.B. Gerber, and V.A. Apkarian, "Vibrational self-consistent field approach to anharmonic spectroscopy of molecules in solids: Application to iodine in argon

- wirth and R.B. Gerber, "Quantum Molecular Dynamics of Ultrafast Processes in Large Polyatomic Systems," *Chem. Rev.* 99, 1583 (1999).
- [29] E.L. Sibert III, J.T. Hynes, and W.P. Reinhardt, "Classical dynamics of highly excited CH and CD overtones in benzene and perdeuterobenzene," *J. Chem. Phys.* 81, 1135 (1984); E.L. Sibert III, W.P. Reinhardt, and J.T. Hynes, "Intramolecular vibrational relaxation and spectra of CH and CD overtones in benzene and perdeuterobenzene," *ibid.* 81, 115 (1984).
- [30] E.B. Wilson, Jr., J.C. Decius, and P.C. Cross, *Molecular Vibrations: The Theory of Infrared and Raman Vibrational Spectra*, Dover (1980).
- [31] A.A. Maradudin and A.E. Fein, "Scattering of neutrons by an anharmonic crystal," *Phys. Rev.* 128, 2589 (1962).
- [32] J. Fabian and P.B. Allen, "Anharmonic Decay of Vibrational States in Amorphous Silicon," *Phys. Rev. Lett.* 77, 3839 (1996).
- [33] B.R. Brooks, R.E. Briccocoleri, B.D. Olson, D.J. States, S. Swaminathan, and M. Karplus, "CHARMM: A Program for Macromolecular Energy, Minimization, and Dynamics Calculations," *J. Comp. Chem.* 4, 187 (1983); A.D. MacKerell, Jr., B.R. Brooks, C.L. Brooks III, L.N. Nilsson, B. Roux, Y. Won, and M. Karplus, "CHARMM: The Energy Function and Its Parameterization with an Overview of the Program," in *The Encyclopedia of Computational Chemistry*, 1, 271, P.v.R. Schleyer et al., editors, John Wiley & Sons: Chichester (1998).
- [34] See, for example, <http://www.rcsb.org/pdb>.
- [35] Y. Mizutani and T. Kitagawa, "Direct observation of cooling of heme upon photodissociation of carbon monoxymyoglobin," *Science* 278, 443 (1997); "Ultrafast structural relaxation of myoglobin following photodissociation of carbon monoxide probed by time-resolved resonance Raman spectroscopy," *J. Phys. Chem. B* 105, 10992 (2001); M.D. Fayer, "Fast protein dynamics probed with infrared vibrational echo experiments," *Ann. Rev. Phys. Chem.* 52, 315 (2001); X. Ye, A. Demidov, and P.M. Champion, "Measurements of the photodissociation quantum yields of MbNO and MbO₂ and the vibrational relaxation of the six-coordinate heme species," *J. Am. Chem. Soc.* 124, 5914 (2002); F. Rosca, A.T.N. Kumar, D. Ionascu, X. Ye, A.A. Demidov, T. Sjödin, D. Wharton, D. Barrick, S.G. Sligar, T. Yonetani, and P.M. Champion, "Investigations of Anharmonic Low-Frequency Oscillations in Heme Proteins," *J. Phys. Chem. A* 106, 3540 (2002).
- [36] D.E. Sagnella, J.E. Straub, T.A. Jackson, M. Lim, and P.A. Anand, "Vibrational population relaxation of carbon monoxide in the heme pocket of photolyzed carbon monoxymyoglobin: Comparison of time-resolved mid-IR absorbance experiments and molecular dynamics simulations," *Proc. Natl. Acad. Sci. U.S.A.* 96, 14324 (1999).
- [37] J.R. Hill, D.D. Dlott, C.W. Rella, K.A. Peterson, S.M. Decatur, S.G. Boxer, and M.D. Fayer, "Vibrational dynamics of carbon monoxide at the active sites of mutant heme proteins," *J. Phys. Chem.* 100, 12100 (1996).
- [38] H.W. Schranz, "Mode to mode energy flow amongst the ring modes in benzene," *J. Mol. Struct. (Theochem)* 368, 119 (1996).
- [39] P.H. Nguyen and G. Stock, "Nonequilibrium molecular-dynamics study of the vibrational energy relaxation of peptides in water," *J. Chem. Phys.* 119, 11350 (2003).

- distribution and secular increase of relative entropy," *J. Phys. Chem. A* 102, 9454 (1998); H. Hasegawa and K. Someda, \Derivative state analysis of intramolecular vibrational energy redistribution of acetylene," *J. Chem. Phys.* 110, 11255 (1999).
- [41] K. Okumura and Y. Tanimura, \Sensitivity of two-dimensional fifth-order Raman response to the mechanism of vibrational mode-mode coupling in liquid molecules," *Chem. Phys. Lett.* 278, 175 (1997).
- [42] T. J. C. Jansen, J. G. Snijder, and K. Duppen, \The third- and fifth-order nonlinear Raman response of liquid CS₂ calculated using a finite field nonequilibrium molecular dynamics method," *J. Chem. Phys.* 113, 307 (2000); \Interaction induced effects in the nonlinear Raman response of liquid CS₂: A finite field nonequilibrium molecular dynamics approach," *ibid.* 114, 10910 (2001).
- [43] K. A. Merchant, W. G. Noid, D. E. Thompson, R. Akiyama, R. F. Loring, and M. D. Fayer, \Structural assignment and dynamics of the A₁ substates of MbCO: Spectrally resolved vibrational echo experiments and molecular dynamics simulations," *J. Phys. Chem. B* 107, 4 (2003).
- [44] J. Edler and P. Hamm, \Two-dimensional vibrational spectroscopy of the amide I band of crystalline acetanilide: Femtosecond resonance, conformational substrates, or vibrational self-trapping?" *J. Chem. Phys.* 119, 2709 (2003).
- [45] D. N. Beratan, J. N. Onuchic, J. E. Winkler, and H. B. Gray, \Electron-Tunneling Pathways in Proteins," *Science* 258, 1740 (1992); J. J. Regan, S. M. Risser, D. N. Beratan, and J. N. Onuchic, \Protein Electron Transport: Single versus Multiple Pathways," *J. Phys. Chem.* 97, 13083 (1993).
- [46] V. Gogonea, D. Suarez, A. van der Vaart, and K. M. Merz, Jr., \New development in applying quantum mechanics to proteins," *Curr. Opin. Struct. Biol.* 11, 217 (2001).
- [47] Y. Komaiji, T. Nakano, K. Fukuzawa, Y. Ueno, Y. Inadomi, T. Nemoto, M. Uebayasi, D. G. Fedorov, and K. Kitaura, \Fragment molecular orbital method: application to molecular dynamics simulation, *ab initio* FMO-MD'," *Chem. Phys. Lett.* 372, 342 (2003).