

Information Processing in Brain Microtubules

Jean Faber¹, Renato Portugal¹, Luiz Pinguelli Rosa²

¹LNCC - National Laboratory of Scientific Computing

Quantum Computing Group, Petropolis, RJ, Brazil.

{faber, portugal}@lncc.br

²Alberto Luiz Coimbra Institute for Graduate Studies and Research in Engineering,

CT, COPPE, Univeristy of Rio de Janeiro, RJ, Brazil,

pr@eletrobras.gov.br

October 28, 2019

Abstract

Models of the mind are based on the possibility of computing in brain microtubules (MT). From this point of view, information processing is the fundamental issue for understanding the brain mechanisms that produce consciousness. The cytoskeleton polymers could store and process information through their dynamic coupling mediated by mechanical energy. We analyze the problem of information transfer and storage in brain microtubules, considering them as a communication channel. We discuss the implications of assuming that consciousness is generated by the subneural process.

1 Introduction

In recent years many papers have addressed the problem of developing a theory of mind [1-13]. R. Penrose and S. Hameroff developed a quantum model of the mind considering the cytoskeleton of neurons cells as the principal component that produces states of mind or consciousness [2,3]. Their model uses the microtubule (MT) structure, the principal part of the cytoskeleton, to produce a quantum computation through tubulins. Tubulins are the units which work as cellular automata in the walls of MTs. The information propagates inside wave guides of the neron MTs.

The microtubule wall could be able to store information and make computation by using combinations of the two possible states (and states) of the tubulins. The MT interior works as an electromagnetic wave guide, filled with water in an organized collective state, transmitting information through

the brain. A gelatinous state of water in brain cells, which was observed by [13], could boost these communication effects.

Using a different approach, Tuszynski et al. are modeling the biophysical aspects of the MTs considering the following questions: What kind of computing do microtubules perform? Can a microtubule store and process information? How do microtubules process information? In order to analyze these questions they start from the classical level, studying the physical properties of the MTs as interacting electric dipoles and move to the quantum level.

According to [6-8,14-17] each tubulin has an electric dipole moment \vec{p} due to an asymmetric charge distribution. The microtubule is thus a lattice of oriented dipoles that can be in random phase, ferroelectric (parallel-aligned) and an intermediate weakly ferroelectric phase like a Spin-Glass phase. It would be natural to consider the electric field of each tubulin as the information transport medium.

Therefore, the dimers tubulin would be considered the information unit in the brain and the MTs sub-processors of the neuron cells. How MTs process information and allow communication inside the brain is the fundamental question of the theory of mind-brain.

In this work we derive some results which were not explicitly obtained in [6-8] and extend the ideas introduced by [6,16] using the point of view of the information theory. We analyze the problem of information transfer and storage in brain microtubules, considering them as a communication channel. The electric field is the mediator of each communicator entity. We discuss the implications in assuming that consciousness is generated by the microtubules as sub-neuronal processors.

2 Biophysical Aspects of the Microtubules

The cytoskeleton has a dynamic structure which reorganizes continually as the cells change their shape, divide, and respond to their environment. The cytoskeleton is composed of intermediate filaments, actin filaments (or microfilaments) and microtubules. The filaments and the microtubules are mutually connected and form a three-dimensional network in the cell. There are many studies [6-8] showing that the cytoskeleton is the main component which organizes the cell, mediates transport of molecules, organelles and synaptic vesicles. The cytoskeleton possibly receives signals from the cellular environment mediated by the membrane of proteins and participates in signal transmission to the outside area of the cell [16,17].

Microtubules are hollow cylinders whose exterior surface cross-section diameter measures 25nm with 13 arrays of protein dimers called tubulines. The interior of the cylinder contains ordered water molecules which implies the existence of an electric dipole moment and an electric field. The MTs can represent a "giant" dipole generating an intrinsic electric field due to individual dipolar charges of each tubulin monomer. The microtubule dipole nature produces a fast growth at the plus end towards the cell periphery and a slow growth at the minus end. The microtubule polarity is closely connected with its functional behavior which can be regulated by phosphorylation and dephosphorylation of microtubule-associated protein (MAP) [6-8,14-17].

Guanosine triphosphate molecules (GTP) are bound to both tubulins in the heterodimer. After polymerization, when the heterodimer is attached to the microtubule, the GTP bound to the α -tubulin is hydrolyzed to the guanosine diphosphate (GDP). On the other hand, the GTP molecule of the β -tubulin is not hydrolyzed. The microtubules present a calm dynamic instability which are their principal feature [6-8].

Many models of conformation (and polarity energy) of the microtubular protofilament have been developed. These models describe the behavior of the pulses generated by the free energy in the GTP hydrolysis. The pulses propagate along of the MTs through an elastic coupling or through electric field propagation between tubulin dimers [5-8,14,15]. Hence, the Hamiltonian model for the dipole moments of a microtubule can be written as

$$H = \sum_{n=1}^N \left[\frac{1}{2} M \left(\frac{du_n}{dt} \right)^2 + \frac{1}{2} K (u_{n+1} - u_n)_{n+1}^2 + \frac{A}{2} u_n^2 + \frac{B}{4} u_n^4 - C u_n \right] ; \quad (1)$$

where u_n represents the dimer's conformational change on the n-protofilament axis - coupled to the dipole moment. The first term represents the kinetic energy, the second term represents the elastic restoring energy and the third term represents the effective nonlinear potential. The last term shows the effect of the external field on the system of coupled dipoles placed on tubulin molecules of mass M.

In figure 1 we plot the effective potential in terms of u_n taking $C = 0$ (no linear term). A and B are parameters of the model, where $A = a(T - T_c)$, T_c is the critical temperature, and $B > 0$ [6,8]

Our assumptions lead us to reconsider these relations taking into account the Information Theory. These models can show us a way to calculate the storage and transference of information considering the MT as a device mediated by the electric field propagating into cellular medium. This propagation of energy can provide a communication channel.

3 Information Theory

Information is related to random events. Entropy measures uncertainty of a physical system. In the classical approach, the Shannon entropy is the principal concept [16,18,20]. The entropy of a random variable is defined by the minus log of the probability that this random variable takes. Therefore, the expectation value of the information in the system, for the discrete case, is given by [18]:

$$h\{i\} = - \sum_i p(x_i) \log p(x_i) \quad (2)$$

This definition shows the amount of physical resources required on average to store the information being produced by the source, in such a way that at a later time the information can be restored.

The principal problem of the information theory is to determine the maximum rate R for reliable communication through the channel N . This maximum rate is given by the capacity of the channel, which can be calculated using the Shannon's theorem. The Shannon's theorem generalizes the case of the binary symmetric channel to the case of a discrete memoryless channel [18]. The memoryless channel is the channel that acts in the same way every time it is used, and different uses are independent of one another.

Consider a channel N with a finite input alphabet A_{in} and a finite output alphabet A_{out} : For the binary symmetric channel, $A_{in} = A_{out} = \{0,1\}$: The action of the channel is described by a set of conditional probabilities, $p(y_j | x_i)$, where $x_i \in A_{in}$ and $y_j \in A_{out}$. The term $p(y_j | x_i)$ is the probability of the different outputs y_j from the channel, given that the input was x_i , and satisfies the rules: $p(y_j | x_i) \geq 0$ and $\sum_j p(y_j | x_i) = 1$ for all i :

The probability of error in a channel is defined by [18]

$$p(C^n \neq D^n) = \min_M \sum_{Y \in M} p(Y \in M | X = C^n(M)) \quad (3)$$

where $C^n(\cdot)$ is the map $C^n : \{1, \dots, 2^{nR}\} \rightarrow A_{in}$ used to code some message M (for n uses of channel) and the output is decoded by using a map $D^n : A_{out}^n \rightarrow \{1, \dots, 2^{nR}\}$.

A rate R is achievable if $p(C^n \neq D^n) \rightarrow 0$ when $n \rightarrow \infty$ for a sequence of encoding-decoding pairs $\{C^n, D^n\}$. Therefore, according to Shannon's theorem,

given a noisy channel N ; its capacity $C(N)$ is defined to be the supremum over all achievable rates for this channel. That is [16,18]

$$C(N) = \max_{p(x_i)} (H(X : Y)); \quad (4)$$

where the maximum is taken over all input distributions $p(x_i)$ of the random variable X , for one use of the channel, and Y is the corresponding induced random variable at the output of the channel. In (4) the term $H(X : Y)$ is the average of the mutual information which is defined by [18]

$$H(X : Y) = H(X) - H(X|Y) = H(Y) - H(Y|X) \quad (5)$$

and

$$H(X|Y) = - \sum_i \sum_k p(x_i, y_k) \log p(x_i|y_k); \quad (6)$$

where $p(x_i, y_k) = p(x_i|y_k)p(y_k)$:

Therefore, the definition of the mutual information can evaluate how much information we have about the value of X given that we know the value of Y . That is, the mutual information measures how much information X and Y have in common [16,18].

The transfer of energy may have many forms and may include the transfer of electrostatic energy, energy of low frequency oscillating fields, energy of light, energy of vibrations, etc. Molecules can contain energy in the chemical bonds, in the excited electron states, in the conformation states, etc. A common measure of the interaction leading to cooperative behaviour is the transfer of information. The electromagnetic field can transfer information through the medium between the systems. And the medium between them could represent a communication channel.

4 Microtubules as Information Processors

Many other features of the cytoskeleton reinforce the idea of computation and information storage. According to [6], the MTs have a charge separation wide enough to store information. Due to its dynamic coupling the information can be stored as mechanical energy and chemical events.

Changes in the opposite direction can be favorable to the SG phase over the F-phase, and according to [6], this change could switch from the growth mode to operational behavior. Our focus is this operational mode. Information processing has been treated by [1-6] considering the highly specialized nature of the functional proteins inside the microtubules.

4.0.1 Storage and information processing on Microtubules

According to [3,5-8,16] the tubulins have a dipole moment net and therefore are sensitive to electric fields. So, at low temperatures, all the dipoles are

oriented at the same direction and the system is in the ferroelectric phase. Hence, in this phase there is an order in the system (figure 2). At around 300K (which is near the body conditions) there is a transition from the ferroelectric (F) to the paraelectric (P) phase, that is, from order to disorder (figure 3). However, at the region of transition the polarization can present a frustrated phase (figure 4) known as spin-glass (SG) [6,8,18].

Following [6] we examine the possible information features of MTs considering the above phases. Assuming an energy approximation dependent on the mean polarity described by the Landau theory of phase transitions, the total energy can be given by [6,16,20]

$$E = \frac{a}{2} \langle \phi \rangle^2 + \frac{b}{4} \langle \phi \rangle^4; \quad (7)$$

where ϕ represents the continuous variable for the mean polarization at each site. The terms a and b are parameters of model, where a has a linear dependence with the temperature $a = \bar{a}(T - T_c)$, T_c is between 200K and 400K and $b > 0$: Then the total energy E will be minimized by $\phi = 0$ for $T > T_c$ and for $T < T_c$ by $\phi = \frac{\bar{a}(T - T_c)}{b}$: We will adopt a Boltzmann distribution function $g(\phi)$ to weight the energy distribution in function of the mean polarity ϕ [6,16]:

$$g(\phi) = Z^{-1} \exp(-\beta E); \quad (8)$$

with $\beta = 1/kT$, where Z is the normalization and k is the Boltzmann constant. Substituting (8) into (7) we get

$$g(\phi) = Z^{-1} \exp\left(-\frac{\bar{a}(T - T_c)}{2kT} \phi^2 - \frac{b}{4kT} \phi^4\right); \quad (9)$$

As ϕ is a continuous variable, we need to use a continuous expression, similar to (2), to calculate the information expectation value of the system. In that sense, $g(\phi)$ is equivalent to $p(x)$ in (2) and therefore we have a relation for the information storage capacity in terms of the mean polarity ϕ :

$$H[\phi] = \frac{\ln Z}{\ln 2} - \frac{\bar{a}(T - T_c)}{2kT \ln 2} \langle \phi^2 \rangle - \frac{b}{4kT \ln 2} \langle \phi^4 \rangle; \quad (10)$$

where the average of ϕ over the whole MT, considering all domains, is obtained from

$$\langle \phi^n \rangle = \frac{1}{Z} \int_{-\infty}^{\infty} \phi^n g(\phi) d\phi; \quad (11)$$

Using (10) we can plot the information capacity against the temperature for some parameter values.

These graphs corroborate the results of [6], which show that, probably at physiological temperature, we can have a mode of information storage in MTs. This is the most important feature for finding another subunit of information

processing inside the brain. Finding another subunit could show us new perspectives for cognitive aspects or memory capacity [4-7].

However, according to these graphs, maximum information storage is obtained at the spin-glass phase, therefore we need to reformulate some assumptions. In this phase, there are domains with many energy levels which can store information [6, 20]. In the next section we will analyze the information processing via the communication channel approach. For each domain in the spin-glass phase, interaction with another by means of its oscillating dipole electric field will be considered. This electric field is emitted to the neighbouring area producing a channel in MT.

4.0.2 Microtubules as a Communication Channel

Given the capacity of information storage presented by the MTs, the subject now is to know if there is some kind of information processing on it. To study any kind of processing, is necessary to describe how the information, on the walls of MT, is linked. That is, we need to understand how the communication happens among different parts along the walls of MT. We saw, that the SG phase supplies the maximum capacity of information storage. Therefore, the subject is reduced for the description of the interaction or communication among domains on MTs in the SG phase. Here, we are assuming that the electric field generated by the MTs dipoles is the main mediator which allows the communication between the domains. We will address this problem from the classical point of view, therefore, we will not consider the quantization of the electric field [1-5].

The previous graphs (figures: (??), (??) and (??)) show that near the critical temperature T_c the information capacity has maximum storage. Following [6], we assume in this phase a partition of lattices by local domains (see figure 4). Therefore, the previous prescription will be valid only on the local domain of coherence. In this way, a domain j has associated a polarization ρ_j with probability distribution $g_j(\rho_j)$. If we make these assumptions the probabilities are independent and are given by the product of local distributions:

$$g = \prod_{j=1}^r g_j(\rho_j); \quad (12)$$

where r is the number of domains [6].

As a consequence of (12) we have for the spin-glass phase:

$$\langle \rho_i \rangle = \prod_j \langle \rho_j \rangle \quad (13)$$

with j in the set of domains.

In order to describe a processing feature we need to measure the amount of information transferred through the channel among the domains [18]. However, there is communication among domains only if they interact. If we consider two

with $S_j = \frac{2^p}{j^3} \frac{4}{j}$; and ω_j is the frequency of the dipole oscillating, ϵ is the permittivity, μ is the permeability of the medium, R_x and R_z are the perpendicular distances from the dipole to the z and x sides, respectively.

Using (5) and (6), and the previous relations, we can derive an expression for the capacity of communication between two domains. The channel between two domains j and k will be denoted by N_{jk} ; hence:

$$(N_{jk}) = hI(E_k) - hI(E_k \oplus_j); \quad (18)$$

where $hI(E_k)$ is given by an expression similar to (10). The conditional information $hI(E_k \oplus_j)$, for a specific polarity ω_j ; can be calculated by a continuous version of (6), that is

$$hI(E_k \oplus_j) = \frac{\ln Z}{\ln 2} \int_0^Z g(\omega_k; \omega_j) E_k + E_j^{\text{signal}} d\omega_k; \quad (19)$$

Through those calculations we can infer that, in the SG phase, there is an inter-dependence among the domains. Each domain communicates to other domain the value of its polarity, transforming the MT in a communication net among their subunits (in this case, the domains). Besides, as each domain possesses a particular polarity, in the context of the information theory, we could interpret each polarity representing a type of symbol, forming along the whole MT a kind of alphabet (with each domain representing a letter). However, as the polarity ω is a continuous variable, the change of a letter in another would be also in a continuous way and not in a discrete way.

The graphs below show the capacity information transference between each domain as a function of the distance.

The transition period observed for the of conformational changes of the tubulin dimer protein is 10^9 to 10^{11} s^{-1} : And the distance between the protein molecules in a cell membrane is of the order of 0.01 m - 0.1 m . The graphs show that its best performance is at the above frequency.

The relative permittivity and permeability of the medium around the oscillating units is assumed to be 1. The capacity of the channel depends on the frequency. At 300K, over a distance of 0.1 m the information transfer at 0.1 THz is suppressed (figure 8). At a distance smaller than 0.1 m the communication starts to become independent of frequency. And finally at a distance greater than 0.1 m the high frequency of the electric field plays a fundamental role in the transfer of information. For the other temperature regimes, the system does not present the SG-phase and the graphs show the loss in performance (figures 9 and 10)

According to [21], biological molecules with dipolar vibrational activity could manifest a coherent mode which could have some effect in isolating the system from thermal environments. The frequency range of the Fröhlich systems is around 10^{11} s^{-1} to 10^{12} s^{-1} [1,4]. Therefore, the high frequency regimes obtained above can work not only to perform communication but also to maintain some quantum coherence.

These results corroborate the possibility of communication between domains (with each energy level corresponding to some kind of symbol). This communication is mediated by the dipole electric field. This interaction is necessary to describe some processing or computing in MT. Through this communication, each domain (or symbol) presents some dependence with another. Therefore there could be storage as well as processing of information.

From the information theory point of view, the formation of domains creates some redundancy for storage or representation of these symbols. This redundancy is important for error correction and information protection.

5 Conclusions

This work confirm the results of [1-8] which consider microtubules as a classical subneuronal information processor. Through the information theory we could calculate the information capacity of the MTs. We conclude that the MTs can store and process information. Utilizing models of the [1-8] we estimated that favorable conditions for storage and processing are found at temperatures close to body temperature.

However there is yet a necessity to make some improvements on the model. For example, the direction of propagation of the information under the environment influence is an interesting issue to be analyzed. According to [1-5] there is some water ordination inside MTs which could increase the quantum processes in MTs. This point deserves to be analyzed using the information theory point of view.

6 References

- [1] S. Hagan, S. R. Hameroff and J. A. Tuszynski, Quantum Computation in Brain Microtubules: Decoherence and Biological Feasibility, *Physical Review E*, volume 65 - 061901, 4 June 2002.
- [2] S. R. Hameroff and R. Penrose, Orchestrated Reduction of Quantum Coherence in Brain Microtubules, in S. Hameroff, A. K. Kaszniak and A. C. Scott, *Toward a Science of Consciousness*, MIT Press, Cambridge, 1996
- [3] S. R. Hameroff and R. Penrose, Conscious Events as Orchestrated Space Time Selections, in J. Shear, *Explaining Consciousness. The Hard Problem*, MIT Press, Cambridge, USA, 1998
- [4] Mari Jibu, Scott Hagan, Stuart R. Hameroff, Karl H. Pribram, Kunio Yasue, Quantum Optical Coherence in Cytoskeletal Microtubules: Implications for Brain Function, *Biosystems* 32, 195-209, 1994.
- [5] Faber, Jean and Rosa, Luiz Pinguelli, Quantum Models of Mind: Are They Compatible with Environmental Decoherence? *accepted for publication in the Phys. Rev. E*.

- [6] J. A. Tuszynski, B. Tripisova, D. Sept, M.V. Sataric, The Enigma of Microtubules and their Self-organization Behavior in the Cytoskeleton. *BioSystems* 42, 153-175, 1997.
- [7] J. A. Tuszynski, J. A. Brown and P. Hawrylak, Dielectric Polarization, Electric Conduction, Information Processing and Quantum Computation in Microtubules. *Are They Plausible?*, The Royal Society, 356, 1897-1926, 1998.
- [8] J.A. Tuszynski, S. Hameroff, M.V. Sataric, B. Trpisova and M. L. A. Nip, Ferroelectric Behavior in Microtubule Dipole Lattices: Implications for Information Processing, Signaling and Assembly/Disassembly. *Journal Theoretical Biology*, 174, 371-380, 1995.
- [9] M. Tegmark, Importance of Quantum Decoherence in brain process, *Phys. Rev. E* 61 , p 4194, 2000
- [10] R. Penrose, *The Emperor New Mind*, Oxford University Press, 1989
- [11] R. Penrose, *Shadows of the Mind*, Vintage, London, 1994
- [12] J. C. Eccles, *How the Self Controls Its Brain*, Springer Verlag, Berlin, 1994
- [13] J. Watterson, *Water Clusters: Pixels of Life*, in S. Hameroff et alli, *Toward a Science of Consciousness*, MIT Press, Cambridge, 1996
- [14] M. V. Sataric, J. A. Tuszynski and R. B. Zakala *Phys. Rev. E*, 48, 589, 1993
- [15] J. A. Brown and J. A. Tuszynski, *Phys. Rev. E* 56, 5, 5834, 1997
- [16] J. Pokoorny and T. Ming Wu, *Biophysics Aspects of Coherence and Biological Order*, Springer, 1998.
- [17] Kandel, Eric R., Schwartz, James H., Jessell Thomas M., *Principles of Neural Science* (Appleton & Lange Norwalk), third edition 1991.
- [18] Michael A. Nielsen and Isaac L. Chuang, *2000 Quantum Computing and Quantum Information* (Cambridge)
- [19] Dotsenko, Viktor, 1994, *An Introduction to The Theory of Spin Glasses and Neural Networks*, (World Scientific Lecture Notes in Physics) - vol. 54.
- [20] Haken, H., 1990. *Synergetics: An Introduction* (Springer, Berlin).
- [21] H. Frohlich, Coherent excitations in active Biological systems, in *Modern Bioelectrochemistry*, edited by F. Gutman and H. Keyzer, Springer Verlag, N York, 1986

This figure "Info1.jpg" is available in "jpg" format from:

<http://arxiv.org/ps/q-bio/0404007v1>

This figure "Info2.jpg" is available in "jpg" format from:

<http://arxiv.org/ps/q-bio/0404007v1>

This figure "Info3.jpg" is available in "jpg" format from:

<http://arxiv.org/ps/q-bio/0404007v1>

This figure "Tub_Pot.jpg" is available in "jpg" format from:

<http://arxiv.org/ps/q-bio/0404007v1>

This figure "ferroel.jpg" is available in "jpg" format from:

<http://arxiv.org/ps/q-bio/0404007v1>

This figure "highT.jpg" is available in "jpg" format from:

<http://arxiv.org/ps/q-bio/0404007v1>

This figure "lowT.jpg" is available in "jpg" format from:

<http://arxiv.org/ps/q-bio/0404007v1>

This figure "lowT1.jpg" is available in "jpg" format from:

<http://arxiv.org/ps/q-bio/0404007v1>

This figure "paramag.jpg" is available in "jpg" format from:

<http://arxiv.org/ps/q-bio/0404007v1>

This figure "spin-glass.jpg" is available in "jpg" format from:

<http://arxiv.org/ps/q-bio/0404007v1>