

UNIDENTIFIABLE DIVERGENCE TIMES IN RATES ACROSS SITES MODELS

STEVEN N. EVANS AND TANDY WARNOW

Abstract. The rates across sites assumption in phylogenetic inference posits that the rate matrix governing the Markovian evolution of a character on an edge of the putative phylogenetic tree is the product of a character-specific scale factor and a rate matrix that is particular to that edge. Thus, evolution follows basically the same process for all characters, except that it occurs faster for some characters than others. To allow estimation of tree topologies and edge lengths for such models, it is commonly assumed that the scale factors are not arbitrary unknown constants, but rather unobserved, independent, identically distributed draws from a member of some parametric family of distributions. A popular choice is the gamma family. We consider an example of a clock-like tree with three taxa, one unknown edge length, and a parametric family of scale factor distributions that contain the gamma family. This model has the property that, for a generic choice of unknown edge length and scale factor distribution, there is another edge length and scale factor distribution which generates data with exactly the same distribution, so that even with infinitely many data it will be typically impossible to make correct inferences about the unknown edge length.

1. Introduction

Statistically-based estimations of phylogenetic trees are popular in molecular systematics, with Bayesian MCMC [HR01] and maximum likelihood methods [Swo96, GG03, PM00, Lew98, OMHO94] used with increasing frequency. Such statistically-based methods assume that the observed sequences are the result of a stochastic process that has operated on a tree, and they make assumptions about the stochastic process (i.e., model) that has produced the data. The models used in phylogenetic inference combine a (typically parameter rich) model of

Date: May 23, 2019.

Key words and phrases. phylogenetic inference, random effects, gamma distribution, identifiability.

SNE supported in part by NSF grant DMS-0071468.

TW supported by NSF grant BCS-0312830.

individual site evolution with the assumption that the different sites evolve under a rates-across-sites model, so that each site c has an associated rate of evolution r_c which is constant across the tree. Thus, sites evolve under essentially the same evolutionary process, and are just scaled up (or down) versions of each other. (Thus, the rates-across-sites assumption implies that if one site is expected to evolve twice as fast as another site on edge e , then it is expected to evolve twice as fast as the other site on every edge.)

Such rates-across-sites models in which each character has its own unknown scale factor are discussed in [SOWH96], but these models still pose difficult inferential problems. As remarked in Chapter 13 of [Fel04]:

As the number of sites increases, the number of parameters being estimated rises correspondingly. This is worrisome: in such "ininitely many parameters" cases maximum likelihood often misbehaves and fails to converge to the correct tree as the number of sites increases.

Indeed, our own example below shows that relative edge lengths are, in general, unidentifiable for such models.

A popular 'fix' that has been proposed for this problem is to adopt a random effects approach and suppose that the successive scale factors r_c are unobserved, independent random draws from a member of some parametric family of distributions. This reduces the dimensionality of the problem by replacing the deterministic sequence of r_c parameters with the small number of parameters that describe the generating distribution (see, for example, [UC71, NCF76, Ols87, HKY87, Yan96]).

A popular choice of distribution for the random scale factors is the two parameter family of gamma distributions. This family has the mathematical advantage that likelihoods still have analytically tractable closed forms. The choice of the gamma family is often supported by claims that it is sufficiently flexible to mimic the variation of rates between characters that is likely to be seen 'in practice'. There also appears to be a general sense among many practitioners that the choice of distributional family for the scale factors is primarily a matter of convenience and that, provided the family is rich enough, substantially correct inferences of relative edge lengths will be possible with sufficient data. To our knowledge, there is no argument in any setting justifying why an assumption of an exact gamma distribution for the scale factors is biologically reasonable. As remarked in [Fel04]:

There is nothing about the gamma distribution that makes it more biologically realistic than any other distribution, such as the lognormal. It is used because of its mathematical tractability.

It was shown in [SSH 94] that the use of random scale factors might not be completely without problems. In their paper they gave an example of a specific choice of edge lengths for each tree topology and a specific choice of discrete distribution for the scale factors (rather than a continuous distribution such as a gamma) such that the resulting distribution for the data under the Neyman two-state model is the same for all tree topologies. This result means that under the discrete distribution model, model trees are not identifiable. However, the result does not show that all model trees (i.e., all trees with particular discrete distribution for the scale factors) are not identifiable.

In this paper we go further, at least in some directions, and provide an example of:

- a rooted tree with three taxa and one unknown edge length (with the remaining edge lengths either known or fixed by the clock-like constraint that all lineages have the same total length),
- a ten parameter family F of scale factor distributions,
- a nine parameter sub-family G of F ;

such that for a generic choice of unknown edge length ℓ and model in $G \subset G$ there is a choice of edge length $\ell \in \mathbb{R}^+$ and model in $F \subset F$ with the property that:

data generated according to the Neyman 2-state model with known ancestral state, scale factor distributed according to G , and edge length ℓ , has the same distribution as data generated according to a Neyman 2-state model with the same ancestral state, scale factor distributed according to F , and edge length ℓ . Thus, even with infinitely many data, it would be impossible to decide whether the unknown edge length is ℓ or ℓ' (even if one somehow knew in advance that the distribution for the scale parameter was one of either F or G).

Here the term generic means that the set of exceptional models G and edge lengths ℓ for which no corresponding F and ℓ' exist is a lower dimensional subset of $G \times \mathbb{R}^+$. In particular, the set of G and ℓ that have corresponding F and ℓ' is an everywhere dense open subset of $G \times \mathbb{R}^+$.

Moreover, the family G consists of distributions with smooth densities and contains all the gamma distributions as a subfamily. Any

gamma distribution and any edge length t will have a distribution $G \pm G$ and ϵ arbitrarily close to them such that there is a corresponding $F \pm F$ and ϵ as above.

Also, our example applies not only to the Neyman model but also to any model such as the binary General Time-Reversible model that contains the Neyman model as a sub-model. Furthermore, our example applies to the General Time-Reversible on an arbitrary finite state-space, because one can choose the substitution rate matrices for such a model to be sufficiently symmetric so that a suitable many-to-one binary encoding of the model is Markovian and evolves according to the Neyman model.

In conclusion, the attempt to achieve identifiability and reasonable inference of edge lengths in the rates across sites model by using random scale factors that come from some common distribution can be problematic.

The rest of the paper is organized as follows. We begin with an introduction of the mathematical terms in Section 2, and we present our example in Section 3. We conclude with a discussion of the ramifications of this result, and directions for future research in Section 4.

2. Basics

In phylogenetic inference, the data are the respective states of an ensemble of characters exhibited by each of a collection of taxa. The most commonly used statistical models in the area are parameterised by a rooted tree with edge lengths (which typically represent the expected number of times a site changes on the edge when the substitution mechanism is in equilibrium) and a set of Markovian stochastic mechanisms for the evolution of successive characters down the tree. It is usually assumed that the observed states for different characters are statistically independent. The goal of phylogenetic inference is to estimate some or all of: the shape (topology) of the tree, the lengths of the edges, and any unknown parameters involved in the specification of the evolution mechanism.

We will restrict attention to the case where each character has the same finite set of possible states. For example, the characters could be nucleotides exhibited at different sites on the genome, and so each character is in one of the four states $\{A; G; C; T\}$. In the example we will give in this paper, we will work with (binary) characters having one of two possible states, 0 or 1. For each character c and each edge e in the tree, one then has a rate matrix $Q_{c,e}$ that describes the evolutionary

process on edge e for character c . Thus, given that the character is in state i at the beginning of the edge, the conditional probability of the (possibly unobserved) event that it is in state j at the end of the edge is the $(i; j)$ entry of the matrix exponential $\exp(tQ_{e;c})$, where t is the length of e . The matrix $Q_{e;c}$ has row sums equal to 0 and non-negative off-diagonal entries: $Q_{e;c}(i; i)$ is the rate at which the character leaves the state i and $Q_{e;c}(i; j) = -Q_{e;c}(i; i)$ is the probability that it jumps to state j when it leaves state i .

Single site substitution models can range from the very simple (e.g. the Jukes-Cantor and Kimura 2-parameter models) to the very complex (e.g. the General Markov Model), which, for a fixed character c , allow the $Q_{e;c}$ matrices to vary significantly from edge to edge, and to have many free parameters. However, the variation between the different matrices obtained by varying the character c is typically more proscribed. The most complex model is where there are no constraints are placed on the $Q_{e;c}$; this is called the "no comm mechanism model" [S97]. Under this no comm mechanism model, it will clearly be difficult to recover any information about edge lengths. The simplest class of models in which it is possible to extract information about edge lengths is the class in which $Q_{e;c}$ is the same for all characters c and edges e . Even for this simple model, there is (as is well-known) a certain lack of identifiability, because the same probability distribution for the data would arise if the comm on rate matrix was multiplied by a comm on scale factor and all edge lengths were divided by that same factor. Thus, even for this model one can only hope to make inferences about relative edge lengths unless at least one edge length is assumed to be known.

The more commonly used models assume that the different $Q_{e;c}$ matrices are themselves the product of a rate matrix specific to the edge e , and a scale factor that is specific to the character c . Thus, the evolutionary process that governs one character is identical, up to a scalar multiple, to that governing another character. This is the rates across sites assumption in molecular phylogenetics, and it has the rather strong implication that if a character c is expected to evolve twice as fast on edge e as character c^0 , then c is expected to evolve twice as fast on every edge in the tree.

Because of the inferential difficulties of allowing the rates for the different sites to be arbitrary, these random scale factors are typically assumed to be drawn from a distribution. Of the many possible distributions, the most popular distributions are the two parameter gamma distributions. In fact, in practice, almost all estimations of phylogenetic trees are based upon the assumption that the rates across sites

are drawn from a gamma distribution, or a discretized gamma distribution.

3. The example

We will present an example of a tree with three taxa, and with sites evolving under the Neyman two-state model (i.e., the two-state version of the Jukes-Cantor model of evolution).

Consider a tree with three taxa x, y , and z ; root v ; and internal node w that is ancestral to x and y . The edges $(w; x)$ and $(w; y)$ have a known length, which we can take as 1. Suppose further that the edge $(v; w)$ has unknown length and that the tree is clock-like, so that the edge $(v; z)$ has length $+ 1$.

Suppose there are $f_0; 1g$ -valued characters labelled $1; 2; \dots$ that have evolved on this tree. The i^{th} character evolves according to the Neyman model with rate r_i . That is, the transition matrix for an edge of length t is

$$e^{-2r_i t} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + \frac{1}{2} (1 - e^{-2r_i t}) \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix} :$$

Assume that successive characters evolve independently.

The probability distribution for the i^{th} character (that is, the marginal likelihood for this character) is thus a linear combination of the terms

$$\begin{aligned} & 1 \\ & \exp(-2r_i) \\ & \exp(-4r_i) \\ & \exp(-2r_i) \\ & \exp(-2r_i(1 + \theta)) \\ & \exp(-2r_i(2 + \theta)) \\ & \exp(-2r_i(3 + \theta)) \\ & \exp(-2r_i(1 + 2\theta)) \\ & \exp(-2r_i(2 + 2\theta)) \\ & \exp(-2r_i(3 + 2\theta)) : \end{aligned}$$

As described in the Introduction, we will adopt the random effects approach and assume that the r_i are, in fact, realisations of a sequence of independent, identically distributed random variables that we will denote by (A_i) .

We are interested in finding such a sequence (A_i) and another independent, identically distributed sequence (B_i) such that the distribution for the data induced by the random choice of scale factors (A_i) is the same as that induced by the (B_i) for another choice of edge length ϵ .

We thus have to find positive random variables A and B with distinct distributions and distinct positive constants α and β with the property that

$$\begin{aligned} E[\exp(-2A)] &= E[\exp(-2B)] \\ E[\exp(-4A)] &= E[\exp(-4B)] \\ E[\exp(-2\alpha A)] &= E[\exp(-2\beta B)] \\ E[\exp(-2(1+\alpha)A)] &= E[\exp(-2(1+\beta)B)] \\ E[\exp(-2(2+\alpha)A)] &= E[\exp(-2(2+\beta)B)] \\ E[\exp(-2(3+\alpha)A)] &= E[\exp(-2(3+\beta)B)] \\ E[\exp(-2(1+2\alpha)A)] &= E[\exp(-2(1+2\beta)B)] \\ E[\exp(-2(2+2\alpha)A)] &= E[\exp(-2(2+2\beta)B)] \\ E[\exp(-2(3+2\alpha)A)] &= E[\exp(-2(3+2\beta)B)]: \end{aligned}$$

Take A to have the distribution which has Laplace transform

$$E[\exp(-sA)] = \prod_{i=1}^7 (1 + d_i)^{-1} (1 + e)^{-1} (1 + f)^{-1}$$

for positive parameters $d_1; \dots; d_7, e, f$. Thus A has the distribution of the sum of 9 independent exponential random variables with respective means $d_1; \dots; d_7, e, f$. Take B to have the distribution which has Laplace transform

$$E[\exp(-sB)] = \prod_{i=1}^7 (1 + g_i)^{-1} (1 + h)^{-2}$$

for positive parameters $g_1; \dots; g_7, h, h$. Thus B has the distribution of the sum of 9 independent exponential random variables with respective means $g_1; \dots; g_7, h, h$.

Define maps $P : \mathbb{R}_+^{10} \rightarrow \mathbb{R}_+^9$ and $Q : \mathbb{R}_+^9 \rightarrow \mathbb{R}_+^9$ by

$$\begin{aligned}
 P_1(d_1; \dots; d_7; e; f) &= \begin{pmatrix} Y^7 \\ (1 + 2d_i) \quad (1 + 2e) \quad (1 + 2f) \\ \vdots \\ (1 + 2d_i) \quad (1 + 2e) \quad (1 + 2f) \end{pmatrix} \\
 P_2(d_1; \dots; d_7; e; f) &= \begin{pmatrix} Y^7 \\ (1 + 4d_i) \quad (1 + 4e) \quad (1 + 4f) \\ \vdots \\ (1 + 4d_i) \quad (1 + 4e) \quad (1 + 4f) \end{pmatrix} \\
 P_3(d_1; \dots; d_7; e; f) &= \begin{pmatrix} Y^7 \\ (1 + 2d_i) \quad (1 + 2e) \quad (1 + 2f) \\ \vdots \\ (1 + 2d_i) \quad (1 + 2e) \quad (1 + 2f) \end{pmatrix} \\
 \vdots \\
 P_9(d_1; \dots; d_7; e; f) &= \begin{pmatrix} Y^7 \\ (1 + 2d_i(3 + 2)) \quad (1 + 2e(3 + 2)) \quad (1 + 2f(3 + 2)) \\ \vdots \\ (1 + 2d_i(3 + 2)) \quad (1 + 2e(3 + 2)) \quad (1 + 2f(3 + 2)) \end{pmatrix}
 \end{aligned}$$

and

$$\begin{aligned}
 Q_1(g_1; \dots; g_7; h) &= \begin{pmatrix} Y^7 \\ (1 + 2g_i) \quad (1 + 2h)^2 \\ \vdots \\ (1 + 2g_i) \quad (1 + 2h)^2 \end{pmatrix} \\
 Q_2(g_1; \dots; g_7; h) &= \begin{pmatrix} Y^7 \\ (1 + 4g_i) \quad (1 + 4h)^2 \\ \vdots \\ (1 + 4g_i) \quad (1 + 4h)^2 \end{pmatrix} \\
 Q_3(g_1; \dots; g_7; h) &= \begin{pmatrix} Y^7 \\ (1 + 2g_i) \quad (1 + 2h)^2 \\ \vdots \\ (1 + 2g_i) \quad (1 + 2h)^2 \end{pmatrix} \\
 \vdots \\
 Q_9(g_1; \dots; g_7; h) &= \begin{pmatrix} Y^7 \\ (1 + 2g_i(3 + 2)) \quad (1 + 2h(3 + 2))^2 \\ \vdots \\ (1 + 2g_i(3 + 2)) \quad (1 + 2h(3 + 2))^2 \end{pmatrix}
 \end{aligned}$$

We want to show that

$$P(d_1; \dots; d_7; e; f) = Q(g_1; \dots; g_7; h)$$

for some choice of parameters with $\epsilon > 0$.

Write $J(d_1; \dots; d_7; e; f)$ for the Jacobian matrix of the mapping

$$(d_1; \dots; d_7; e; f) \mapsto P(d_1; \dots; d_7; e; f)$$

(thus, J is a 9×9 matrix). Write $K(g_1; \dots; g_7; h)$ for the Jacobian matrix of Q . A straightforward check with a computer algebra package such as Mathematica shows that the polynomials $\det J$ and $\det K$ are not identically 0. (While the determinants could possibly be computed symbolically, it is easier to compute the matrices symbolically, substitute in appropriate integer values for the parameters, and use exact integer arithmetic to compute the determinant.)

for those values: For example, $\det J(2;3;4;5;6;7;8;9;10;11) \notin 0$ and $\det K(2;3;4;5;6;7;8;9;10) \notin 0$.) Because these determinants are polynomials, the set of values where J (resp. K) is non-singular is a relatively open subset of \mathbb{R}_+^{10} (resp. \mathbb{R}_+^9) with a closure that is all of \mathbb{R}_+^{10} (resp. \mathbb{R}_+^9) (that is, they are everywhere dense).

We can therefore find a point $(\gamma; g_1; \dots; g_7; h)$ in the interior of \mathbb{R}_+^9 such that:

- (i) the matrix $K(\gamma; g_1; \dots; g_7; h)$ is non-singular,
- (ii) in any open neighbourhood of $(\gamma; g_1; \dots; g_7; h)$ in \mathbb{R}_+^9 there are points $(d_1; \dots; d_7; e; f)$ such that the matrix $J(\gamma; d_1; \dots; d_7; e; f)$ is non-singular.

By assumption (i) and the implicit function theorem (see, for example, [KP02]), the range of Q contains an open neighbourhood of $Q(\gamma; g_1; \dots; g_7; h)$. Note that

$$P(\gamma; g_1; \dots; g_7; h; h) = Q(\gamma; g_1; \dots; g_7; h);$$

and so for all points $(\gamma; d_1; \dots; d_7; e; f)$ in some open neighbourhood of $(\gamma; g_1; \dots; g_7; h; h)$ we can find $(\gamma; g_1; \dots; g_7; h)$ such that

$$P(\gamma; d_1; \dots; d_7; e; f) = Q(\gamma; g_1; \dots; g_7; h);$$

We will be done if we can show that it is not always the case that $\gamma = \dots = g_7 = h$ for such a solution. To see this, we will fix $\gamma = \dots = g_7 = h$ and let $(d_1; \dots; d_7; e; f)$ vary. By assumption (ii) and the implicit function theorem, the image of any open neighbourhood of $(\gamma; g_1; \dots; g_7; h; h)$ by the map

$$(d_1; \dots; d_7; e; f) \mapsto P(\gamma; d_1; \dots; d_7; e; f)$$

has non-empty interior. However, the range of the map

$$(\gamma; g_1; \dots; g_7; h) \mapsto Q(\gamma; g_1; \dots; g_7; h)$$

is at most 8-dimensional, and, in particular, has empty interior. Therefore, there certainly exists $(d_1; \dots; d_7; e; f)$ such that

$$P(\gamma; d_1; \dots; d_7; e; f) = Q(\gamma; g_1; \dots; g_7; h)$$

for some $(\gamma; g_1; \dots; g_7; h)$ with $\gamma \neq \dots \neq g_7 = h$.

Remark. Note that if we take $g_1 = \dots = g_7 = h$, then we have a gamma distribution with shape parameter 9. Also, we could still produce the unidentifiability phenomenon witnessed above if we raised all the Laplace transforms to the same power $c > 0$. In that case, setting $g_1 = \dots = g_7 = h$ would give a gamma distribution with shape

parameter ρ . Since the unidentifiability occurs on a dense set of parameters (g_1, \dots, g_7, h) , any gamma distribution will have distributions arbitrarily close to it that exhibit the phenomenon.

4. Conclusions and Future Research

The example we have given shows that the attempt to achieve identifiability and reasonable inference of edge lengths in the rates across sites model by using random scale factors that come from some common distribution can be problematic.

The gamma distributions work, but comparably simple distributions arbitrarily close to them don't. Using the gamma family is thus not just a matter of working with distributions that have enough flexibility to capture reasonable variation in rates. Rather, identifiability of edge lengths for the gamma family relies on quite specific features of that family that are not shared by a somewhat larger class of equally reasonable distributions.

The result has consequences for the estimation of times at internal nodes, since if edge lengths cannot be estimated, then neither can the dates (since the edge length is a product of the elapsed time on the edge, and the equilibrium expected rate of evolution for on that edge). It is also reasonable to consider the question of whether model trees are identifiable under generic cases of these simple distributions of rates across sites. Our result does not answer this, and so further research is necessary.

Finally, this is a theoretical result, and the consequences of this theoretical result can be tested in simulation. To date, few (if any) such studies have been done that have not presumed that the rates are distributed by a gamma distribution, or a distribution consisting of some invariable sites, and the remaining sites evolving under a gamma distribution. This also reflects the implicit belief that the assumption of a gamma distribution is acceptable. We hope this paper will help encourage researchers to reconsider this assumption.

References

- [Fel04] J. Felsenstein, *Inferring phylogenies*, Sinauer Associates, Sunderland, Massachusetts, 2004.
- [GG03] Stéphane Guindon and Olivier Gascuel, A simple, fast, and accurate algorithm to estimate large phylogenies by maximum likelihood, *Syst. Biol.* 52 (2003), no. 5, 696{704.
- [HKY87] M. Hasegawa, H. Kishino, and T. Yano, Man's place in Homonoidea as inferred from molecular clocks of DNA, *J. Mol. Evol.* 26 (1987), 132{147.

- [HR01] J.P. Huelsenbeck and R. Ronquist, MrBayes: Bayesian inference of phylogeny., *Bioinformatics* 17 (2001), 754{755.
- [KP02] Steven G. Krantz and Harold R. Parks, *The implicit function theorem*, Birkhauser Boston Inc., Boston, MA, 2002, History, theory, and applications. MR 2003f:26001
- [Lew98] P. Lewis, A genetic algorithm for maximum likelihood phylogeny inference using nucleotide sequence data, *Mol. Biol. Evol.* 15 (1998), 277{283.
- [NCF76] M. Nei, R. Chakraborty, and P.A. Fuerst, Infinite allele model with varying mutation rate, *Proc. Natl. Acad. Sci. USA* 73 (1976), 4164{4168.
- [Ol87] G.J. Olsen, Earliest phylogenetic branchings: Comparing rRNA-based evolutionary trees inferred with various techniques, *Cold Spring Harbor Symposia on Quantitative Biology* 52 (1987), 825{837.
- [OMHO94] G. Olsen, H. Matsuda, R. Hagstrom, and R. Overbeek, FastDNAml: A tool for construction of phylogenetic trees of DNA sequences using maximum likelihood, *Computations in Applied Biosciences* 10 (1994), no. 1, 41{48.
- [PM00] S. L. Kosakovsky Pond and S. Muse, Hyphy package distribution and documentation page, 2000, Distributed by the authors at <http://www.hyphy.org>.
- [SOWH96] D.L. Swofford, G.J. Olsen, P.J. Waddell, and D.M. Hillis, *Phylogenetic inference*, *Molecular Systematics* (D.M. Hillis, C. Moritz, and B.K. Mable, eds.), Sinauer Associates, Sunderland, Massachusetts, 1996.
- [SSH94] M.A. Steel, L.A. Szekely, and M.D. Hendy, Reconstructing trees when sequence sites evolve at variable rates, *J. Comp. Biol.* 1 (1994), 153{163.
- [Swo96] D. Swofford, PAUP*: phylogenetic analysis using parsimony (and other methods), version 4.0.
- [TS97] C. Tuley and M. Steel, Links between maximum likelihood and maximum parsimony under a simple model of site substitution, *Bulletin of Mathematical Biology* 59 (1997), 581{607.
- [UC71] T. Uzzell and K.W. Corbin, Fitting discrete probability distributions to evolutionary events, *Science* 72 (1971), 1089{1096.
- [Yan96] Z. Yang, Maximum-likelihood estimation of phylogeny from DNA sequences when substitution rates differ over sites, *Mol. Biol. Evol.* 10 (1996), 1396{1401.

E-mail address: evans@stat.Berkeley.EDU

Department of Statistics # 3860, University of California at Berkeley, 367 Evans Hall, Berkeley, CA 94720-3860, U.S.A

E-mail address: tandy@cs.utexas.edu

Department of Computer Sciences, University of Texas at Austin, Austin, TX 78712, U.S.A.