

The Mystery of Two Straight Lines in Bacterial Genome Statistics

Alexander Gorban

Centre for Mathematical Modelling, University of Leicester, U.K.

Andrey Zinovyev^y

Institut des Hautes Etudes Scientifiques, Bures-sur-Yvette, France

Abstract

In special coordinates (codon position {specific nucleotide frequencies}) bacterial genomes form two straight lines in 9-dimensional space: one line for eubacterial genomes, another for archaeal genomes. All the 143 known bacterial genomes belong to these lines with high accuracy, and these two lines are certainly different.

For each genome we can define the codon position {specific nucleotide frequencies} in the coding part of this genome. The coding part of the genome is separated into codons. The codon position {specific nucleotide frequencies} are 12 numbers p^i , where i is a nucleotide symbol (A, C, G, or T), and $i = 1; 2; 3$ is a position number of nucleotide in codon. Among 12 frequencies p^i only 9 are independent because normalization conditions $p_A^1 + p_C^1 + p_G^1 + p_T^1 = 1$. The possibility of non-uniqueness of separation of the coding part into codons is here neglected. The vectors with all possible coordinates p^i fill a 9-dimensional polyhedron (a direct sum of three 3-dimensional simplices). In this 9-dimensional space bacterial genomes form two straight lines: one line for eubacterial genomes, another for archaeal genomes. All the 143 known bacterial genomes (from Genbank release August 2004) belong to these lines with high accuracy, and these two lines are certainly different. Both these lines can be parameterized by G+C genomic content in a very natural way, because p^i prove to be linear functions of (non position {specific}) G+C content with high accuracy. These functions are different for different lines. The results of statistical analysis (the regression lines together with experimental points) are presented on Fig. 1. Significant differences between eubacteria and archaea are observed for $p_A^1, p_C^1, p_G^1, p_G^3$, and p_T^3 functions.

The average codon usage in genome is represented by 64 frequencies p^M , where M is a nucleotide symbol, p^M is the frequency of the codon M in coding part of genome. The mean-field approximation for codon usage is $p^M = p^1 p^2 p^3$. The frequencies p^M for bacterial genomes belong to lines of third order (two such lines: one for eubacterial genomes, another for archaeal genomes) in the 63-dimensional simplex of codon usage (64 frequencies minus one normalization condition) and the codon usage for known genomes form the clouds near these lines. These clouds of codon usage have a distinctive horseshoe form [1].

ag153@le.ac.uk

^yzinovyev@ihes.fr




Figure 1: Codon position specific nucleotide frequencies as functions of average GC-content. Solid line and empty points correspond to 124 completed eubacterial genomes, broken line and filled points correspond to 19 completed archaeal genomes.

Some hints to observed structure were reported recently in studies on multivariate analysis of bacterial codon usage (for example, see Figure 6 from [1], or in codon bias study in [2]). More illustration can be found in e-print [3] and on the web{site [4].

There are even two mysteries:

1. Why bacterial genomes form straight lines?
2. Why these lines are certainly different for eubacteria and archaea?

References

- [1] Lobry, J.R., Chessel, D., Internal correspondence analysis of codon and amino-acid usage in thermophilic bacteria. *J. Appl. Genet.* 44 (2) (2003), 235{261.
- [2] Carbone, A., Kepes, F., Zinovyev, A., Codon Bias Signatures, Organisation of Microorganisms in Codon Space and Lifestyle. *Mol Biol Evol.* 2004 Nov 10.
- [3] Gorban, A.N., Popova, T.G., Zinovyev, A.Yu., Four basic symmetry types in the universal 7-cluster structure of 143 complete bacterial genomic sequences, e-print: <http://arxiv.org/abs/q-bio.GN/0410033>.
- [4] Cluster structures in genomic word frequency distributions, Web{site: <http://www.jhes.fr/zinovyev/7clusters>.