The Logic Backbone of a Transcription Network

M. Cosentino Lagomarsino

UMR 168 / Institut Curie, 26 rue d'Ulm 75005 Paris, France*

P. Jona

Politecnico di Milano, Dip. Fisica, Pza Leonardo Da Vinci 32, 20133 Milano, Italy[†]

B. Bassetti

Università degli Studi di Milano, Dip. Fisica, and I.N.F.N. Via Celoria 16, 20133 Milano, Italy[‡]

(Dated: February 9, 2020)

Abstract

A great part of the effort in the study of coarse grained models of transcription networks is directed to the analysis of their dynamical features. In this letter, we consider the *equilibrium* properties of such systems, showing that the logic backbone underlying all dynamic descriptions has the structure of a computational optimization problem. It involves variables, which correspond to gene expression levels, and constraints, which describe the effect of *cis*-regulatory signal integration functions. In the simple paradigmatic case of Boolean variables and signal integration functions, we derive and discuss phase diagrams. Notably, the model exhibits a connectivity transition between a regime of simple, but uncertain, gene control, to a regime of complex combinatorial control.

^{*}e-mail address: mcl@curie.fr

[†]e-mail address: patrizia.jona@fisi.polimi.it

[‡]e-mail address: bassetti@mi.infn.it

a. Introduction. Identity, response and architecture of a living system are central topics of molecular biology. Presently, they are largely seen as a result of the interplay between a gene repertoire and the regulatory machinery [1, 2]. Gene transcription in mRNA form is an important step in this process. At this level, the fundamental regulatory machinery is embodied by the transcription factors, proteins that bind to the *cis* regulatory sites along DNA and control the activity of RNA polymerase [3, 4] (Fig. 1). This process is referred to as signal integration. Together, the *cis*-regulatory regions establish a set of interdependencies between transcription factors and genes, including other transcription factors: a "transcription network" [1]. Some of the transcription networks of living organism are now being explored experimentally [2]. Studies on the structure have shown a modularity that has important biological implications [5].

Characterizing and predicting the gene expression patterns determined by these networks remains an enormous challenge. The problem is that transcription networks are fairly large. While detailed models or simulations work well on small (sub-)systems [6, 7], typically a coarse grained approach is needed. Microscopically, it is well accepted that the Gillespie algorithm [8], while disregarding spatial correlations, correctly describes the stochastic asynchronous events of chemical kinetics involved. On the other hand, with a mesoscopic average in time, it is not clear what the emergent time scales might be. In particular, the pioneering approach of Kauffman [9] suggesting a synchronous deterministic dynamics for a Boolean (i.e. on/off) representation of the network is still being debated [10].

We approach this problem with a model that focuses, rather than on pure dynamics, on the compatibility between gene expression patterns and signal integration functions. Simply put, a cell with N genes can express them in exponentially many ways, 2^N in the Boolean representation. However, the cell never explores all the patterns of expression. It only knows clusters of correlated configurations. An elementary example is the cI-cro switch of λ -phage [3]. Looking at the system one could observe the states 10, 01 or perhaps 00, but not 11. In a cell, one can think that many of the states are not observable for the same compatibility reasons.

This formulates the problem as an optimization problem, which we call GR, from Gene Regulation. The simplest case (GR1) assumes Boolean variables and functions. It maps to a Satisfiability problem [11] (Sat), an optimization problem where N Boolean variables are constrained by M conjunctive normal form (CNF) constraints (i.e. by a Boolean polynomial constructed as a product (\land) of M disjunctive monomials (\lor)). We investigate theoretically the behavior of GR1 for large N, and for random realizations of the constraints [12].




FIG. 1: Schematics of our representation of a transcription network. Each signal integration function at the *cis*-regulatory region of a gene is transformed in a constraint on the gene expression variables. Bottom: example of factor graph of GR1 for K = 2. Each diamond node represents a signal integration function, while each black circle is a variable. The directionality of the constraint is represented graphically by connecting the input variables to two edges of the diamonds and the output variables to a vertex.

The phase diagrams in Fig. 2 to 4 resume our main findings. An important premise is the fact, which is evident looking at the available data [5], that some genes are essentially "free" from the point of view of transcription. These are mainly controllers and are connected to external stimuli. We introduce the "complex combinatorial core" (CCC) of the network, the subset of genes that determine a choice of an expression program. We define order parameters related to the number of free genes in the CCC and the connectivity of the constraints. There are three distinct regimes of gene control. In the first (UNSAT), there are no free genes in the core, and the system cannot control the simultaneous expression of all its genes. In the second regime ("complex control" or HARD-SAT), the core contains free genes that control, both directly and indirectly, many others. The general dynamics is residual (many variables are fixed, the others can change). In the third regime, the core is empty. Each free gene (which is external to the core) controls the state of a small number of genes ("simple control", or SAT phase). By varying the connectivity and the number of constraints, a hypothetical CCC can "tune" its transition from the simple to the complex control phase.

b. Model. Mapping on a Satisfiability Problem. The two main ingredients of our representation of a transcription network are: (i) A set of N discrete variables $\{x_i\}_{i=1..N}$ associated to genes or operons (identified with their transcripts and protein products). These variables represent the expression levels and take values in $\{0, ..., q\}$. (ii) A set of interactions, or constraints, $\{I_b(x_{b_0}, x_{b_1}, ..., x_{b_{K_b}})\}_{b=1..M}$ between the genes, representing the signal integration. It is useful to represent variables and interactions in a so-called factor graph, as illustrated in Fig. 1. Note that the constraints I_b contain implicitly the topology of the graph. Considering ensembles of random graphs, their probability distribution needs to be specified through these constraints. We refer to this as q-level GR, or GRq.

The x_i represent coarse-grained expression levels. It is possible to use the Shea and Ackers model of gene activation by recruitment [4, 15] to construct a local free energy associated to each signal integration node, that generates the constraints [16]. Here we consider the simplest possible scenario, GR1, treating the expression levels as Boolean variables, setting q = 1, and the signal integration functions as Boolean functions $\{f_b(x_{i(b,1)}, ..., x_{i(b,K_b)})\}_{b=1..M}$. Here $k_b = 1 + K_b$ can be seen as the local connectivity of a function node. Note that the factor graph is also defined by a variable connectivity c_i , the number of functions connected to x_i .

The expression

$$x_{i(b,0)} = f_b(x_{i(b,1)}, .., x_{i(b,K_b)}),$$
(1)

imposes that the variable $x_{i(b,0)}$ is the output of the function f_b . It translates into the XOR Boolean constraint $\neg(x_{i(b,0)} \dot{\lor} f_b)$. This can be easily recast in CNF, as 2^{K_b} clauses involving k_b variables, each clause corresponding with a trivial map to a line in the truth table of f_b .

The formula

$$I = \bigwedge_{b=1..M} I_b = \bigwedge_{b=1..M} \neg (x_{i(b,0)} \dot{\lor} f_b)$$
(2)

defines a Satisfiability problem on the variables $x_1, ..., x_N$. From the biological viewpoint, this is a logic representation of the computational tasks encoded in the transcription network by evolution, i.e. which sets of genes have to be switched on at any given condition. If we restrict to the case of fixed in-ward connectivity, $K_b = K$, $\forall b$, these conditions, defining K-GR1, are also found in Kauffman networks [9, 10] and define a k-Sat problem whose connectivity is k = K + 1.

Having established a link between GR1 and a particular optimization problem we set out to look at random instances for the signal integration functions. For fixed connectivity one can expect a similar behavior for K-GR1 as random k-Sat, or k-XORSAT (a Sat problem with clauses containing only XOR clauses), with the presence of a phase transition in the number of satisfying states. The suitable order parameter for such a transition is $\gamma = M/N$. Notably, the space of functions of the three models have different dimensions. Furthermore, differently from Sat or XORSAT, in GR1 each gene expression variable is regulated at most by one signal integration function, so that $\gamma \leq 1$.

c. Phase Diagram To compute the number of satisfying solutions \mathcal{N} , for random instances of the constraints, each Boolean variable $x_i \in \{0, 1\}$ is transformed into a spin $\sigma_i \in \{-1, 1\}$. Under the restrictions defining GR1, the total energy of the system is simply the number of violated constraints. A zero-energy configuration satisfies all the constraints, and is therefore able to comply to all the logic functions encoded by the network.

The conventional average of \mathcal{N} on the realizations might be biased by the weight of exceptions [17]. The correct quantity to compute is the "quenched average" of the system's free energy, $\overline{\log \mathcal{N}}$, which is usually accessed with the replica, or similar methods [13]. However, in the case under examination we will use a simpler method, based on the "leaf removal" [14] algorithm, which allows to compute only the *annealed* average $\log \overline{\mathcal{N}}$. For the case of random XORSAT, Mezard and collaborators [14] have shown that the annealed average on the core variables coincides with the quenched one. In general $\overline{\log \mathcal{N}} \leq \log \overline{\mathcal{N}}$. For GR1, we performed estimates that indicate the presence of the same self-averaging property, which, if confirmed, would render our calculations exact.

For a given realization of the constraints $\{\vec{I}, \vec{f}\}$, the number of satisfying states \mathcal{N} can be written as

$$\mathcal{N}(\vec{I}, \vec{f}) = \sum_{\vec{\sigma}} \prod_{b=1}^{M} \delta(1; f_b(\sigma_{i(b,1)}, ..., \sigma_{i(b,K_b)}) \sigma_{i(b,0)}).$$

Here, the randomness is contained: (i) in the specification of the network structure, $\vec{I} = (I_1, ..., I_M)$, i.e. in the coordinates i(b, l), which point at the variable occupying place l in the *b*th constraint; (ii) in the specification of the functions $\vec{f} = (f_1, ..., f_M)$ with a certain probability distribution in the class \mathcal{F} . An overbar (⁻) indicates an average on both distributions, $p(\vec{I})$ and $p(\vec{f})$.

Let us first concentrate on the case with fixed in-ward connectivity K. We consider a set functions in a family \mathcal{F}_K which satisfy the condition $\frac{1}{2^{2K}} \sum_{f \in \mathcal{F}_K} p(\vec{f}) f(\vec{x}) = \rho$, as will always be the case if the outputs of the functions are uncorrelated, even in the presence of bias. It is then easy to verify that

$$\overline{\mathcal{N}} = \sum_{\vec{\sigma},\vec{I}} p(\vec{I}) \prod_{b=1}^{M} \left(\rho \delta(1; \sigma_{i(b,0)}) + (1-\rho) \delta(-1; \sigma_{i(b,0)}) \right) ;$$

so that, as for the XORSAT model,

$$\overline{\mathcal{N}} = 2^{N-M} . \tag{3}$$

Interestingly, the same procedure is valid for GRq, where $\overline{\mathcal{N}} = q^{N-M}$.




FIG. 2: Phase diagram of 4-GR1. For $\gamma > \gamma_c$ no solutions exist in the typical realization (UNSAT phase). For $\gamma < \gamma_d$ the system is paramagnetic typically a satisfying state exists (SAT phase, or simple gene control). For $\gamma_d < \gamma < \gamma_c$ there is complex gene control.

The leaf removal [14] algorithm makes use of the fact that not all the M equations and N variables are meaningful to calculate \mathcal{N} . Indeed, for every variable that appears in only one constraint, both the constraint and the variable can be eliminated without affecting the number of solutions. This is a nonlinear procedure, as more variables can disappear together with a single constraint, and leads to the definition of a "core", the CCC, of significant variables and constraints, in numbers of N_C and M_C respectively. In the CCC, M_C genes are controlled, and $\Delta_C = N_C - M_C$ are the "free" genes with an essential role in controlling the expression states, as a function of an input signal. We can use $\frac{\Delta_C}{N_C} \leq 0$, there are no free genes in the core, and the system cannot comply to all the expression programs encoded into its DNA. This is a UNSAT phase from the computational point of view. (ii) For $0 \leq \frac{\Delta_C}{N_C} < 1$, the Δ_C genes of the CCC control O(N) genes each, and therefore are able to determine an expression state. This is a "complex gene control"

phase, or HARD-SAT (iii) For $\frac{\Delta_C}{N_C} = 1$, which in the thermodynamic limit $N \to \infty$, γ const., implies $\Delta_C = 0$, the free genes control O(1) genes to generate a satisfying state ("simple control", or SAT phase).

These three phases depend on the value of γ . In general the phase diagram can be explored studying the rank and the kernel of the connectivity matrix. Following the analysis of Mezard and collaborators [14], in the case of Poisson variable connectivity $p(c) = \frac{(k\gamma)^c}{c!}e^{-k\gamma}$, the phase diagram as a function of γ is identical to the random XORSAT problem. It is illustrated in Fig. 2. For $\gamma > \gamma_c$ no solutions exist in the typical realization (UNSAT phase). For $\gamma < \gamma_d$ the system is paramagnetic (SAT phase). For $\gamma_d < \gamma < \gamma_c$ exponentially many satisfying states exist. Here, the space of solutions breaks down into clusters separated by free energy barriers. The typical dynamics in a cluster will be residual, in the sense that a block of genes are fixed (on or off) and the rest may move. The number of clusters is controlled by the (computational) complexity Σ of the system. The number of observed configurations is $\mathcal{N}^* \sim \exp[N\Delta_C]$. Thus, by definition Σ is directly related to the order parameter Δ_C/N_C , i.e. to the partitioning of the core genes. How the system explores (or not) the clusters depends on details of its dynamics.

A biologically more interesting case is when both the inward connectivity K and the outward connectivity c vary along the network. Considering $p(k|c) = \frac{(k\gamma)^c}{c!}e^{-(k\gamma)}$, the conditioned probability that a variable is in c clauses of the k kind, we have $p(c) = \sum_k \frac{(k\gamma)^c}{c!}e^{-(k\gamma)} \cdot p(k)$. The leaf removal algorithm can be applied separately to sets of clauses with a given connectivity, defining $N_C \equiv \langle N_C \rangle_k$ and $M_C \equiv \langle M_C \rangle_k$, where $\langle X \rangle_k = \sum_p p(k) \cdot X(k)$. Choosing $p(k) = Z^{-1}(\nu)e^{-\nu k}$ does not affect the exponential asymptotic decay of p(c) for large c. We can call this case multi-Poisson, as the graph is a superimposition of graphs with fixed clause connectivity and Poisson variable connectivity, which in turn follows a Poisson distribution.

The behavior of GR1 on such a topology is nontrivially different from the fixed connectivity case. The main reason for this is that, while $\Delta_C(\gamma)$ is still locally decreasing, many new discontinuities emerge, due to the influence of clauses with different connectivities. This gives rise to two phenomena. Firstly, Δ_C can increase globally with increasing γ . Indeed, it does increase after γ_d , to decrease again before γ_c (Fig. 3). After the onset of the complex control phase, the complexity initially increases (step-wise), reaches a maximum, and then decreases monotonically. This has an influence on the number of observed states as a function of γ . Secondly, Δ_C can become negative and then jump back to a positive state, creating a reentrant UNSAT phase (Fig. 4).




FIG. 3: Δ_C as a function of γ for different values of ν in the multi-Poisson case. The discrete jumps are due to the onset of HARD phases for the different values of k. Δ_C can become negative many times, giving rise to reentrant UNSAT phases. The figure refers to a connectivity distribution with a cutoff at k = 12

d. Discussion and Conclusions To conclude, we established a simple framework for the equilibrium modeling of large scale transcription networks. It is a compatibility analysis on the constraints established by the signal integration functions. Its advantage is that, while giving non-trivial results, it avoids to deal directly with issues related to the dynamics. In absence of an explicit knowledge of the emergent time scales involved in the dynamics, we feel this is an appropriate approach. Particularly in the Boolean approximation, which we treat here.

From a general, qualitative standpoint, our model shows that the "biological" complexity is not simply measured by the number of genes. A more proper indicator is Δ_C which depends on the order parameter γ , or - roughly - on the number of transcription factors per gene. At fixed number of genes, it is known that this quantity increases in bacteria that need to react to more environments [18]. Imagining that prokaryotes, being unicellular, "live" in a simple control phase, our phase diagram predicts an intrinsic limit to this process, represented by the phase boundary with the HARD-SAT, and the connected slowing down of the dynamics. Considering varying




FIG. 4: Phase diagram $\gamma - \nu$ for the multi-Poisson case. The dashed line represents the mean value of the numerically evaluated critical parameter $\gamma_d(\nu)$ for the SAT-HARD transition of network realizations with $N = 3 \times 10^4$.

N, one may wonder why, in real organisms, a small γ is correlated with a small N [19]. A possible answer to this question is the following. With large N and small γ , the system is shifted to the SAT phase, and therefore needs to explore a very big configuration space without sufficient "guidelines". In other words, the available configurations are too many to be reached in reasonable time by the dynamics.

Looking at the phase diagram in Fig. 2, the complex control phase, having general residual dynamics, matches a qualitative feature of many cells, where some genes are constantly expressed, and the rest vary. On the other hand, the dynamical slowing down characteristic of any glassy phase raises an issue that must be solved by the chemical dynamics of the cell. In analogy with Kauffman's ideas, the breakdown in many different attraction basins might be interpreted as epigenesis. That is, in the HARD-SAT phase there will be typically many cell types. How many, is determined by the complexity Σ , which is directly related to our Δ_C . While for fixed K this quantity simply decreases with increasing γ , its behavior is more interesting in the multi-Poisson case. However, this remains an open issue which has to be regarded with more detail. The experimental scaling of the number of cell types is sub-linear in the number of genes N [9, 20]. In the fixed K case, GR1 gives exponential scaling with N at fixed γ in the complex control phase. On the other hand, the results of random-GR1 are perhaps more easily related to the number of species times the number of cell types at equal number of genes.

The approach presented here is new, and remains largely unexplored. It is naturally fit to study networks with non-Boolean variables and non-fixed in-ward connectivities. It can be easily extended to nonzero temperature (probabilistic constraints). It can be of use for models that include other regulation mechanisms than just transcription. More far-reached extensions include evolutionary models, with a fitness landscape on the space of constraints. It is not clear to us exactly how useful it can become for the study of concrete biological networks. A biologically significant model has to be able to deal with the details of an individual realization the system. This is, we think, the main challenge to our approach. For example, it will be interesting to identify whether the CCC is meaningful in known transcription networks. Furthermore, there exist techniques able to analyze single realizations of optimization problems which might be applicable to sufficiently characterized genetic networks. We are aware of work in this direction, which is now in course of publication [21]. For example, a typical experiment in molecular biology involves a local interaction with a regulation network with overexpression or deletion of a gene. Such methods of analysis could be used to generate predictions of the effects that keep into account the network connections. Perhaps a deeper understanding of these interdependent pathways might help to interact more effectively with them. Hopefully with a constructive purpose.

Acknowledgments

We would like to acknowledge useful discussions with A. Sportiello, J. Berg, M. Leone, A. Pagnani, M. Caselle, P.R. ten Wolde, R. Zecchina.

M. Babu, N. Luscombe, L. Aravind, M. Gerstein, and S. Teichmann, Curr Opin Struct Biol 14, 283 (2004).

^[2] M. Herrgard, M. Covert, and B. Palsson, Curr Opin Biotechnol 15, 70 (2004).

^[3] M. Ptashne, A Genetic Switch (Cell Press, MA, 1992), second edition ed.

- [4] N. Buchler, U. Gerland, and T. Hwa, Proc Natl Acad Sci U S A 100, 5136 (2003).
- [5] S. Shen-Orr, R. Milo, S. Mangan, and U. Alon, Nat Genet 31, 64 (2002).
- [6] H. McAdams and A. Arkin, Proc Natl Acad Sci U S A 94, 814 (1997).
- [7] A. Arkin, J. Ross, and H. McAdams, Genetics 149, 1633 (1998).
- [8] D. Gillespie, J. Phys. Chem. 81, 2340 61 (1977).
- [9] S. Kauffman, The Origins of Order (Oxford Univ. Press, New York, 1993).
- [10] C. Gershenson, in Artificial Life IX Workshops and Tutorials (2004).
- [11] M. Mezard and R. Zecchina, Phys Rev E Stat Nonlin Soft Matter Phys 66, 056126 (2002).
- [12] S. Mertens, Computing in Science and Engineering 4, 31 (2002).
- [13] M. Mezard, G. Parisi, and R. Zecchina, Science 297, 812 (2002).
- [14] M. Mezard, F. Ricci-Tersenghi, and R. Zecchina, J. Stat. Phys 505 (2003).
- [15] M. Shea and G. Ackers, J Mol Biol 181, 211 (1985).
- [16] M. Cosentino Lagomarsino, B. Bassetti, and P. Jona (2004), in Preparation.
- [17] M. Mezard, G. Parisi, and M. Virasoro, *Spin Glass Theory and Beyond* (World Scientific, Singapore, 1987).
- [18] I. Cases, V. de Lorenzo, and C. Ouzounis, Trends Microbiol 11, 248 (2003).
- [19] E. van Nimwegen, Trends Genet **19**, 479 (2003).
- [20] S. Kauffman, J Theor Biol 230, 581 (2004).
- [21] L. Correale, M. Leone, A. Pagnani, M. Weigt, and R. Zecchina (2004), in Preparation.