

Introns Restructure tRNA Genes of Archaea

Zhumur Ghosh ^(a), Smarajit Das ^(a), Jayprokas Chakrabarti ^(a, b, *),
Bibekanand Mallick ^(a) and Satyabrata Sahoo ^(a)

(a) Computational Biology Group (CBG)

Theory Department
Indian Association for the Cultivation of Science
Calcutta 700032 India

(b) Biogyan

BF 286, Salt Lake
Calcutta 700064 India

*** Author for correspondence**

Telephone: +91-33-24734971, ext. 281(Off.)
Fax: +91-33-24732805
E-mail: tpjc@iacs.res.in ; biogyan@vsnl.net

ABSTRACT

Motivation: Introns in tRNAs are suspected to play many roles. Analysis of the bulge-helix-bulge structural motifs at the intron-exon boundaries provide some insights. The splice-sites on these structural motifs suggest that a single intron-containing-tRNA-gene can give rise to more than one tRNA product.

Results: Partially overlapped mitochondrial tRNA genes are known to exist. But cytoplasmic tRNA genes overlap in archaeal methanogens in ways that are different. We present here bioinformatic evidence that in these methanogens the domain of overlap is far wider, encompassing the entire tRNA genes. tRNA genes are embedded within one another. Intron of one acts as exon of the other and vice-versa. We propose alternate intron splice-sites on bulge-helix-bulge motifs.

INTRODUCTION

In our recent work (Mallick *et al.*, 2005) we analysed cytoplasmic tRNA genes (tDNAs) of 22 species of 12 orders of three phyla of archaea for identity elements of aminoacylation. As these investigations progressed we observed some tDNAs missing in archaea. On closer examination some of these missing tDNAs were found embedded in other tDNAs. In this communication we present these observations and explain the rationale behind our conjecture of intron splicing at alternate positions in archaeal tDNAs.

Introns are found in tDNAs in all three domains of life (Diener and Moore, 1998). These were first discovered in yeast's nuclear tDNA (Valenzuela *et. al.*, 1978); later found in several archaeal tDNAs between nucleotide-positions 37 and 38, located in anticodon (AC) loop (Daniels *et al.*, 1985). These are the canonical introns. Archaea, an intermediate between Eukarya and Bacteria, have tRNAs that share many similarities with either or both these domains (Marck and Grosjean 2002). Archaeal tDNAs also have introns at positions other than canonical. These unusually located noncanonical introns in archaeal tDNAs were observed in 1987 (Wich *et al.*, 1987). Presence of intron in tDNA at canonical position is detected by the algorithm tRNAscan-SE (Lowe & Eddy, 1997). But detecting the unusually located noncanonical introns could be difficult. This is due to the lack of prior knowledge regarding their lengths and exact locations. An incomplete set of tDNAs within the genome is generally an indication that the missing tDNAs contain one or two unusually located introns.

Eukaryotic and archaeal tDNA splicing chemistry share deep similarities (Kleman-Leyer *et al.*, 1997). Intron splicing in both these domains occur through endonucleolytic cleavage of tDNAs producing 5' hydroxyl and 2', 3' cyclic phosphate termini at intron-exon boundaries (Kjems and Garrett, 1988). There are at least three forms of tRNA endonuclease in archaea: homotetrameric, homodimeric, and heterotetrameric. Crenarchaeal and nanoarchaeal endonucleases are heterotetrameric. Euryarchaeal endonucleases are either of the first two with an exception in *M. kandleri* where heterotetrameric endonuclease is found (Tocchini-Valentini *et al.*, 2005). Splicing mechanism in most of the cases depend on structures or specific sequences that lie primarily within the introns themselves and occurs by transesterification reactions (Negri *et al.*, 1997). Archaeal splicing endonucleases recognize and cleave a conserved secondary structure motif called the bulge-helix-bulge (BHB) that appears at the intron-exon boundaries. Archaeal splicing machinery cleaves introns at variable positions in pre-tRNAs within the BHB motif (Li *et al.*, 1998). Splicing of introns hence is a RNA-protein interaction which requires mutual recognition of two complementary tertiary structures. tDNA introns contribute to the specificity of splice-site recognition (Bufardecì *et al.*, 1993). Mutational analysis, secondary structure probing, and sequence analysis have shown that the conformation of the BHB motif is more important for archaeal endonuclease recognition than its sequence (Lykke-Andersen and Garrett, 1994, 1997).

In our previous study (Mallick *et al.*, 2005) some common tRNA genes were missed in the following three euryarchaeal methanogens: In *Methanococcus maripaludis* S2 (NC_005791) -37 tRNA genes were identified; in *Methanopyrus kandleri* AV19 (NC_003551) -35 tRNA and in *Methanocaldococcus jannaschii* (NC_000909)- 36 tRNA genes were identified. Note all other archaea have about 46 tRNA genes.

In our present work we re-investigated the genomes of the above in search for the missing tRNA genes. Interestingly we found evidence that some of these missing tDNAs are embedded in other tDNAs. Mitochondrial tRNA genes were found earlier to overlap by one to six nucleotides with downstream genes on the same strand (Yokobori and Pääbo, 1995). For instance, tRNA^{Tyr} and tRNA^{Cys} of human mitochondrial genome overlap with one another by one nucleotide at the last base of tRNA^{Cys}. This nucleotide is the discriminator base of tRNA^{Tyr} (Reichert *et al.*, 1998). But tDNA-overlaps in archaeal methanogens is an altogether new phenomena. Here the domain of overlap is far wider, encompassing the entire tDNAs. These tDNAs are embedded within one another. We propose intron- splicing at alternate positions on the bulges of the BHB motif.

ALGORITHM

We identified tRNA-genes from archaeal genomes by tRNAscan-SE and ARAGORN. These tRNA search programs key on primary sequence patterns and or secondary structure specific to tRNA. A few loop-holes exist in these algorithms. It is the inability of these existing routines to identify tRNA genes if it harbours noncanonical introns. Some of the tRNAs are misidentified; some are missed out by these search algorithms. We developed a computational approach to search tRNA genes having introns at positions other than canonical and identified some missing tRNA-genes in the three methanogens. About one thousand tRNA-genes from archaea were studied for this purpose. From this database of 1000 tRNA-genes we fine-tuned the strategy to locate noncanonical introns. The salient features were : (i) sequences were considered that gave rise to the regular cloverleaf secondary structure. (ii) conserved elements : T8 (except Y8 in *M. kandleri*), G18, R19, R53, T44, Y55, and A58 were considered as conserved bases for all archaeal tRNA. Further there were tRNA-specific conserved or identity elements (Mallick *et al.*, 2005) of archaea, (iii) the promoter ahead of the 5' end looked for, (iv) the constraints of lengths of stems of regular tRNA A-arm, D-arm, AC-arm and T-arm were 7, 4, 5 and 5 bp respectively. In few cases the constraints on lengths of D-arm and AC-arm were relaxed. (v) Base positions optionally occupied in D-loop were 17, 17a, 20a and 20b. (vi) the extra arm or V-arm was considered for type I and II tRNAs. The constraint on length of V-arm: less than 21 bases (vii) Noncanonical introns were considered at any position in tRNA-genes. The introns constrained to harbour the Bulge-Helix-Bulge (BHB) secondary structure for splicing out during tRNA maturation. The minimum length of intron allowed was 6 bases. (viii) The locations of the noncanonical intron and of the splice-sites on the bulges fixed to have the right conserved and identity bases for the two embedded tDNAs. With these features in our algorithm, we extracted some tRNA genes missed out in the three archaea.

STRUCTURAL SIGNATURES AND SPLICE-SITES

M. maripaludis

tRNA^{Ala} / tRNA^{Trp} Embedded genes :

From our database of 22 archaeal genomes we observed that three alanine tRNAs function to read all the alanine codons in archaea. But in *M. maripaludis* tRNA^{Ala}(CGC) was missing. Using our

identification algorithm we identified it, lying in the range c1627676-1627513. It has two introns. A noncanonical intron, of length 45 bases, between 31/32 of the tRNA and a canonical intron of length 44 bases. The sequence, along with introns, is in fig 1. It has a stable cloverleaf secondary structure. All the conserved bases and base-pairs of archaeal tRNA^{Ala} are there in this tRNA. This tRNA has G3:U70 and A73. These are unique to tRNA^{Ala} and key elements recognized by alanyl-tRNA-synthetase. In tRNA^{Ala}(CGC) gene of *M. maripaludis* the Bulge-Helix-Bulge(BHB) secondary structural motif (at the intron-exon boundaries) along with the cloverleaf of tRNA is in fig 2a. The five base-pair long exonic helix (h_e) is followed by a bulge (b) wherein lies the NCI start site (one of the intron splice sites). At the central position is the common bulge (B). From B three bifurcated arms shoot out. The end position of the NCI, along with the start and end positions of canonical intron are in these arms. In the figure the splice sites are denoted by arrows. The entire intronic structure along with the anticodon arm is $h_e b h' B [(h_1' b h_1' L_1) (h_2' L_2) (h_3' L_3)]$ (h_e is the exonic helix, b denotes bulge ; h' is purely intronic helix or a mixed helix).

Interestingly, we observed that tRNA^{Trp}(CCA) gene is embedded in tRNA^{Ala}(CGC) . Considered as tDNA^{Trp}(CCA) the above sequence (fig 1) has a canonical intron of length 89 bases. This tRNA has A73, C34, C35 and A36, the key aminoacylating elements of archaeal tRNA^{Trp} . The BHB motif remains the same. We propose alternate splice-sites on the bulges in fig 2a. Splicing positions for this pre-tRNA shows that the 3'-splice-site for its canonical intron is the same as that of tRNA^{Ala}(CGC) gene. Thus the proposed intron splice-sites give rise to two tRNAs embedded in the composite tDNA^{Ala} / tDNA^{Trp} .

M. kandleri

tRNA^{Arg} / tRNA^{Trp} Embedded genes :

Arginine is a six codon amino acid. Normally four archaeal arginine tRNAs read the arginine codons in all archaea. In this species only two tRNAs with anticodons UCG and UCU are present. This is clearly unusual.

We find tDNA^{Arg}(CCT) gene between 55221-55068. It has A73, U36 and C35, important elements for proper aminoacylation. A20 is another important identity element for archaeal tRNA^{Arg} (exception: *N. equitans* has U20). Interestingly, this tRNA of *M. kandleri* has U20. It has all the conserved bases and base-pairs of archaeal arginine tRNA. In fig 1 we show the sequence with its introns. Fig 2b shows the secondary structure of this tRNA. Notice the stable cloverleaf structure. It has noncanonical intron of length 38 at 33/34 and canonical intron of length

38. In fig the BHB structure and the splice-sites corresponding to these introns are clearly marked on the bulges.

The same range (fig 1) of the genome sequence of *M. kandleri* embeds tRNA^{Trp} gene. This tRNA has all conserved nucleotides (A73, C34, C35 and A36) for a proper archaeal tryptophan tRNA. It has a canonical intron (fig 1) of length 77 bases. Introns of both of these tRNAs have the BHB motif of type h_ebh/bh/bh/L with different splice sites. Notice again, the 3' splice-sites of the canonical intron of arginine and tryptophan tDNAs on the bulge coincide (fig 2b).

tRNA^{Ile} / tRNA^{Phe} Embedded genes

In archaea tRNA^{Ile}(UAU) gene was first observed in *N. equitans* (Mallick *et al.*,2005). In eukaryotes this tDNA is present may be in multiple copies in several species. Interestingly we found that the sequence ranging from c1639188-1639080 which has the signatures of tRNA^{Ile}(UAU) gene. It has a noncanonical intron of 16 bases at 33/34. There is a canonical intron of length 16 bases. Both the introns have proper BHB motifs of type h_ebh/L. The splicing site is found therein (see figure 2c). The secondary structure shows the presence of A73, A35, T36 base and C27:G43 bp which are important elements for aminoacylation.

On the contrary the same tDNA sequence has the features of tRNA^{Phe}. It has a canonical intron of length 32 having different splice sites within the same BHB motif (figure 2c). A73, G34, A35, A36 and bp C13:G22 are also present which are important for proper aminoacylation. Also 3' splice site of the CIs of the two tRNAs are similar.

M. jannaschii

tRNA^{Pro} / tRNA^{Trp} Embedded genes :

Two of the four Proline tRNAs are present in this species .Again this is unusual. We identify tRNA^{Pro}(CGG) gene lying between 1189945-1190052. This gene has a 26 length long noncanonical intron between bases 31 and 32 and a canonical intron of length 7 (fig 1). Apart from the presence of all the conserved bases and base pairs of archaeal Proline tRNA it has the following important elements for aminoacylation: A73, G36, and three GC at 5' end of A-arm.

We observe that same sequence embeds tRNA^{Trp}(CCA) gene of *M. jannaschii* . It has a 33 base long canonical intron (fig 1). Notice it also has all the signatures and aminoacylation elements (described earlier) of archaeal tRNA^{Trp} gene.

An active intron consists of two domains, one extending from the 5' splice site to the position of the base participating in the A-I (anticodon-intron) interaction (Baldi *et al.*, 1992), the other including the latter base and extending up to the 3' splice site. tRNA precursors with introns 6-10 bases long are efficiently cleaved only at one of the two domains. If an A-I pair is present, they have domain I; if the A-I pair is absent and the intron is at least 8 bases long, they have domain II (Negri *et al.*, 1997). tRNA^{Pro}(CGG) gene of *M. jannaschii* has a short length CI is of 7 nt which should be cleaved at one of the two domains. At its 32nd position 'C' is present which can pair with 'G' at the 3' end of the intron. A-I interaction is possible in this case. Hence it has only domain I having splice site at 3' end. Both the CIs of the two tRNA genes and NCI of Proline tRNA gene have proper BHB motif of type h_ebh^hB [(h₁' L₁) (h₂' L₂)]. The splice sites are marked by arrow in figure 2d where the 3'-splice site of the CIs of the two tDNAs are same.

ANSATZ:

In some of the primary transcripts of mitochondrial tRNAs of animals, tRNA occasionally abuts another tRNA/mRNA and overlaps by one to several bases (Yokobori and Pääbo, 1995). In archaea we studied the availability of all tRNAs over the entire set of sequenced genomes. We identified the tRNAs missing in the three methanogens. Studying then the structural motif, the bulge-helix-bulge, at the exon-intron boundaries, we found we could get a few of the missing ones if introns are spliced at alternate sites on the bulges. Such a process would create two tRNAs from one composite tDNA containing introns. Exon of one of the tRNA is intron of the other and vice-versa. The two tRNAs would embed onto each other through this intron-exon coupling. Bioinformatic evidence led us to the ansatz of alternate splice-sites on the bulge-helix-bulge motifs. Introns of these embedded tDNAs have nontrivial secondary structure. They are relatively stable and, probably, have further unknown functions. Assuming this alternate intron-splicing ansatz we are led to speculate on how this works. There are at least two possibilities. One is to assume that tRNA- endonuclease recognition sites are not uniquely determined. Coded into the tDNA there are sequence/structural signatures that lead to one or the other of the splice-sites. An alternate possibility is to assume that there exist two competing almost equally stable structural motifs leading to two alternate splicing modes.

Acknowledgements

We acknowledge useful discussions with Chanchal Dasgupta, Siddhartha Roy, Samit Adhya, Shaon Raychaudhuri and Ashok Thakur.

References:

Baldi, M. I., Mattoccia, E., Bufardecì, E., Fabbri, S. and TocchiniValentini, G.P. (1992) Participation of the intron in the reaction catalyzed by the *Xenopus* tRNA splicing endonuclease. *Science*, **255**, 1404-1408.

Bufardecì, E., Fabbri, S., Baldi, M. I., Mattoccia, E. and Tocchini-Valentini, G.P. (1993) In vitro genetic analysis of the structural features of the pre-tRNA required for determination of the 3' splice site in the intron excision reaction. *EMBO J.*, **12**, 4697-4704.

Daniels, C.J., Gupta, R. and Doolittle, W.F. (1985) Transcription and excision of a large intron in the tRNA Trp gene of an archaebacterium, *Halobacterium volcanii*. *J Biol Chem*, **260**, 3132-3134.

Diener, J. L. and Moore, P. B. (1998) Solution Structure of a Substrate for the Archaeal Pre-tRNA Splicing Endonucleases: The Bulge-Helix-Bulge Motif. *Molecular Cell* ,**1** , 883-894.

Kjems, J. and Garrett, R. (1988) Novel splicing mechanism for the ribosomal RNA intron in the archaebacterium *Desulfurococcus mobilis*. *Cell* , **54**, 693-703.

Kleman-Leyer, K., Armbruster, D.W. and Daniels, C.J. (1997) Properties of *H. volcanii* tRNA Intron Endonuclease Reveal a Relationship between the Archaeal and Eukaryal tRNA. *Cell*, **89**, 839-847.

Li, H., Trotta, C.R. and Abelson, J.(1998) Crystal structure and evolution of a transfer RNA splicing enzyme. *Science*, **280**, 279-284.

Lowe, T.M. and Eddy, S.R. (1997) tRNAscan-SE: A program for improved detection of transfer RNA genes in genomic sequence. *Nucleic Acids Res.*, **25**, 955-964.

Lykke-Andersen, J. and Garrett, R.A. (1994) Structural characteristics of the stable RNA introns of archaeal hyperthermophiles and their splicing junctions. *J. Mol. Biol.*, **243**, 846-855.

Mallick, B., Chakrabarti,J., Sahoo, S., Ghosh, Z. and Das, S.(2005) Identity Elements of Archaeal tRNA. *DNA Research* (article in press).

Marck, C. and Grosjean, H. (2002) tRNomics: Analysis of tRNA genes from 50 genomes of Eukarya, Archaea, and Bacteria reveals anticodon-sparing strategies and domain-specific features. *RNA*, **8**, 1189-1232.

Negri, E. D. N., Fabbri, S., Bufardecì, E., Baldi, M.I., Attardi, D.G., Mattoccia, E. and Tocchini-Valentini, G.P. (1997) The Eukaryal tRNA Splicing Endonuclease Recognizes a Tripartite Set of RNA Elements. *Cell.*, **89**, 859-866.

Reichert, A., Rothbauer, U. and Mörl, M. (1998) Processing and Editing of Overlapping tRNAs in Human Mitochondria. *J. Biol. Chem.*, **278**, 31977-31984.

Tocchini-Valentini, G.D., Fruscoloni, P. and Tocchini-Valentini, G.P. (2005) Structure, function, and evolution of the tRNA endonucleases of Archaea: An example of subfunctionalization. *Proc. Natl. Acad. Sci. USA*, **102**, 8933-8938.

Valenzuela, P., Venegas, A., Weinberg, F., Bishop, R. and Rutter, W.J. (1978) Structure of yeast phenylalanine-tRNA genes: An intervening DNA segment within the region coding for the tRNA. *Proc Natl Acad Sci USA*, **75**, 190-194.

Wich, G., Leinfelder, W., and Böck, A. (1987) Genes for stable RNA in the thermophile *Thermoproteus tenax*: introns and transcription signal. *EMBO J.* ,**6**, 523-528.

Yokobori, S. -i. and Pääbo, S. (1995) Transfer RNA Editing in Land Snail Mitochondria. *Proc Natl Acad Sci USA* ,**92** , 10432-10435.

M. mairipaludis (NC_005791)

tRNA^{Ab} c1627676-1627513

GGGAGTATGGTGTAGTCTGGTCTATCATCGGGG**ACTCCAAGTGCCTATATGGCCTT**
CTTAGAAATCTAAGAATTTTTAAGGCGCGCAAACACCAGTTTGTGCGTGGACTAAA
GCCOGATGCAGGAGAAATCCCTGGACCTGGGTTCAAATCCAGTACTTCCA

tRNA^{Ab} c1627676-1627513

GGGAGTATGGTGTAGTCTGGTCTATCATCGGGGACTCCA**AGTGCCTATATGGCCTT**
CTTAGAAATCTAAGAATTTTTAAGGCGCGCAAACACCAGTTTGTGCGTGGACTAAA
GCCOGATGCAGGAGAAATCCCTGGACCTGGGTTCAAATCCAGTACTTCCA

M. kandleri (NC_003551)

tRNA^{As} 55221-55068

GGGCCCGCGTGTAGCCAGGTCTATCATGCGGGGCT**CCAGGGACCCGCCGATGACT**
TCCATGGGGTGTGCCGGACCCTG**TGGGGAGACCCTGGAGCGACGAGGCGGGTCCG**
AAGACACCCCGTGACCCGGGTTCAAATCCCGGCGGGCCACCA

tRNA^{As} 55221-55068

GGGCCCGCGTGTAGCCAGGTCTATCATGCGGGGCTCCAG**GGACCCGCCGATGACT**
TCCATGGGGTGTGCCGGACCCTG**TGGGGAGACCCTGGAGCGACGAGGCGGGTCCG**
AAGACACCCCGTGACCCGGGTTCAAATCCCGGCGGGCCACCA

tRNA^{Al} c1639188-1639080

GCCGCGGCAGCTCAGCCTGGGAGAGCGCOGACT**G AAGAGATCCCGCGGGTATTCC**
CGCGGGATCGCOGCCATCCGGTGGGCCCGGGTTCAAATCCCGGCCGCGGCACCA

tRNA^{Al} c1639188-1639080

GCCGCGGCAGCTCAGCCTGGGAGAGCGCOGACTGAAG**AGATCCCGCGGGTATTCC**
CGCGGGATCGCOGCCATCCGGTGGGCCCGGGTTCAAATCCCGGCCGCGGCACCA

M. jannaschii (NC_000909)

tRNA^{As} 1189945-1190052

GGGGGTGTGGTGTAGCCAGGTCTATCATCGGGGACT**CTCCAGTTGCCTATAGGGCGCT**
GGACCACGC**AGGAGAA**ATCCCTGGACCTGGGTTCAAATCCAGCACCCCA

tRNA^{As} 1189945-1190052

GGGGGTGTGGTGTAGCCAGGTCTATCATCGGGGACTCCAG**TTGCCTATAGGGCGCT**
GGACCACGC**AGGAGAA**ATCCCTGGACCTGGGTTCAAATCCAGCACCCCA

Figure 1




Figure 2a

Figure Legends

Figure 1: Sequences of the embedded tRNA genes.

Blue coloured region denotes noncanonical intron, brown canonical intron and black the exonic region.

Figure 2a : Secondary structure of tRNA^{Ala}(CGC) / tRNA^{Trp}(CCA) along with the BHB of their introns of *M. maripaludis*.

NCIs: Noncanonical intron start position; NCIE: Noncanonical intron end position; CIs: Canonical intron start position; CIE: Canonical intron end position

—————> This signifies splicing sites of the introns in pre-tRNAs

h_e: Exonic helix; h[/]: mixed helix (part of it is intronic and part of it is exonic) (h₁[/]: mixed helix in the 1st branch; h₂[/]: mixed helix in the 2nd branch; h₃[/]: mixed helix in the 3rd branch); b: bulge; L: loop (L₁: loop in the 1st branch; L₂: loop in the 2nd branch; L₃: loop in the 3rd branch); B: common bulge.

Figure 2b: Secondary structure of tRNA^{Arg}(CCT) / tRNA^{Trp}(CCA) along with the BHB of their introns of *M. kandleri*.

NCIs: Noncanonical intron start position; NCIE: Noncanonical intron end position; CIs: Canonical intron start position; CIE: Canonical intron end position

—————> This signifies splicing sites of the introns in pre-tRNAs

h_e: Exonic helix; h[/]: mixed helix; L : loop; b: bulge.

Figure 2c: Secondary structure of tRNA^{Ile}(UAU) / tRNA^{Phe}(GAA) along with the BHB of their introns of *M. kandleri*.

NCIs: Noncanonical intron start position; NCIE: Noncanonical intron end position; CIs: Canonical intron start position; CIE: Canonical intron end position

—————> This signifies splicing sites of the introns in pre-tRNAs

h_e: Exonic helix; h[/]: mixed helix; L : loop; b: bulge.

Figure 2d: Secondary structure of tRNA^{Pro}(CGG) / tRNA^{Trp}(CCA) along with the BHB of their introns of *M. jannaschii*.

NCIs: Noncanonical intron start position; NCie: Noncanonical intron end position; CIs: Canonical intron start position; Cie: Canonical intron end position

————▶ This signifies splicing sites of the introns in pre-tRNAs

h_e : Exonic helix; h' : mixed helix (part of it is intronic and part of it is exonic) (h_1' : mixed helix in the 1st branch; h_2' : mixed helix in the 2nd branch); b: bulge; L: loop (L_1 : loop in the 1st branch; L_2 : loop in the 2nd branch); B: common bulge.