

Network growth approach to macroevolution

Shao-Meng Qin, Yong Chen[†], and Pan Zhang

Institute of Theoretical Physics, Lanzhou University, Lanzhou 730000, China

Abstract. We propose a novel coarse-grained network growth model coupled with the competition interaction to simulate the macroevolution. Our work shows that the competition plays an important role in the macroevolution and it is more rational to describe the relationship between species by network structures. Our model presents a complete picture of the development of phyla and associated splitting process. It was found that periodic mass extinction occurred in our networks without any extraterrestrial factors and the lifetime distribution of species was very close to that seen in the fossil record. We also perturbed the networks with two scenarios of mass extinctions on different hierarchic levels to study the recovery.

PACS numbers: 87.23.-n, 89.75.Hc, 07.05.-t

[†] To whom correspondence should be addressed. Email: ychen@lzu.edu.cn

1. INTRODUCTION

The ecosystem, which is formed from a myriad of interactions between various species, is one of the best-known example of complexity. During last decade, theoretical research on the coevolution of species and the statistics of extinctions has been strongly influenced by the pioneering interdisciplinary works of Per Bak and his collaborators [1, 2, 3, 4]. Many simple and delicate models have explained large numbers of the phenomena exhibited in the fossil record. Notably, with the development of nonlinear dynamics and complex networks, it was found that a food web model is the most suitable way to describe the ecosystem. More recent models based on network structure and various types of interaction have presented convincing results not only for macroevolution but also for microevolution. Although microevolution, which focuses on the influence one species has on others, is more important in protecting the environment, macroevolution, which focuses on species coevolution and periodic species extinction, is more interesting and more important for species diversity.

Many remarkable phenomena in the history have been found in the fossil record but until now no satisfactory explanation has been presented. These phenomena include the Cambrian explosion that gave rise to most of all the known animal phyla, the periodicity of mass extinctions and characteristics of the evolution, such as extinction rate, origination rate, percent extinction, standing diversity, survivorship and lifetime distribution.

Recently, there have been attempts to study macroevolution using models equipped with dynamics that operate at the level of individuals [5, 6, 7]. Lotka-Volterra models are relatively successful in describing many aspects of population dynamics. Coppex introduced a simple two-trophic network to describe an ecosystem with N species of predators and a single prey species [8]. His numerical simulations show that there is a power-law distribution of intervals between extinctions. Since the Lotka-Volterra model is a differential equation model, it is hard to simulate the global-level processes with a large value of N . Furthermore, his model does not describe the macroevolution following the beginning of the Cambrian. Luz-Burgoa and his cooperators investigated the process of sympatric speciation in a simple food web model [9]. They found that sympatric speciation is obtained for the top species in both cases, and their results suggested that the speciation velocity depends on how far up in the food chain the focus population is feeding. However, this model investigated only the cause of species speciation and cannot explain the splitting process of a phylum.

B.F De Blasio and F.V. De Blasio have introduced competition interaction into the computer model of the influence of ecospace colonization on adaptive radiation designed by J.W. Valentine and T.D. Walker [10, 11]. This macroevolution model presents a clear picture of the simultaneous appearance of so many phyla in the Cambrian almost at the same time and post-explosion evolution of the ecosystem. However this model cannot show an accurate picture of macroevolution. It lacks periodicity extinction and the results presented did not describe whether this model is consistent with the characteristics of the real fossil record. Moreover, this competition rule seems too simple to describe the intricate interaction between different species.

In fact, such interspecies competitive interactions play a key role in macroevolution

whether considered as the sum of predator-prey interaction, host-parasite interaction or competitions between species using the same resource. Competition also appears to be essential for species proliferation. Laboratory bacterial experiments have suggested that species branching is promoted by competition [12, 13]. Competition among higher taxonomical groups may also play a major role in macroevolution [14].

In this paper we present a macroevolution model also based on competitive interactions. In contrast to the model in Ref. [11], our model is built on the growth of network structures. From the simulation results of the properties of our model, such as degree distribution, clustering coefficients, and especially the modularity dynamics, one can gain more insight into the processes of creating and splitting phyla and the possible effects of mass extinction. Additionally, we observe the periodicity of mass extinctions without any extraterrestrial causes (not mentioned in the model of Ref. [11]). This phenomenon is suggested by Raup and Sepkoski in 1984 [15]. Our model indicates that this phenomenon, periodicity of mass extinctions, might be a natural consequence of the macroevolution and not the result of any extraterrestrial causes as predicted by the model of Lipowski [16, 17]. Furthermore, the simulation results of lifetime distribution are very similar to the fossil records and the normal experiences of human beings. Finally, in order to study the effects of mass extinctions, large numbers of species were removed from the network.

This article is organized as follows. We will explain how to build our model in Sec. 2. In Sec. 3, we show and discuss the simulation results, including network structure properties, lifetime distributions, extinction dynamics, and perturbations of the model. In the last Section, we summarize our conclusions and suggest further extensions of our work.

2. NETWORK MODEL

We take every species as a point in 2-dimensional morphospace and let each dimension be independent. Each dimension morphospace represents a phenotypic character of species such as shape, form and structure and so on. We select 2-dimensional morphospace is an compromise between integrated description of species and fast computation. Every species is characterized by a point in the morphospace. Species who belong to the same phylum are similar in phenotypic character, and they are close in morphospace. Phenotypic similarity between species implies that they are likely to exploit the same resources [18]. However, even species that are far morphologically may compete for common resources, like the competition for light among plants and water among animals. Long-range competition is accounted for by allowing a finite tail in the possible competitive interaction for morphologically distant species. The model dynamics are described subsequently.

(1) *Architecture of the network.* Every node in the network represents one species and the weight of the connecting line edge indicates the competition strength between two species. The evolution of the network represents evolution of all species. We give the weight on the edges following a distribution like gaussian form. Its mean is 0.5 and variance is 0.5, but we cast the random numbers larger than 1 and negative. Once an edge is built, it does not change. However, when a species becomes extinct, we remove this node and all edges connected to it

from the network.

(2) *Initialization.* At the beginning, three species with the same phenotype are connected as the initial network. Then we assign the weights of three edges.

(3) *Competition.* Since not all species could compete with each other even if they are very close in morphospace, the network model should not be a fully interconnected graph. The competition strength is equal to the weight of the edges. The extinction probability per step p_{ext} of species s_i depends on the total weight of edges connected with it. So the extinction probability of one species is defined by,

$$p_{ext}(s_i) = \varepsilon \sum_j f_{ij}, \quad (1)$$

where the f_{ij} is the weight of the edge between species s_i and s_j , and the fraction of competition ε is constant. If one species becomes extinct, it will be erased from the network immediately and other species will no longer experience competition from the extinct species.

(4) *Speciation.* In the short term, it appears to be approximately correct to say that the whole world can support a certain number of species. Modern-day ecological data on island biogeography support this view [19]. It is reasonable to assume that new species are more difficult to evolve as the species number approaches to this upper limit than that when the species is rare. When a new species originate from an old species, the distance in the morphospace between this new species and its parents is taken to be Gaussian distributed with a small variance σ . Here, the speciation range $\sigma = 0.1$ is same for all simulations. The new species are formed at each time step with a speciation rate,

$$p_{sp} = \beta \left(1 - \frac{N}{N'}\right), \quad (2)$$

where β is the fraction of the speciation rate, N' is the maximal species number in the world and N is the current species number or node number in the network. In this work, the maximal species is set to $N' = 5500$ for all simulations.

After a new species born, it becomes a part of the network, but it is isolated from other nodes. The probability $f(d_{ij})$ that the new node i will be connected to the old node j depends on the distance between them in the morphospace,

$$f(d_{ij}) = \begin{cases} f_0, & d_{ij} < R; \\ f_\infty, & d_{ij} \geq R. \end{cases} \quad (3)$$

Here the d_{ij} is the distance between species s_i and s_j . We set short-range radius $R = 2$ for all simulations, which was chosen larger than the speciation range σ so that speciation takes place within the range of the short-range competition. The f_0 should be much larger than f_∞ .

In this model, the reason of species alive is not the intrinsic morphological advantages but the morphological disparity of a species. In other words, the fitness landscape is flat. In each time step, we chose a species randomly and decide whether it would evolve into a new species based on the speciation rate. Then, we randomly selected another species and decided whether it should be removed from the network at that time step. This procedure is repeated millions of times in our simulations.

3. Simulation Result and Discussion

In this section we present our simulation results of our network model, such as, degree distribution, cluster coefficient, and community structure. Moreover, to study macroevolution we consider the lifetime distribution, the extinction dynamics, and perturbation of network growth in our simulations.

3.1. Time Evolution of Network Structure

In general, two statistical properties, degree distribution and clustering coefficient, are used to measure the structures of complex networks. However, for studying phylum splitting, we calculate the community structure of networks.

Community structure is a mesoscopic description of networks. Newman and Girvan [20, 21] refers to the fact that nodes in many real networks appear to group in subgraphs. The community in the network represent phyla. In our model, there is only one taxonomic grouping. So we can also believe community present genera, families, orders, classes, or any other taxonomic grouping. We can use this property to measure the split of the phylum. Newman and Girvan also proposed a divisive algorithm that used edge betweenness as a metric to identify the boundaries of communities and introduced a measure, based on the concept of modularity Q . This algorithm has been applied successfully to a variety of networks. Theoretically, the concept of modularity Q cannot be used in weighted networks. Even then, it is still observed the structure evolution from the different aspects via community structure in our simulations. The main reason for this observation is that in our model the connections among the nodes play a major role in the interspecies competition. Therefore we focus our attention on the structures and ignore the differences between the edges.

It should be noted that our model is a network with dynamic node number. The macroevolution is described by the process of network growth. As we will show, the main characters of our model include the species outbreak in the beginning phase, the split of the phylum, and the final saturation state. In additional, we used the algorithm to find community, as in Ref. [22] which is based on an extremal optimization of the value of modularity and is feasible for the accurate identification of community structure in large complex networks.

In Fig. 1, we present the simulation results of the time evolution of the main properties of our network growth model. The biosphere gradually stabilizes after an initial period of strong diversification. The early phase of the simulation reveals that the increase of species number is a step-like behavior. The insets of Fig. 1(b) and Fig. 1(c) show these similar behaviors of the network structure at the beginning stage. The species number in the saturation state is in direct ratio to speciation rate β and in inverse ratio to fraction of competition ε . Note that the discrete transitions for species number, clustering coefficients, and modularity occur at the same time.

Note that the definition of the modularity given by Newman, $Q = \sum_i (e_{ii} - a_i^2)$, where e_{ij} is the fraction of all edges in network that link nodes in community i to nodes in community j and $a_i = \sum_j e_{ij}$. The parameter $f_0 = 0.9$ is much larger than $f_\infty = 0.03$ in our network




Figure 1. Temporal evolution of the structural properties of the network with parameters $\beta = 0.6$, $\varepsilon = 0.015$, $f_0 = 0.9$, and $f_\infty = 0.03$. The insets show the magnification of the first 40000 steps. A step-like increase is observed in the inset picture and corresponds to the split of the phyla.

growth. So one can assume that in every community i the number of edges z_i^{in} that link nodes in the same community is much larger than the number of edges z_i^{ex} between the different communities, or the internal structure of different communities is similar. As a result, we make an estimate of the value of Q ,

$$Q = \sum_i \left(\frac{z_i^{in}}{2m} - \frac{2z_i^{in} + z_j^{ex}}{(2m)^2} \right), \quad (4)$$

where $m = \frac{1}{2} \sum_i (2z_i^{in} + z_i^{ex})$ is the number of edges in the network. Using the $z_i^{in} \gg z_i^{ex}$, and $z_i^{in} = z_j^{in}$, $z_i^{ex} = z_j^{ex}$ with $i \neq j$, we get

$$Q(n) \approx 1 - 1/n, \quad (5)$$




Figure 2. Species distribution in morphospace at four different times. Top left, 10,000 time steps after the beginning of simulation; top right, after 20,000 time steps; bottom left, after 100,000 time steps; bottom right, after 1,000,000 time steps. The figure shows the dynamics of phylum split and the phylum branching process in morphospace. The parameters of simulation are identical to those in Fig. 1.

where n is the community number in the network. As shown in Fig. 1(b), we find empirically found that at the beginning of the simulation the modularity Q at the plateau after the step increase follows a similar form to Eq. (4), where the n is the number of the sidesteps or the number of the communities.

Thus, we can present a phenomenological picture of the phylum such split that the species split into two different groups when the number of species or modularity ascends to the next plateau. As the modularity Q approaches 1, the presence of many communities in this network implies many phyla. Each phylum has a large number of species and the main competition of each species is from other species in the same phylum.

After the rapid increase in the initial phase, species proliferation decelerates to a constant value with little fluctuation. The modularity and cluster also decrease gradually while the species number increases slowly. This suggests that the number of community is increasing




Figure 3. (Color online) Degree distribution of our network model with saturation node number $N \approx 2600$ after enough steps simulation. Parameter values are $\beta = 1.0$, $\varepsilon = 0.015$, $f_0 = 0.5$, and $f_\infty = 0.01$. The solid line is a Gaussian fitting result, $P(k) = 0.00149 + \frac{1.88322}{18.47469 \times \sqrt{\pi/2}} \times \exp(-2 \times (\frac{k-68.26007}{18.47469})^2)$.

and the long-term competition surmounts the interaction of short-term. In other words, the number of phyla is increasing. In Fig. 2, we present the simulation results of the species distribution in morphospace at different times. Every point in the morphospace denotes a species. Obviously, species congregate around to form a new phylum. Furthermore we find that every species belongs to a phylum.

In order to understand the properties of the network model for biology, one must have some knowledge about the degree distribution $P(k)$, which is the probability that a node chosen uniformly at random has degree k . Note that the number of vertices of this network model is dynamic. To study $P(k)$, it should be assigned a large β relative and small other parameters to generate a network with large number of nodes. In fact, the network growth and the final saturation state of network structure are decided by two factors. One is the short-term competition, which has a large connecting probability f_0 and small number of nodes. Another is the long-term competition, which has a small connecting probability f_∞ and a large number of nodes (it is almost equal to the sum of the number of the nodes of network). In Fig. 3, we present the degree distribution of the network with $N \approx 2600$ at the final saturation state. It was found that $P(k)$ is very close to a Gaussian distribution with mean value $k_0 \approx 68$. This suggests that the degree distribution of our network model is similar to the random network with large modularity because the network growth has conspicuous stochasticity. We know that in random networks, the modularity is very small and even close to 0 in most cases. Therefore, it is difficult to categorize our network model as a type of network such as random,




Figure 4. (Color online) Histogram of lifetimes of marine genera during the Phanerozoic (black circles). The black dash line is the best power-law fit to the points between 10 and 100 My and the black solid line is the best exponential fit to all the points [23, 24]. The red and blue solid lines are our simulation results from the network growth model using two different parameters.

scale-free, or small-world.

3.2. Lifetime Distribution

One of the properties frequently studied in macroevolution models is lifetime distribution of species. Newman and Sibani mathematically derived a number of relations between the normal quantities (extinction rates, diversity, lifetime distribution, etc.) and show how these different trends are inter-related [24]. Since the lifetime distribution is easy to calculate, it becomes a canonical simulation result in most models of multispecies ecosystems.

Using simulations based on various models, many scholars have measured the lifetimes of competing species and suggested that their distribution is well approximated by a power-law form [24]. Similar estimations demonstrate that this distribution is equivalent to the power-law distribution of genus lifetimes, since the longer lived genera give rise on average to larger numbers of species [25]. However, the real fossil records are fit equally well by an exponential form [24].

In Fig. 4 we show a histogram of species lifetime distributions for the network growth model with two different parameters β . As the plot shows, the distribution closely follows an exponential law. Note that two simulation results of the lifetime distribution of species for different β are very similar. We conjecture that the lifetime distribution of species is not under the influence of the fraction of the speciation rate β , but depends on the correlation between f_0 and f_∞ (or the relation between competition within a phylum and competition for resources).




Figure 5. (Color online) Periodicity of mass extinctions without an extraterrestrial cause. Parameter values are $\beta = 0.9$, $\varepsilon = 0.05$, $f_0 = 0.9$, and $f_\infty = 0.14$. (a) Time series of the cluster coefficient, modularity Q , and species number N . (b) Fourier transform of the data in (a) after filtering the high frequency component. (c) The maximal periodicity of oscillation τ from the time series of species number N as a function of f_∞ . The other parameters are same as in(a).

Moreover, this observation emphasizes that the competitive interactions play a key role in ecological dynamics.

3.3. Extinct Dynamics

Of the estimated one to four billion species which have existed on the Earth since life first appeared here, less than 50 million are still alive today. Paleontological data, which

show broad distributions of the extinct events in the Earth's history, suggest the existence of strong correlations between extinctions [26]. Normally, the majority of researchers prefer the alternative explanation that the extinctions appear because of external stresses imposed on the ecosystems by the environment. Recently, the occurrence of extinctions in the absence of periodic external perturbation was suggested by Lipowski in a lattice model [16, 17].

Fig. 5 shows temporal evolution of N , Q , and the cluster coefficient at the saturation state of the network with parameters $\beta = 0.9$, $\varepsilon = 0.05$, $f_0 = 0.9$, and $f_\infty = 0.14$. Note that we omit the initial phase of the network growth in Fig. 5(a). The Fourier transform of the data is presented in Fig. 5(b). In order to give prominence to oscillation and compare the periodicity of N , Q and the cluster coefficient, we subtract their means and normalize the data before Fourier transform. We observe that three curves have the same peak at frequency $3.35693(\times 10^{-6}\text{Hz})$. The maximal periodicity of oscillations τ for various f_∞ is shown in Fig. 5(c). The red dashed line is a fitted result $\tau = 351410 - 375.074\exp(38.81 \times f_\infty)$.

The peak in Fig. 5(b) shows that species number N and the modularity Q and cluster coefficient of the network have same periodicity and amplitude. As is well known, the oscillation of the clustering coefficients and modularity indicates changing network structures. Note that in our network these parameters denote the phylum number in the whole world and the average species number in one phylum. Therefore this oscillation indicates an extinction at the phylum level. The oscillation of the network structure parameter reflects phylum changing of the number and the scale. Phylum which bears intense competition will decrease its scale or even that this phylum extinct. In most instances, large numbers of phyla do not routinely go extinct, and they may belong to extant groups rather than constituting distinct phyla.

It should be noted that the species number in Fig. 5(a) is very small, $N \approx 140$, due to the larger long-distance competition $f_\infty = 0.14$. In fact, by enlarging the simulation results of the saturation phase in Fig. 1 with $f_\infty = 0.03$, it was found that the similar oscillation behaviors take place in the large networks. However, the oscillation in Fig. 1 appears to be small-amplitude fluctuations.

It is clear from Fig. 5(c) that the periodicity of oscillation is a function of f_∞ . Moreover, our extensive simulation results demonstrate that there are no obvious correlation between the periodicity and other parameters. Therefore we conjecture that the long-distance competitions for public resources induce the oscillation of the species number. In the case of larger f_∞ , which means the resources are scarce or lacking, this species number oscillation is strong enough to produce similar oscillation of the modularity Q and the cluster coefficient. This demonstrates the existence of periodic extinctions of the phyla. However, smaller f_∞ or a scenario with abundant public resources would not only increase the system size N and the phylum number, but also would not only increase the system size and the phylum number, but would also decrease the probability of extinction of species and phyla. As a result, only the small fluctuation of species and phyla exists.




Figure 6. (Color online) The network growth with a random extinction at 3×10^6 steps. (a) Time series of the species number. The inset shows a magnification of the extinction region. (b) The modularity and cluster in the extinction region. The dotted line is the time of extinction. The parameters are $\beta = 0.5$, $\varepsilon = 0.015$, $f_0 = 0.7$, and $f_\infty = 0.015$.

3.4. Perturbing the Model

In order to study the effect of mass extinctions on the macroevolution within the network growth approach, a large number of species were removed from the network at the saturation state. Such a scenario is designed to mimic the influence of an external factor, such as the change of climate, the impacts of comets or meteorites, etc. We discuss the following two cases.

Case I: Random perturbation on species. In this case, half of the species were removed randomly. Such a situation is appropriate for biosphere in which the species become extinct because of 'bad luck'. Clearly, the dynamics of the network should be reduced to the saturation state quickly since each species has the same probability of extinction.

Case II: Random perturbation on community. In this case, the artificial perturbations act on the community. We remove half of the community in the network randomly. The




Figure 7. (Color online) The network growth with a random community extinction at 3×10^6 steps. (a) The time evolution of the species number. The inset shows a magnification of the extinction region. (b) The modularity and cluster dynamics in the extinction region. The dotted line is the time of extinction. The parameters are identical to those used in Fig. 6.

motivation for this case is based on the idea that some phyla have 'bad genes' and cannot adapt to the sudden change of environment.

Figs. 6-7 show the simulation results of the time series of the species number and the network structure for both cases. Obviously, even though both scenarios of mass extinction act on different hierarchic levels, the species in the world will eventually recover and the response to an extinction will include a rapid expansion of species diversity. However, there is a noticeable difference, which can be seen in Fig. 6-7, in the detailed recovery process.

In case I, after removing the nodes in the network, the species number quickly returned to the pre-perturbation level (see Fig. 6(a)). Fig. 6(b) depicts the response of network structure to extinction. There is not any significant change of the modularity and cluster and both are still in normal fluctuation range.

In contrast to case I, the recovery in case II is slow and difficult. Many more steps are necessary for the species number to reach the saturation stage (see Fig. 7 (a)). However, note that there is also a sharply rising stage (as shown in the inset of Fig. 7(a)). The initial stage

of the response to the extinction, when compared with the inset of Fig. 6(a), is similar for both cases. In Fig. 7(b), we present the responses of the modularity and clustering coefficient to this extinction. These curves represent a sudden increase preceding the decrease of the diversity at the phylum level.

As demonstrated by the arguments given above, it is clear that the response of the biological evolution for two scenarios of mass extinction is completely different. Although both perturbations erase almost the same ratio of the species number, in Case I, the perturbation does not break down the structure of the network. However, in Case II, a number of whole communities are removed, or some phyla become extinct. Therefore the network structure is destroyed and the perturbation of Case II is more deleterious to the recovery of the .

The statement above, implies that forming phyla is a way for species to avoid or replace the short-term competition in order to increase the species number in the long-term. Therefore it is much easier for species to conquer short-term competition for speciation than to develop long-term competition. In fact, in Case I, removing the nodes in the network decreases both long- and short- term competition. In fact, in Case I, removing the nodes in the network decreases both long- and short- term competition. So Case I exactly takes the place for new species. These new species did not need to develop new phyla, which explains why the recovery after disturbance is so fast. In contrast, Case II only decreases the long-term competition. The new species will develop in their phylum because the absence of long-term competition decreases the short-term competition below saturation and the small σ make the new species remain within their phylum. After this stage, when every phylum is full of species and reaches saturation, the new species must keep away from the phylum to decrease short-term competition. These new species will develop new phyla which are booming and forcing old phyla until they have the same scale with the existed phyla. Thus, the behaviors of the modularity and cluster are different in the two scenarios following disturbances, and the species number increases slowly to the saturation level after the initial sharply rising stage.

4. Conclusion

In this paper, we present a simple network growth model with competitive interactions to approximate the biological evolution. This model is characterized by four tunable parameters: the fraction of competition ε , the fraction of speciation rate β , the short-term competition f_0 , and the long-term competition f_∞ . We start with a few species, and then let our model grow in diversity and complexity until it reaches the saturation state. In our simulation, the species number is much smaller than the limitation. It should be noted that the limitation plays an important role in the evolution. Without this parameter, periodical extinction can not be displayed. These simulation results are not presented in this paper.

Based on the simulation results of our network growth model, we demonstrate several different aspects of biological evolution, such as the species number, phyla, cluster, the lifetime distribution, etc. With simulations up to sufficiently long times (at least several 10^6 steps), one can observe a semi-periodic mass extinction from the combination of the close

phyla. Finally, the network model is perturbed in two different ways: random perturbation species and random perturbation on community. The effects of these different extinction scenarios on the two taxonomic hierarchic levels, species and phyla, may be useful for interpretation of the causes of historical mass extinction.

Our model is aimed at emphasizing the importance of the competition and network structure in the biological evolution. Competition not only decides the species number in the whole system, but also makes the species split to different phyla. As there are two kinds of different competitions, the response of the biological evolution for two mass extinctions is completely different. Competition coupled with network structure drive the number of species to fluctuate periodically.

Although many of the properties shown in our model are similar to the fossil records especially the lifetime distribution, it is difficult to compare the model with the actual history of phyla diversities and real world. Since this model is a macroevolution model, it cannot include all the properties and details of the biology.

Further extensions of our model offer interesting opportunities. For example, one can change the form of the competing function or differently implement the interaction among species. This could complicate the simulation tremendously, and we believe that the results would have the similar qualitative properties. In the real world, the climate switches between the ice age and warmth with a period of about 10,000 years[27]. Comparably, our model could also use the periodic change parameters instead of the constant form. Such a modification would likely result in other thought-provoking phenomena. However this modification is not currently reasonable because we cannot assess whether the changing climate is strong enough to affect the parameters in our model or whether a relationship exists between the climate and the model.

Acknowledgments

This work was supported in part by the National Natural Science Foundation of China under Grant No. 10305005 and by the Special Fund for Doctor Programs at Lanzhou University.

References

- [1] P. Bak and K. Sneppen, Phys. Rev. Lett. **71**, 4083(1993); M. Paczuski, S. Maslov, and P. Bak, Phys. Rev. E **53**, 414(1996).
- [2] R. V. Solé, S. C. Manrubia, M. Benton, and P. Bak, Nature **388**, 764(1997).
- [3] P. Bak and S. Boettcher, Physica D **107**, 143(1997).
- [4] R. V. Solé, S. C. Manrubia, M. Benton, S. Kauffman, and P. Bak, *Trends in Ecology and Evolution*, **14**, 156(1999).
- [5] D. Chowdhury, D. Stauffer, and A. Kunwar, Phys. Rev. Lett. **90**, 068101(2003).
- [6] P. A. Rikvold and R. K. P. Zia, Phys. Rev. E. **68**, 031913(2003).
- [7] M. Hall, K. Christensen, S. A. di Collobiano, and H. J. Jensen Phys. Rev. E **66**, 011904(2002).
- [8] F. Coppex, M. Droz, and A. Lipowski, Phys. Rev. E **69**, 061901(2004).
- [9] K. Luz-Burgoa, T. Dell, and S. M. de Oliveira, Phys. Rev. E **72**, 011914(2005).
- [10] J. W. Valentine, Paleobiology **6**, 444(1980); J. W. Valentine and T. D. Walker, Physica D **22**, 31(1986).

- [11] B. F. De Blasio and F. V. De Blasio, *Phy. Rev. E* **72**, 031916(2005).
- [12] R. E. Lenski and M. Travisano, *Proc. Natl. Acad. Sci. U.S.A.* **91**, 6808(1994).
- [13] D. Schluter, *Science* **266**, 798(1994).
- [14] M. J. Benton, *Biol. Rev. Cambridge Philos. Soc.* **62**, 305(1987); J. J. Sepkoski, in *Paleobiology II: A Synthesis*, edited by D.E. G. Briggs and P. R. Crowther (Blackwell Science, Malden, 2000).
- [15] D. M. Raup and J. J. Sepkoski, *Proc. Natl. Acad. Sci. U.S.A.* **81**, 801(1984).
- [16] A. Lipowski, *Phy. Rev. E* **71**, 052902(2005).
- [17] A. Lipowski and D. Lipowska, *Theor. Biosci.* **125**, 67(2006).
- [18] A. R. H. Swan, in *Paleobiology II: A Synthesis*.
- [19] M. L. Rosenzweig *Species Diversity in Space and Time*. Cambridge University Press (Cambridge).(1995)
- [20] M. E. J. Newman and M. Girvan, *Phy. Rev. E* **69**, 026113(2004).
- [21] M. Girvan and M. E. J. Newman, *Proc. Natl. Acad. Sci. U.S.A.* **99**, 7821(2002).
- [22] J. Duch and A. Arenas, *Phy. Rev. E* **72**, 027104(2005).
- [23] J. J. Jr Sepkoski, *Milwaukee Public Museum Contributions in Biology and Geology* **83**, (1992).
- [24] M. E. J. Newman and P. Sibani, *Proc. R. Soc. Lond. B* **266**, 1593(1999). B. Burlando, *J. Theor. Biol.* **146**, 99(1990); M. E. J. Newman and K. Sneppen, *Phys. Rev. E* **54**, 6226(1996).
- [25] C. Adami, *Phys. Lett. A* **203**, 29(1995); K. Sneppen, P. Bak, H. Flyvbjerg, and M. H. Jensen, *Proc. Nat. Acad. Sci.* **92**, 5209(1995); M. E. J. Newman, *Physica D* **107**, 293(1997).
- [26] M. E. J. Newman and R. G. Palmer, *Modelling Extinction* (Oxford University Press, New York, 2003); M. E. J. Newman and R. G. O. Palmer, e-print adap-org/9908002.
- [27] P. M. Grootes and M. Stuiver, *J. Geophys. Res.* **102**, 26455(1997).