# SUPPLEMENTAL MATERIAL

TABLE I: parameters in G-protein cycle

parameter	value	References and Notes
k1	$0.12nM^{-1}min^{-1}$	$2 \times 10^6 M^{-1} s^{-1} $ (12)
k2	$0.6min^{-1}$	$0.01 \ s^{-1} \ (12)$
k3	$0.24min^{-1}$	$4 \times 10^{-3} s^{-1} $ (12)
k4	$320nM \cdot min^{-1}$	fit to the time-course of G protein activation
k5	7 nM	(12).(See Fig.2A )
k6	$30nM \cdot min^{-1}$	$4(molecules/cell)s^{-1}$ (12)
k7	$0.024min^{-1}$	$4 \times 10^{-4} s^{-1} $ (12)
k8	$0.0048min^{-1}$	$10^{-5} (molecules/cell)^{-1} s^{-1} $ (12)
k9	$115nM \cdot min^{-1}$	fit to the time-course of G protein activation
k10	60 nM	(12)(See Fig.2A),
k11	$20nM \cdot min^{-1}$	and experimental data that G protein increased from
k12	$0.08min^{-1}$	8000 molecules/cell to 12000 molecules/cell (8)
k13	$0.24min^{-1}$	$4 \times 10^{-3} s^{-1} $ (12)
k14	$0.025nM^{-1}min^{-1}$	$0.11s^{-1}$ (12)
k15	$480nM^{-1}min^{-1}$	$1(molecule/cell)^{-1}s^{-1}(12)$

TABLE II: parameters in recruitment of scaffolds

parameter	value	References and Notes
k16	$0.05nM^{-1}min^{-1}$	$t_{1/2} = 8.22s$ for Ste5s' recovery
k17	$5min^{-1}$	at shmoo tips (49)
k18	$0.00007nM^{-1}min^{-1}$	fit to (24)
k19	$0.001 min^{-1}$	(See Fig.2C. )
k23	$30min^{-1}$	$t_{1/2} = 2s$ for the recovery of ste5 in the nucleus.
		(49)

TABLE III: dephosphorylation

protein	a1	d1	p1	a2	d2	p2
	$(nM^{-1}min^{-1})$	$(min^{-1})$	$(min^{-1})$	$(nM^{-1}min^{-1})$	$(min^{-1})$	$(min^{-1})$
Ste11(MAPKKK)	0.5	24	10	1	24	10
Ste7(MAPKK)	0.5	24	10	0.5	24	10
Fus3(MAPK)	0.2	10	30	0.4	20	30

TABLE IV: phosphorylation in cytosol

protein	a3	d3	р3	a4	d4	p4
	$(nM^{-1}min^{-1})$	$(min^{-1})$	$(min^{-1})$	$(nM^{-1}min^{-1})$	$(min^{-1})$	$(min^{-1})$
Ste7(MAPKF	<ol> <li>1</li> </ol>	36	10	1	36	10
Fus3(MAPK	0.1	36	10	0.1	36	10

Notes: Equations of reaction in this part are all in the following form:  $S + E \stackrel{a_n}{\rightleftharpoons} SE \stackrel{p_n}{\Longrightarrow} S^* + E$ .

### As for the underset n:

1: dephosphorylation of the once phosphorylated kinase; 2: dephosphorylation of the twice phosphorylated kinase; 3: once phosphorylation; 4: twice phosphorylation. Parameters are estimated in analogy to Ref.11.

TABLE V: Other reactions in the cytosol

parameter	value	References
k24	$1.2 \ nM^{-1}min^{-1}$	Ref.23
k25	$24min^{-1}$	Ref.23
k26	$0.0015 \ nM^{-1}min^{-1}$	Ref.40
k27	$0.1 \ min^{-1}$	Ref.41




FIG. 1: 27 kinds of scaffold-kinase complexes in solution and 27 kinds of scaffold-kinase complexes at the membrane

$B1 + Ste11 \rightleftharpoons B2$	$B1 + Stellpp \rightleftharpoons B3$	$B1 + Ste7 \rightleftharpoons B4$
$B1 + Ste7pp \leftarrow B5$	$B1 + Fus3 \rightleftharpoons B6$	$B1 + Fus3pp \leftarrow B7$
$B2 + Ste7 \rightleftharpoons B8$	$B2 + Ste7pp \leftarrow B11$	$B2 + Fus3 \rightleftharpoons B12$
$B2 + Fus3pp \leftarrow B15$	$B3 + Ste7 \rightleftharpoons B9$	$B3 + Ste7pp \leftarrow B10$
$B3 + Fus3 \rightleftharpoons B13$	$B3 + Fus3pp \leftarrow B14$	$B4 + Ste11 \rightleftharpoons B8$
$B4 + Stellpp \rightleftharpoons B9$	$B4 + Fus3 \rightleftharpoons B16$	$B4 + Fus3pp \leftarrow B19$
$B5 + Fus3pp \rightleftharpoons B10$	$B5 + Ste11 \rightleftharpoons C11$	$B5 + Fus3 \rightleftharpoons B17$
$B5 + Fus3pp \leftarrow B18$	$B6 + Ste11 \rightleftharpoons B12$	$B6 + Ste11pp \rightleftharpoons B13$
$B6 + Ste7 \rightleftharpoons B16$	$B6 + Ste7pp \leftarrow B17$	$B7 + Ste11pp \rightleftharpoons B14$
$B7 + Ste11 \rightleftharpoons B15$	$B7 + STe7pp \leftarrow B18$	$B7 + Ste7 \rightleftharpoons B19$
$B8 + Fus3 \rightleftharpoons B20$	$B8 + Fus3pp \leftarrow B23$	$B9 \to B10$
$B9 + Fus3 \rightleftharpoons B21$	$B9 + Fus3pp \leftarrow B25$	$B10 + Fus3 \rightleftharpoons B24$
$B10 + Fus3pp \leftarrow B27$	$B11 + Fus3 \rightleftharpoons B22$	$B11 + Fus3pp \leftarrow B26$
$B12 + Ste7 \rightleftharpoons B20$	$B12 + Ste7pp \leftarrow B22$	$B13 + Ste7 \rightleftharpoons B21$
$B13 + Ste7pp \leftarrow B24$	$B14 + Ste7 \rightleftharpoons B25$	$B14 + Ste7pp \leftarrow B27$
$B15 + Ste7 \rightleftharpoons B23$	$B15 + Ste7pp \leftarrow B26$	$B16 + Ste11pp \rightleftharpoons B21$
$B16 + Ste11 \rightleftharpoons B20$	$B17 \rightarrow B18$	$B17 + Ste11pp \rightleftharpoons B24$
$B17 + Ste11 \rightleftharpoons B22$	$B18 + Stel1pp \rightleftharpoons B27$	$B18 + Ste11 \rightleftharpoons B26$
$B19 + Stel1pp \rightleftharpoons B25$	$B19 + Ste11 \rightleftharpoons B23$	$B21 \rightarrow B24$
$B22 \rightarrow B26$	$B24 \rightarrow B27$	$B25 \rightarrow B27$
$C1 + Ste11 \rightleftharpoons C2$	$C1 + Stel11pp \rightleftharpoons C3$	$C1 + Ste7 \rightleftharpoons C4$
$C1 + Ste7pp \leftarrow C5$	$C1 + Fus3 \rightleftharpoons C6$	$C1 + Fus3pp \leftarrow C7$
$C2 \rightarrow C3$	$C2 + Ste7 \rightleftharpoons C8$	$C2 + Ste7pp \leftarrow C11$
$C2 + Fus3 \rightleftharpoons C12$	$C2 + Fus3pp \leftarrow C15$	$C3 + Ste7 \rightleftharpoons C9$
$C3 + Ste7pp \leftarrow C10$	$C3 + Fus3 \rightleftharpoons C13$	$C3 + Fus3pp \leftarrow C14$
$C4 + Ste11 \rightleftharpoons C8$	$C4 + Ste11pp \rightleftharpoons C9$	$C4 + Fus3 \rightleftharpoons C16$
$C4 + Fus3pp \leftarrow C19$	$C5 + Fus3pp \rightleftharpoons C10$	$C5 + Ste11 \rightleftharpoons C11$
$C5 + Fus3 \rightleftharpoons C17$	$C5 + Fus3pp \leftarrow C18$	$C6 + Ste11 \rightleftharpoons C12$
$C6 + Stellpp \rightleftharpoons C13$	$C6 + Ste7 \rightleftharpoons C16$	$C6 + Ste7pp \leftarrow C17$
$C7 + Ste11pp \rightleftharpoons C14$	$C7 + Ste11 \rightleftharpoons C15$	$C7 + STe7pp \leftarrow C18$
$C7 + Ste7 \rightleftharpoons C19$	$C8 \rightarrow C9$	G0 . E 0 G00
	00 / 00	$C8 + Fus3 \rightleftharpoons C20$
$C8 + Fus3pp \leftarrow C23$	$C9 \rightarrow C10$	$C8 + Fus3 \rightleftharpoons C20$ $C9 + Fus3 \rightleftharpoons C21$
$C8 + Fus3pp \leftarrow C23$	$C9 \rightarrow C10$	$C9 + Fus3 \rightleftharpoons C21$
$C8 + Fus3pp \leftarrow C23$ $C9 + Fus3pp \leftarrow C25$	$C9 \rightarrow C10$ $C10 \leftarrow C11$	$C9 + Fus3 \rightleftharpoons C21$ $C10 + Fus3 \rightleftharpoons C24$
$C8 + Fus3pp \leftarrow C23$ $C9 + Fus3pp \leftarrow C25$ $C10 + Fus3pp \leftarrow C27$	$C9 \rightarrow C10$ $C10 \leftarrow C11$ $C11 + Fus3 \rightleftharpoons C22$	$C9 + Fus3 \rightleftharpoons C21$ $C10 + Fus3 \rightleftharpoons C24$ $C11 + Fus3pp \leftarrow C26$
$C8 + Fus3pp \leftarrow C23$ $C9 + Fus3pp \leftarrow C25$ $C10 + Fus3pp \leftarrow C27$ $C12 \leftarrow C13$	$C9 \rightarrow C10$ $C10 \leftarrow C11$ $C11 + Fus3 \rightleftharpoons C22$ $C12 + Ste7 \rightleftharpoons C20$	$C9 + Fus3 \rightleftharpoons C21$ $C10 + Fus3 \rightleftharpoons C24$ $C11 + Fus3pp \leftarrow C26$ $C12 + Ste7pp \leftarrow C22$
$C8 + Fus3pp \leftarrow C23$ $C9 + Fus3pp \leftarrow C25$ $C10 + Fus3pp \leftarrow C27$ $C12 \leftarrow C13$ $C13 + Ste7 \rightleftharpoons C21$	$C9 \rightarrow C10$ $C10 \leftarrow C11$ $C11 + Fus3 \rightleftharpoons C22$ $C12 + Ste7 \rightleftharpoons C20$ $C13 + Ste7pp \leftarrow C24$	$C9 + Fus3 \rightleftharpoons C21$ $C10 + Fus3 \rightleftharpoons C24$ $C11 + Fus3pp \leftarrow C26$ $C12 + Ste7pp \leftarrow C22$ $C14 \leftarrow C15$
$C8 + Fus3pp \leftarrow C23$ $C9 + Fus3pp \leftarrow C25$ $C10 + Fus3pp \leftarrow C27$ $C12 \leftarrow C13$ $C13 + Ste7 \rightleftharpoons C21$ $C14 + Ste7 \rightleftharpoons C25$	$C9 \rightarrow C10$ $C10 \leftarrow C11$ $C11 + Fus3 \rightleftharpoons C22$ $C12 + Ste7 \rightleftharpoons C20$ $C13 + Ste7pp \leftarrow C24$ $C14 + Ste7pp \leftarrow C27$	$C9 + Fus3 \rightleftharpoons C21$ $C10 + Fus3 \rightleftharpoons C24$ $C11 + Fus3pp \leftarrow C26$ $C12 + Ste7pp \leftarrow C22$ $C14 \leftarrow C15$ $C15 + Ste7 \rightleftharpoons C23$ $C16 + Ste11 \rightleftharpoons C20$
$C8 + Fus3pp \leftarrow C23$ $C9 + Fus3pp \leftarrow C25$ $C10 + Fus3pp \leftarrow C27$ $C12 \leftarrow C13$ $C13 + Ste7 \rightleftharpoons C21$ $C14 + Ste7 \rightleftharpoons C25$ $C15 + Ste7pp \leftarrow C26$	$C9 \rightarrow C10$ $C10 \leftarrow C11$ $C11 + Fus3 \rightleftharpoons C22$ $C12 + Ste7 \rightleftharpoons C20$ $C13 + Ste7pp \leftarrow C24$ $C14 + Ste7pp \leftarrow C27$ $C16 + Ste11pp \rightleftharpoons C21$	$C9 + Fus3 \rightleftharpoons C21$ $C10 + Fus3 \rightleftharpoons C24$ $C11 + Fus3pp \leftarrow C26$ $C12 + Ste7pp \leftarrow C22$ $C14 \leftarrow C15$ $C15 + Ste7 \rightleftharpoons C23$ $C16 + Ste11 \rightleftharpoons C20$
$C8 + Fus3pp \leftarrow C23$ $C9 + Fus3pp \leftarrow C25$ $C10 + Fus3pp \leftarrow C27$ $C12 \leftarrow C13$ $C13 + Ste7 \rightleftharpoons C21$ $C14 + Ste7 \rightleftharpoons C25$ $C15 + Ste7pp \leftarrow C26$ $C17 \rightarrow C18$	$C9 \rightarrow C10$ $C10 \leftarrow C11$ $C11 + Fus3 \rightleftharpoons C22$ $C12 + Ste7 \rightleftharpoons C20$ $C13 + Ste7pp \leftarrow C24$ $C14 + Ste7pp \leftarrow C27$ $C16 + Ste11pp \rightleftharpoons C21$ $C17 + Ste11pp \rightleftharpoons C24$	$C9 + Fus3 \rightleftharpoons C21$ $C10 + Fus3 \rightleftharpoons C24$ $C11 + Fus3pp \leftarrow C26$ $C12 + Ste7pp \leftarrow C22$ $C14 \leftarrow C15$ $C15 + Ste7 \rightleftharpoons C23$ $C16 + Ste11 \rightleftharpoons C20$ $C17 + Ste11 \rightleftharpoons C22$
$C8 + Fus3pp \leftarrow C23$ $C9 + Fus3pp \leftarrow C25$ $C10 + Fus3pp \leftarrow C27$ $C12 \leftarrow C13$ $C13 + Ste7 \rightleftharpoons C21$ $C14 + Ste7 \rightleftharpoons C25$ $C15 + Ste7pp \leftarrow C26$ $C17 \rightarrow C18$ $C18 + Ste11pp \rightleftharpoons C27$	$C9 \rightarrow C10$ $C10 \leftarrow C11$ $C11 + Fus3 \rightleftharpoons C22$ $C12 + Ste7 \rightleftharpoons C20$ $C13 + Ste7pp \leftarrow C24$ $C14 + Ste7pp \leftarrow C27$ $C16 + Ste11pp \rightleftharpoons C21$ $C17 + Ste11pp \rightleftharpoons C24$ $C18 + Ste11 \rightleftharpoons C26$	$C9 + Fus3 \rightleftharpoons C21$ $C10 + Fus3 \rightleftharpoons C24$ $C11 + Fus3pp \leftarrow C26$ $C12 + Ste7pp \leftarrow C22$ $C14 \leftarrow C15$ $C15 + Ste7 \rightleftharpoons C23$ $C16 + Ste11 \rightleftharpoons C20$ $C17 + Ste11 \rightleftharpoons C22$ $C19 + Ste11pp \rightleftharpoons C25$
$C8 + Fus3pp \leftarrow C23$ $C9 + Fus3pp \leftarrow C25$ $C10 + Fus3pp \leftarrow C27$ $C12 \leftarrow C13$ $C13 + Ste7 \rightleftharpoons C21$ $C14 + Ste7 \rightleftharpoons C25$ $C15 + Ste7pp \leftarrow C26$ $C17 \rightarrow C18$ $C18 + Ste11pp \rightleftharpoons C27$ $C19 + Ste11 \rightleftharpoons C23$ $C22 \rightarrow C24$ $C24 \rightarrow C27$	$C9 \rightarrow C10$ $C10 \leftarrow C11$ $C11 + Fus3 \rightleftharpoons C22$ $C12 + Ste7 \rightleftharpoons C20$ $C13 + Ste7pp \leftarrow C24$ $C14 + Ste7pp \leftarrow C27$ $C16 + Ste11pp \rightleftharpoons C21$ $C17 + Ste11pp \rightleftharpoons C24$ $C18 + Ste11 \rightleftharpoons C26$ $C20 \rightarrow C21$ $C22 \rightarrow C26$ $C25 \rightarrow C27$	$\begin{array}{l} C9 + Fus3 \rightleftharpoons C21 \\ C10 + Fus3 \rightleftharpoons C24 \\ C11 + Fus3pp \leftarrow C26 \\ C12 + Ste7pp \leftarrow C22 \\ C14 \leftarrow C15 \\ C15 + Ste7 \rightleftharpoons C23 \\ C16 + Ste11 \rightleftharpoons C20 \\ C17 + Ste11 \rightleftharpoons C22 \\ C19 + Ste11pp \rightleftharpoons C25 \\ C21 \rightarrow C24 \\ C23 \rightarrow C25 \\ C26 \rightarrow C27 \\ \end{array}$
$C8 + Fus3pp \leftarrow C23$ $C9 + Fus3pp \leftarrow C25$ $C10 + Fus3pp \leftarrow C27$ $C12 \leftarrow C13$ $C13 + Ste7 \rightleftharpoons C21$ $C14 + Ste7 \rightleftharpoons C25$ $C15 + Ste7pp \leftarrow C26$ $C17 \rightarrow C18$ $C18 + Ste11pp \rightleftharpoons C27$ $C19 + Ste11 \rightleftharpoons C23$ $C22 \rightarrow C24$ $C24 \rightarrow C27$ $B1 \rightleftharpoons C1$	$C9 \rightarrow C10$ $C10 \leftarrow C11$ $C11 + Fus3 \rightleftharpoons C22$ $C12 + Ste7 \rightleftharpoons C20$ $C13 + Ste7pp \leftarrow C24$ $C14 + Ste7pp \leftarrow C27$ $C16 + Ste11pp \rightleftharpoons C21$ $C17 + Ste11pp \rightleftharpoons C24$ $C18 + Ste11 \rightleftharpoons C26$ $C20 \rightarrow C21$ $C22 \rightarrow C26$ $C25 \rightarrow C27$ $B2 \rightleftharpoons C2$	$\begin{array}{l} C9 + Fus3 \rightleftharpoons C21 \\ C10 + Fus3 \rightleftharpoons C24 \\ C11 + Fus3pp \leftarrow C26 \\ C12 + Ste7pp \leftarrow C22 \\ C14 \leftarrow C15 \\ C15 + Ste7 \rightleftharpoons C23 \\ C16 + Ste11 \rightleftharpoons C20 \\ C17 + Ste11 \rightleftharpoons C22 \\ C19 + Ste11pp \rightleftharpoons C25 \\ C21 \rightarrow C24 \\ C23 \rightarrow C25 \\ C26 \rightarrow C27 \\ B3 \rightleftharpoons C3 \\ \end{array}$
$C8 + Fus3pp \leftarrow C23$ $C9 + Fus3pp \leftarrow C25$ $C10 + Fus3pp \leftarrow C27$ $C12 \leftarrow C13$ $C13 + Ste7 \rightleftharpoons C21$ $C14 + Ste7 \rightleftharpoons C25$ $C15 + Ste7pp \leftarrow C26$ $C17 \rightarrow C18$ $C18 + Ste11pp \rightleftharpoons C27$ $C19 + Ste11 \rightleftharpoons C23$ $C22 \rightarrow C24$ $C24 \rightarrow C27$ $B1 \rightleftharpoons C1$ $B4 \rightleftharpoons C4$	$C9 \rightarrow C10$ $C10 \leftarrow C11$ $C11 + Fus3 \rightleftharpoons C22$ $C12 + Ste7 \rightleftharpoons C20$ $C13 + Ste7pp \leftarrow C24$ $C14 + Ste7pp \leftarrow C27$ $C16 + Ste11pp \rightleftharpoons C21$ $C17 + Ste11pp \rightleftharpoons C24$ $C18 + Ste11 \rightleftharpoons C26$ $C20 \rightarrow C21$ $C22 \rightarrow C26$ $C25 \rightarrow C27$ $B2 \rightleftharpoons C2$ $B5 \rightleftharpoons C5$	$C9 + Fus3 \rightleftharpoons C21$ $C10 + Fus3 \rightleftharpoons C24$ $C11 + Fus3pp \leftarrow C26$ $C12 + Ste7pp \leftarrow C22$ $C14 \leftarrow C15$ $C15 + Ste7 \rightleftharpoons C23$ $C16 + Ste11 \rightleftharpoons C20$ $C17 + Ste11 \rightleftharpoons C22$ $C19 + Ste11pp \rightleftharpoons C25$ $C21 \rightarrow C24$ $C23 \rightarrow C25$ $C26 \rightarrow C27$ $B3 \rightleftharpoons C3$ $B6 \rightleftharpoons C6$
$C8 + Fus3pp \leftarrow C23$ $C9 + Fus3pp \leftarrow C25$ $C10 + Fus3pp \leftarrow C27$ $C12 \leftarrow C13$ $C13 + Ste7 \rightleftharpoons C21$ $C14 + Ste7 \rightleftharpoons C25$ $C15 + Ste7pp \leftarrow C26$ $C17 \rightarrow C18$ $C18 + Ste11pp \rightleftharpoons C27$ $C19 + Ste11 \rightleftharpoons C23$ $C22 \rightarrow C24$ $C24 \rightarrow C27$ $B1 \rightleftharpoons C1$ $B4 \rightleftharpoons C4$ $B7 \rightleftharpoons C7$	$C9 \rightarrow C10$ $C10 \leftarrow C11$ $C11 + Fus3 \rightleftharpoons C22$ $C12 + Ste7 \rightleftharpoons C20$ $C13 + Ste7pp \leftarrow C24$ $C14 + Ste7pp \leftarrow C27$ $C16 + Ste11pp \rightleftharpoons C21$ $C17 + Ste11pp \rightleftharpoons C24$ $C18 + Ste11 \rightleftharpoons C26$ $C20 \rightarrow C21$ $C22 \rightarrow C26$ $C25 \rightarrow C27$ $B2 \rightleftharpoons C2$ $B5 \rightleftharpoons C5$ $B8 \rightleftharpoons C8$	$C9 + Fus3 \rightleftharpoons C21$ $C10 + Fus3 \rightleftharpoons C24$ $C11 + Fus3pp \leftarrow C26$ $C12 + Ste7pp \leftarrow C22$ $C14 \leftarrow C15$ $C15 + Ste7 \rightleftharpoons C23$ $C16 + Ste11 \rightleftharpoons C20$ $C17 + Ste11 \rightleftharpoons C22$ $C19 + Ste11pp \rightleftharpoons C25$ $C21 \rightarrow C24$ $C23 \rightarrow C25$ $C26 \rightarrow C27$ $C3 \rightleftharpoons C3$ $C3 \rightleftharpoons C3$ $C4 \rightleftharpoons C6$ $C5 \rightleftharpoons C9$
$C8 + Fus3pp \leftarrow C23$ $C9 + Fus3pp \leftarrow C25$ $C10 + Fus3pp \leftarrow C27$ $C12 \leftarrow C13$ $C13 + Ste7 \rightleftharpoons C21$ $C14 + Ste7 \rightleftharpoons C25$ $C15 + Ste7pp \leftarrow C26$ $C17 \rightarrow C18$ $C18 + Ste11pp \rightleftharpoons C27$ $C19 + Ste11 \rightleftharpoons C23$ $C22 \rightarrow C24$ $C24 \rightarrow C27$ $B1 \rightleftharpoons C1$ $B4 \rightleftharpoons C4$ $B7 \rightleftharpoons C7$ $B10 \rightleftharpoons C10$	$C9 \rightarrow C10$ $C10 \leftarrow C11$ $C11 + Fus3 \rightleftharpoons C22$ $C12 + Ste7 \rightleftharpoons C20$ $C13 + Ste7pp \leftarrow C24$ $C14 + Ste7pp \leftarrow C27$ $C16 + Ste11pp \rightleftharpoons C21$ $C17 + Ste11pp \rightleftharpoons C24$ $C18 + Ste11 \rightleftharpoons C26$ $C20 \rightarrow C21$ $C22 \rightarrow C26$ $C25 \rightarrow C27$ $B2 \rightleftharpoons C2$ $B5 \rightleftharpoons C5$ $B8 \rightleftharpoons C8$ $B11 \rightleftharpoons C11$	$C9 + Fus3 \rightleftharpoons C21$ $C10 + Fus3 \rightleftharpoons C24$ $C11 + Fus3pp \leftarrow C26$ $C12 + Ste7pp \leftarrow C22$ $C14 \leftarrow C15$ $C15 + Ste7 \rightleftharpoons C23$ $C16 + Ste11 \rightleftharpoons C20$ $C17 + Ste11 \rightleftharpoons C22$ $C19 + Ste11pp \rightleftharpoons C25$ $C21 \rightarrow C24$ $C23 \rightarrow C25$ $C26 \rightarrow C27$ $B3 \rightleftharpoons C3$ $B6 \rightleftharpoons C6$ $B9 \rightleftharpoons C9$ $B12 \rightleftharpoons C12$
$C8 + Fus3pp \leftarrow C23$ $C9 + Fus3pp \leftarrow C25$ $C10 + Fus3pp \leftarrow C27$ $C12 \leftarrow C13$ $C13 + Ste7 \Rightarrow C21$ $C14 + Ste7 \Rightarrow C25$ $C15 + Ste7pp \leftarrow C26$ $C17 \rightarrow C18$ $C18 + Ste11pp \Rightarrow C27$ $C19 + Ste11 \Rightarrow C23$ $C22 \rightarrow C24$ $C24 \rightarrow C27$ $B1 \Rightarrow C1$ $B4 \Rightarrow C4$ $B7 \Rightarrow C7$ $B10 \Rightarrow C10$ $B13 \Rightarrow C13$	$C9 \rightarrow C10$ $C10 \leftarrow C11$ $C11 + Fus3 \rightleftharpoons C22$ $C12 + Ste7 \rightleftharpoons C20$ $C13 + Ste7pp \leftarrow C24$ $C14 + Ste7pp \leftarrow C27$ $C16 + Ste11pp \rightleftharpoons C21$ $C17 + Ste11pp \rightleftharpoons C24$ $C18 + Ste11 \rightleftharpoons C26$ $C20 \rightarrow C21$ $C22 \rightarrow C26$ $C25 \rightarrow C27$ $B2 \rightleftharpoons C2$ $B5 \rightleftharpoons C5$ $B8 \rightleftharpoons C8$ $B11 \rightleftharpoons C11$ $B14 \rightleftharpoons C14$	$C9 + Fus3 \rightleftharpoons C21$ $C10 + Fus3 \rightleftharpoons C24$ $C11 + Fus3pp \leftarrow C26$ $C12 + Ste7pp \leftarrow C22$ $C14 \leftarrow C15$ $C15 + Ste7 \rightleftharpoons C23$ $C16 + Ste11 \rightleftharpoons C20$ $C17 + Ste11 \rightleftharpoons C22$ $C19 + Ste11pp \rightleftharpoons C25$ $C21 \rightarrow C24$ $C23 \rightarrow C25$ $C26 \rightarrow C27$ $B3 \rightleftharpoons C3$ $B6 \rightleftharpoons C6$ $B9 \rightleftharpoons C9$ $B12 \rightleftharpoons C12$ $B15 \rightleftharpoons C15$
$C8 + Fus3pp \leftarrow C23$ $C9 + Fus3pp \leftarrow C25$ $C10 + Fus3pp \leftarrow C27$ $C12 \leftarrow C13$ $C13 + Ste7 \rightleftharpoons C21$ $C14 + Ste7 \rightleftharpoons C25$ $C15 + Ste7pp \leftarrow C26$ $C17 \rightarrow C18$ $C18 + Ste11pp \rightleftharpoons C27$ $C19 + Ste11 \rightleftharpoons C23$ $C22 \rightarrow C24$ $C24 \rightarrow C27$ $B1 \rightleftharpoons C1$ $B4 \rightleftharpoons C4$ $B7 \rightleftharpoons C7$ $B10 \rightleftharpoons C10$ $B13 \rightleftharpoons C13$ $B16 \rightleftharpoons C16$	$C9 \rightarrow C10$ $C10 \leftarrow C11$ $C11 + Fus3 \rightleftharpoons C22$ $C12 + Ste7 \rightleftharpoons C20$ $C13 + Ste7pp \leftarrow C24$ $C14 + Ste7pp \leftarrow C27$ $C16 + Ste11pp \rightleftharpoons C21$ $C17 + Ste11pp \rightleftharpoons C24$ $C18 + Ste11 \rightleftharpoons C26$ $C20 \rightarrow C21$ $C22 \rightarrow C26$ $C25 \rightarrow C27$ $B2 \rightleftharpoons C2$ $B5 \rightleftharpoons C5$ $B8 \rightleftharpoons C8$ $B11 \rightleftharpoons C11$ $B14 \rightleftharpoons C14$ $B17 \rightleftharpoons C17$	$C9 + Fus3 \rightleftharpoons C21$ $C10 + Fus3 \rightleftharpoons C24$ $C11 + Fus3pp \leftarrow C26$ $C12 + Ste7pp \leftarrow C22$ $C14 \leftarrow C15$ $C15 + Ste7 \rightleftharpoons C23$ $C16 + Ste11 \rightleftharpoons C20$ $C17 + Ste11 \rightleftharpoons C22$ $C19 + Ste11pp \rightleftharpoons C25$ $C21 \rightarrow C24$ $C23 \rightarrow C25$ $C26 \rightarrow C27$ $B3 \rightleftharpoons C3$ $B6 \rightleftharpoons C6$ $B9 \rightleftharpoons C9$ $B12 \rightleftharpoons C12$ $B15 \rightleftharpoons C18$
$C8 + Fus3pp \leftarrow C23$ $C9 + Fus3pp \leftarrow C25$ $C10 + Fus3pp \leftarrow C27$ $C12 \leftarrow C13$ $C13 + Ste7 \rightleftharpoons C21$ $C14 + Ste7 \rightleftharpoons C25$ $C15 + Ste7pp \leftarrow C26$ $C17 \rightarrow C18$ $C18 + Ste11pp \rightleftharpoons C27$ $C19 + Ste11 \rightleftharpoons C23$ $C22 \rightarrow C24$ $C24 \rightarrow C27$ $B1 \rightleftharpoons C1$ $B4 \rightleftharpoons C4$ $B7 \rightleftharpoons C7$ $B10 \rightleftharpoons C10$ $B13 \rightleftharpoons C13$ $B16 \rightleftharpoons C16$ $B19 \rightleftharpoons C19$	$C9 \to C10$ $C10 \leftarrow C11$ $C11 + Fus3 \rightleftharpoons C22$ $C12 + Ste7 \rightleftharpoons C20$ $C13 + Ste7pp \leftarrow C24$ $C14 + Ste7pp \leftarrow C27$ $C16 + Ste11pp \rightleftharpoons C21$ $C17 + Ste11pp \rightleftharpoons C24$ $C18 + Ste11 \rightleftharpoons C26$ $C20 \to C21$ $C22 \to C26$ $C25 \to C27$ $B2 \rightleftharpoons C2$ $B5 \rightleftharpoons C5$ $B8 \rightleftharpoons C8$ $B11 \rightleftharpoons C11$ $B14 \rightleftharpoons C14$ $B17 \rightleftharpoons C20$	$C9 + Fus3 \rightleftharpoons C21$ $C10 + Fus3 \rightleftharpoons C24$ $C11 + Fus3pp \leftarrow C26$ $C12 + Ste7pp \leftarrow C22$ $C14 \leftarrow C15$ $C15 + Ste7 \rightleftharpoons C23$ $C16 + Ste11 \rightleftharpoons C20$ $C17 + Ste11 \rightleftharpoons C22$ $C19 + Ste11pp \rightleftharpoons C25$ $C21 \rightarrow C24$ $C23 \rightarrow C25$ $C26 \rightarrow C27$ $B3 \rightleftharpoons C3$ $B6 \rightleftharpoons C6$ $B9 \rightleftharpoons C9$ $B12 \rightleftharpoons C12$ $B18 \rightleftharpoons C18$ $B21 \rightleftharpoons C21$
$C8 + Fus3pp \leftarrow C23$ $C9 + Fus3pp \leftarrow C25$ $C10 + Fus3pp \leftarrow C27$ $C12 \leftarrow C13$ $C13 + Ste7 \rightleftharpoons C21$ $C14 + Ste7 \rightleftharpoons C25$ $C15 + Ste7pp \leftarrow C26$ $C17 \rightarrow C18$ $C18 + Ste11pp \rightleftharpoons C27$ $C19 + Ste11 \rightleftharpoons C23$ $C22 \rightarrow C24$ $C24 \rightarrow C27$ $B1 \rightleftharpoons C1$ $B4 \rightleftharpoons C4$ $B7 \rightleftharpoons C7$ $B10 \rightleftharpoons C10$ $B13 \rightleftharpoons C13$ $B16 \rightleftharpoons C16$	$C9 \rightarrow C10$ $C10 \leftarrow C11$ $C11 + Fus3 \rightleftharpoons C22$ $C12 + Ste7 \rightleftharpoons C20$ $C13 + Ste7pp \leftarrow C24$ $C14 + Ste7pp \leftarrow C27$ $C16 + Ste11pp \rightleftharpoons C21$ $C17 + Ste11pp \rightleftharpoons C24$ $C18 + Ste11 \rightleftharpoons C26$ $C20 \rightarrow C21$ $C22 \rightarrow C26$ $C25 \rightarrow C27$ $B2 \rightleftharpoons C2$ $B5 \rightleftharpoons C5$ $B8 \rightleftharpoons C8$ $B11 \rightleftharpoons C11$ $B14 \rightleftharpoons C14$ $B17 \rightleftharpoons C17$	$C9 + Fus3 \rightleftharpoons C21$ $C10 + Fus3 \rightleftharpoons C24$ $C11 + Fus3pp \leftarrow C26$ $C12 + Ste7pp \leftarrow C22$ $C14 \leftarrow C15$ $C15 + Ste7 \rightleftharpoons C23$ $C16 + Ste11 \rightleftharpoons C20$ $C17 + Ste11 \rightleftharpoons C22$ $C19 + Ste11pp \rightleftharpoons C25$ $C21 \rightarrow C24$ $C23 \rightarrow C25$ $C26 \rightarrow C27$ $B3 \rightleftharpoons C3$ $B6 \rightleftharpoons C6$ $B9 \rightleftharpoons C9$ $B12 \rightleftharpoons C12$ $B15 \rightleftharpoons C18$

TABLE VI: parameters in MAPK cascade on scaffold

Protein	on	off	**on	**off	phosphorylation	References and notes
	$(nM^{-1})$	$(min^{-1}]$	$(nM^{-1})$	$(min^{-1}]$	$(min^{-1})$	
	$\min^{-1}$ )		$\min^{-1}$ )			
Ste11	1.2	10	1	10	20	Stellpp can rebind to the scafold (27)
(MAP						
KKK)						
Ste7	1.2	0.1	-	0.5;	46	Ste7pp in the scaffold undergoes hyper-
(MAP				20		phosphorylation by Fus3pp, which accel-
KK)						erates Ste7's dissociation from the scaffold
Fus3	1.2	10	-	200	250	$t_{1/2} = 0.3s$ for the recovery of Fuspp in
(MAPK)						shmoo tips $(10,57)$

TABLE VII: parameters in downstream reactions

protein	value	References and Notes
k28	$8nMmin^{-1}$	
k29	40nM	
k30	$0.01min^{-1}$	
k31	$0.5nMmin^{-1}$	
k32	10nM	
k33	$2nMmin^{-1}$	
k34	7nM	
k35	$0.002min^{-1}$	
k36	$8nM^{-1}min^{-1}$	
k37	$1min^{-1}$	
k38	$10min^{-1}$	
k39	$10min^{-1}$	
k40	$0.1nM^{-1}min^{-1}$	
k41	$0.01min^{-1}$	
k42	$0.01nM^{-1}min^{-1}$	
k43	$0.1 min^{-1}$	
k44	$32nMmin^{-1}$	fit the time course of Sst2 determined in experiment (8)
k45	1nM	the same as k44
k46	$0.052min^{-1}$	the sam as k44
k47	$10min^{-1}$	$t_{1/2} = 4s$ for the recovery of Fus3in nucleus (57)
k48	$14min^{-1}$	$[Fus3_{in}]/[Fus3_{out}] = 1.4 \text{ at } t = 0 \text{ (45)}$
k49	$8min^{-1}$	$[Fus3_{in}]/[Fus3_{out}] = 2$ after 60 min treatment of
k50	$15min^{-1}$	$\alpha$ -factor (45)

TABLE VIII: initial amount of proteins

Protein	value(nM)	notes and references
Ste2	1000	8000 mole/cell (8)
Sst2	250	2000 mole/cell (8)
G protein	1000	8000 mole/cell (8)
Ste20	1000	
Ste11	200	less than 2000 mole/cell(1)
Fus3Ste7	200	Total amount of Ste7 should be less than 2000 mole/cell, and about 95% of
Ste7	10	Ste7 is in the form of Fus3Ste7 before pheromone induction (1)
$Fus3_{in}$	700	about 5000 mole/cell (1), and before pheromone
$Fus3_{out}$	300	induction, $[Fus3_{in}]/[Fus3_{out}] = 1.4 (45)$
MAPKKK-P	100	
MAPKK-P	100	
MAPK-Pin	100	
MAPK-Pout	100	
Ste5in	125	(1)
Ste5out	0.1	
Cdc28	100	

TABLE IX: mutants

No.	Mutant	Behavior	Implementation	Reference
1	$Ste2^{300\triangle}$ (endocytosis of receptor Ste2 is impaired)	G protein activation levels up	k3=0.08	Ref.12
2	treated with cyclo- heximide(synthesis inhibitor)	G protein cycle is closed down	k31=k32=k44= k4=k9=0	Ref.12
3	SST2∆	super-sensitivity upon pheromone induction	$k44=0, [Sst2]_{t=0}=0$ $nM$	Ref.8
4	$2 \times SST2$	response upon pheromone reduces	$k44=64,$ $[Sst2]_{t=0}=500 \text{ nM}$	Ref.8
5	$2 \times G\beta\gamma$	super-sensitivity to pheromone induction	$[Sst2]_{t=0}$ =500 nM $[G\beta\gamma]_{t=0}$ = 1000nM	Ref.8
6	negative feedbacks are cut down	activation of Fus3pp doesn't attenuate with time	k26=k27=k31= k44=0	
7	dissociating speed for hyper- phosphorylated Ste7pp is slowed down	activation of Fus3pp doesn't attenuate with time	$**off'_{KK} = 0.5$	
8	dissociating speed for normal Ste7pp is enhanced	activation of Fus3pp decreases	$**off_{KK} = 20$	
9	shuttle of the scaffold Ste5 is cut down	super-sensitivity to pheromone induction	k22=k23=0	
10	Ste11 is continuously activated	Ste7pp is activated while Fus3pp is re- pressed down	$ \begin{array}{cccc} [Ste11pp]_{t=0} & = & \\ 200nM, [Ste11]_{t=0} & = & \\ 0, [MAPKKK & - & \\ P]_{totoal} & = & \\ 0, [\alpha - factor] = 0 & \end{array} $	
11	Ste7 is continuously activated	Fus3pp isn't activated	$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	Ref.46

```
ODE functions:
   \frac{d[Ste2]}{\cdot \cdot} = -k1[\alpha - factor][Ste2] + k2[Ste2_{active}] - k7[Ste2] +
                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    k4[Ste12_{active}]^2
                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                 \overline{k5^2 + [Ste12_{active}]^2}
   \frac{d[Ste2_{active}]}{..} = k1[\alpha - factor][Ste2] - k2[Ste2_{active}] - k3[Ste2_{active}]
     d[Sst2_{active}]
                                                                                                                                                                                                                                                                                                                                                                    -k46[Sst2_{active}]
                                                                                                                                                         \overline{k45^2 + [Ste12_{active}]^2}
                                                   dt
   \frac{d[G]}{..} = -k8[Ste2_{active}][G] + k15[G_{\alpha}d][G_{\beta}\gamma] +
                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                        k12[G] + k11
                                                                                                                                                                                                                                                                                                                                                                                                                                             \overline{k10^2 + [Ste12_{active}]^2}
   \frac{d[G_{\alpha}t]}{\cdot \cdot} = k8[Ste2_{active}][G] - k13[G_{\alpha}t] - k14[G_{\alpha}t][Sst2_{active}]
   \frac{dt}{d[G_{\alpha}d]} = k13[G_{\alpha}t] + k14[G_{\alpha}t][Sst2_{active}] - k15[G_{\alpha}d][G_{\beta}\gamma]
   \overline{d^t_{G_\beta\gamma]}}
                                                                         = k8[Ste2_{active}][G] - k15[G_{\alpha}d][G_{\beta}\gamma] - k40[G_{\beta}\gamma][Far1pp_{out}] + k41[Far1pp_{out}G_{\beta}\gamma]20 - k18[G_{\beta}\gamma][Ste20] + k19[G_{\beta}\gamma Ste20] + 
   \frac{dt}{d[Ste20]} = -k18[G_{\beta}\gamma][Ste20] + k19[G_{\beta}\gamma Ste20]
   \frac{d}{dt} = -k18[G_{\beta}\gamma][Ste20] + k16[G_{\beta}\gamma Ste20] - k16[G_{\beta}\gamma Ste20]B1 + k17C1 - k16[G_{\beta}\gamma Ste20]B2 + k17C2 - k16[G_{\beta}\gamma Ste20]B3 + k17C3 - k16[G_{\beta}\gamma Ste20]B7 + k17C7 - k16[G_{\beta}\gamma Ste20]B8 + k17C7 - k16[G_{\beta}\gamma 
\frac{dt}{dt} = k18[G_{\beta}\gamma][Ste20] - k19[G_{\beta}\gamma Ste20] - k10[G_{\beta}\gamma Ste20]B1 + k17C1 - k10[G_{\beta}\gamma Ste20]B2 + k17C7 - k16[G_{\beta}\gamma Ste20]B8 + k17C4 - k16[G_{\beta}\gamma Ste20]B5 + k17C5 - k16[G_{\beta}\gamma Ste20]B6 + k17C6 - k16[G_{\beta}\gamma Ste20]B7 + k17C7 - k16[G_{\beta}\gamma Ste20]B8 + k17C8 - k16[G_{\beta}\gamma Ste20]B9 + k17C9 - k16[G_{\beta}\gamma Ste20]B10 + k17C10 - k16[G_{\beta}\gamma Ste20]B11 + k17C11 - k16[G_{\beta}\gamma Ste20]B12 + k17C12 - k16[G_{\beta}\gamma Ste20]B13 + k17C13 - k16[G_{\beta}\gamma Ste20]B14 + k17C14 - k16[G_{\beta}\gamma Ste20]B15 + k17C15 - k16[G_{\beta}\gamma Ste20]B16 + k17C16 - k16[G_{\beta}\gamma Ste20]B15 + k17C19 - k16[G_{\beta}\gamma Ste20]B10 + 
   k16[G_{\beta}\gamma Ste20]B17 + k17C17 - k16[G_{\beta}\gamma Ste20]B18 + k17C18 - k16[G_{\beta}\gamma Ste20]B19 + k17C19 - k16[G_{\beta}\gamma Ste20]B20 + k17C20 - k16[G_{\beta}\gamma Ste20]B20 + k16[G_{
   k16[G_{\beta}\gamma Ste20]B21 + k17C21 - k16[G_{\beta}\gamma Ste20]B22 + k17C22 - k16[G_{\beta}\gamma Ste20]B23 + k17C23 - k16[G_{\beta}\gamma Ste20]B24 + k17C24 - k16[G_{\beta}\gamma Ste20]B24 + k17C24 - k16[G_{\beta}\gamma Ste20]B25 + k17C25 + k16[G_{\beta}\gamma Ste20]B25 + k16[G_{\beta}\gamma Ste20]B
   k16[G_{\beta}\gamma Ste20]B25 + k17C25 - k16[G_{\beta}\gamma Ste20]B26 + k17C26 - k16[G_{\beta}\gamma Ste20]B27 + k17C27
     \frac{d[Ste11]}{dt} = p1_{KKK}[Ste11pMAPKKK-P] + off_{KKK}(C2 + C8 + C11 + C12 + C15 + C20 + C22 + C23 + C26 + B2 + B8 + B11 + B12 + B15 + B20 + B22 + C26 + B2 + B11 + B12 + B15 + B20 + B22 + B11 + B12 + B15 + B20 + B22 + B11 + B12 + B13 + B14 + B1
     B23 + B26) - on_{KKK}[Ste11](C4 + C6 + C7 + C1 + C16 + C17 + C18 + C19 + C5 + B4 + B6 + B7 + B1 + B16 + B17 + B18 + B19 + B5) - k26[Ste11][Fus3pp_{out}] + B16 + B17 + B18 + B19 +
   \frac{d[Ste11p]}{dt} = -a1_{KKK}[Ste11p]([MAPKKK-P]_0 - [Ste11pMAPKKK-P] - [Ste11ppMAPKKK-P]) + d1_{KKK}[Ste11pMAPKKK-P] + d1_{KKK}[Ste11pMAPKK-P] + d1_{KKK}[Ste11pMAPKK-P] + d1_{KKK}[Ste11pMAPKK-P] + d1_{KKK}[Ste11pMAPKK-P] + d1_{KKK}[Ste11pMAPKK-P] + d1_{KKK}[Ste11pMAPK-P] + d
 p2_{KKK}^{at}[Stel1ppMAPKKK - P]
     d[Ste11pMAPKKK - P]
 a1_{KKK}[Ste11p]([MAPKKK-P]_0 - [Ste11pMAPKKK-P] - [Ste11ppMAPKKK-P]) - (d1_{KK} + p1_{KK})[Ste11pMAPKKK-P] - [Ste11ppMAPKKK-P] - [Ste11ppMAPKK-P] - [Ste11ppMAPK-P] - [S
   d[Stellpp]
                                                                                                                                  -a2KKK[Ste11pp]([MAPKKK-P]_0-[Ste11pMAPKKK-P]-[Ste11ppMAPKKK-P])+d2KKK[Ste11ppMAPKKK-P])+d2KKK[Ste11ppMAPKKK-P]+d2KKK[Ste11ppMAPKKK-P]+d2KKK[Ste11ppMAPKKK-P]+d2KKK[Ste11ppMAPKKK-P]+d2KKK[Ste11ppMAPKKK-P]+d2KKK[Ste11ppMAPKKK-P]+d2KKK[Ste11ppMAPKKK-P]+d2KKK[Ste11ppMAPKKK-P]+d2KKK[Ste11ppMAPKKK-P]+d2KKK[Ste11ppMAPKKK-P]+d2KKK[Ste11ppMAPKKK-P]+d2KKK[Ste11ppMAPKKK-P]+d2KKK[Ste11ppMAPKKK-P]+d2KKK[Ste11ppMAPKKK-P]+d2KKK[Ste11ppMAPKKK-P]+d2KKK[Ste11ppMAPKKK-P]+d2KKK[Ste11ppMAPKKK-P]+d2KKK[Ste11ppMAPKKK-P]+d2KKK[Ste11ppMAPKKK-P]+d2KKK[Ste11ppMAPKKK-P]+d2KKK[Ste11ppMAPKKK-P]+d2KKK[Ste11ppMAPKKK-P]+d2KKK[Ste11ppMAPKKK-P]+d2KKK[Ste11ppMAPKKK-P]+d2KKK[Ste11ppMAPKKK-P]+d2KKK[Ste11ppMAPKKK-P]+d2KKK[Ste11ppMAPKKK-P]+d2KKK[Ste11ppMAPKKK-P]+d2KKK[Ste11ppMAPKKK-P]+d2KKK[Ste11ppMAPKKK-P]+d2KKK[Ste11ppMAPKKK-P]+d2KKK[Ste11ppMAPKKK-P]+d2KKK[Ste11ppMAPKKK-P]+d2KK[Ste11ppMAPKKK-P]+d2KK[Ste11ppMAPKKK-P]+d2KK[Ste11ppMAPKKK-P]+d2KK[Ste11ppMAPKKK-P]+d2KK[Ste11ppMAPKKK-P]+d2KK[Ste11ppMAPKKK-P]+d2KK[Ste11ppMAPKKK-P]+d2KK[Ste11ppMAPKKK-P]+d2KK[Ste11ppMAPKKK-P]+d2KK[Ste11ppMAPKKK-P]+d2KK[Ste11ppMAPKKK-P]+d2KK[Ste11ppMAPKKK-P]+d2KK[Ste11ppMAPKKK-P]+d2KK[Ste11ppMAPKKK-P]+d2KK[Ste11ppMAPKKK-P]+d2KK[Ste11ppMAPKKK-P]+d2KK[Ste11ppMAPKKK-P]+d2KK[Ste11ppMAPKKK-P]+d2KK[Ste11ppMAPKKK-P]+d2KK[Ste11ppMAPKKK-P]+d2KK[Ste11ppMAPKKK-P]+d2KK[Ste11ppMAPKKK-P]+d2KK[Ste11ppMAPKKK-P]+d2KK[Ste11ppMAPKKK-P]+d2KK[Ste11ppMAPKKK-P]+d2KK[Ste11ppMAPKK-P]+d2KK[Ste11ppMAPKK-P]+d2KK[Ste11ppMAPKK-P]+d2KK[Ste11ppMAPKK-P]+d2KK[Ste11ppMAPKK-P]+d2KK[Ste11ppMAPKK-P]+d2KK[Ste11ppMAPKK-P]+d2KK[Ste11ppMAPKK-P]+d2KK[Ste11ppMAPK-P]+d2KK[Ste11ppMAPK-P]+d2KK[Ste11ppMAPKK-P]+d2KK[Ste11ppMAPK-P]+d2KK[Ste11ppMAPK-P]+d2KK[Ste11ppMAPK-P]+d2KK[Ste11ppMAPK-P]+d2KK[Ste11ppMAPK-P]+d2KK[Ste11ppMAPK-P]+d2KK[Ste11ppMAPK-P]+d2KK[Ste11ppMAPK-P]+d2KK[Ste11ppMAPK-P]+d2KK[Ste11ppMAPK-P]+d2KK[Ste11ppMAPK-P]+d2KK[Ste11ppMAPK-P]+d2KK[Ste11ppMAPK-P]+d2KK[Ste11ppMAPK-P]+d2KK[Ste11ppMAPK-P]+d2KK[Ste11ppMAPK-P]+d2KK[Ste11ppMAPK-P]+d2KK[Ste11ppMAPK-P]+d2K(Ste11ppMAPK-P]+d2K(Ste11ppMAPK-P]+d2K(Ste11ppMAPK-P]+
 C6+C7+C16+C17+C18+C19+B1+B4+B5+B6+B7+B16+B17+B18+B19)
     d[StellppMAPKKK - P]
   a2_{KKK}[Ste11pp]([MAPKKK-P]_0 - [Ste11pMAPKKK-P] - [Ste11ppMAPKKK-P]) - (d2_{KKK} + p2_{KKK})[Ste11ppMAPKKK-P])
   \frac{d[Ste7]}{...} = -a3_{KK}[Ste7][Ste11pp] + d3_{KK}[Ste11ppSte7] + p1_{KK}[Ste7pMAPKK - P] + off_{KK}(C4 + C8 + C9 + C16 + C19 + C20 + C21 + C
 C^{23} + C^{25} + B^{4} + B^{8} + B^{9} + B^{16} + B^{19} + B^{20} + B^{21} + B^{23} + B^{25}) - on_{KK}[Ste7](C^{1} + C^{2} + C^{3} + C^{6} + C^{7} + C^{12} + C^{13} + C^{14} + C^{15} + B^{1} + C^{14} + C^{15} + B^{1} + C^{14} + C^{15} + B^{1} + C^{15} + C^{14} + C^{15} + C^{15}
   B2 + B3 + B6 + B7 + B12 + B13 + B14 + B15) - k24[Ste7][Fus3_{out}] + k25[Fus3_{out}Ste7]
   \frac{d[Ste7Ste11pp]}{d} = a3_{KK}[Ste11pp][Ste7] - (d3_{KK} + p3_{KK})[Ste11ppSte7]
   \frac{d[Ste7p]}{J_L} = -a1_{KK}([MAPKK-P]_0 - [Ste7pMAPKK-P] - [Ste7ppMAPKK-P])[Ste7p] + d1_{KK}[Ste7pMAPKK-P] + d1_{KK}[Ste7pMAPKC-P] + d1_{KK}[Ste7pMA
   p3_{KK}^{ac}[Ste11ppSte7] - a4_{KK}[Ste7p][Ste11pp] + d4_{KK}[Ste11ppSte7p] + p2_{KK}[Ste7ppMAPKK - P]
   \frac{d[Ste7pMAPKK-P]}{d} = a1_{KK}[Ste7p]([MAPKK-P]_0 - [Ste7pMAPKK-P] - [Ste7ppMAPKK-P]) - (d1_{KK} + p1_{KK})[Ste7pMAPKK-P]
   \frac{dt}{d[Ste7pSte11pp]} = a4_{KK}[Ste11pp][Ste7p] - (d4_{KK} + p4_{KK})[Ste11ppSte7p]
     d[Ste7pp]
                                                                                                    =-a2_{KK}[Ste7pp]([MAPKK-P]_0-[Ste7pMAPKK-P]-[Ste7ppMAPKK-P])+d2_{KK}[Ste7ppMAPKK-P]+d2_{KK}[Ste7ppMAPKK-P]+d2_{KK}[Ste7ppMAPKK-P]+d2_{KK}[Ste7ppMAPKK-P]+d2_{KK}[Ste7ppMAPKK-P]+d2_{KK}[Ste7ppMAPKK-P]+d2_{KK}[Ste7ppMAPKK-P]+d2_{KK}[Ste7ppMAPKK-P]+d2_{KK}[Ste7ppMAPKK-P]+d2_{KK}[Ste7ppMAPKK-P]+d2_{KK}[Ste7ppMAPKK-P]+d2_{KK}[Ste7ppMAPKK-P]+d2_{KK}[Ste7ppMAPKK-P]+d2_{KK}[Ste7ppMAPKK-P]+d2_{KK}[Ste7ppMAPKK-P]+d2_{KK}[Ste7ppMAPKK-P]+d2_{KK}[Ste7ppMAPKK-P]+d2_{KK}[Ste7ppMAPKK-P]+d2_{KK}[Ste7ppMAPKK-P]+d2_{KK}[Ste7ppMAPKK-P]+d2_{KK}[Ste7ppMAPKK-P]+d2_{KK}[Ste7ppMAPKK-P]+d2_{KK}[Ste7ppMAPKK-P]+d2_{KK}[Ste7ppMAPKK-P]+d2_{KK}[Ste7ppMAPKK-P]+d2_{KK}[Ste7ppMAPKK-P]+d2_{KK}[Ste7ppMAPKK-P]+d2_{KK}[Ste7ppMAPKK-P]+d2_{KK}[Ste7ppMAPKK-P]+d2_{KK}[Ste7ppMAPKK-P]+d2_{KK}[Ste7ppMAPKK-P]+d2_{KK}[Ste7ppMAPKK-P]+d2_{KK}[Ste7ppMAPKK-P]+d2_{KK}[Ste7ppMAPKK-P]+d2_{KK}[Ste7ppMAPKK-P]+d2_{KK}[Ste7ppMAPKK-P]+d2_{KK}[Ste7ppMAPKK-P]+d2_{KK}[Ste7ppMAPKK-P]+d2_{KK}[Ste7ppMAPKK-P]+d2_{KK}[Ste7ppMAPKK-P]+d2_{KK}[Ste7ppMAPKK-P]+d2_{KK}[Ste7ppMAPKK-P]+d2_{KK}[Ste7ppMAPKK-P]+d2_{KK}[Ste7ppMAPKK-P]+d2_{KK}[Ste7ppMAPKK-P]+d2_{KK}[Ste7ppMAPKK-P]+d2_{KK}[Ste7ppMAPKK-P]+d2_{KK}[Ste7ppMAPKK-P]+d2_{KK}[Ste7ppMAPKK-P]+d2_{KK}[Ste7ppMAPKK-P]+d2_{KK}[Ste7ppMAPKK-P]+d2_{KK}[Ste7ppMAPKK-P]+d2_{KK}[Ste7ppMAPKK-P]+d2_{KK}[Ste7ppMAPKK-P]+d2_{KK}[Ste7ppMAPKK-P]+d2_{KK}[Ste7ppMAPKK-P]+d2_{KK}[Ste7ppMAPK-P]+d2_{KK}[Ste7ppMAPK-P]+d2_{KK}[Ste7ppMAPK-P]+d2_{KK}[Ste7ppMAPK-P]+d2_{KK}[Ste7ppMAPK-P]+d2_{KK}[Ste7ppMAPK-P]+d2_{KK}[Ste7ppMAPK-P]+d2_{KK}[Ste7ppMAPK-P]+d2_{KK}[Ste7ppMAPK-P]+d2_{KK}[Ste7ppMAPK-P]+d2_{KK}[Ste7ppMAPK-P]+d2_{KK}[Ste7ppMAPK-P]+d2_{KK}[Ste7ppMAPK-P]+d2_{KK}[Ste7ppMAPK-P]+d2_{KK}[Ste7ppMAPK-P]+d2_{KK}[Ste7ppMAPK-P]+d2_{KK}[Ste7ppMAPK-P]+d2_{KK}[Ste7ppMAPK-P]+d2_{KK}[Ste7ppMAPK-P]+d2_{KK}[Ste7ppMAPK-P]+d2_{KK}[Ste7ppMAPK-P]+d2_{KK}[Ste7ppMAPK-P]+d2_{KK}[Ste7ppMAPK-P]+d2_{KK}[Ste7ppMAPK-P]+d2_{KK}[Ste7ppMAPK-P]+d2_{KK}[Ste7ppMAPK-P]+d2_{KK}[Ste7ppMAPK-P]+d2_{KK}[Ste7ppMAPK-P]+d2_{KK}[Ste7ppMAPK-P]+d2_{KK}[Ste7ppMAPK-P]+d2_{KK}[Ste7ppMAPK-P]+d2_{KK}[Ste7
 p4_{KK}[Ste11ppSte7p] - a3_{K}[Ste7pp][Fus3_{out}] + (d3_{K} + p3_{K})[Ste7ppFus3_{out}] - a4_{K}[Ste7pp][Fus3p_{out}] + (d4_{K} + p4_{K})[Ste7ppFus3p_{out}] + **
 off_{KK}(C5+C10+C11+C17+C22+C24+B5+B10+B11+B17+B22+B24) + **off_{KK}'(C18+C26+C27+B18+B26+B27) - k27[Ste7pp] \\ = (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0.5) + (1.51+0
   \frac{d[Ste7ppMAPKK-P]}{d[Ste7ppMAPKK-P]} = a2_{KK}[Ste7pp]([MAPKK-P]_0 - [Ste7pMAPKK-P] - [Ste7ppMAPKK-P]) - (d2_{KK} + p2_{KK})[Ste7ppMAPKK-P] - [Ste7ppMAPKK-P] - [Ste7ppMAPK
   \frac{d[Fus3_{out}]}{dt} = -a3_K[Ste7pp][Fus3_{out}] + d3_K[Ste7ppFus3_{out}] + p1_K[Fus3p_{out}MAPK - P_{out}] + off_K(C6 + C12 + C13 + C16 + C17 + C20 + C10 + C10)
 C21 + C22 + C24 + B6 + B12 + B13 + B16 + B17 + B20 + B21 + B22 + B24) - on_{K}[Fus3_{out}](C1 + C2 + C3 + C4 + C5 + C8 + C9 + C10 + C11 
                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         k32[Ste12_{active}]^2
     B1 + B2 + B3 + B4 + B5 + B8 + B9 + B10 + B11) - k24[Ste7][Fus3_{out}] + k25[Fus3_{out}Ste7] + k47[Fus3_{in}] - k48[Fus3_{out}] + k47[Fus3_{in}] + k47[Fus3_{i
   \frac{d[Fus3_{out}Ste11pp]}{\cdot \cdot} = a3_K[Ste7pp][Fus3_{out}] - (d3_K + p3_K)[Ste7ppFus3_{out}]
     d[Fus3\underline{p_{out}}]
                                                                                                                           =-a1_K[Fus3p_{out}][MAPK-P_{out}]+d1_K[Fus3p_{out}MAPK-P_{out}]+p3_K[Ste7ppFus3_{out}]-a4_K[Ste7pp][Fus3p_{out}]+d1_K[Ste7pp][Fus3p_{out}]+d1_K[Ste7pp][Fus3p_{out}]+d1_K[Ste7pp][Fus3p_{out}]+d1_K[Ste7pp][Fus3p_{out}]+d1_K[Ste7pp][Fus3p_{out}]+d1_K[Ste7pp][Fus3p_{out}]+d1_K[Ste7pp][Fus3p_{out}]+d1_K[Ste7pp][Fus3p_{out}]+d1_K[Ste7pp][Fus3p_{out}]+d1_K[Ste7pp][Fus3p_{out}]+d1_K[Ste7pp][Fus3p_{out}]+d1_K[Ste7pp][Fus3p_{out}]+d1_K[Ste7pp][Fus3p_{out}]+d1_K[Ste7pp][Fus3p_{out}]+d1_K[Ste7pp][Fus3p_{out}]+d1_K[Ste7pp][Fus3p_{out}]+d1_K[Ste7pp][Fus3p_{out}]+d1_K[Ste7pp][Fus3p_{out}]+d1_K[Ste7pp][Fus3p_{out}]+d1_K[Ste7pp][Fus3p_{out}]+d1_K[Ste7pp][Fus3p_{out}]+d1_K[Ste7pp][Fus3p_{out}]+d1_K[Ste7pp][Fus3p_{out}]+d1_K[Ste7pp][Fus3p_{out}]+d1_K[Ste7pp][Fus3p_{out}]+d1_K[Ste7pp][Fus3p_{out}]+d1_K[Ste7pp][Fus3p_{out}]+d1_K[Ste7pp][Fus3p_{out}]+d1_K[Ste7pp][Fus3p_{out}]+d1_K[Ste7pp][Fus3p_{out}]+d1_K[Ste7pp][Fus3p_{out}]+d1_K[Ste7pp][Fus3p_{out}]+d1_K[Ste7pp][Fus3p_{out}]+d1_K[Ste7pp][Fus3p_{out}]+d1_K[Ste7pp][Fus3p_{out}]+d1_K[Ste7pp][Fus3p_{out}]+d1_K[Ste7pp][Fus3p_{out}]+d1_K[Ste7pp][Fus3p_{out}]+d1_K[Ste7pp][Fus3p_{out}]+d1_K[Ste7pp][Fus3p_{out}]+d1_K[Ste7pp][Fus3p_{out}]+d1_K[Ste7pp][Fus3p_{out}]+d1_K[Ste7pp][Fus3p_{out}]+d1_K[Ste7pp][Fus3p_{out}]+d1_K[Ste7pp][Fus3p_{out}]+d1_K[Ste7pp][Fus3p_{out}]+d1_K[Ste7pp][Fus3p_{out}]+d1_K[Ste7pp][Fus3p_{out}]+d1_K[Ste7pp][Fus3p_{out}]+d1_K[Ste7pp][Fus3p_{out}]+d1_K[Ste7pp][Fus3p_{out}]+d1_K[Ste7pp][Fus3p_{out}]+d1_K[Ste7pp][Fus3p_{out}]+d1_K[Ste7pp][Fus3p_{out}]+d1_K[Ste7pp][Fus3p_{out}]+d1_K[Ste7pp][Fus3p_{out}]+d1_K[Ste7pp][Fus3p_{out}]+d1_K[Ste7pp][Fus3p_{out}]+d1_K[Ste7pp][Fus3p_{out}]+d1_K[Ste7pp][Fus3p_{out}]+d1_K[Ste7pp][Fus3p_{out}]+d1_K[Ste7pp][Fus3p_{out}]+d1_K[Ste7pp][Fus3p_{out}]+d1_K[Ste7pp][Fus3p_{out}]+d1_K[Ste7pp][Fus3p_{out}]+d1_K[Ste7pp][Fus3p_{out}]+d1_K[Ste7pp][Fus3p_{out}]+d1_K[Ste7pp][Fus3p_out]+d1_K[Ste7pp][Fus3p_out]+d1_K[Ste7pp]+d1_K[Ste7pp][Fus3p_out]+d1_K[Ste7pp]+d1_K[Ste7pp]+d1_K[Ste7pp]+d1_K[Ste7pp]+d1_K[Ste7pp]+d1_K[Ste7pp]+d1_K[Ste7pp]+d1_K[Ste7pp]+
 d4_{K}[Ste7ppFus3p_{out}] + p2_{K}[Fus3pp_{out}MAPK - P_{out}]
   \frac{d[Fus3p_{out}MAPK-P_{out}]}{d} = a1_K[Fus3p_{out}][MAPK-P_{out}] - (d1_K + p1_K)[Fus3p_{out}MAPK-P_{out}]
   \frac{d[Fus3p_{out}STe7pp]}{..} = a4_K[Ste7pp][Fus3p_{out}] - (d4_K + p4_K)[Ste7ppFus3p_{out}]
     d[Fus3pp_{\underline{out}}]
                                                                                                                                          = -a2_K[Fus3pp_{out}][MAPK - P_{out}] + d2_K[Fus3pp_{out}MAPK - P_{out}] + p4_K[Ste7ppFus3p_{out}] + **off_K(C7 + C14 + C15 + C15 + C14 + C15 
 \frac{dt}{C18 + C19 + C23 + C25 + C26 + C27 + B7 + B14 + B15 + B18 + B19 + B23 + B25 + B26 + B27) + k49[Fus3pp_{in}] - k50[Fus3pp_{out}]}
   \frac{d[Fus3pp_{out}MAPK-P_{out}]}{d[Fus3pp_{out}MAPK-P_{out}]} = a2_K[Fus3pp_{out}][MAPK-P_{out}] - (d2_K + p2_K)[Fus3pp_{out}MAPK-P_{out}]
     d[MAPK - P_{out}]
                                                                                                                                                                               = -a1_{K}[Fus3p_{out}][MAPK - P_{out}] + (d1_{K} + p1_{K})[Fus3p_{out}MAPK - P_{out}] - a2_{K}[Fus3p_{out}][MAPK - P_{out}] + (p2_{K} + p2_{K})[Pus3p_{out}][MAPK - P_{out}] + (p2_{K} + p2_{K})[Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p_{out}][Pus3p
                                                                                                                                                                                                                                                                                                                                            k31[Ste12_{active}]^2
 d2_K)[Fus3pp_{out}MAPK-P_{out}]+\\
                                                                                                                                                                                                                                                                                                                               \overline{k5^2 + [Ste12_{active}]^2}
     d[Fus3_{ou}\underline{t}Ste7]
                                                                                                                                                        =k24[Ste7][Fus3_{out}]-k25[Fus3_{out}Ste7]
                                                               dt
```

```
\frac{dB1}{\cdot \cdot} = -B1(on_{KKK}[Ste11] + **on_{KKK}[Ste11pp] + on_{KK}[Ste7] + on_{K}[Fus3_{out}]) + off_{KKK}B2 + **off_{KK}B3 + off_{KK}B4 + **off_{KK}B4 + *
\frac{dt}{off_{KK}B5} - bff_{K}B6 + **off_{K}B7 - k16[G_{\beta}\gamma Ste20]B1 + k17C1 + k22[Ste5_{in}] - k23B1
       \frac{dB2}{dt} = on_{KKK}B1[Ste11] - off_{KKK}B2 - on_{KK}[Ste7]B2 - on_{K}[Fus3_{out}]B2 + off_{KK}B8 + **off_{KK}B11 + off_{K}B12 + **off_{K}B15 - off_{KK}B11 + off_{KK}B
  k16[G_{\beta}\gamma Ste20]B2 + k17C2
       \frac{dB3}{...} = **on_{KKK}[Ste11pp]B1 - on_{KK}[Ste7]B3 - on_{K}[Fus3_{out}]B3 + off_{KK}B9 + **off_{KK}B10 + off_{K}B13 + **off_{K}B14 - **off_{KK}B10 + off_{KK}B10 + of
  \frac{dt}{dt} = ** on_{KK} [Stellpp]B1 - on_{K
     \frac{d\vec{B4}}{d\vec{B4}} = on_{KK}B1[Ste7] - off_{KK}B4 - on_{KKK}[Ste11]B4 - **on_{KKK}B4[Ste11pp] - on_{K}[Fus3_{out}]B4 + off_{KKK}B8 + **off_{KKK}B9 + on_{KKK}B4[Ste11pp] + on_{KKK}B4[Ste11pp] - on_{K}[Fus3_{out}]B4 + off_{KKK}B8 + **off_{KKK}B9 + on_{KKK}B4[Ste11pp] - on_{K}[Fus3_{out}]B4 + off_{KKK}B8 + **off_{KKK}B9 + on_{KKK}B4[Ste11pp] - on_{K}[Fus3_{out}]B4 + off_{KKK}B8 + **off_{KKK}B9 + on_{KKK}B4[Ste11pp] - on_{K}[Fus3_{out}]B4 + off_{KKK}B8 + on_{KKK}B4[Ste11pp] - on_{K}[Fus3_{out}]B4 + off_{KKK}B4[Ste11pp] - on_{KKK}B4[Ste11pp] - on_{KKK}B4[Ste11pp] - on_{KKK}B4[Ste11pp] - on_{KKK}B4[Ste11pp] - on_{KKK}B4[Ste11
       off_K B16 + ** off_K B19 - k16[G_{\beta} \gamma Ste20]B4 + k17C4
       dB5
                                                                                                            -**on_{KKK}[Ste11pp]B5 - on_{KKK}[Ste11]B5 - on_{K}[Fus3_{out}]B5 + **off_{KKK}B10 + off_{KKK}B11 + off_{K}B17 + **off_{K}B18 - **off_{KKK}B11 + off_{K}B18 - off_{KKK}B11 + off_{K}B18 - 
  off_{KK}B5 - k16[G_{\beta}\gamma Ste20]B5 + k17C5
                                                                 = - **on_{KKK}[Ste11pp]B6 + on_KB1[Fus3_{out}] - off_KB6 - on_{KKK}[Ste11]B6 - on_{KK}[Ste7]B6 + off_{KKK}B12 + **off_{KKK}B13 + off_{KKK}B12 + **off_{KKK}B13 + off_{KKK}B13 + off_{KKK
  off_{KK}B16 + ** off_{KK}B17 - k16[G_{\beta}\gamma Ste20]B6 + k17C6
       \frac{dB7}{dB7} = - **on_{KKK}[Ste11pp]B7 - on_{KK}[Ste11]B7 - on_{KK}[Ste7]B7 + **off_{KKK}B14 + off_{KKK}B15 + **off_{KK}'B18 + off_{KK}'B19 - **off_{KK}'B18 + off_{KK}'B19 - **off_{KK}'B18 + off_{KK}'B19 - **off_{KK}'B18 + off_{KK}'B19 - **off_{KK}'B19 - **off
  off_K B7 - k16[G_{\beta}\gamma Ste20]B7 + k17C7
     \frac{dB8}{dB8} = -on_K[Fus3_{out}]B8 - (off_{KK} + off_{KK})B8 + on_{KK}[Ste11]B4 + on_{KK}[Ste7]B2 + off_{K}B20 + **off_{K}B23 - k16[G_{\beta}\gamma Ste20]B8 + k17C8 + k16[G_{\beta}\gamma Ste20]B8 + k16[G_{\beta}\gamma St
       _{dB9}^{dt}
                                                            = **on_{KK}B4[Ste11pp] - p_{KK}B9 - (**off_{KK} + off_{KK})B9 - on_{K}[Fus3_{out}]B9 + on_{KK}[Ste7]B3 + off_{K}B21 + **off_{K}B25 - off_{KK}B4[Ste11pp] + off_{KK}B4[Ste11pp]
     k16[G_{\beta}\gamma Ste20]B9 + k17C9
       **on_{KKK}[Ste11pp]B5 - (**off_{KKK} + **off_{KK})B10 - on_{K}[Fus3_{out}]B10 + off_{K}B24 + **off_{K}B27 + p_{KK}B9 - k16[G_{\beta}\gamma Ste20]B10 + k17C10 + k17
       \frac{dB11}{dB11} = -(off_{KKK} + **off_{KK})B11 - on_K[Fus3_{out}]B11 + on_{KKK}[Ste11]B5 + off_KB22 + **off_KB26 - k16[G_{\beta}\gamma Ste20]B11 + k17C11 + 
     dt
dB12
                    dt
          -on_{KK}[Ste7]B12 - (off_{KKK} + off_K)B12 + on_{KKK}[Ste11]B6 + on_K[Fus3_{out}]B2 + off_{KK}B20 + **off_{KK}B22 - k16[G_{\beta}\gamma Ste20]B12 + k17C12 + k17C12
       dB13
                      *on_{KKK}[Ste11pp]B6 - (**off_{KKK} + off_{K})B13 - on_{KK}[Ste7]B13 + on_{K}[Fus3_{out}]B3 + off_{KK}B21 + **off_{KK}B24 - k16[G_{\beta}\gamma Ste20]B13 + k17C13
                                                                           = **on_{KKK}[Ste11pp]B7 - (**off_{KKK} + **off_{K})B14 - on_{KK}[Ste7]B14 + off_{KK}B25 + **off_{KK}'B27 - k16[G_{\beta}\gamma Ste20]B14 + k17C14 + k17C14
     \frac{dt}{dB15} = -(off_{KKK} + **off_K)B15 - on_{KK}[Ste7]B15 + on_{KKK}[Ste11]B7 + off_{KK}B23 + **off'_{KK}B26 - k16[G_{\beta}\gamma Ste20]B15 + k17C15
       dB^{ai}
                                                                                   = - **on_{KKK}[Ste11pp]B16 - on_{KKK}[Ste11]B16 - (off_{KK} + off_{K})B16 + on_{KK}[Ste7]B6 + on_{K}[Fus3_{out}]B4 + off_{KKK}B20 + **on_{KKK}[Ste11pp]B16 - on_{KKK}[Ste11pp]B16 - o
  off_{KKK}B21 - k16[G_{\beta}\gamma Ste20]B16 + k17C16
       \frac{dB17}{dB17} = - **on_{KKK}[Ste11pp]B17 - p_KB17 - (**off_{KK} + off_{K})B17 - on_{KKK}[Ste11]B17 + on_{K}[Fus3_{out}]B5 + off_{KKK}B22 + **off_{KKK}B22 + *
       off_{KKK}B24 - k16[G_{\beta}\gamma Ste20]B17 + k17C17
       dB18
               dt
                           **on_{KKK}[Ste11pp]B18 - (**off'_{KK} + **off_{K})B18 - on_{KKK}[Ste11]B18 + off_{KKK}B26 + **off_{KKK}B27 + p_{K}B17 - k16[G_{\beta}\gamma Ste20]B18 + k17C18 + c_{KK}B27 + c_{KK}B2
                                                                           = - **on_{KKK}[Ste11pp]B19 - (off_{KK} + **off_K)B19 - on_{KKK}[Ste11]B19 + on_{KK}[Ste7]B7 + off_{KKK}B23 + **off_{KKK}B25 - on_{KKK}[Ste7]B7 + on_{KKK}[Ste7]B7 +
     k16[G_{\beta}\gamma Ste20]B19 + k17C19
       dB20
                                                                         = -(off_{KKK} + off_{KK} + off_{K})B20 + on_{KKK}[Ste11]B16 + on_{KK}[Ste7]B12 + on_{K}[Fus3_{out}]B8 - k16[G_{\beta}\gamma Ste20]B20 + k17C20 + 
       dt
dB21
                                                                                   = **on_{KKK}[Ste11pp]B16 - (**off_{KKK} + off_{KK} + off_{K})B21 - p_{KK}B21 + on_{KK}[Ste7]B13 + on_{K}[Fus3_{out}]B9 - k16[G_{\beta}\gamma Ste20]B21 + k17C21
       d_{B22}^{dt}
                                                                                   = -(off_{KKK} + **off_{KK} + off_{K})B22 - (p_{K})B22 + on_{KKK}[Ste11]B17 + on_{K}[Fus3_{out}]B11 - k16[G_{\beta}\gamma Ste20]B22 + k17C22 + k17
     \frac{dt}{dB23} = -(off_{KKK} + off_{KK} + **off_{K})B23 + on_{KKK}[Ste11]B19 + on_{KK}[Ste7]B15 - k16[G_{\beta}\gamma Ste20]B23 + k17C23
                                                                         = **on_{KKK}[Ste11pp]B17 - (**off_{KKK} + **off_{KK} + off_{K})B24 - p_{K}B24 + on_{K}[Fus3_{out}]B10 + p_{KK}B21 - k16[G_{\beta}\gamma Ste20]B24 + k17C24 + on_{K}[Fus3_{out}]B10 + p_{KK}B21 - k16[G_{\beta}\gamma Ste20]B24 + k17C24 + on_{K}[Fus3_{out}]B10 + p_{KK}B21 - k16[G_{\beta}\gamma Ste20]B24 + k17C24 + on_{K}[Fus3_{out}]B10 + p_{KK}B21 - k16[G_{\beta}\gamma Ste20]B24 + k17C24 + on_{K}[Fus3_{out}]B10 + on_{K}[Fus3_
       \frac{dt}{dB25}
                                                                                = **on_{KKK}[Ste11pp]B19 - (**off_{KK} + off_{KK} + off_{KK})B25 - p_{KK}B25 + on_{KK}[Ste7]B14 - k16[G_{\beta}\gamma Ste20]B25 + k17C25 + on_{KK}[Ste7]B14 - k16[G_{\beta}\gamma Ste20]B25 + on_{KK}[Ste7]B14 - on_{KK}[Ste7]B1
       dt dB26
                                                                                   = -(off_{KKK} + **off_{KK}' + **off_{K})B26 + on_{KKK}[Ste11]B18 + p_{K}B22 - k16[G_{\beta}\gamma Ste20]B26 + k17C26
     _{dB27}^{\phantom{dt}dt}
                                                                           = **on_{KKK}[Ste11pp]B18 - (**off_{KK} + **off_{KK}' + **off_{K}')B27 + p_{KK}B25 + p_{K}B24 - k16[G_{\beta}\gamma Ste20]B27 + k17C27 + p_{KK}\beta Steam + k16[G_{\beta}\gamma Ste20]B27 + p_{KK}\beta Steam + k16[G_{\beta}\gamma Steam + k16[G_{\beta}\gamma
```

```
\frac{dC1}{..} = -C1(on_{KKK}[Ste11] + **on_{KKK}[Ste11pp] + on_{KK}[Ste7] + on_{K}[Fus3_{out}]) + off_{KKK}C2 + **off_{KK}C3 + off_{KK}C4 + **on_{KKK}C4 + **
\frac{dt}{off_{KK}C5} - Off_{KK}C6 + **off_{K}C7 + k16[G_{\beta}\gamma Ste20]B1 - k17C1
      \frac{dC2}{dt} = on_{KK}C1[Ste11] - off_{KK}C2 - p_{KK}C2 - on_{K}[Ste7]C2 - on_{K}[Fus3_{out}]C2 + off_{KK}C8 + **off_{KK}C11 + off_{K}C12 + **off_{KK}C11 + off_{KK}C11 + 
  off_{K}C15 + k16[G_{\beta}\gamma Ste20]B2 - k17C2
      \frac{d\vec{C3}}{..} = **on_{KKK}[Ste11pp]C1 + p_{KKK}C2 - on_{KK}[Ste7]C3 - on_{K}[Fus3_{out}]C3 + off_{KK}C9 + **off_{KK}C10 + off_{K}C13 + **off_{K}C14 - **off_{KK}C10 + off_{KK}C13 + **off_{KK}C14 - **off_{KK}C14 + **off_{K
  off_{KKK}C3 + k16[G_{\beta}\gamma Ste20]B3 - k17C3
    \frac{d\vec{C4}}{\vec{C4}} = on_{KK}C1[Ste7] - off_{KK}C4 - on_{KKK}[Ste11]C4 - **on_{KKK}C4[Ste11pp] - on_{K}[Fus3_{out}]C4 + off_{KKK}C8 + **off_{KKK}C9 + on_{KKK}C4[Ste11pp] - on_{K}[Fus3_{out}]C4 + off_{KKK}C8 + on_{KKK}C4[Ste11pp] - on_{K}[Fus3_{out}]C4 + off_{KKK}C8 + on_{KKK}C4[Ste11pp] - on_{K}[Fus3_{out}]C4 + off_{KKK}C8 + on_{KKK}C9 + on_{KKK}C4[Ste11pp] - on_{K}[Fus3_{out}]C4 + off_{KKK}C8 + on_{KKK}C9 + on_{KKK}C4[Ste11pp] - on_{K}[Fus3_{out}]C4 + off_{KKK}C8 + on_{KKK}C9 + on_{KKK}C9 + on_{KKK}C9] - on_{KKK}[Ste11pp] - on_{K}[Fus3_{out}]C4 + off_{KKK}C8 + on_{KKK}C9 + on_{KKK}C9] - on_{KKK}[Ste11pp] - on_{K}[Ste11pp] - on_{K}[Ste11p
      off_KC16 + ** off_KC19 + k16[G_{\beta}\gamma Ste20]B4 - k17C4
                                                                                                         **on_{KKK}[Ste11pp]C5 - on_{KKK}[Ste11]C5 - on_{K}[Fus3_{out}]C5 + **off_{KKK}C10 + off_{KKK}C11 + off_{K}C17 + **off_{K}C18 - **off_{KKK}C11 + off_{KKK}C11 + off_{KKK}C
  off_{KK}C5 + k16[G_{\beta}\gamma Ste20]B5 - k17C5
                                                   = - **on_{KKK}[Ste11pp]C6 + on_{K}C1[Fus3_{out}] - off_{K}C6 - on_{KKK}[Ste11]C6 - on_{KK}[Ste7]C6 + off_{KKK}C12 + **off_{KKK}C13 + off_{KKK}C13 + off_{KKK}C12 + off_{KKK}C13 + off_{K
  o\bar{f}f_{KK}C16 + **off_{KK}C17 + k16[G_{\beta}\gamma Ste20]B6 - k17C6
      \frac{dC7}{..} = - **on_{KKK}[Ste11pp]C7 - on_{KK}[Ste11]C7 - on_{KK}[Ste7]C7 + **off_{KKK}C14 + off_{KK}C15 + **off_{KK}'C18 + off_{KK}C19 - **off_{KK}C19 + **
    off_K C7 + k16[G_{\beta}\gamma Ste20]B7 - k17C7
      \frac{dC8}{dt} =
                  P_{KKK}C8 - on_{K}[Fus_{3out}]C8 - (off_{KKK} + off_{KK})C8 + on_{KKK}[Ste11]C4 + on_{KK}[Ste7]C2 + off_{K}C20 + **off_{K}C23 + k16[G_{\beta}\gamma Ste20]B8 - k17C8 + off_{KKK}C8 + on_{KK}[Ste7]C2 + off_{KKC}C8 + on_{KK}[Ste7]C8 + on_{
                                                   = **on_{KK}C4[Ste11pp] - p_{KK}C9 - (**off_{KK} + off_{KK})C9 - on_{K}[Fus3_{out}]C9 + on_{KK}[Ste7]C3 + off_{K}C21 + **off_{K}C25 + off_{KK}C4[Ste11pp] + off_{KK}C4[Ste11pp]
  p_{KKK}C8 + k16[G_{\beta}\gamma Ste20]B9 - k17C9
      \frac{dC10}{dC10} = **on_{KKK}[Ste11pp]C5 - (**off_{KKK} + **off_{KK})C10 - on_{K}[Fus3_{out}]C10 + off_{K}C24 + **off_{K}C27 + p_{KKK}C11 + p_{KK}C9 + p_{KKK}C11 + p_{KK}C11 +
    k1\overline{6}[G_{\beta}\gamma Ste20]B10 - k17C10
      dC11
                                                              \frac{dt}{dC12} = -p_{KKK}C12 - on_{KK}[Ste7]C12 - (off_{KKK} + off_K)C12 + on_{KKK}[Ste11]C6 + on_K[Fus3_{out}]C2 + off_{KK}C20 + ** off_{KK}C22 + off_{KK}C20 + ** off_{KK}C20 +
    k16[G_{\beta}\gamma Ste20]B12 - k17C12
    \frac{dC\bar{1}3}{..} = **on_{KKK}[Ste11pp]C6 - (**off_{KK} + off_{K})C13 - on_{KK}[Ste7]C13 + on_{K}[Fus3_{out}]C3 + off_{KK}C21 + **off_{KK}C24 + p_{KKK}C12 + off_{KK}C24 + 
      k16[G_{\beta}\gamma Ste20]B13 - k17C13
      \frac{dC14}{dt} =
      **on_{KK}[Ste11pp]C7 - (**off_{KK} + **off_K)C14 - on_{KK}[Ste7]C14 + off_{KK}C25 + **off'_{KK}C27 + p_{KKK}C15 + k16[G_{\beta}\gamma Ste20]B14 - k17C14
      \frac{dC15}{r} = -p_{KKK}C15 - (off_{KKK} + **off_K)C15 - on_{KK}[Ste7]C15 + on_{KKK}[Ste11]C7 + off_{KK}C23 + **off_{KK}'C26 + k16[G_{\beta}\gamma Ste20]B15 - k17C15
      dC^{dt}_{C16}
                                                                                         -**on_{KKK}[Ste11pp]C16 - on_{KKK}[Ste11]C16 - (off_{KK} + off_{K})C16 + on_{KK}[Ste7]C6 + on_{K}[Fus3_{out}]C4 + off_{KKK}C20 + **C10_{KKK}C20 + **C10_{KKK}
  off_{KKK}C21 + k16[G_{\beta}\gamma Ste20]B16 - k17C16
      \frac{dC17}{dC17} = - **on_{KKK}[Ste11pp]C17 - p_{K}C17 - (**off_{KK} + off_{K})C17 - on_{KKK}[Ste11]C17 + on_{K}[Fus3_{out}]C5 + off_{KKK}C22 + **off_{KKK}C22 +
  off_{KKK}C24 + k16[G_{\beta}\gamma Ste20]B17 - k17C17
      \frac{d\ddot{C}18}{d} =
                    **on_{KKK}[Ste11pp]C18 - (**off_{KK}' + **off_{K})C18 - on_{KKK}[Ste11]C18 + off_{KKK}C26 + **off_{KKK}C27 + p_{K}C17 + k16[G_{\beta}\gamma Ste20]B18 - k17C18 + off_{KKK}C27 + of
    \frac{dC19}{dt} = - ** on_{KK} [Ste11pp]C19 - (off_{KK} + ** off_{K})C19 - on_{KK} [Ste11]C19 + on_{KK} [Ste7]C7 + off_{KKK}C23 + ** off_{KKK}C25 + off_{KKK}C25 + off_{KKK}C25]
    k16[G_{\beta}\gamma Ste20]B19 - k17C19
      \frac{dC20}{dC20} = -(off_{KKK} + off_{KK} + off_{K})C20 - p_{KKK}C20 + on_{KKK}[Ste11]C16 + on_{KK}[Ste7]C12 + on_{K}[Fus3_{out}]C8 + k16[G_{\beta}\gamma Ste20]B20 - k17C20 + con_{KK}C20 + on_{KKK}C20 + on_{K
      dCB21
                                                                                    = **on_{KKK}[Ste11pp]C16 - (**off_{KKK} + off_{KK} + off_{K})C21 - p_{KK}C21 + on_{KK}[Ste7]C13 + on_{K}[Fus3_{out}]C9 + p_{KKK}C20 + on_{KKK}[Ste7]C13 + on_{KKK}[S
   dt \atop k16[G_{\beta}\gamma Ste20]B21 - k17C21 
    \frac{dC\dot{2}2}{dt} = -(off_{KKK} + **off_{KK} + off_{K})C22 - (p_{KKK} + p_{K})C22 + on_{KKK}[Ste11]C17 + on_{K}[Fus3_{out}]C11 + k16[G_{\beta}\gamma Ste20]B22 - k17C22
      \frac{dt}{dC23} = -(off_{KKK} + off_{KK} + ** off_{K})C23 - p_{KKK}C23 + on_{KKK}[Ste11]C19 + on_{KK}[Ste7]C15 + k16[G_{\beta}\gamma Ste20]B23 - k17C23
      \frac{dt}{dC24} =
      **on_{KKK}[Ste11pp]C17 - (**off_{KK} + **off_{KK} + off_{K})C24 - p_{K}C24 + on_{K}[Fus_{3out}]C10 + p_{KKK}C22 + p_{KK}C21 + k16[G_{\beta}\gamma Ste20]B24 - k17C24
      \frac{dC25}{r} = **on_{KK}[Ste11pp]C19 - (**off_{KK} + off_{KK} + **off_{K})C25 - p_{KK}C25 + on_{KK}[Ste7]C14 + p_{KK}C23 + k16[G_{\beta}\gamma Ste20]B25 - k17C25 + con_{KK}[Ste7]C14 + p_{KK}C23 + k16[G_{\beta}\gamma Ste20]B25 - k17C25 + con_{KK}[Ste7]C14 + p_{KK}C23 + k16[G_{\beta}\gamma Ste20]B25 - k17C25 + con_{KK}[Ste7]C14 + p_{KK}C23 + k16[G_{\beta}\gamma Ste20]B25 - k17C25 + con_{KK}[Ste7]C14 + p_{KK}C23 + k16[G_{\beta}\gamma Ste20]B25 - k17C25 + con_{KK}[Ste7]C14 + p_{KK}C23 + k16[G_{\beta}\gamma Ste20]B25 - k17C25 + con_{KK}[Ste7]C14 + p_{KK}C23 + k16[G_{\beta}\gamma Ste20]B25 - k17C25 + con_{KK}[Ste7]C14 + p_{KK}C23 + k16[G_{\beta}\gamma Ste20]B25 - k17C25 + con_{KK}[Ste7]C14 + p_{KK}C23 + k16[G_{\beta}\gamma Ste20]B25 - k17C25 + con_{KK}[Ste7]C14 +
      dt dC26
                                                          = -(off_{KKK} + **off_{KK}' + **off_{K})C26 - p_{KKK}C26 + on_{KKK}[Ste11]C18 + p_{K}C22 + k16[G_{\beta}\gamma Ste20]B26 - k17C26 + on_{KKK}[Ste11]C18 + p_{K}C22 + k16[G_{\beta}\gamma Ste20]B26 - k17C26 + on_{KKK}[Ste11]C18 + p_{K}C22 + k16[G_{\beta}\gamma Ste20]B26 - k17C26 + on_{KKK}[Ste11]C18 + p_{K}C22 + k16[G_{\beta}\gamma Ste20]B26 - k17C26 + on_{KKK}[Ste11]C18 + p_{K}C22 + k16[G_{\beta}\gamma Ste20]B26 - k17C26 + on_{KKK}[Ste11]C18 + on_{KK
      \frac{dt}{dC27}
                                                              = **on_{KKK}[Ste11pp]C18 - (**off_{KK} + **off_{KK}' + **off_{KK}')C27 + p_{KKK}C26 + p_{KK}C25 + p_{K}C24 + k16[G_{\beta}\gamma Ste20]B27 - k17C27 + p_{KKK}C26 + p_{KK}C25 + p_{KK}C26 + p_{KK}
```

```
\frac{d[Ste5_{in}]}{d[Ste5_{in}]} = -k22[Ste5_{in}] + k23B1
\frac{dt}{d[Fus3_{in}]}
                                              = -k47[Fus3_{in}] + k48[Fus3_{out}] + p1_{K}[Fus3p_{in}MAPK - P_{in}]
\frac{dt}{d[Fus3p_{in}]} = p2_K[Fus3pp_{in}MAPK - P_{in}] - a1_K[Fus3p_{in}][MAPK - P_{in}] + d1_K[Fus3p_{in}MAPK - P_{in}]
\frac{d[Fus3p_{in}MAPK - P_{in}]}{d[Fus3p_{in}MAPK - P_{in}]} = a1_{K}[Fus3p_{in}][MAPK - P_{in}] - d1_{K}[Fus3p_{in}MAPK - P_{in}] - p1_{K}[Fus3p_{in}MAPK - P_{in}]
\frac{d[Fus3pp_{in}]}{d[Fus3pp_{in}]} = -k49[Fus3pp_{in}] + k50[Fus3pp_{out}] - a2_K[Fus3pp_{in}][MAPK - P_{in}] + d2_K[Fus3pp_{in}MAPK - P_{in}MAPK - P_{in}MAP
\frac{d[Fus3pp_{in}MAPK - P_{in}]}{d[Fus3pp_{in}MAPK - P_{in}]} = a2_K[Fus3pp_{in}][MAPK - P_{in}] - d2_K[Fus3pp_{in}MAPK - P_{in}] - p2_K[Fus3pp_{in}MAPK - P_{in}]
\overline{d[MAPK - \overset{dt}{P_{in}}]}
\frac{dt}{-a1_{K}[Fus3p_{in}][MAPK-P_{in}] + (d1_{K} + p1_{K})[Fus3p_{in}MAPK-P_{in}] - a2_{K}[Fus3pp_{in}][MAPK-P_{in}] + (p2_{K} + d2_{K})[Fus3pp_{in}MAPK-P_{in}]}
\frac{d[Stel2_{active}]}{dt} = \frac{k28[Fus3pp_{in}]}{k29 + [Fus3pp_{in}]} - k30[Stel2_{active}]
\frac{d[Far1]}{dt} = \frac{k33[Ste12_{active}]^2}{k34^2 + [Ste12_{active}]^2} - k36[Far1][Fus3pp_{in}] - k35[Far1] + k37[Far1pp_{in}]
\frac{d[Far1pp_{in}]}{d[Far1pp_{in}]} = k36[Far1][Fus3pp_{in}] - k37[Far1pp_{in}] - k38[Far1pp_{in}] + k39[Far1pp_{out}] - k43[Far1pp_{in}][Cdc28] + k42[Far1pp_{in}Cdc28] + k42
\frac{dt}{d[Cdc28]} = k42[Far1pp_{in}Cdc28] - k43[Far1pp_{in}][Cdc28]
\frac{dt}{d[Far1pp_{in}Cdc28]} = -k42[Far1pp_{in}Cdc28] + k43[Far1pp_{in}][Cdc28]
\frac{d[Far1pp_{out}]}{d[Far1pp_{out}]} = -k40[Far1pp_{out}][G_{\beta}\gamma] + k41[Far1pp_{out}G_{\beta}\gamma] + k38[Far1pp_{in}] - k39[Far1pp_{out}]
\frac{d[Far1pp_{out}G_{\beta}\gamma]}{..}=k40[G_{\beta}\gamma][Far1pp_{out}]-k41[Far1pp_{out}G_{\beta}\gamma]
```

# Dynamic Studies of Scaffold-dependent Mating Pathway in Yeast

Danying Shao\*†, Wen Zheng\*†, Wenjun Qiu\*†, Qi Ouyang \*†, and Chao Tang\*†‡

\*Center for Theoretical Biology and †Department of Physics Peking University, Beijing, 100871, China

<sup>‡</sup>California Institute for Quantitative Biomedical Research
Departments of Biopharmaceutical Sciences and Biochemistry and Biophysics
UCSF Box 2540, University of California at San Francisco, San Francisco, CA 94143-2540

Danying Shao and Wen Zheng contributed equally to this work.

Correspondence: Address reprint requests to Qi Ouyang (qi@pku.edu.cn) or Chao Tang

(chao.tang@ucsf.edu).

Running Title: Dynamics of yeast mating pathway

**keywords:** Mating pathway, Scaffold, Shuttling, Phosphatase, Concentration effect, MAPK Cascade

Abstract: The mating pathway in Saccharomyces cerevisiae is one of the best understood signal transduction pathways in eukaryotes. It transmits the mating signal from plasma membrane into the nucleus through the G-protein coupled receptor and the mitogenactivated protein kinase (MAPK) cascade. According to the current understandings of the mating pathway, we construct a system of ordinary differential equations to describe the process. Our model is consistent with a wide range of experiments, indicating that it captures some main characteristics of the signal transduction along the pathway. Investigation with the model reveals that the shuttling of the scaffold protein and the dephosphorylation of kinases involved in the MAPK cascade cooperate to regulate the response upon pheromone induction and to help preserving the fidelity of the mating signaling. We explored factors affecting the dose-response curves of this pathway and found that both negative feedback and concentrations of the proteins involved in the MAPK cascade play crucial role. Contrary to some other MAPK systems where signaling sensitivity is being amplified successively along the cascade, here the mating signal is transmitted through the cascade in an almost linear fashion.

### INTRODUCTION

Cells have to respond to changes in the environment and/or to the external stimuli. This is accomplished by signal transduction pathways which sense the signal, transduce it and induce necessary changes in the cell, e.g. in gene expression. One of the best understood signaling pathways in eukaryotes is the mating pathway in the budding yeast *Saccharomyces cerevisiae* (1,2). Extensive studies of the mating pheromone response have contributed much to the understanding of the mechanisms of several conservative biological modules (3), such as the G protein cycle (2,4) and the mitogen-activated protein kinase (MAPK) cascade (2,5). Genetic, biochemical and molecular analysis of the response have combined to establish basic principles of the signaling and regulation. Many important discoveries are made in the study of this pathway, for instance, the concept of a kinase-scaffold protein (6) and the role of regulator of G protein signaling (RGS) proteins in the pathway (7).

The budding yeast can exist in either of the two types, MATa or  $MAT\alpha$ . These two types of cells will mate when each one receives the mating signal, a peptide pheromone, secreted by the opposite type (a-factor by MATa and  $\alpha$ -factor by  $MAT\alpha$ ). Once the pheromone binds to the seven-transmembrane-segment receptor in the plasma membrane (Ste2 in MATa and Ste3 in  $MAT\alpha$ ), the receptor is activated, which then activates the heterotrimeric G protein that couples to it (Fig. 1). The activated G protein transmits signal to multiple effectors, resulting in the beginning of the MAPK cascade, which is embedded in a scaffold protein Ste5. This cascade consists of three kinases: Ste11 (MAPKKK), Ste7 (MAPKK), and Fus3 (MAPK). The activation of the cascade finally leads to the phosphorylation of Fus3. The phosphorylated MAPK then travels into the nucleus, and transmits the signal to downstream effectors, leading to preparation for mating, including the cell cycle arrest in G1 phase to assure synchronism of the mating partners, the induction of new gene expression necessary for mating, and the polarized growth in the direction of the pheromone source.

Much qualitative and quantitative information in this pathway have been documented. With the increasing amount of experimental data and information, it is now possible to study this pathway quantitatively at a systems level. Several mathematical models have been employed to study this (8-12) and some other related systems (13-16), showing that mathematical modeling and simulation can be a powerful method in the analysis of functional and structural characteristics of biological pathways.

We set up an Ordinary Differential Equation (ODE) model to describe the mating pathway in budding yeast. Although several models have been employed to illustrate mechanisms in the pathway, there has not been one that integrates all the known essential features with a comprehensive analysis of its dynamic properties. Some models were constrained to a single step (8,12), while others oversimplified the regulations and functions of the scaffold (10). In our model, biochemical interactions, induced gene expressions which feed back to the pathway, and translocations of key components such as the scaffold protein Ste5 are all considered. Results from our model are consistent with a wide range of experimental data. We then tested the current understanding of regulations of cellular responses and further explored the intrinsic mechanisms in the

pathway, with special interest in the role of the scaffold protein Ste5. We find that the shuttling of the scaffold and dephosphorylation of the MAP kinases cooperate to regulate the responses upon pheromone induction, and to help keeping the fidelity of the mating pathway. We further explored the mechanisms of the dose-response curves of this pathway, and elucidated the role of enzyme concentration. We found that instead of an ultrasensitive response as in some other MAPK cascade (17), the mating signal here is transmitted through the cascade in an almost linear manner due to negative feedback.

## THE MODEL

We choose a mutant (TMY101), a MATa type of Saccharomyces cerevisiae, as the main modeling subject. In this type of cells, the gene BAR1 is deleted. In a wild type MATa cell, the product of BAR1 can be excreted from the cells and cleave the  $\alpha$ -factor. In order to simulate a continuous and constant  $\alpha$ -factor treatment, we use this mutant in our model.

The mating response can be divided into three modules in a temporal order: the activation of G protein cycle, the scaffold-depended MAPK cascade, and the downstream effects of activated MAPK (See Fig. 1). Viewing the response as a series of modules arranged in the temporal order can help to better understand the signaling process. Yet in our model, couplings and feedback between these modules are also taken into account.

The activation of G protein cycle.  $\alpha$ -factor secreted by  $MAT\alpha$  binds to and hence activates the seven-transmembrane-segment receptor (Ste2) on the plasma membrane surface of MATa. Pheromone binding enhances monoubiquitination of the receptor, and the ubiquitination in this case serves as a signal for endocytosis and delivery to the vacuole (18). This comprises a negative feedback loop (at short time scales). In our model, this process is treated as a process of accelerated degradation for simplicity. The synthesis of receptor Ste2 is included; the downstream effecter Ste12 is responsible for the gene expression of Ste2. Thus, it comprises a positive feedback (at long time scales).

The interaction between the activated receptor and  $G\alpha$  leads to some conformational changes, which enable  $G\alpha$  to release GDP and to bind GTP (19).  $G\alpha$ ·GTP can not interact with  $G\beta\gamma$ , resulting in a release of  $G\beta\gamma$  from the receptor. The  $G\gamma$  unit fixes the heterodimer on the plasma membrane surface, while the  $G\beta$  unit can interact with several effectors to transmit the signal. In this sense,  $G\alpha$  unit is a negative regulator of the pathway; it plays a role in an adaptational response to pheromone through preventing the availability of  $G\beta\gamma$  when there is no signal (20). The  $G\alpha$ ·GTP can be hydrolyzed into  $G\alpha$ ·GDP, which can re-associate with  $G\beta\gamma$  into a heterotrimer. The cycle of G protein is thus closed. Regulators of G protein signaling (RGS) proteins can accelerate the hydrolyzation of  $G\alpha$ ·GTP (21). In this pathway, the most important RGS protein is Sst2, which is considered in our model. The gene expression of Sst2 is also regulated by Ste12 in the downstream. Therefore, Sst2 is part of a negative feedback loop that leads to the adaptation (8,22). Since there is experimental evidence that the amount of  $G\alpha$  increases significantly when the cells are treated with pheromone (8), we add G protein synthesis in our model. It is commonly accepted that Ste12 transcripts  $G\alpha$  genes. According to the above description, we formulate the reactions

in G protein cycle as follows:

$$Ste2 + \alpha - factor \underset{k2}{\overset{k1}{\rightleftharpoons}} Ste2_{active}$$

$$Ste2_{active} \xrightarrow{k3} \qquad (degradation)$$

$$\xrightarrow{k4,k5,ste12a} Ste2 \qquad (synthesis)$$

$$Ste2 \xrightarrow{k7}$$

$$G \xrightarrow{Ste2_{active}} G\alpha \cdot GTP + G\beta\gamma$$

$$\xrightarrow{k9,k10,ste12a} G \xrightarrow{k12}$$

$$G\alpha \cdot GTP \xrightarrow{k13} G\alpha \cdot GDP$$

$$G\alpha \cdot GTP + G\beta\gamma \xrightarrow{k15} G$$

where k's are the kinetic parameters, the protein above or below the arrow is enzyme or transcription factor of the reaction.

The scaffold depended MAPK pathway. The released  $G\beta\gamma$  has several effectors (23). One effector for mating is Ste20, the first p21-activated protein kinase to be identified in any eukaryote (24). Ste20 is also activated by Cdc42, which is regulated by Cdc24. However, this process is not included in our model because the Cdc42 binding domain of Ste20 has been shown to be dispensable for pheromone signaling in yeast (25,26), and there should be enough active Cdc24<sup>GEF</sup> and Cdc42 constitutively at the membrane to activate the amount of Ste20 required for initial signaling. Besides, mutants in Cdc24 do not have much influence on the pathway (27,28).

Another effector of  $G\beta\gamma$  is the scaffold protein Ste5. The correlation between the disruption of the Ste4( $G\beta$ )-Ste5 interaction and sterility confirms the importance of this interaction in signal transduction (29).  $G\beta\gamma$  can bind to Ste5 on the LIM domain of Ste5, which is required for Ste11 (MAPKKK) activation (30), probably through inducing a conformational change that enhances Ste20-dependent activation of Ste11. Also it interacts with Ste5 in the RING-H2 domain which is essential for Ste5 oligemerization (31).

Most scaffolds are contained in the nucleus during vegetative growth. Upon pheromone induction they undergo enhanced exportation from the nucleus and localize at the shmoo tip (1). Although the detailed controlling mechanism of exportation of Ste5 is not clear, it is plausible that mating pheromone increases the rate of Ste5 export (57). Here, we utilize an active control mechanism where the import rate is kept constant, while the export rate is dependent on the total concentration of the released  $G\beta\gamma$ . When there is no signal, the export rate is very low, keeping most scaffolds in the nucleus. When the mating signal opens the G-protein cycle, released  $G\beta\gamma$  enhances the export rate, driving scaffolds to the shmoo tip. In this way, the localization of the scaffolds can be regulated by G protein cycle.

The mating pathway is highly dependent on the scaffold protein Ste5. First, Ste5 functions as an adapter protein. It recruits Ste11 to the plasma membrane, where Ste20 is also tethered, to facilitate Ste11's activation (28), triggering the MAPK cascade. Another

function for Ste5 is scaffolding. Ste5 tethers Ste11 (MAPKKK), Ste7 (MAPKK) and Fus3 (MAPK) to form a complex (32), keeping the kinases and their substrates in proximity, as well as preventing the influence of phosphatases. This function is supposed to be important in enzyme regulation and in preventing cross talk (33,34).

When Ste5 is in the cytosol, it can form scaffold-kinase complexes with Ste11, Ste7 and Fus3. Every kinase binding site on the scaffold is in one of the three possible states: without a kinase, with an unphosphorylated kinase, or with a dual phosphorylated kinase. So for scaffold-kinase complexes in solution, there are  $3 \times 3 \times 3 = 27$  states: B1, B2, ...B27 (See Fig. 2A).  $G\beta\gamma$  can bind to Ste20 and Bi (i = 1, 2...27). Because Ste20 is already on the plasmid membrane through the interaction with Cdc42 before signaling, and scaffold must shuttle out from the nucleus to bind to  $G\beta\gamma$ , we assume that  $G\beta\gamma$  first binds to Ste20, then binds to Bi. Once Bi binds to  $G\beta\gamma$ Ste20 complex, it is fixed at the plasmid membrane and the whole complex is denoted Ci (see Fig. 2B). Ci and Bi are the same in the interaction with MAPK kinases, except that Ci can phosphorylate Ste11 while Bi can not.

$$\begin{split} G\beta\gamma + Ste20 & \stackrel{k18}{\rightleftharpoons} G\beta\gamma Ste20 \\ Bi + G\beta\gamma Ste20 & \stackrel{k16}{\rightleftharpoons} Ci (i=1,2,...27) \\ Ste5_{in} & \stackrel{k22}{\rightleftharpoons} B1 \end{split}$$

where " $Ste5_{in}$ " denotes Ste5 in the nucleus, B1 denotes Ste5 outside of the nucleus, and k22 is dependent on the total concentration of released  $G\beta\gamma$ :  $k22 = 0.0003 + 0.3 \frac{G_{act}}{G_{act} + 2500}$ ,  $G_{act} = [G_{\alpha} \cdot GTP] + [G_{\alpha} \cdot GDP]$ .

Ste11 is the MAPKKK of yeast pheromone pathway, which consists of an N-terminal regulatory domain and a C-terminal kinase region (1). The interaction of these two domains keeps Ste11 in an anti-self state. CBD domain in N-terminal contains serine and threonine residues that can be phosphorylated by Ste20. Ste20-mediated phosphorylation of these residues activates Stell (35). In addition to recruiting Stell to a pool of its activators Ste20, Ste5 also binds to Ste11 in its N-terminal, making the CBD domain in N-terminal more accessible for Ste20. Ste50 also helps to make the CBD domain more accessible to Ste20 by a direct interaction between the SAM domain of itself and the SAM domain of Ste11 (36), but is less essential than Ste5. Cells lacking Ste50 are not truly sterile, thus we do not include Ste50 in our model. Ste11pp phosphorylates the target residues in the activation loops of Ste7 (MAPKK), and activates it (37). The activated Ste7pp then phosphorylates, and activates its targets, the MAPKs Fus3 and Kss1 (38). In the mating pathway, Fus3 plays a much more important role, while Kss1 is the main MAPK in the filamentation-invasion pathway in nitrogen starved cells (5). Thus we do not consider Kss1 in our model, although analysis about crosstalk will be given in the Discussion. While Ste11 and Ste7 are predominantly cytoplasmic proteins, Fus3 can shuttle between the nucleus and the cytoplasm. It concentrates in the nucleus after activation, thus bringing the signal to the nucleus (7,35,39). There are also several feedback loops: Ste11 (MAPKKK) undergoes ubiquitination and MAPK dependent degradation (40); Ste7 (MAPKK) is assumed to undergo enhanced degradation after phosphorylation (41). In addition, Ste7pp in the scaffold is assumed to be hyper-phosphorylated by activated Fus3pp, which reduces the binding efficiency between Ste7pp and the scaffold sharply (42).

All the kinases can be categorized into two pools: on the scaffold and in the solution. We assume that phosphorylation on scaffold employs a processive mechanism, while phosphorylation in solution is distributive(11). Processive mechanism means that the active kinase collides with and binds to a substrate, phosphorylates it once, then it may slides to align the second phosphorylation site of the substrate with the active site of the kinase, phosphorylates the substrate a second time before finally dissociates. Distributive mechanism means that the active kinase collides with and binds to a substrate, phosphorylates it once and releases the monophosphrylated product, which then collides with a second molecule of the active kinase, and is phosphorylated a second time (14). In our model, we assume that a kinase is activated when and only when it is dual phosphorylated, while a partial phosphorylated kinase possesses no activity. Dual phosphorylation in a distributive manner could lead to a sharp, sigmodial stimulus-response curve (14,17), leading an all-or-none cell fate (14). However, the scaffold might diminish this property if phosphorylations on the scaffold occur in a processive manner (11). The dephosphorylations in the solution employ the distributive mechanism while dephosphorylations in scaffolds are precluded in our model due to sterical obstruction of the phosphatases groups. The proteins responsible for dephosphorylation of Ste11pp and Ste7pp are not clear. In the model, we add two proteins with constant concentration to dephosphorylate Ste11pp and Ste7pp respectively. There are several phosphatases for Fus3pp: the dual-specificity phosphatase Msg5 (equally distributed in nucleus and cytoplasm), and the tyrosine phosphatases Ptp3 (cytoplasm) and Ptp2 (nucleus), all of which can result in the inactivation of Fus3pp (7,43,44). The basal level of Fus3 phosphorylation is controlled mainly by Ptp3, the amount of which is constant during the stimulation (44). Pheromone treatment induces the expression of Msg5 through the effects of Ste12 (43), which then acts together with Ptp3 to inactivate Fus3pp. In our model, we use MAPK-P with an initial concentration and with a synthesis rate regulated by Ste12 to represent these three phosphatases. A recent experiment shows that different inputs by Ste5 and Msg5 phosphatase lead MAPK cascade to multiple outcomes (45), indicating that MAPK-P is a key regulator in the network. Reactions of MAPK cascade in cytosol are formulated as follows:

```
Ste11p + MAPKKK - P \rightleftharpoons Ste11pMAPKKK - P \rightarrow Ste11 + MAPKKK - P Ste11pp + MAPKKK - P \rightleftharpoons Ste11ppMAPKKK - P \rightarrow Ste11p + MAPKKK - P Ste7 + Ste11pp \rightleftharpoons Ste7Ste11pp \rightarrow Ste7p + Ste11pp Ste7p + MAPKK - P \rightleftharpoons Ste7pMAPKK - P \rightarrow Ste7 + MAPKK - P Ste7p + Ste11pp \rightleftharpoons Ste7pSte11pp \rightarrow Ste7pp + Ste11pp Ste7pp + MAPKK - P \rightleftharpoons Ste7ppMAPKK - P \rightarrow Ste7p + MAPKK - P Ste7pp + MAPKK - P \rightleftharpoons Ste7ppMAPKK - P \rightarrow Ste7p + MAPKK - P Fus3_{out} + Ste7pp \rightleftharpoons Fus3_{out}Ste7pp \rightarrow Fus3_{out} + Ste7pp Fus3_{out} + MAPK - P_{out} \rightleftharpoons Fus3_{out}MAPK - P_{out} \rightarrow Fus3_{out} + MAPK - P_{out}
```

```
\begin{split} Fus3p_{out} + Ste7pp &\rightleftharpoons Fus3p_{out}Ste7pp \rightarrow Fus3pp_{out} + Ste7pp \\ Fus3pp_{out} + MAPK - P_{out} &\rightleftharpoons Fus3pp_{out}MAPK - P_{out} \rightarrow Fus3p_{out} + MAPK - P_{out} \\ Ste7 + Fus3_{out} &\stackrel{k24}{\rightleftharpoons} Ste7Fus3_{out} \\ Ste11 &\stackrel{k26,Fus3pp_{out}}{\longrightarrow} \\ Ste7pp &\stackrel{k27}{\longrightarrow} \end{split}
```

where p indicates once phosphorylation, and pp indicates twice phosphorylation.

As for the scaffolds, we made the following assumptions in the model: (a) Inactive kinase can bind to Bi and Ci. On the scaffold, this inactive kinase can either dissociate from the scaffold without phosphorylation or undergo processive phosphorylation before getting off the scaffold if its upstream kinase happens to be on the same scaffold and in the active state. (b) Dephosphorylations on scaffolds are precluded due to sterical obstruction. (c) There is no binding of partially activated kinases to the scaffold proteins. For free fully activated kinases, only Ste11pp can bind to the scaffold. Experimental evidence indicates that active Ste5 can also accept Ste11pp activated by other pathways and channel those signals to Fus3 (46), and that a greater amount of scaffold proteins interact with Ste11 rather than the other two kinases. As for Ste7, it undergoes hyper-phosphorylation by activated Fus3pp, which accelerates its dissociation from the scaffold (42). Fus3pp dissociates rapidly from the scaffold after phosphorylation (47) and travels into the nucleus. Thus, the reassociation of Ste7pp and Fus3pp to scaffold seems to be unlikely. (d) Scaffold molecules do possess some catalytic properties (33), so that the reaction rates within a scaffold complex are greater than in the solution. Moreover, Ste7 and Fus3 can bind firmly (23), and the residues for Ste7 binding in Fus3 are the same as the residues for Ste5 binding (48), so it is reasonable to assume that Ste7 competes with Ste5 for binding to Fus3. Fig. 3 illustrates the scaffolddependent reactions of Ste11.

**Downstream effects**: After activation, Fus3pp dissociates rapidly from the scaffold, while the scaffold remains tethered to the plasma membrane and partly in the solution (47,49), acting as a platform for activation of many molecules of Fus3 and leading to the propagation of the signal. The activated Fus3pp transmits the signal into the nucleus, resulting in the activation of transcription and the induction of cell cycle arrest. Fus3pp is assumed to mediate the pheromone-induced transcription of PRE-containing genes through phosphorylation and activation of at least three nuclear proteins: Dig1, Dig2 and Ste12 (1,2). In unstimulated cells, Dig1 and Dig2 bind to and thus repress Ste12 (50). Fus3pp phosphorylates Dig1, Dig2 and Ste12, and induces the release of Ste12 from the complex (50,51). The free Ste12 then interacts with other proteins of the transcription machinery and thereby activates transcription of many different genes. Among the products of these genes are proteins that activate (e.g. Fus3, receptor) or inhibit (e.g. Msg5, Sst2) the pathway (2). Therefore, the transcription affords several feedback loops in the pathway. Another important substrate of Fus3pp is Far1. Activated Ste12 increases the transcription of Far1, and Fus3pp is able to phosphorylate Far1 and thus to stabilize it (52). Far1 is a bi-functional scaffold protein. In the cytoplasm, Far1 is involved in polarized growth; while in the nucleus, it has a key function in controlling cell cycle (53,54). Far1 inhibits Cln-Cdc28 complex, the master regulator of the yeast cell cycle in G1 phase. In our model, the binding of Far1 to Cln-Cdc28 is treated as a symbol for cell cycle arrest. Ste12 is also the transcription factor of Bar1, which is then excreted from the cell and inactivates  $\alpha$ -factor (53). It is deleted in our model because most experiment results that we use were from  $bar1\triangle$  mutants. However,  $bar1\triangle$  could cause more than 100-fold sensitivity increase in downstream transcription response (12). Reactions in the downstream are:

$$Fus3pp_{in},k28,k29 \atop k30} Ste12_{active}$$

$$\frac{Ste12_{active},k31,k5}{Ste12_{active},k33,k5} MAPK - P_{out}$$

$$\frac{Ste12_{active},k33,k34}{Ste12_{active},k35} Fus3_{out}$$

$$Ste12_{active},k33,k34 Far1 \xrightarrow{Fus3pp_{in},k36} Far1pp_{in} \xrightarrow{k38} Far1pp_{out}$$

$$Far1pp_{out} + G\beta\gamma \xrightarrow{k40} Far1pp_{out}G\beta\gamma$$

$$Far1pp_{in} + Cdc28 \xrightarrow{k42} Far1pp_{in}Cdc28$$

$$Ste12_{active},k44,k45 \atop k46} Sst2_{active}$$

$$Fus3_{in} \xrightarrow{k47} Fus3_{out}$$

$$Fus3pp_{in} \xrightarrow{k49} Fus3pp_{out}$$

$$Fus3pp_{in} + MAPK - P_{in} \rightleftharpoons Fus3pp_{in}MAPK - P_{in}$$

$$\Rightarrow Fus3p_{in} + MAPK - P_{in} \rightleftharpoons Fus3p_{in}MAPK - P_{in}$$

$$\Rightarrow Fus3p_{in} + MAPK - P_{in} \rightleftharpoons Fus3p_{in}MAPK - P_{in}$$

$$\Rightarrow Fus3p_{in} + MAPK - P_{in} \rightleftharpoons Fus3p_{in}MAPK - P_{in}$$

$$\Rightarrow Fus3_{in} + MAPK - P_{in} \rightleftharpoons Fus3p_{in}MAPK - P_{in}$$

$$\Rightarrow Fus3_{in} + MAPK - P_{in} \rightleftharpoons Fus3p_{in}MAPK - P_{in}$$

Spatial location: we consider two compartments in the cell: the nucleus and the shmoo tip (a projection towards the direction of pheromone formed as a result of polarized growth). The nucleus is where downstream effects take place. The shmoo tip is where the many signaling proteins are concentrated (55) and the main place for the upstream reactions, including the G-protein cycle and Ste5-related reactions. Thus in our model we neglected the rest part of the cytosol. In other words, we "restricted" the cytosol to the shmoo tip. The scaffold protein Ste5, the MAPK Fus3 (both activated and inactivated), and Far1 are shuttling between the nucleus and the shmoo tip. Ste5 is mostly sequestered in the nucleus in the absence of pheromone while pheromone enhances nuclear exportation of Ste5 (56). Nuclear localization of Fus3 is slightly enhanced by the pheromone treatment (7,39,57,58).

**The mathematical model**: we employ a set of ordinary differential equations (ODEs) to describe the changes in the concentration of proteins involved in the mating pathway. Generally, in a system of l biochemical species with the concentration  $c_i$  (i=1,2,···,l) and m biochemical reactions with the rates  $v_j$  (j=1,2,···,m), the following series of equations can be utilized to describe the biochemical mechanism in the system:

$$\frac{dc_1}{dt} = f_1(c_1, c_2, \dots, c_l) = n_{11}v_1 + n_{12}v_2 + \dots + n_{1m}v_m$$

$$\frac{dc_2}{dt} = f_2(c_1, c_2, \cdots, c_l) = n_{21}v_1 + n_{22}v_2 + \cdots + n_{2m}v_m$$

$$\vdots$$

$$\frac{dc_l}{dt} = f_l(c_1, c_2, \cdots, c_l) = n_{l1}v_1 + n_{l2}v_2 + \cdots + n_{lm}v_m$$

The quantity  $n_{ij}$  denotes the stoichiometric coefficient. The rate of a reaction is a function of the concentrations of substrates, products and probable effectors (10). If we treat the gene expression as a special kind of reaction which can be described with Hill functions, the equations listed above can be employed to describe the dynamics of our system. In our model, all the unbound substances in various phosphorylation states and complexes formed by them are viewed as individual species. All complex formations, dissociations, degradations, phosphorylations, and dephosphorylations are treated as reactions. The parameters and initial concentrations in the model are derived from experiments whenever possible. For the remaining parameters, some are determined by fitting the results of the model to indirect experiments; others are estimated according to the mechanisms and similar reactions in other organisms. The list of the model parameters as well as the detailed ODEs are presented in  $SUPPLEMENTAL\ MATERIAL$ . For simulation, we use Matlab, version 6.5 (The MathWorks, Natick, MA).

## RESULTS

## A. Temporal characteristics

**G-protein cycle.** Fig. 4 summarizes the dynamics of the G-protein cycle. Upon saturated pheromone induction (1  $\mu$ M  $\alpha$ -factor), the level of the activated G-protein climbs up rapidly, reaches its peak at about 30 s, and then gradually declines to a bottom at about 7.5 minutes before it gradually increases again, as shown in Fig. 4A. The simulation result (solid line) fits quite well with the experiment data (12) (circles with error-bars). One crucial factor that might contribute to enhancing the closure of G-protein cycle is the endocytosis of activated receptor Ste2. This hypotheses is supported by experiment with mutant  $Ste2^{300\triangle}$  (the C-terminal tail of the  $\alpha$ -factor receptor gene STE2 is removed to impair its endocytosis) (12). We slowed down the degradation rate of the active Ste2 to simulate the  $Ste2^{300\triangle}$  mutant. Consistent with the experiment data, the closure of G-protein cycle is apparently impaired and the amount of the activated G-protein levels off after reaching its peak, as shown in Fig. 4B. The behavior of  $Ste2^{300\triangle}$  cells (dashed line for simulation, and up-triangle with error-bars for experimental data) indicates that endocytosis is a key factor that causes G protein cycle to close up. Fig. 4A shows that after about t=10 minutes, the activated G-protein continues to rise steadily. We attribute it to protein synthesis, because that is the time scale for gene expression. To test this hypothesis, we delete protein synthesis of all proteins considered in our model, and find that the level of the activated G-protein does not rise in the simulation. The behavior of cycloheximide treated cells, as shown in Fig. 4B (dotted line for simulation and square with error-bars for experiment data (12)), supports this hypotheses. For comparison, time-course for TMY101 cells is also shown in the figure (solid line for simulation, and circles with error-bars for experiment data).

Binding of Ste20 to  $G\beta\gamma$ : After activation,  $G\beta\gamma$  activates two effectors: Ste20 (MAP-KKKK) and Bi (Scaffold in the solution), hence transmitting the signal downwards. In our model, the time-course for the pheromone induced binding of Ste20 to  $G\beta\gamma$  fits well with experimental data (24), (Fig. 4C). It shows that Ste20 binds quickly to  $G\beta\gamma$  during the first 5 minutes. The binding slows down afterwards and then speeds up. This time-course seems to follow the activation of G-protein cycle upstream (Fig. 4A), consistent with the presumption that  $G\beta\gamma$ -dependent activation rather than Cdc42-dependent activation of Ste20 is critical in the mating pathway.

**Activation of MAPK pathway**: Following the recruitment of the scaffold protein Ste5 to the membrane, signal passes down through the MAPK cascade (Fig. 4D). Note that except for the gradual recruitment of Ste5, the signal transduction is very fast.

**Downstream effects**: Fig. 4E shows the activation of Ste12, Far1ppG $\beta\gamma$  and Far1ppCdc28 to illustrate the downstream effects of Fus3pp (MAPK). The formation of complex Far1pp-Cln-Cdc28 is responsible for the cell cycle arrest, and Far1pp-G $\beta\gamma$  causes the polarized growth and the formation of the shmoo tip. Since G $\beta\gamma$  is a part of the upstream complex involving Ste5 to provide a scaffold for the MAPK pathway, excess G $\beta\gamma$  is not available until the pathway is attenuated to some extent. Thus the curve for Far1-G $\beta\gamma$  begins to rise at 20 minutes after pheromone treatment, relatively late compared to other downstream effectors.

# B. Features of the pathway

Scaffold shuttling and dephosphorylation cooperate to regulate MAPK cascade quantitatively and to keep its fidelity to mating signal: One of the most distinctive features of the mating pathway is its dependence on scaffold. Interestingly, we found that the amount of Ste5 localizing out of nuclear upon pheromone induction coincides with activated Fus3pp (MAPK) in the dose-response curve, as shown in Fig. 5A. Thus we speculate that the mating pathway is tightly controlled by scaffold protein Ste5 and its shuttling. It is highly possible that with different concentrations of scaffolds out of nuclear, the efficiency of MAPK cascade varies. When scaffold concentration is relative low, the pathway efficiency should increase with scaffold concentration because of the catalytical function and spacial protection function of the scaffold. When scaffold concentration is too high, it may reduce the mating efficiency because effective complex suitable for signal transduction is hardly found (9). To investigate this possibility, we shut down the shuttling of the scaffold protein Ste5 and varied the concentration of total Ste5 at the shmoo tip from 1nM to 1000 nM. For the levels of pheromone induction, we varied the concentration from 0.1 nM to 1000 nM. Consistent with our expectation, the intensity of the output of MAPK cascade, indicated by the amount of the activated Fus3pp, first increases with scaffold concentration, then decrease, with the optimal scaffold concentration around 100 nM, as shown in Fig. 5B. Thus, within our simulation range, pathway efficiency monotonously increases with scaffold available in the shmoo tip, and it is reasonable to suggest that scaffold proteins have the ability to quantitatively control the strength of signal transmission through the MAPK cascade. This function of Ste5 could be achieved by facilitating Ste11's activation by Ste20 through binding to  $G\beta\gamma$ , concentrating MAPK cascade components, and sequestering them from inhibition by phosphatases (See below).

Dephosphorylation is the other mechanism we speculate that might contribute to controlling the activation of the mating pathway. To test this hypothesis, we varied MAPK-P's concentration (both in the nucleus and at the shmoo tip) from 100 nM down to 0 nM. As expected, Fus3pp (MAPK) shows super-sensitivity upon pheromone induction, see Fig. 5C. How does dephoshorylation control activation of MAPK cascade? Since the kinases are exposed to phosphatase only when they are in the cytosol and the scaffold could help to prevent the influence of the phosphatase on the kinases bound to it, we suggest that certain level of phosphatase concentration can keep the kinase phosphorylation in the cytosol at a very low level, and thus constrain the signal transduction on the scaffold. Hence, when the phosphatases are attenuated, a large amount of activated Ste11pp, Ste7pp, and Fus3pp could be accumulated in the cytosol even at lower level of scaffold protein recruited to the shmoo tip, bypassing the control of scaffold protein Ste5. In short, the shuttling of the scaffold and the dephosphorylation of the MAP kinases cooperate to control the activation of MAPK cascade quantitatively.

The cooperation between the scaffold shuttling and the MAPK dephosphorylation is also crucial to the specificity of the pathway. There are at least 5 MAPK signal transduction pathways in budding yeast (5), some of which share the same proteins, such as Ste11 and Ste7. A big puzzle is how specificity is achieved. To investigate the mating pathway's ability to isolate inappropriate signals leaking in from other pathways such as the filamentationinvasion pathway, we tested the behavior of some mutants. We set all Ste11 molecules in the dual phosphorylated state at t=0 and shut off MAPKKK-P to simulate the constitutive activation of Stellpp in invasive growth in the absence of mating pheromone. From the time-course curve of Fig. 6A, we observed that although Ste7pp (MAPKK) is activated to a relatively low extent, little Fus3pp (MAPK) is stimulated. We then set all Ste7 molecules in the dual phosphorylated state at t=0 and shut off MAPKK-P to simulate constitutive Ste7 activation, and found that Fus3pp (MAPK) could only be activated transiently in cytosol the activation dropped down immediately (within ten seconds) (inset in Fig. 6B). This result is consistent with the experiments, revealing that persistent activation by constitutive Ste7pp fails to support Fus3-dependent mating in the yeast (42). Further simulation by removing MAPK-P indicates that the later deactivation of Fus3pp is caused by dephosphorylation (Fig. 6B). Note that only the kinases in the solution are exposed to phosphatases, and the scaffold could help to shield from the influence of phosphatases. Thus, phosphatases constrain the signal on the scaffold. While on the scaffold, Ste7pp prefers to phosphorylate Fus3 instead of Kss1, the signal constrained on the scaffold will lead to the activation of Fus3, rather than Kss1, the main MAPK in the filamentation-invasion pathway in nitrogenstarved cells. That means that the activation of Fus3 strictly relies on the scaffold protein. When there is no pheromone induction, no scaffold protein is recruited to the shmoo tip, and the phosphatases inhibit the basal activation of Fus3. With pheromone treatment, the active scaffold proteins (the activation means recruitment to the plasma membrane in our model) help to assemble MAP Kinases cascade components and accelerate the activation of Fus3pp, the main MAPK in the mating pathway. When activation exceeds dephosphorylation, the signal will be transmitted into the nuclear.

To conclude, when there is no pheromone induction, phosphatases repress the activation of the mating pathway and prevent inappropriate signals from leaking in. When pheromone exists, scaffolds are recruited to the shmoo tip by the activated G protein, gathering MAPK cascade components and sequestering them from phosphatases so that the mating signal can be transmitted downwards. Thus, mating pathway is highly depended on scaffolds. This conclusion is consistent with experiments (45). Although there are other factors that contribute to suppress the crosstalk between the two pathways (34), the mechanism outlined above could also play an important role.

Desensitization to pheromone induction: The amount of activated Fus3pp (MAPK) decreases with time even when the cells are exposed to prolonged  $\alpha$ -factor induction, as shown in Fig. 7A our wild type cell simulation (solid line). This indicates desensitization effect. Desensitization is a key feature of the pathway, which enables cells to reenter the cell cycle to resume vegetative growth. We investigated the possible factors that might contribute to this desensitization, and found that multiple negative feedback loops, such as the degradation of Ste11 (MAPKKK) and Ste7pp (MAPKK), the synthesis of Msg5 (MAPK phosphatase) and Sst2, should be the major cause. As shown in Fig. 7A, a wild type cell with all the negative feedback shows desensitization, but a mutant without these negative feedback does not (dashed line): the activation of Fus3pp does not decrease even after one hour treatment with saturating pheromone.

Another important cause is the negative regulation of Ste7's (MAPKK) binding ability to scaffold by Fus3pp (MAPK). In the scaffold, Ste7pp, which undergoes feedback phosphorylation by activated Fus3pp, dissociates more quickly from the scaffold (42), hence exposes itself to ubiquitination and degradation (41,59). This feedback can also accelerate the disassembling of scaffold complexes. In our model, we assume that the dissociation rate for Ste7pp on scaffolds with both Ste7pp and Fus3pp is larger than that for Ste7pp on scaffolds without Fus3pp. If we change former parameter to be the same as the latter one, activation of Fus3pp continues to rise up after prolonged stimulation, as shown in the dashed line in Fig. 7B. On the other hand, the dissociation rate for Ste7pp from the scaffold without Fus3pp has to be relatively slow to keep the intensity of the signal transduction. A mutant in which the dissociation rate for the normal phosphorylated Ste7pp (with no Fus3pp on the scaffold) is enhanced (dotted line in Fig. 7B) results in a very low intensity of the signal transduction. Thus our model shows that differentiated binding abilities of Ste7 to the scaffold ensure the right behavior of the MAPK cascade.

**Sensitivity to pheromone**: Aside from the temporal characteristics, the sensitivity to different levels of pheromone induction is another key feature of the signal transduction

pathway. We compared the dose-response curves predicted by our model to those observed in experiments (12,45), and there is a quantitative agreement (Fig. 8). While our model specifically simulates the  $\Delta Bar1$  strain, our results are also consistent with experiments of the wild type, taking into account the 100 fold sensitivity shift (8,60). In our study, cells are assumed to be treated with indicated concentration (0.001nM-1000 nM) of  $\alpha$ -factor for 20 min. The response of every component to a certain concentration of  $\alpha$ -factor is represented by the maximum amount of that component. For the activated Ste2, the peak is obtained within seconds; for G protein activation (measured by the sum of  $G\alpha$ -GTP and  $G\alpha$ -GDP), the maximum amount is achieved at about 30 seconds; for Ste11pp, Ste7pp and Fus3pp, the peak value appears at about 20 minutes. These different time scales are consistent with experimental observations and illustrate the characteristics of different components' activation. All these curves are normalized. We fit the dose-response curves of Ste2a, G protein activation, scaffold recruitment, Ste11pp, Ste7pp and Fus3pp with the Hill Function:

$$[out] = \frac{A \times [in]^n}{[in]^n + th^n},$$

where n is the Hill coefficient and th the threshold where the response reaches half its maximum (Table I). Note that the sensitivity to  $\alpha$ -factor is well conserved throughout the whole pathway, from receptor Ste2 at the very beginning throughout the MAPK cascade.

As stated before, our model separates the whole mating pathway into different modules. It is interesting to explore the dose-response curve of each module. The first module is ligand binding. The reason why the dose-response curve of the activated Ste2 is a Hill function with  $n \approx 1$  is the reaction it takes. Consider the reaction  $Ste2 + \alpha \stackrel{k1}{\rightleftharpoons} Ste2_{act}$ , with the input concentration of  $\alpha$  (on the left hand side) fixed at  $[\alpha]$  and the total concentration of Ste2 fixed at  $[Ste2]_0$ . At steady state:

$$[Ste2_{act}] = \frac{A_1 \times [\alpha]}{[\alpha] + th_1},\tag{1}$$

with  $A_1 = [Ste2]_0$  and  $th_1 = k2/k1$ , which takes the value  $th_1 = 5.0$  with our choice of parameters k1 and k2. The coefficients derived from our simulation with our whole model are n = 0.9, th = 9.1. The differences come from protein synthesis, degradation and signal dependent feedback. When these effects are deleted from the whole model, the simulation results agree perfectly with the analysis.

We further explored the dose-response curve for G protein activation. Again, for simplicity, we do not take into account in the calculation the signal-dependent production, degradation and the feedback of the RGS protein (we treat the amount of Sst2 as a constant) and look for the steady state solution. The result of the calculation is:

$$[G\beta\gamma] = \frac{[G]_0 \times [Ste2_{act}]}{[Ste2_{act}] + \frac{k13'}{k8}},\tag{2}$$

where k13' = k13 + k14[Sst2]. To get this result, we made the assumption that the hydrolysis is a relatively slow process, so that  $\frac{k13'}{k15} \ll [G]_0$ . If we substitute Eq.(1) into Eq.

(2), we get:

$$[G\beta\gamma] = \frac{A_2 \times [\alpha]}{[\alpha] + th_2},\tag{3}$$

where  $A_2 = [G]_0 \frac{A_1}{A_1 + \frac{k_1 3'}{k_8}}$  and  $th_2 = th_1 \frac{k_1 3'}{k_8 \times (A_1 + \frac{k_1 3'}{k_8})}$ . Eq. (3) is a Hill function with n = 1.0 and th = 2.9. The simulation result of our whole model is n = 1.0 and th = 4.8. Again, if the protein synthesis, degradation and feedbacks are deleted, the simulation results of n = 1.0 and th = 2.9 agree well with the analysis.

We see that the curve of  $[G\beta\gamma] \sim [\alpha - factor]$  employs the same type of function as the curve of  $[Ste2_{act}] \sim [\alpha$ -factor], with the same Hill coefficient. The only difference is that there is a shift in threshold.

The above analysis suggests that the amount of Sst2 can affect the threshold of the dose-response curve. Fig. 9A indicates that Sst2, which can accelerate the closure of the G protein cycle, is indeed a key regulator of mating pathway's sensitivity. Comparison of the dose-response curves for mutants SST2 $\triangle$  (dotted line for simulation, up-triangle with errorbars for experiment data), wild type cells (solid line for simulation, squares for experiment data (8)) and  $2 \times SST2$  (dashed line for simulation, down-triangle for experiment data) clearly shows that the system is sensitive to the amount of Sst2, which is consistent with the theoretical analysis above. Another mutant we studied in this module is one with excess  $G\beta\gamma$  copies. Fig. 9B compares dose-response curves for TMY101 cells (solid line for simulation, and squares for experiment data) and cells with  $2 \times G\beta\gamma$  (dashed line for simulation, and up-triangles for experiment data(8)), indicating that  $G\beta\gamma$  alone is sufficient to switch on the downstream signal transduction.

The next module in the pathway is the scaffold-dependent phosphorylation cascade. Again, we do not take into account protein synthesis and degradation, nor all the feedback. Here,  $G\beta\gamma$  is the input, while the concentration of Ste11pp, Ste7pp and Fus3pp is chosen as the output.  $G\beta\gamma$  first binds to Ste20, then binds to scaffold in the solution to generate C (we use C to indicate the ensemble of C1, C2,...C27) which then initiates MAPK cascade. We calculate the Hill coefficients of Ste11pp, Ste7pp, and Fus3pp using  $G\beta\gamma$  as input, and find that each of them shows ultrasensitivity (Numerical results:  $n_{ste11pp} = 2.3$ ,  $n_{ste7pp} = 2.2$ ,  $n_{fus3pp} = 1.8$ ). Since  $G\beta\gamma$  experiences two simple reversible reactions to generate C, the relation between C and  $G\beta\gamma$  must be a hill function with n = 1. Therefore, the ultrasensitivity in this module must arise from C to Ste11pp, Ste7pp and Fus3pp. Let us first consider the phosphorylation and dephosphorylation cycle of Ste11. Ste11 can be dual phosphorylated on the scaffold in a processive mechanism and dual dephosphorylated in the cytosol in a distributive mechanism:

$$C + Ste11 \stackrel{on}{\rightleftharpoons} CSte11 \stackrel{p}{\Rightarrow} C + Ste11pp.$$
 
$$Ste11pp + MAPKKK - P \stackrel{a_2}{\rightleftharpoons} Ste11ppMAPKKK - P \stackrel{p_2}{\Rightarrow} Ste11p + MAPKKK - P$$
 
$$Ste11p + MAPKKK - P \stackrel{a_1}{\rightleftharpoons} Ste11pMAPKKK - P \stackrel{p_1}{\Rightarrow} Ste11 + MAPKKK - P$$

where we assume that Ste11pp leaves the scaffold so fast that CSte11pp can be neglected in the process. Introduce Michaelis constant  $K = \frac{off+p}{on}$ ,  $K_1 = \frac{d_1+p_1}{a_1}$ ,  $K_2 = \frac{d_2+p_2}{a_2}$ , and define  $\alpha_1 = \frac{pK_1}{p_1K}$ ,  $\alpha_2 = \frac{pK_2}{p_2K}$ ,  $p' = \frac{p_1p_2}{p_1+p_2}$ ,  $K' = \frac{p_1K_2+p_2K_1}{p_1+p_2}$ , we get:

$$[Ste11pp] = \frac{\alpha_2}{\alpha_1 + \alpha_2} G\left(pC_{total}, p'[MAPKKK - P]_0, \frac{K}{[Ste11]_0}, \frac{K'}{[Ste11]_0}\right) [Ste11]_0, \quad (4)$$

where

$$G(u, v, M, N) = \frac{2uN}{v - u + vM + uN + \sqrt{(v - u + vM + uN)^2 - 4(v - u)uN}},$$

$$u = p[Ste11]_0, \quad v = p'[MAPKKK - P]_0, \quad M = \frac{K}{[Ste11]_0}, \quad N = \frac{K'}{[Ste11]_0}.$$

The above G function can be fitted to a sigmoidal curve (Hill function)(61), with the Hill coefficient and the threshold value to be

$$n = \frac{1}{\log_{81} \frac{81(M+0.1)(N+0.1)}{(M+0.9)(N+0.9)}},\tag{5}$$

$$th = \frac{p'(1+2M)}{p(1+2N)}[MAPKKK - P]_0.$$
 (6)

According to the parameters in the Ste11 cycle,  $M = K/[Ste11]_0 \approx 0.125$  and  $N = K'/[Ste11]_0 = 0.255$ . So Eq. (7) approximates a Hill function with  $n \approx 2.6$  and  $th \approx 21nM$ . Therefore, it is the zeroth-order ultrasensitivity that leads to the ultrasensitive response of Ste11pp. Similarly, in the Ste7 cycle,  $M \approx 0.183$  and N = 0.324; in the Fus3 cycle,  $M \approx 0.135$  and N = 0.181. Therefore, Ste7 and Fus3 are also located in the zeroth-order region, making them ultrasensitive.

The analysis above together with Eq. (6) indicates that the relation between Ste11pp, Ste7pp, Fus3pp and  $\alpha$  should exhibit ultrasensitivity (numerical result:  $n_{ste11pp} = 2.1$ ,  $n_{ste7pp} = 1.8$ ,  $n_{fus3pp} = 1.9$ ). This appears to contradict the results in Table I where all the dose-response curves overlap with Hill coefficient  $n \approx 1$ . The reason for this contradiction is that in the analysis we cut off all the feedback in the original model. There are all together nine feedback in the whole model, six of which are negative feedback: the transcription of Sst2, MAPK-P, the degradation of activated Ste2, Ste11, Ste7pp, and the hyper-phosphorylation of Ste7pp. When  $[\alpha]$  is low, all these negative feedback are kept low, leaving the output nearly unaffected. When  $[\alpha]$  is high, the negative feedback will also be strong, significantly reducing the output. Therefore, negative feedback can make the dose-response curve less steep. It may be the counterbalance between the zeroth-order ultrasensitivity and the negative feedback that keeps the Hill coefficients of Ste11pp, Ste7pp, and Fus3pp remain 1. In order to test this view, we add each of the six negative feedback into the simplified model where all feedback are cut off (Table II. Numerical result shows that each feedback can reduce the Hill coefficients, and if all the negative feedback are added

together, the Hill coefficients can decrease to nearly 1, indicating that negative feedback indeed can reduce the Hill coefficients.

We further study how the Hill coefficients of Ste11pp, Ste7pp, Fus3pp depend on the scaffold, the substrate and the phosphatase concentrations (Table III. First, we increase Ste11, Ste7 or Fus3 concentration by 10 times, and find that the Hill coefficients only change a little. The probable reason is that the original kinase concentration is already much larger than the scaffold concentration, so when Ste11, Ste7 or Fus3 concentration increases, the added part could not get to the scaffold to be phosphorylated, and thus does not contribute to response. Then, we increase kinase concentration and scaffold concentration together, and find that Hill coefficients have a substantial increase. This is because when scaffold and kinase concentration increase together, effective substrate concentration increases. Therefore, the zeroth-order ultrasensitivity becomes more significant, while feedback, which is mainly dependent on the downstream regulation, does not increase as fast. Table III also shows that when each phosphatase concentration is decreased by 10 times, Hill coefficients arise as well. This is because low phosphatase concentration allow more substrate to be phosphorylated in the cytosol in a distributive mechanism, which is regarded as another mechanism to generate ultrasensitivity aside from the zeroth-order ultrasensitivity (17). Thus, the scaffold, the substrate and the phosphatase concentration play an important role in determining the Hill coefficients of the MAPK pathway.

**Analysis of parameters**: Due to the lack of experimental data to determine all the parameters, it is necessary to analyze the sensitivity of the system to changes of the parameters. To do this, we define a quantity:

$$D = \sqrt{(\frac{([Fus3pp]_{cal} - [Fus3pp]_{ori})^2}{[Fus3pp]_{ori}})^2}$$

where  $[Fus3pp]_{cal}$  denotes the calculated output when a parameter is changed and  $[Fus3pp]_{obs}$  denotes the original output, and the bar denotes the average of the relative variance of the output over an input ( $\alpha$ -factor) range of  $10^{-3}$  nM to  $10^{3}$  nM. We multiply and divide one parameter by 2 at each time, calculate D, and then take the average for the two situations of increasing and decreasing the parameter. The most sensitive parameters are listed in Table IV. Note that these parameters all have direct or indirect experimental support, which is reassuring. These parameters also give some clues about the mechanisms of the mating pathway. The parameters which influence the outcome of the pathway most are those involved in the receptor activation, which is consistent with our finding that the shape of dose-response curve is determined by the first step in the pathway—the receptor activation. The parameter in the production of receptor Ste2 is also essential to the outcome of the system. The other two influential parameters are the rates of the kinases to get off the scaffold, so the scaffold protein is also an important factor in this pathway.

# **DISCUSSION**

Our model describes the whole mating pathway comprising of the G protein cycle, the scaffold-dependent MAPK cascade and the downstream effects in the nucleus. We have in-

vestigated multiple features of the pathway, including its various characteristic times scales, desensitization, scaffold's effect, specificity, sensitivity to different levels of pheromone induction, the role of feedback, and sensitivity amplification. Although many the parameters in our model do not have solid experimental support and the detailed mechanisms of some steps are still not clear, the results given by our model are consistent with the current understanding of the pathway and with a wide range of experimental data.

The duration and sensitivity of the mating pathway have to be tightly regulated; an inappropriate activation of Fus3 will block the normal invasive growth. Our model shows that activation of the mating pathway attenuates with time even when the pathway is exposed to prolonged pheromone induction. This desensitization is attributed to several feedback such as the enhanced degradation of Ste7 (MAPKK). These feedback enable the cell to recover from mating and continue their vegetative growth upon prolonged pheromone induction.

Evidence shows that oligomerization of the scaffold protein is required for its activation (31,57,62,63). However, oligomerization is not included in our model for simplicity. Since we consider all possible complexes involving the scaffold protein, taking the oligomorization into account would make the model extremely complicated (with  $27 \times 27 = 729$  possible complexes involving the scaffold protein). Moreover, there are few experimental data concerning the interactions between the scaffold protein and the kinases. Therefore, we have to make some simplifications. A clue for the simplification is the experimental evidence that nuclear export and shmoo tip recruitment of Ste5 are coordinated with oligomarization (57,61). Thus we use nuclear export as a controlling step. In our model, the recruitment of the scaffold protein to the shmoo tip implies its activation, including the effect of nuclear transportation and oligomorization. However, oligomarization may have other effect in the pathway besides the activation of scaffold protein. Due to the lack of experimental data in this process, it is difficult to consider it in detail in our model. Experimental progress about this process is much needed to further improve the model.

The scaffold protein undergoes continuous shuttling and enhanced exportation upon pheromone induction. The obvious question is why yeast cells take so much trouble to shuttle a huge protein through the nucleus when it functions predominantly in cytoplasm. Our model indicates that nuclear shuttling might be a key step controlling the availability of the scaffold protein to the pathway. Since the activation of the MAPK cascade in the mating pathway is dependent on the scaffold protein, whether and how many scaffold proteins are available determine whether and how efficient the MAPK pathway is stimulated. However, the mechanism of scaffold shuttling are still not clear. The complex Msn5p/Ste21 is suggested to be responsible for the export of Ste5p. Further work is required to establish the accurate and detailed mechanism of this controlling step. In our model, we employ an active control mechanism. That is, the scaffold export rate is dependent on the concentration of the separated  $G\beta\gamma$ , which is released by the phemorone. The more the pheromone, the more the activated  $G\beta\gamma$ , and the higher the export rate. In other words, G protein cycle controls the shuttling of the scaffold and the concentration of scaffold at the shmoo tip.

Another function of the scaffold is its role in keeping the pathway's fidelity to the signal. The capability of the scaffold protein to prevent kinases from dephosphorylation assures the mating pathway's dependence on the scaffold protein, and the availability of scaffold in the shmoo tip is further controlled by the G protein cycle. Recently the specificity of different pathways in yeast is under intensive study (34). It is interesting to see that different cellular signals, which can be transmitted by the same components, result in distinct responses. Especially, the haploid invasive growth pathway employs the same MAPK components (Ste11 as MAPKKK, Ste7 as MAPKK, Fus3 and Kss1 as MAPK) as the mating pathway, except that Fus3 is more active during mating while Kss1 is preferentially activated during invasive growth. Then how are the different outputs controlled? Our model suggests that dephosphorylation and scaffolds work in coordination to prevent improper signal from leaking in, and thus contributing to the mating pathway's fidelity to pheromone induction. Further work is needed to include the parallel pathway of the invasive growth, through which a more comprehensive understanding of specificity might be obtained.

MAPK cascade which is conserved in all eukaryotic cells is composed of both phosphory-lation and dephosphorylation. Our model reveals that dephosphorylation has several roles in the mating pathway. It is obvious that it contributes to the desensitization of the pathway which enables the cell to reenter the cell cycle. Furthermore, it cooperates with the shuttling of the scaffold proteins, to realize other important features of the signaling pathway. First, it helps to preserve the consistence of the sensitivity from the G-protein cycle to the MAPK cascade. Second, the amount of MAPK-P, together with the scaffold protein, contributes to the pathway fidelity.

Notably, we suggest that negative feedback plays an important role in the experimentally observed preservation of sensitivity along the MAPK cascade. Whether or not the sensitivity is amplified as the MAPK cascade descends is also determined by the concentrations of the kinases and phosphatases involved. We call for new experiments to test this hypothesis.

### ACKNOWLEDGEMENT

We thank members of the Peking University Center for Theoretical Biology for their valuable suggestions. This research is supported by National Key Basic Research Project of China, Chinese Natural Science foundation, and the Chun-Tsung endowment at Peking University. C.T. acknowledges support from the Sandler Family Supporting Foundation.

### REFERENCES

- 1. Bardwell, L. 2004. A walk-through of the yeast mating pheromone response pathway. *Peptides.* 25:1465-1476.
- 2. Dohlman, H. G., and J. W. Thorner. 2001. Regulation of G protein-initiated signal transduction in yeast: paradigms and principles. *Annu. Rev. Biolchem.* 70:703-754.
- 3. Wang, Y., and H. G. Dohlman. 2004. Pheromone signaling mechanisms in yeast: a prototypical sex machine. *Science*. 306:1508-1509.

- 4. Dohlman, H. G. 2002. G protein and pheromone signaling. *Annu. Rev. Physiol.* 64:129-152.
- 5. Gustin, M. C., J. Albertyn, M. Alexander, and K. Davenport. 1998. MAP kinase pathways in the yeast *Saccharomyces cerevisiae*. *Microbiol*. *Mol. Biol. Rev.* 62:1264-1300.
- 6. Choi, K., B. Satterberg, D. M. Lyons, and E. A. Eloin. 1994. Ste5 tethers multiple protein kinases in the MAp kinase cascade required for matin in *S. cerevisiae*. *Cell*. 78:499-512.
- 7. Blackwell, E., I. M. Halatek, H. N. Kim, A. T. Ellicott, A. A. Obukhov, and D. E. Stone. 2003. Effect of the pheromone-responsive  $G\alpha$  and phosphatase proteins of saccharomyces cerevisiae on the subcellular localization of the Fus3 Mitogen-Activated Protein Kinase. Mol. Cell. Biol.23:1135-1150.
- 8. Hao, N., N. Yildirim, Y. Wang, T. C. Elston, and H. G. Dohlman. 2003. Regulators of G protein signaling and transient activation of signaling. *J. Biol. chem.* 278:46506-46515.
- 9. Kholodenko, B. N.. 2000. Negative feedback and ultrasensitivity can bring about oscillation in the mitogen-activated protein kinase cascades. *Eur.J. Biochem.* 267:1583-1588.
- 10. Kofahl, B., and E. Klipp. 2004. Modeling the dynamics of the yeast pheromone pathway. *Yeast.* 21:831-850.
- 11. Levchenko, A., J. Bruck, and P. W. Sternberg. 2000. Scaffold proteins may biphasically affect the levels of Mitogen-activates protein signaling and reduce its threshold properties. *PNAS*. 97:5818-5823.
- 12. Yi,T., H. Kitano, and M. Simon. 2003. A quantitative characterization of the yeast heterotrimeric G protein cycle. *PNAS*. 100:10764-10769.
- 13. Bhalla, U. S., P. T. Ram, and R. Lyengar. 2002. MAP kinase phosphatase as a locus of flexibility in a mitogen-activated protein kinase signaling network. *Science Reports*. 297:1018-1023.
- 14. Ferrell, J. E., and R. R. Bhatt. 1997. Mechanistic studies of the dual phosphorylation of mitogen-activated protein kinase. *J. Biol. Chem.* 272:19008-19016.
- 15. Ferrell, J. E., and E. M. Machleder. 1998. The biochemical basis of an all-or-none cell fate ewith in *Xenopus Oocytes*. *Science Reports*. 280:895-898.
- 16. Xiong, W., and E. Ferrell. 2003. A positive-feedback-based bistable 'memory module' that governs a cell fate decision. *Nature*. 426:460-465.
- 17. Huang, C. F., and J. E. Ferrell. 1996. Ultrasensitivity in the mitogen-activated protein kinase cascade. *Proc. Natl. Acad. Sci. USA*. 93:10078-10083.
- 18. Hicke, L., and H. Reizman. 1996. Ubiquitination of a yeast plasma membrane receptor signals its ligand-stimulated endocytosis. *Cell.* 84:277-288.
- 19. Conklin, B. R., and H. R. Bourne. 1993. Structural elements of  $G\alpha$  subunits that interact with  $G\beta\gamma$ , receptors and effectors. *Cell.* 73:631-641.
- 20. Cole, G. M., D. E. Stone, and S. I. Reed. 1990. Stoichiometry of G protein subunits affects the *Saccharomyces cerevisiae* mating pheromone signal transduction pathway. *Mol. Cell. Biol.* 10:510-517.

- 21. Dohlman, H. G., and J. Thorner. 1997. RGS proteins and signaling by heterotrimeric G protein. *J. Biol. Chem.* 272:3871-3874.
- 22. Dohlman, H. G., D. Apaniesk, Y. Chen, J. Song, D. Nusskern. 1995. Inhibition of G-protein signaling by dominant gain-of-function mutations in Sst2p, a pheromone desensitization factor in *Saccharomyces cerevisiae*. *Mol. Cell. Biol.* 15:3635-3643.
- 23. Bardwell, L., J. G, Cook, E. C. Chang, B. R. Cairns, and J.Thorner. 1996. Signaling in the yeast pheromone response pathway: specific and high-affinity interaction of the mitogen-activated protein (MAP) kinases Kss1 and Fus3 with the upstream MAP kinase kinase Ste7. *Mol. Cell. Biol.*16:3637-3650.
- 24. Leeuw, T., C. Wu, J.D. Schrag, M. Whiteway, D. Y. Thomas, and E. Leberer. 1998. Interaction of a G-protein  $\beta$ -subunit with a conserved sequence in Ste20/PAK family protein kinases. *Nature*. 391:191-195.
- 25. Leberer, D.. 1997. Functional characterization of the Cdc42p binding domain of yeast Ste20p protein kinase.  $EMBO\ J\ 16:83-97.$
- 26. Peter, M., A. M. Neiman, H. O. Park, M. Van Lohuizen, and I. Herskowitz. 1996. Functional analysis of the interaction between the small GTP binding Cdc42 and the Ste20 protein kinase in yeast. *EMBO J.* 15:7046-7059.
- 27. Lamson, R. E., M. J. Winters, and P. M. Pryciak. 2002. Cdc42 regulation of kinase activity and signaling by the yeast p21-activates kinase Ste20. *Mol. Cell. Biol.* 22:2939-2951.
- 28. Pryciak, P. M., and F. A. Huntress. 1998. Membrane recruitment of the kinase cascade scaffold protein Ste5 by the  $G\beta\gamma$  complex underlies activation of the yeast pheromone response pathway. *Genes Dev.* 12:2684-2697.
- 29. Dowell, S. J., A. L. Bishop, S. L. Dyos, A. J. Brown, and M. S. Whiteway. 1998. Mapping of a yeast G  $\beta\gamma$  signaling interaction. *Genetics*. 150:1407-1417.
- 30. Feng, Y., L. Y. Song, E. Kincaid, S. K. Mahanty, and E. A. Elion. 1998. Functional binding between  $G\beta$  and LIM domain of Ste5 is required to activate the MEKK Ste11. *Curr. biol.* 8:267-278.
- 31. Sette, C., C. J. Inoyue, S. L. Stroschein, P. J. Iaquinta, and J. Thorner. 2000. Mutational analysis suggests that activation of the yeast pheromone response mitogen-activated protein kinase pathway involves conformational changes in the Ste5 scaffold protein. *Mol. Biol. Cell.* 11:4033-4049.
- 32. Printen, J. A., and G. F. Sprague. 1994. Protein-protein interaction in the yeast pheromone response pathway: Ste5 interacts with all members of the MAP kinase cascade. *Genetics*. 138:609-619.
- 33. Park, S., A. Zarrinpar, and W. A. Lim. 2003. Rewiring MAP kinase pathways using alternative scaffold assembly mechanisms. *Science*. 299:1061-1064.
- 34. Schwartz, M.A., and H.D.Madhani. 2004. Principles of MAP kinase signaling specificity in *Saccharomyces Cerevisiae*. *Annu.Rev.Genet*. 38:725-48.
- 35. Van Drogen, F., S. M. O'Rourke, V. M. Stucke, M. Jaquenoud, A. M. Neiman, and M. Peter. 2000. Phosphorylation of the MEKK Ste11p by the PAK-like kinase Ste20p is

- required for MAP kinase signaling in vivo. Curr. Biol. 10:630-639.
- 36. Kwan, J. J., N. warner, T. Pawson, and L. W. Donaldson. 2004. The solution structure of the *S. cerevisiae* Ste11 MAPKKK SAM domain and its partnership with Ste50. *J. Mol. Biol.* 342:681-693.
- 37. Neiman, A. M., and I. Herskowitz. 1994. Reconstitution of a yeast protein kinase cascade in vitro: Activation of the yeast MEK homolog STE7 by STE11. *Proc. Natl. Acad. Sci. USA*. 91:3398-4402.
- 38. Errede, B., A. Gartner, Z. ZHou, K. Nasmyth, and G. Ammerer. 1993. MAP kinase-related Fus3 from *S. cerevisiae* is activated by Ste7 in vitro. *Nature*. 362:261-264.
- 39. Choi, K., J. E. Kranz, S. K. Mahanty, K. Park, and E. A. Elion. 1999.
- Characterization of Fus3 localization: active Fus3 localizes in complexes of varying size and specific activity. *Mol. Bio. Cell.* 10:1553-1568.
- 40. Esch, R. K., and B. Errede. 2002. Pheromone induction promotes Ste11 degradation through a MAPK feedback and ubiquitin-dependent mechanism. *PNAS*. 99:9160-9165.
- 41. Wang, Y., and H. G. Dohlman. 2002. Pheromone-dependent ubiquitination of the mitogen-activated protein kinase kinase Ste7. *J. Biol. Chem.* 277:15766-15772.
- 42. Maleri, S., Q. Ge, E. A. Hackett, Y. Wang, H. G. Dohlman, and B. Errede. 2004. Persistent activation by constitutive Ste7 promotes kss1-mediated invasive growth but fails to support Fus3-dependent mating in yeast. *Mol. Cell. Biol.* 24:9221-9238.
- 43. Doi, K., A. Gartner, G. Ammerer, B. Errede, H. Shinkawa, K. Sugimoto. 1994. MSG5, a novel protein phosphatase promotes adaption to pheromone response in *S. cerevisiae*. *EMBO J.* 13:61-70.
- 44. Zhan, X. L., R. J. Deschenes, and K. L. Guan. 1997. Differential regulation of FUS3 MAP kinase by tyrosine-specific phosphatases PTP2/PTP3 and dual-specificity phosphatase MSG5 in *Saccharomyces cerevisiae*. *Gene. Dev.* 11:1690-1702.
- 45. Andersson, J., D. M. Simpson, M. Qi, Y. Wang, and E. A. Elion. 2004. Differential input by Ste5 scaffold and Msg5 phosphatase route a MAPK cascade to multiple outcomes. *J. EMBO*. 23:2564-2576.
- 46. Flatauer, L. J., S. F. Zadeh, and L. Bardwell. 2005. Mitogen-Activated Protein Kinase with distinct requirements for Ste5 scaffolding influence signaling specificity in *Saccharomyces cerevisiae*. *Mol. Cell. Biol.* 25;1793-1803.
- 47. Van Drogen, F., and M. Peter. 2001. MAP kinase dynamics in yeast. *Biol. Cell.* 93:63-70.
- 48. Kusari, A. B., D. M. Molina, W. Sabbagh, C. S. Lau, and L. Bardwell. 2004. A conserved protein interaction network involving the yeast MAP kinases Fus3 and Kss1. *J. Cell. Biol.* 164:267-277.
- 49. Van Drogen, F., V. M. Stucke, G. Jorritsma and M. Peter. 2001. MAP kinase dynamics in response to pheromone in budding yeast. *Nature Cell Biol.* 3:1051-1059.
- 50. Tedford, K., S. Kim, D. Sa, K. Stevens, and M. Tyers. 1997. Regulation of the mating pheromone and invasive growth responses in yeast by two MAP kinase substrates. *Curr. Biol.* 7:228-238.

- 51. Cook, J. G., L. Bardwell, S. J. Kron, and J. Thorner. 1996. Two novel targets of the MAP kinase Kss1 are negative regulators of invasive growth in the yeast *Saccharomyces cerevisiae*. *Genes. Dev.* 10:2831-2848.
- 52. Gartner, A., A. Jovanovic, D. I. Jeoung, S. Bourlat, F. R. Cross, and G. Ammerer. 1998. Pheromone-dependent G1 cell cycle arrest requires Far1 phosphorylation, but may not involve inhibition of Cdc28-Cln2 kinase, in vivo. *Mol. Cell. Biol.* 18:3681-3691.
- 53. Ballensiefen, W., and H. D. Schmitt. 1997. Periplasmic *Bar1* protease of *Saccharomyces cerevisiae* is active before reaching its extracellular destination. *Eur. J. Biochem.* 247:142-147.
- 54. Chang, F., and I. Herskowitz. 1990. Identification of a gene necessary for cell cycle arrest by a negative growth factor of yeast: FAR1 is an inhibitor of a Gly cyclin, CLN2. *Cell.* 63:999-1011.
- 55. Bagnat, M., and K. Simon. 2002. Cell surface polarization during yeast mating. *Proc. Natl. Acad. Sci. USA*. 99:14183-14188.
- 56. Mahanty, S. K., Y. Wang, F. W. Farley, and E. A. Elion. 1999. Nuclear shuttling of yeast scaffold Ste5 is required for its recruitment to the plasma membrane and activation of the mating MAPK cascade. *Cell.* 98:501-512.
- 57. Elion, E. A.. 2001. The Ste5p scaffold. J. Cell. Sci. 114:3967-3978.
- 58. Elion, E. A.. 2000. Pheromone response, mating and cell biology. *Current opinion in cell. biol.* 3:573-581.
- 59. Wang, Y., Q. Ge, D. Houston, J. Thorner, B. Errede, and H. G. Dohlman. 2003. Regulation of Ste7 ubiquitination by Ste11 phosphorylation and the Skp1-Cullin-F-box complex. *J. Biol. Chem.* 278:22284-22289.
- 60. Garrison, T. R., Y. Zhang, M. Pausch, D. Apanovitch, R. Aebersold, and H. G. Dohlman. 1999. Feedback phosphorylation of an RGS protein by MAP kinase in yeast. *J. Biol. Chem.* 274:36387-36391.
- 61. Goldbeter, Albert and Danniel E. Koshland. 1981. An amplified sensitivity arising from covalent modification in biological systems. *Proc. Nati Acad. Sci. USA*. 78:6840-6844.
- 62. Wang, Y., and E. A. Elion. 2003. Nuclear export and plasma membrane recruitment of the Ste5 scaffold are coordinated with oligomerization and association with signal transduction components. *Mol. Biol. cell.* 14:2543-2558.
- 63. Yablonski, D., I. Marbach, and A. Levitzki. 1996. Dimerization of Ste5, a mitogen-activated protein kinase cascade scaffold protein, is required for signal transduction. *Proc. Natl. Acad. USA*. 93:13864-13869.

TABLE I: Coefficients in Hill function

Coefficient	Ste2a	G activation	Ste11pp	Ste7pp	Fus3pp
n	0.9	1.0	1.2	1.2	1.2
th (nM)	9.1	4.8	4.0	4.8	4.8

Note: simulation results with the whole model.

TABLE II: Hill coefficient n when negative feedback are added

added feedback	Ste11pp	Ste7pp	Fus3pp
none	2.1	1.8	1.9
transcription of Sst2	1.8	1.5	1.7
transcription of MAPK-P	2.0	1.7	1.8
degradation of $Ste2_{act}$	1.8	1.4	1.8
degradation of Ste11	1.3	1.2	1.8
degradation of Ste7pp	2.1	1.7	1.7
hyper-phosphorylation of Ste7pp	1.8	1.6	1.4
all the six feedback	1.2	1.2	1.2

TABLE III: Hill Coefficient **n** when concentration changes

concentration changes       Ste11pp       Ste7pp       Fus3pp $[Ste11] \times 10$ 1.2       1.2       1.3 $[Ste7] \times 10$ 1.3       1.6       1.2 $[Fus3] \times 10$ 1.2       1.2       1.2 $[Ste11] \times 10, [Ste5] \times 10$ 1.7       1.8       1.4 $[Ste7] \times 10, [Ste5] \times 10$ 1.8       2.5       2.8 $[Fus3] \times 10, [Ste5] \times 10$ 1.4       1.7       1.7
$[Ste7] \times 10$ 1.3       1.6       1.2 $[Fus3] \times 10$ 1.2       1.2       1.2 $[Ste11] \times 10, [Ste5] \times 10$ 1.7       1.8       1.4 $[Ste7] \times 10, [Ste5] \times 10$ 1.8       2.5       2.8
$[Fus3] \times 10$ 1.2 1.2 1.2 [Ste11] $\times$ 10, [Ste5] $\times$ 10 1.7 1.8 1.4 [Ste7] $\times$ 10, [Ste5] $\times$ 10 1.8 2.5 2.8
$[Ste11] \times 10, [Ste5] \times 10$ 1.7 1.8 1.4 $[Ste7] \times 10, [Ste5] \times 10$ 1.8 2.5 2.8
$[Ste7] \times 10, [Ste5] \times 10$ 1.8 2.5 2.8
$[Fus3] \times 10 \ [Ste5] \times 10$ 14 17 17
[1 460] × 10, [5 600] × 10
[MAPKKK - P]/10 1.7 1.4
[MAPKK - P]/10 1.2 1.7 1.2
[MAPK - P]/10 1.2 1.3 2.1

TABLE IV: Influential Parameters

Parameter	$D = \sqrt{\frac{\left(\frac{\triangle[Fus3pp]}{[Fus3pp]}\right)^2}{[Fus3pp]}}$	Related reaction
k1	0.519	$Ste2 + \alpha \stackrel{k1}{\underset{k2}{\rightleftharpoons}} Ste2_{act}$
k2	0.266	
k6	0.167	$\xrightarrow{k4,k5,ste12a} Ste2 \xrightarrow{k7}$
$**off_K$	0.156	Fus3pp gets off from the scaffold protein
$**off'_{KK}$	0.150	Ste7pp gets off from the scaffold protein

### FIGURE LEGENDS

- Figure 1. Spacial structure of the mating pathway.
- Figure 2. A. 27 solution-located scaffold complexes. B. 27 membrane-located scaffold complexes. The diamond on the upleft corner indicates the  $G\beta\gamma$ Ste20 complex
  - Figure 3. Scaffold-dependent reactions.
- Figure 4. Time course of the G protein cycle activation. A. G protein activation. The values are normalized to the maximum concentration around 30 s. Result from simulation is shown in solid line and experiment data (12) are plotted in circles with error-bars. B. G protein activation in  $Ste2^{300\triangle}$  cells, the wild types and cycloheximide treated cells (experiment data are from (12)). C. The time-course of Binding Ste20 to  $G\beta\gamma$ . The values are normalized to the maximum concentration (experiment data are from (24)). D. The recruitment of scaffold protein Ste5 (circles) which is the sum of scaffolds in the solution and scaffolds at the membrane, the activation of MAPK cascade components Ste11pp (MAPKKKK)(dotted line), Ste7pp (MAPKK) (dashed line) and Fus3pp (MAPK)(solid line). E. Downstream responses to  $\alpha factor$  induction: activated Ste12 (solid line),  $Far1pp \cdot Cdc28$  (dashed line), and  $Far1pp \cdot G\beta\gamma$ .
- Figure 5. A. Predicted dose-response curves of recruitment of scaffold protein Ste5 (dashed line) and activation of Fus3 (MAPK) (solid line). B. The dependence of Fus3pp on the concentration of scaffold proteins Ste5, with different concentration of  $\alpha$ -factor. C. Predicted dose-response curves for mutants in which MAPK-P is under-expressed. All the values are normalized. In all above simulation, Cells are treated with indicated with  $\alpha factor$  for 20 min.
- Figure 6. Dephosphorylation prevents improper signal to leak in. A. All Ste11 is dual phosphorylated at t=0, and MAPKKK-P is shut off. This simulates the condition in which signal in invasive growth pathway is on. B. All Ste11 is dual phosphorylated at t=0, and MAPKK-P is shut off. The activation of Fus3 is still repressed down except for the small pulse at the very beginning (Inset graph). Dashed line indicates the activation of Fus3 when MAPK-P is eliminated.
- Figure 7. Desensitization. A. Time-course of activation of Fus3 (MAPK) in a wild type cell (solid line) and in a mutant (dashed line). B. Effect of feedback hyper-phosphorylation of Ste7 in the scaffold. Predicted time-course activation of Fus3 for wild type (solid line) and for mutants (dashed line and dotted line).
- Figure 8. Dose-response curves for key components in mating pathway in TMY101 cells: activated Ste2 (dotted line for simulation, down-triangles for experiment data from (12));

G protein activation, i.e. the sum of  $G\alpha$ -GTP and  $G\alpha$ -GDP (dashed line for simulation, circles for experiment data from (12)); Fus3pp (MAPK) (solid line for simulation and crosses for experiment data from (45)).

Figure 9. Response curve of Ste12 to  $\alpha$ -factor for: A. different Sst2 expression level; B. different  $G\beta\gamma$  expression level. To simulate the  $2 \times G\beta\gamma$  cells, we separately add another 1000nM  $G\beta\gamma$  at t=0 min to double the total concentration of  $G\beta\gamma$ . Cells are treated with  $\alpha$ -factor for 60 min. Note that the experiment data are shifted left by more than 100-fold because Bar1 is deleted in TMY101 cells. Experiment data are from (8).




FIG. 1:




FIG. 2:




FIG. 3:




FIG. 4:





FIG. 5:






FIG. 6:




FIG. 7:





FIG. 8:






FIG. 9: