

Function Constrains Network Architecture and Dynamics: A Case Study on the Yeast Cell Cycle Network

Kai-yeung Lau,¹ Surya Ganguli,² and Chao Tang^{1,3},

¹Departments of Biopharmaceutical Sciences and Biochemistry and Biophysics, University of California San Francisco, 1700 4th Street, San Francisco, CA 94143-2540, USA

²Sloan-Swartz Center for Theoretical Neurobiology, University of California, 513 Parnassus Avenue, San Francisco, CA 94143-0444, USA

³Center for Theoretical Biology, Peking University, Beijing 100871, China
(Dated: March 16, 2019)

We develop a general method to explore how the function performed by a biological network can constrain both its structural and dynamical network properties. This approach is orthogonal to prior studies which examine the functional consequences of a given structural feature, for example a scale free architecture. A key step is to construct an algorithm that allows us to efficiently sample from a maximum entropy distribution on the space of boolean dynamical networks constrained to perform a specific function, or cascade of gene expression. Such a distribution can act as a "functional null model" to test the significance of any given network feature, and can aid in revealing underlying evolutionary selection pressures on various network properties. Although our methods are general, we illustrate them in an analysis of the yeast cell cycle cascade. This analysis uncovers strong constraints on the architecture of the cell cycle regulatory network as well as significant selection pressures on this network to maintain ordered and convergent dynamics, possibly at the expense of sacrificing robustness to structural perturbations.

PACS numbers: 87.10.+e, 87.17.Aa

I. INTRODUCTION

A central problem in biology involves understanding the relationship between structure and function. This problem becomes especially intricate in the era of systems biology in which the objects of study are biological networks composed of large numbers of interacting molecules. To what extent does the structure of a biological network constrain the range of functions, or types of dynamical behaviors, that the network is capable of producing? Conversely, to what extent does the requirement of carrying out a specific function constrain the structural and more general dynamical properties of a network?

There already exists a large body of theoretical work addressing the former question. For example Kauzmann [1] and others performed an extensive study of ensembles of simplified boolean networks with fixed structural properties, such as the number of nodes and the mean degree of connectivity. A principal finding was a phase transition in the resulting dynamical behavior, from ordered to chaotic, as the connectivity increased [2]. More recently, the observation that many biological networks are scale free spurred a flurry of research into the dynamical consequences of the scale free structural feature [3-6]. A principal finding was that the scale-free architecture is more robust to random failures and dynamic fluctuations.

Alternatively, the latter question involving the struc-

tural and dynamical consequences of performing a specific function remains relatively unexplored [7, 8]. It is an important question because many biological functions are performed by relatively small network modules for which gross statistical properties such as mean connectivity, or any kind of degree distribution, scale free or not, do not have a clear significance. A key example of such a module is the yeast cell cycle control network, whose essential function was reduced to the boolean dynamics of a set of 11 nodes by Li, et. al. [9]. Despite the small network size, a dynamical analysis of this simple model demonstrated a great deal of robustness of the cell cycle trajectory to both fluctuations in protein states and perturbations of network structure. Is this robustness carefully selected for through evolution and encoded somehow in the topological structure of the cell cycle network, or does it arise for free, simply as a consequence of the functional constraint of having to produce the long cascade of gene expression that controls the cell cycle?

In this paper we develop techniques to address this question, and more generally to address the consequences of specific functional, rather than structural constraints. A key step in the above work exploring the dynamical consequences of fixed structural constraints was the ability to efficiently sample from the maximum entropy distribution on the space of networks constrained to have a fixed structural feature, such as a given degree distribution. We call such an ensemble a structural ensemble. Similarly, in order to address the structural consequences of fixed functional constraints, we develop an efficient algorithm to sample from a maximal entropy distribution on the space of biological networks constrained to perform a specific function. We call the resulting ensemble

To whom correspondence should be addressed. E-mail: Chao.Tang@ucsf.edu.

of networks a functional ensemble.

Since we use a maximum entropy distribution, which introduces no further assumptions about the network other than the fact that it performs a given function, we can use such functional ensembles to test whether any dynamical or structural property of a given biological network is simply a consequence of the function it performs, or rather a consequence of further selection. Under this functional null model, statistically significant properties of a real biological network can give us insight into additional selective pressures that have operated on that network above and beyond the baseline functional requirements. As an illustration of this general method, we focus on the specific case of the cell cycle network of [9], uncovering deeper insight into the evolutionary pressures on its structural and dynamical properties.

II. METHODS

A. The Dynamical Model

We consider a simplified, boolean model of biological network dynamics, in which each degree of freedom, or node s_i , $i = 1 :: N$ takes one of two values at any given time: either 0 (inactive) or 1 (active). For example, the two values could signify whether a protein is expressed or not, or whether a kinase is activated or not. Thus the full state at time t is captured by a column vector

$$S(t) = (s_1(t); s_2(t); \dots; s_N(t))^T \quad (1)$$

that can take one of 2^N values. Time progresses in discrete steps, and the nodes can either activate or inhibit each other at the next step. These interactions are captured by the network connectivity matrix C with elements c_{ij} representing an interaction arrow from node j to node i . The allowed values of c_{ij} are given by

$$c_{ij} \in \begin{cases} \{1; 0; -1\} & \text{if } i \neq j; \\ \{1; 0\} & \text{if } i = j; \end{cases} \quad (2)$$

For two different nodes, if c_{ij} is nonzero, it can be either activating (+1) or inhibiting (-1). If a diagonal element $c_{ii} = -1$ we say node i is self-degrading. This terminology is justified by the dynamical rule

$$s_i(t+1) = f_i(C; S(t)); \quad (3)$$

where

$$f_i(C; S(t)) = \begin{cases} 1; & \text{if } \sum_{j \neq i} c_{ij} s_j(t) > 0 \\ 0; & \text{if } \sum_{j \neq i} c_{ij} s_j(t) < 0 \\ s_i(t); & \text{if } \sum_{j \neq i} c_{ij} s_j(t) = 0; c_{ii} = 0 \\ 0; & \text{if } \sum_{j \neq i} c_{ij} s_j(t) = 0; c_{ii} = -1; \end{cases} \quad (4)$$

Essentially, if the total input to a node is positive (negative), it will be on (off) at the next time step. In the case of zero input, the node either turns off, or maintains its state, depending on whether or not it is self-degrading.

B. Generating Functional Ensembles

A biological network achieves its function by successfully taking the values of its nodes through a sequence of states. Thus we will equate the notion of function with a specified state trajectory

$$S(0) \rightarrow S(1) \rightarrow \dots \rightarrow S(T); \quad (5)$$

In our example of the cell cycle network, the state sequence is simply the natural cell cycle trajectory. We wish to either enumerate or uniformly sample from the space of networks, or equivalently connectivity matrices C , that can successfully perform the above sequence of T state transitions. We can think of each of the T transitions as providing one constraint on the connectivity matrix C via the dynamical rule given by equations (3) and (4). Assuming the nodes are distinguishable, the number of networks, given by the number of allowed connectivity matrices is $M = (2 - 3^{N-1})^N$. Even for small, mesoscopic scale networks such as the cell cycle with 11 nodes, $M \approx 6.23 \times 10^{55}$ and so it is computationally infeasible to iterate through all of these networks and find those for which the T constraints corresponding to the T transitions in (5) are satisfied. Even if one were to sample from these M networks, one would rarely find a network that could perform the function in (5).

However, it is important to note that the constraint on the network connectivity C imposed by a given transition actually decouples across the rows of the connectivity matrix. That is for each node i , the dynamical rule in Eq. (4) depends only on the i 'th row $c_{ij}; j = 1 :: N$ of C , or equivalently on the N incoming interaction arrows whose target is node i . Thus we can check that the T constraints induced by the target sequence (5) are satisfied for each row, independently of the other rows. For any given i , the number of possible rows is $Z = 2 - 3^{N-1}$, which is 118,098 for the yeast cell cycle network. Thus it becomes computationally feasible to exhaustively iterate through all possible row values, or incoming arrow combinations, for each node, and check that the T constraints are satisfied for each such combination.

After following this procedure, let $\mathcal{I}_i \subset Z$ index the set of allowed incoming arrow combinations to node i that satisfy all T constraints. If for each node i , we uniformly choose a particular i_i , and assemble the corresponding N allowed rows into a matrix C , we will have constructed a network that can successfully carry out the state trajectory in (5). This is essentially our sampling procedure. It produces a functional ensemble; a maximum entropy distribution on the space of networks constrained to produce the function represented by (5).

C. Combined structural and functional ensembles.

In order to perform a more fine scale study of the properties of the yeast cell cycle network, we wish to constrain more than just a predefined function. We would

also like to understand how various properties depend on the number of nonzero interaction arrows in the network. Thus we need to develop a method to uniformly sample from the space of networks that both perform a fixed function and have a fixed number of arrows. We do so as follows.

In the course of sampling from the functional ensemble above, in each step where we select a particular combination of incoming connections for a node, we compute the probability of choosing such a combination from the corresponding set by calculating the convolution of (i) the probability for the node to have a certain number of inward arrows and (ii) the probability of choosing a combination with that number of arrows. Mathematically, let H be the number of arrows in the network. It can be distributed among the nodes in different ways. Let W_{hi} be the number of times h inward arrows being assigned to node i among all possible ways of dividing the H arrows. If there are Q_{hi} inward connection combinations selected for node i that have h inward arrows, then the probability for assigning h inward arrows to node i is

$$p = \frac{W_{hi} Q_{hi}}{\sum_{l=1}^N W_{li} Q_{li}} : \quad (6)$$

Once the number of arrows is assigned, an inward connection combination can then be selected randomly from the Q_{hi} combinations. This process is repeated for each node, each time using the remaining number of arrows (i.e. $H - h$) to calculate W_{hi} and Q_{hi} . For each network, we randomize the order of nodes for which we select the connection combination.

We used this algorithm to generate the ensemble of "cell cycle" networks described below. In order to compare this ensemble to a set of more random networks that serve no particular function, we generated this "random network" ensemble by randomly rewiring the connections in each "cell cycle" network under the constraint that all nodes must be connected to the same network, i.e. no isolated nodes or sub-networks.

III. THE YEAST CELL-CYCLE NETWORK

The simplified yeast cell-cycle Boolean network [see Fig. 1] given in Li et al. [9] contains 11 proteins, or nodes, and 1 checkpoint. There are 34 arrows connecting the nodes: 15 activating, 14 deactivating and 5 "self-degrading." Using the above dynamical model, this network can produce the cell-cycle process, as shown in Table I. Starting from the "excited" G_1 state, the process goes through the S phase, the G_2 phase, the M phase, and finally returns to the biological G_1 stationary state. The network also has 7 x-points, with the G_1 stationary state being the biggest, having a basin size of 1764 (86% of all protein states).




FIG. 1: Simplified yeast cell-cycle network from Li et al. [9]. Solid, dashed, and dotted arrows are activating, deactivating, and "self-degrading" interactions, respectively. The black solid and black dashed arrows are absolutely required for a network to produce the cell-cycle process [see Table I].

IV. RESULTS

We used the same 11 nodes and the types of connections in the simplified yeast cell-cycle Boolean network to construct our ensembles of networks. Using our technique to generate purely functional ensembles, we were able to select 11 sets of inward connection combinations (one for each node) that produce the cell-cycle process [see Table I]. Figure 2 shows the compositions of different types of connections in the sets. The number of selected connection combinations for each node (shown in parenthesis in Fig. 2) varies for two orders of magnitude. The number of networks that can realize the cell-cycle process is $6.55 \cdot 10^{32}$ and the distribution against the number of arrows is shown in Fig. 3.

A. Constraints on Structure from Function.

From Fig. 2 we can deduce that there are 10 core connections (shown as black solid and black dashed arrows in Fig. 1) that are absolutely required in order to produce the cell-cycle process. These required connections become obvious once we look closer into the cell-cycle process. For example, comparing the G_1 stationary state and the "excited" G_1 state, Cln3 is the only node that is turned on; this implies that MBF and SBF, which are turned on in the next time step, can only be activated by Cln3. The remaining required connections can all be deduced using the same logic. The compositions of different

TABLE I: The cell-cycle process generated by the simplified yeast cell-cycle network in Fig. 1.

Time	Cln3	MBF	SBF	Cln1,2	Clb5,6	Clb1,2	Mcm1	Cdc20	Swi5	Sic1	Cdh1	Phase
1	1	0	0	0	0	0	0	0	0	1	1	\Excited" G ₁
2	0	1	1	0	0	0	0	0	0	1	1	G ₁
3	0	1	1	1	0	0	0	0	0	1	1	G ₁
4	0	1	1	1	0	0	0	0	0	0	0	G ₁
5	0	1	1	1	1	0	0	0	0	0	0	S
6	0	1	1	1	1	1	1	0	0	0	0	G ₂
7	0	0	0	1	1	1	1	1	0	0	0	M
8	0	0	0	0	0	1	1	1	1	0	0	M
9	0	0	0	0	0	1	1	1	1	1	0	M
10	0	0	0	0	0	0	1	1	1	1	0	M
11	0	0	0	0	0	0	0	1	1	1	1	M
12	0	0	0	0	0	0	0	0	1	1	1	G ₁
13	0	0	0	0	0	0	0	0	0	1	1	Stationary G ₁




FIG. 2: Fractions of different types of inward connections to each node from all other nodes (including itself) that produce the cell-cycle process [see Table I]. Numbers in parenthesis are the number of connection combinations selected by our algorithm [see Methods].

ent connection types follow a common trend where there is a higher chance for a node to be positively regulated by nodes that are active earlier in the cell-cycle process (positive feed-forward) and negatively regulated by nodes that are active later in the process (negative feedback). This trend seems to be general for networks that produce cascades of activation. To check this, we looked at the connection compositions for two simple activation cascades, where 11 nodes are activated in turn for 4 or 5 time steps, and they indeed show the same trend [See Table II and Fig. 4 for 5 time steps activation cascade].




FIG. 3: Distribution of number of networks that can realize the cell-cycle process over the number of arrows in the networks.

Next, we generated two network ensembles for each number of arrows allowed in the space of cell cycle networks. This number of arrows varied from 24 to 117 as seen in Fig. 3. The first ensemble is a combined structural/functional ensemble [see Methods] consisting of 1,000 networks that both realize the cell cycle and have a fixed number of arrows. These networks will be referred to as the "cell-cycle networks" (CN). The second ensemble was generated by randomly reconnecting the arrows for each network in the first ensemble [see Methods]. This ensemble will be referred to as the "random networks" (RN).

TABLE II: A simple activation cascade with 11 nodes activated in turn for 5 time steps

Time	A	B	C	E	F	G	H	I	J	K	L
1	1	0	0	0	0	0	0	0	0	0	0
2	1	1	0	0	0	0	0	0	0	0	0
3	1	1	1	0	0	0	0	0	0	0	0
4	1	1	1	1	0	0	0	0	0	0	0
5	1	1	1	1	1	0	0	0	0	0	0
6	0	1	1	1	1	1	0	0	0	0	0
7	0	0	1	1	1	1	1	0	0	0	0
8	0	0	0	1	1	1	1	1	0	0	0
9	0	0	0	0	1	1	1	1	1	0	0
10	0	0	0	0	0	1	1	1	1	1	0
11	0	0	0	0	0	0	1	1	1	1	1
12	0	0	0	0	0	0	0	1	1	1	1
13	0	0	0	0	0	0	0	0	1	1	1
14	0	0	0	0	0	0	0	0	0	1	1
15	0	0	0	0	0	0	0	0	0	0	1
16	0	0	0	0	0	0	0	0	0	0	0




FIG. 4: Fractions of different types of inward connections to each node from all other nodes (including itself) that produce the activation cascade in Table II.

B. Analysis of Attractors: Large Basins for Free

We studied the time evolution of protein states of the two ensembles by using the dynamical model described in the Methods and initiating the networks from each of the $2^{11} = 2;048$ states. We found that the CN networks have fewer attractors and larger attractor basin sizes compared to the RN networks [see Fig. 5]. The number of attractors decreases and the size of attractor basins increase as the number of arrows increases [see




FIG. 5: Number of attractors and basin sizes of attractors for the two ensembles of networks. (a) Distribution of number of attractors. (b) Number of attractors averaged over networks with the same number of arrows. (c) Distribution of size of basins of attractions. (d) Basin size averaged over networks with the same number of arrows.




FIG. 6: (a) Distribution of convergence value w_n for network states in each ensemble. (b) Overall convergence averaged over networks with the same number of arrows. Error bars represent one standard error.

Fig. 5 (b), (d)]. In the CN ensemble, the probabilities for a network with 34 arrows to have 7 attractors and to have the biggest attractor basin size 1764 (as in the case of yeast cell-cycle network) are 7.0% and 14.4% respectively. In the RN ensemble the corresponding percentages are only 2.7% and 1.6% respectively. Thus we see that the constraint of having to perform the yeast cell cycle cascade alone can, to a certain extent, help explain the origins of these two measures of dynamical robustness; a large basin essentially arises for free as a consequence of the cell cycle function. Indeed, the average basin size of the biggest attractors for the CN ensemble is 1705.53 compared to 1452.21 for the RN ensemble. In addition, 97.5% of the networks in the CN ensemble have the G_1 stationary state as the biggest attractor and the average basin size of these attractors is 1674.07.




FIG. 7: Distribution of transient times.

C. Convergence of Trajectories.

Following [9], we define a measure w_n that quantifies the "degree of convergence" of the dynamical network trajectories onto each network state n where $n = 1; \dots; 2048$. Let $T_{j;k}$ denote the number of trajectories starting from all 2048 initial network states that travel from state j to state k in one time step. Let L_n denote the number of steps it takes to get from state n to its attractor, so that we can index the states along the outward trajectory by $k = 1; \dots; L_n$. Then $w_n = \frac{1}{L_n} \sum_{k=1}^{L_n} T_{k-1;k} = L_n$. The overall convergence, or overlap W of trajectories is given by the average of w_n over all states n .

The distribution of the w_n values is shown in Fig. 6 (a). The result shows that there are more states in the CN ensemble having larger w_n values indicating a higher degree of convergence in the network dynamics. The local maxima at $w_n = 620$ for this ensemble should be a result of the requirement that networks in this ensemble must produce the cell-cycle process. The overall overlap W [see Fig. 6 (b)], which is the average of w_n over all network states, for the yeast cell-cycle network is 743 and the probability for a network with 34 arrows in the CN ensemble to have $W \geq 743$ is 4.4%. Such a result is highly unlikely in the RN ensemble. In fact no networks in the RN ensemble with 34 arrows had an overlap $W \geq 743$. Thus a higher degree of convergence is a dynamical consequence of performing the cell cycle function, but nevertheless, the actual cell cycle network in Fig. 1 still displays a relatively high degree of convergence even within the CN ensemble.

D. Dynamical Order: Transients and Sensitivity.

To compare the degree of order between the networks in the two ensembles, we calculated the transient time for all network states for all networks [see Fig. 7]. The transient time is defined as the amount of time for a net-




FIG. 8: Network sensitivity averaged over networks with the same number of arrows. A asterisk (*) shows the network sensitivity of the yeast cell-cycle network. Error bar represent two times standard error.

work state to evolve to its attractor, which is equal to the length of its trajectory [10]. The result shows that CN networks have longer transient times and thus are more chaotic than RN networks (unless the RN networks have long attracting limit cycles). The average transient time for the yeast cell-cycle network is 7.47, and the probability for a cell-cycle network with 34 arrows to have ≥ 7.47 average transient time is 7.2%.

We then calculated the network sensitivity s [11] for all networks [see Fig. 8]. Network sensitivity is the average expected number of node state changes in the output given a one node state change in the input. In other words, s calculates the average hamming distance of the output states of the network for all hamming neighbor input states (i.e. hamming distance = 1). If $s < 1$, the network is ordered, where fluctuations in node states die out quickly and remain localized. If $s > 1$, the network is chaotic, where fluctuations spread across the entire network. When $s = 1$, the network is critical. The result indicates that networks in both ensembles were chaotic on average with any number of arrows within the range we studied. s increases monotonically with the number of arrows in a network. The values of s for RN ensemble remain smaller than those for the CN ensemble demonstrating that CN networks are indeed more chaotic than RN networks. The yeast cell-cycle network has a network sensitivity of 1.27, which is more ordered than the average, and the probability for a network with 34 arrows in the CN ensemble to have $s \geq 1.27$ is only 3.0%. Thus the actual cell cycle is remarkably ordered despite the fact that the functional constraint of performing the cell cycle drives networks to be more chaotic.




FIG. 9: Response to perturbations to network structure. (a) Fractions of perturbed networks, for each number of arrows, that retain the original biggest attractor. A sterisk () shows the fraction for the perturbed yeast cell-cycle network. Error bar represent one standard error. (b) Distribution of relative changes of the size of the basin of attraction for the biggest attractor.

E. Dynamical response to structural perturbations.

To check and compare how the networks respond to structural perturbations, we performed the same kinds of alterations described in Li et al. [9] on all networks in the two ensembles. The alterations included deleting arrows from, adding arrows to and switching the signs of arrows in the networks. We assessed the response by calculating the percentage of perturbed networks that retain their original biggest attractor [see Fig. 9 (a)] and also the relative change in the size B of the basin of attraction for the original biggest attractors [see Fig. 9 (b)]. The percentages of perturbed networks in the two ensembles that retain their original biggest attractor both increase initially when the number of arrows in the network is small. The percentages are similar when there are about 34 to 54 arrows in the networks but as the number of arrows exceeds 60, the percentage for the RN ensemble drops quickly and the percentage for the CN ensemble remain high. The percentage for yeast cell-cycle network (68%) is smaller than the average for CN networks with the same number of arrows. The probability to obtain an equal or higher percentage is 82.3%, indicating that the yeast cell-cycle network has a worse than average robustness with respect to such structural perturbations.

We noticed from the distributions of $\Delta B/B$ that there is a higher chance for perturbations to have a deleterious effect to networks in the CN ensemble where the change in the sizes of basins of attraction is usually negative. However, there is a much higher chance for networks in the RN ensemble to completely lose the original biggest attractor ($\Delta B/B = 1$), which is even more unfavorable. The above effects should be attributed to the smaller basins of attraction for networks in the RN ensemble. The average $\Delta B/B$ for yeast cell-cycle network is -0.326 and the probability for a CN network with 34 arrows




FIG. 10: Fractions of trajectories of the perturbed networks starting from the "excited" G_1 state that still evolve to the biological G_1 stationary state and fraction of cell-cycle processes of the perturbed networks remain unchanged. A sterisk () and cross (x) show the fraction of trajectories reaching G_1 stationary state and fraction of cell-cycle processes unchanged, respectively, for perturbed yeast cell-cycle networks. Error bar represent one standard error.

to have average $\Delta B/B$ value -0.326 is 90.9%. This again signifies a worse than average robustness for the yeast cell-cycle network.

F. Stability of the cell cycle process.

Finally, we checked how many perturbed networks from the CN ensemble could still maintain the cell-cycle process [see Fig. 10]. Starting from the "excited" G_1

state, the fraction of trajectories reaching the G_1 stationary state and the fraction of cell-cycle processes unchanged increase as the number of arrows in the network increases. The fraction of trajectories reaching the G_1 stationary state and the fraction of cell-cycle processes unchanged for the yeast cell-cycle network are 0.54 and 0.23 respectively. The probability for a CN network with 34 arrows to maintain 54% of trajectories reaching the G_1 stationary state and 23% of cell-cycle processes unchanged are 80.6% and 62.1% respectively.

V. DISCUSSION

We presented a maximum entropy analysis method that can reveal the underlying structural constraints, as well as the statistical significance of various dynamical properties, of networks that perform a certain function. We applied this method to the yeast cell-cycle network and the accompanying cell-cycle process [9].

We demonstrated that requiring a network to produce an activation cascade, e.g. the cell-cycle process, requires the network to have positive feed-forward and negative feedback interactions between their nodes. It is not just the case that this is a good design principle to realize a long transient cascade; it is essentially the only way to achieve it generically, and yields an example of how network function constrains network structure.

We also showed that certain dynamical features arise purely as a consequence of performing the cell-cycle function. Compared to the random (RN) ensemble, networks in the cell-cycle (CN) ensemble had much larger basins of attraction, a higher degree of convergence of trajectories, longer transient times, and more chaotic behavior, as measured by the network sensitivity s . These properties may be essential for networks to produce a long sequence of state transitions. The long trajectory of the cell-cycle process provides many possible merge points for other trajectories, which certainly contribute to the high degree of convergence in the network dynamics and the large basin of attraction for G_1 stationary state. Thus the existence of this globally attracting trajectory of the dynamics alone can explain to a certain extent the observed robustness against dynamical perturbations.

On the other hand, with respect to structural perturbations, the actual yeast cell cycle is relatively less robust compared to other networks in the CN ensemble. This is in stark contrast to the high degree of dynamical order displayed by the cell cycle network, which suggests that there may be a trade off between ordered dynamics

and structural robustness. The network sensitivity [11], which measures the degree of order, calculates how sensitive a network is to fluctuations in the states of the nodes, which is a major source of variation in a cell population [12][15]. Evolution may have favored a design for the yeast cell-cycle network that is ordered and less sensitive to fluctuations in the states of the nodes (e.g. it has been reported that there is on average only 1 copy of SWI5 mRNA per cell in yeast [16]), by sacrificing robustness against perturbations to the network structure. However, we expect that the complete yeast cell-cycle network is more robust against such perturbations since it has "redundant" components and connections that perform similar jobs.

In any case, the observation that only 3% of randomly chosen cell cycle networks with 34 arrows are more ordered (as measured by the sensitivity s) than the real cell cycle network reveals an unsuspected but significant selection pressure for ordered dynamics that cannot be explained by the functional requirement of maintaining the cell cycle process; indeed the functional requirement of maintaining the cell cycle process forces the dynamics in the opposite direction, i.e. to be more chaotic. Similarly, we have seen that simply requiring a long cell cycle to occur automatically raises the average degree of convergence of trajectories. However, even after accounting for this increase within the functional ensemble, only 4.4% of all cell cycle networks with 34 arrows have a greater degree of convergence, reflecting an evolutionary pressure for convergent dynamics above and beyond that necessitated by the requirements of the cell cycle alone.

Although we have focused on a single biological example, the cell cycle, our analysis method is quite general. It would be interesting to perform it on other mesoscopic scale networks that have a comparable number of components to uncover their structural and dynamical constraints. More generally, we believe these techniques of simultaneously analyzing both structural and functional ensembles of networks will prove useful in the larger quest to deduce general principles governing relations between structure, dynamics, function, robustness and evolution.

VI. ACKNOWLEDGEMENTS

We would like to thank Morten K lobster for suggesting the sampling method. C.T. acknowledges support from the Sandler Family Supporting Foundation and from the National Key Basic Research Project of China (2003CB715900).

[1] S. A. Kauffman, *The Origins of Order.*, Oxford Univ. Press, Oxford.

[2] M. Adakana, S.N. Coppersmith, L. Kadano, *Boolean Dynamics with Random Coupling*, Perspectives and Problems in Nonlinear Science. Springer Applied Mathematical

cal Sciences Series. E. Kapan, J.E. Marsden, and R.R. Screenivasan, Eds. (Springer Verlag, 2003).

[3] H. Jeong, B. Tombor, R. Albert, Z.N. Oltvai, and A.L. Barabasi, *Nature* 407, 651 (2000).

[4] M. Adakana and P. Cizel, *Proc. Natl. Acad. Sci. U.S.A.*

- 100, 8710 (2003).
- [5] H. Jeong, S.P. Mason, A.L. Barabasi, and Z.N. Oltvai, *Nature* 411, 41 (2001).
- [6] S.M. Aslov and K. Sneppen, *Science* 296, 910 (2002).
- [7] A.W. Agner, *Proc. Natl. Acad. Sci. U.S.A.* 102, 11775-11780 (2005).
- [8] W. Ma, L. Lai, Q. Ouyang, and C. Tang, to be published.
- [9] F. Li, T. Long, Y. Lu, Q. Ouyang, and C. Tang, *Proc. Natl. Acad. Sci. U.S.A.* 101, 4781 (2004).
- [10] A. Wuensche, *Pac. Sym. Biocomput.*, 89 (1998).
- [11] I. Shmulevich and S.A. Kauffman, *Phys. Rev. Lett.* 93, 048701 (2004).
- [12] M.B. Elowitz, A.J. Levine, E.D. Siggia, and P.S. Swain, *Science* 297, 1183 (2002).
- [13] J.M. Raser and E.K. O'Shea, *Science* 304, 1811 (2004).
- [14] J.M. Raser and E.K. O'Shea, *Science* 309, 2010 (2005).
- [15] P.S. Swain, M.B. Elowitz, and E.D. Siggia, *Proc. Natl. Acad. Sci. U.S.A.* 99, 12795 (2002).
- [16] M.J. Holland, *J. Biol. Chem.* 277, 14363 (2002).