

Two exactly-solvable problems in one-dimensional quantum mechanics on circle

L.G.Mardoyan¹, G.S.Pogosyan^{1,2,3} and A.N.Sissakian²

¹*International Center for Advanced Studies, Yerevan State University, Yerevan, Armenia*

²*Laboratory of Theoretical Physics, Joint Institute for Nuclear Research, Dubna, 141980, Moscow Region, Russia*

³*Centro de Ciencias Físicas, Universidad Nacional Autónoma de México Apartado Postal 48-3, Cuernavaca, Morelos 62251, México*

Abstract

In this note we establish a relation between two exactly-solvable problems on circle, namely singular Coulomb and singular oscillator systems.

1. In a recent paper [1] have constructed a series of complex mappings $S_{2C} \rightarrow S_2$, $S_{4C} \rightarrow S_3$ and $S_{8C} \rightarrow S_5$, which extend to spherical geometry the Levi-Civita, Kustaanheimo-Stiefel and Hurwitz transformations, well known for Euclidean space. We have shown that these transformations establish a correspondence between Coulomb and oscillator problems in classical and quantum mechanics for dimensions (2,2), (3,4) and (5,8) on the spheres. A detailed analysis of the real mapping on the curved space has been done in [2]. It was remarked that in the stereographic projection the relation between Coulomb and oscillator problems functionally coincide with the flat space Levi-Civita and Kustaanheimo-Stiefel transformations. The relation between the quasiradial Schrödinger equations for Coulomb and oscillator problems on the n -dimensional spheres and one- and two-sheeted hyperboloids for $n \geq 2$ was found in the article [3].

The present note are devoted to two one-dimensional exactly-solvable potentials on circle S_1 : $s_0^2 + s_1^2 = R^2$

$$V^{so}(\vec{s}) = \frac{\omega^2 R^2 s_1^2}{2 s_0^2} + \frac{1}{2} \frac{k_1^2 - \frac{1}{4}}{s_1^2}, \quad V^c(\vec{s}) = -\frac{\mu}{R} \frac{s_0}{|s_1|} + \frac{1}{2} \frac{p^2 - \frac{1}{4}}{s_1^2}, \quad (1)$$

where s_0, s_1 are Cartesian coordinates in ambient Euclidean space E_2 .

2. The Schrödinger equation describing the nonrelativistic quantum motion on the circle S_1 in polar coordinate $\varphi \in [-\pi, \pi]$

$$s_0 = R \cos \varphi, \quad s_1 = R \sin \varphi$$

has the following form ($\hbar = m = 1$)

$$\frac{d^2 \Psi}{d\varphi^2} + 2R^2 [E - V(\varphi)] \Psi = 0 \quad (2)$$

Substituting the singular oscillator potential $V^{so}(\varphi)$ in eq. (2), we obtain Pöschl-Teller-type equation

$$\frac{d^2\Psi}{d\varphi^2} + \left[\epsilon - \frac{k_0^2 - \frac{1}{4}}{\cos^2 \varphi} - \frac{k_1^2 - \frac{1}{4}}{\sin^2 \varphi} \right] \Psi = 0 \quad (3)$$

where $\epsilon = 2R^2E + \omega^2R^4$ and $k_0^2 = \omega^2R^4 + \frac{1}{4}$. The regular (at points $\varphi = 0$ and $\pi/2$) solution of above equation maybe chosen in following form [5]

$$\Psi_n(\varphi) = \sqrt{\frac{2(2n + k_0 \pm k_1 + 1)\Gamma(n + k_0 + k_1 + 1)\Gamma(n \pm k_1 + 1)}{R[\Gamma(1 \pm k_1)]^2\Gamma(n + k_0 + 1)(n)!}}$$

$$(\sin \varphi)^{\frac{1}{2} \pm k_1} (\cos \varphi)^{\frac{1}{2} + k_0} {}_2F_1(-n, n + k_0 \pm k_1 + 1; 1 \pm k_1; \sin^2 \varphi), \quad (4)$$

with

$$\epsilon = (2n \pm k_1 + k_0 + 1)^2, \quad n = 0, 1, 2, \dots \quad (5)$$

Then the energy spectrum of the one-dimensional singular oscillator is given by

$$E_n(R) = \frac{1}{2R^2} \left[(2n \pm k_1 + \frac{1}{2})^2 + (2k_0 + 1)(2n \pm k_1 + 1) \right]. \quad (6)$$

Let us remark that the wave-functions have been normalized in the domain $[0, \pi/2]$. The positive sign at the k_1 has to taken whenever $k_1 > \frac{1}{2}$, i.e. the additional term to oscillator potential is repulsive at the origin and the motion take place only in domain $\varphi \in [0, \pi/2]$. If $0 < |k_1| \leq \frac{1}{2}$, i.e. the additional term is attractive at the origin, both the positive and negative sign must be taken into account in the solution. The motion in this case take place in $\varphi \in [-\pi/2, \pi/2]$. This is indicated by the notion $\pm k_1$, in the formulas.

3. Let us to write the Schrödinger equation (2) for singular Coulomb potential $V^c(\vec{s})$

$$\frac{d^2\Psi}{d\varphi^2} + \left(2R^2E + 2\mu R \cot |\varphi| + \frac{p^2 - \frac{1}{4}}{\sin^2 \varphi} \right) \Psi = 0. \quad (7)$$

First we will consider the region $\varphi \in [0, \pi]$. We make now a transformation to the new variable $\theta \in [0, \frac{\pi}{2}]$

$$e^{i\varphi} = \cos \theta, \quad (8)$$

which is possible if we continue the variable φ in the complex domain: $\text{Re } \varphi = 0, 0 \leq \text{Im } \varphi < \infty$ (see Fig. 1). We complexify also the coupling constant μ by putting $k = i\mu$ such that

$$\mu \cot \varphi = k(1 - 2 \sin^{-2} \theta). \quad (9)$$

As result we obtain the equation

$$\frac{d^2W}{d\theta^2} + \left[\epsilon - \frac{k_0^2 - \frac{1}{4}}{\cos^2 \theta} - \frac{k_1^2 - \frac{1}{4}}{\sin^2 \theta} \right] W = 0, \quad (10)$$

where $W(\theta) = (\cot \theta)^{\frac{1}{2}}Z(\theta)$ and

$$\epsilon = 2R^2E + 2kR, \quad k_0^2 = 2R^2E - 2kR, \quad k_1^2 = 2 - 4p^2. \quad (11)$$

From the above equation we see that, up to the substitution (11) the equation (10) for one-dimensional singular Coulomb problem (with restriction $0 < p^2 \leq \frac{1}{2}$) coincides to the one-dimensional singular oscillator equation (3).




Figure 1: Domain $\mathbf{G} = \{ 0 \leq \text{Re } \varphi \leq \pi; 0 \leq \text{Im } \varphi < \infty \}$ on the complex plane of φ .

The regular, for $\theta \in [0, \pi/2]$ and $k_0 \geq 1/4$, solution of this equation according to (4) is

$$\Psi(\theta) = \frac{W(\theta)}{\sqrt{\cot \theta}} = C_n (\sin \theta)^{1 \pm k_1} (\cos \theta)^{k_0} {}_2F_1(-n, n + k_0 \pm k_1 + 1; 1 \pm k_1; \sin^2 \theta) \quad (12)$$

where C_n is the normalization constant. To compute the constant C_n we require that the wave function (12) satisfy the condition

$$R \int_0^\pi \Psi_n \Psi_n^\diamond d\varphi = \frac{1}{2}, \quad (13)$$

where the symbol “ \diamond ” means the complex conjugate together with the inversion $\varphi \rightarrow -\varphi$, i.e. $\Psi^\diamond(\varphi) = \Psi^*(-\varphi)$. [We choose the scalar product as Ψ^\diamond because for $\varphi \in \mathbf{G}$ and real μ , and ϵ the function $\Psi^\diamond(\varphi)$ also belongs to the solution space of (7)]. By analogy to the work

[1, 3], we consider the integral over contour G in the complex plane of variable φ (see Fig.1)

$$\begin{aligned} \oint \Psi_n(\varphi) \Psi_n^\diamond(\varphi) d\varphi &= \int_0^\pi \Psi_n(\varphi) \Psi_n^\diamond(\varphi) d\varphi + \int_\pi^{\pi+i\infty} \Psi_n(\varphi) \Psi_n^\diamond(\varphi) d\varphi \\ &+ \int_{\pi+i\infty}^{i\infty} \Psi_n(\varphi) \Psi_n^\diamond(\varphi) d\varphi + \int_{i\infty}^0 \Psi_n(\varphi) \Psi_n^\diamond(\varphi) d\varphi. \end{aligned} \quad (14)$$

Using the facts that the integrand vanishes as $e^{2ik_0\varphi}$ and that $\Psi_n(\varphi)$ is regular in the domain \mathbf{G} (see Fig.1), then according to the Cauchy theorem we have

$$\int_0^\pi \Psi_n(\varphi) \Psi_n^\diamond(\varphi) d\varphi = (1 - e^{2i\pi k_0}) \int_0^{i\infty} \Psi_n(\varphi) \Psi_n^\diamond(\varphi) d\varphi. \quad (15)$$

Making the substitution (8) in the right integral of eq. (15), we find

$$\int_0^\pi \Psi_n(\varphi) \Psi_n^\diamond(\varphi) d\varphi = i (1 - e^{2i\pi k_0}) \int_0^{\frac{\pi}{2}} [\Psi_n] \tan \theta d\theta. \quad (16)$$

and after integration over the angle θ we finally get [6]

$$C_n = \sqrt{\frac{(-ik_0)(2n + k_0 \pm k_1 + 1)\Gamma(n + 1 \pm k_1)\Gamma(n + k_0 \pm k_1 + 1)}{R[1 - e^{2i\pi k_0}](2n \pm k_1 + 1)n![\Gamma(1 \pm k_1)]^2\Gamma(n + k_0 + 1)}}. \quad (17)$$

Let us now consider two most interesting cases.

(i). The case when $p^2 = \frac{1}{4}$. Then the duality transformation (8) establish the connection between pure Coulomb problem and singular oscillator with $k_1^2 = 1$. Comparing the eqs. (5) with (11) and putting $k = i\mu$, we get

$$k_0 = -(n + 1) + i\sigma, \quad \sigma = \frac{\mu R}{n + 1} \quad (18)$$

and for energy spectrum

$$E_n(R) = \frac{(n + 1)^2}{2R^2} - \frac{\mu^2}{2(n + 1)^2}, \quad n = 0, 1, 2, \dots \quad (19)$$

Returning to the variable φ , we obtain that wave function at $0 \leq \phi \leq \pi$ has the form

$$\Psi_{n\sigma}(\varphi) = C_n(\sigma) \sin \varphi e^{-i\varphi(n-i\sigma)} {}_2F_1(-n, 1 + i\sigma; 2; 1 - e^{2i\varphi}), \quad (20)$$

where the normalization constant $C_n(\sigma)$ is

$$C_n(\sigma) = e^{\frac{\sigma\pi}{2}} |\Gamma(1 + i\sigma)| \sqrt{\frac{(n + 1)^2 + \sigma^2}{\pi R}}. \quad (21)$$

The wave function in the region $-\pi \leq \varphi < 0$ ($s_1 < 0$) maybe determined from the eq. (20) by the reflection $\varphi \rightarrow -\varphi$. Therefore the general solution of the Schrödinger equation for $\varphi \in [-\pi, \pi]$ can be presented in form of even and odd function

$$\begin{aligned} \Psi_{n\sigma}^{(+)}(\varphi) &= C_n(\sigma) \sin |\varphi| e^{-i(n-i\sigma)|\varphi} F(-n, 1 + i\sigma; 2; 1 - e^{2i|\varphi|}), \\ \Psi_{n\sigma}^{(-)}(\varphi) &= C_n(\sigma) \sin \varphi e^{-i(n-i\sigma)|\varphi} F(-n, 1 + i\sigma; 2; 1 - e^{2i|\varphi|}). \end{aligned}$$

Thus by using the relation between Coulomb and singular oscillator systems we have constructed the wave functions and energy spectrum for a Coulomb system on the one-dimensional sphere.

(ii). Choosing now $k_1^2 = \frac{1}{4}$ or the same $p^2 - \frac{1}{4} = \frac{3}{16}$. In this case the centrifugal potential term is repulsive at the origin for singular Coulomb systems and the motion take place only in one of the domains $\varphi > 0$ or $\varphi < 0$. For oscillator system this term equal zero and therefore the duality transformation (8) connect singular Coulomb and pure oscillator systems.

Let us introduce the quantity ν which takes two values $\nu = \frac{1}{4}$ and $\nu = \frac{3}{4}$. Making all calculations by analogy to previous case, it is easy to obtain the energy spectrum

$$E_n^\nu(R) = \frac{(n + \nu)^2}{2R^2} - \frac{\mu^2}{2(n + \nu)^2}, \quad n = 0, 1, 2, \dots \quad (22)$$

and wave functions

$$\begin{aligned} \Psi_{n\sigma}^\nu(\varphi) &= e^{\frac{\sigma\pi}{2}} 2^\nu \frac{|\Gamma(\nu + i\sigma)|}{\Gamma(2\nu)} \sqrt{\frac{[(n + \nu)^2 + \sigma^2]\Gamma(n + 2\nu)}{4\pi R(n + \nu)n!}} \\ &\times (\sin \varphi)^\nu e^{-i\varphi(n - i\sigma)} {}_2F_1(-n, \nu + i\sigma; 2\nu; 1 - e^{2i\varphi}), \end{aligned}$$

where $\sigma = \mu R / (n + \nu)$.

In the contraction limit $R \rightarrow \infty$, $\varphi \rightarrow 0$ and $R\varphi \sim x$ - fixed, we see that

$$\epsilon_n = \lim_{R \rightarrow \infty} E_n^\nu(R) = -\frac{\mu^2}{2(n + \nu)^2}, \quad n = 0, 1, \dots \quad (23)$$

and

$$\Phi_n^\nu(y) = \lim_{\substack{R \rightarrow \infty \\ \varphi \rightarrow 0}} \Psi_{n\sigma}^\nu(\varphi) = \frac{\sqrt{\mu}}{\Gamma(2\nu)} \frac{1}{(n + \nu)} \sqrt{\frac{\Gamma(n + 2\nu)}{2n!}} y^\nu e^{-\frac{|y|}{2}} {}_1F_1(-n; 2\nu; y), \quad (24)$$

where $y = 2\mu x / (n + \nu)$. Formulas (23) and (24) coincides with the formulas for energy levels ϵ_n and up to the factor $\sqrt{2}$ for wave functions $\Phi_n^\nu(y)$ for two type one-dimensional Coulomb anyons with $\nu = \frac{1}{4}$ and $\nu = \frac{3}{4}$ respectively [4].

Acknowledgments

The research of L.G.M. and G.S.P. was partially supported by the NATO Collaborative Linkage Grant No. 978431 and ANSEF Grant No. PS81.

References

- [1] E.G.Kalnins, W.Miller Jr. and G.S.Pogosyan. Coulomb-oscillator duality in spaces of constant curvature, *J.Math.Phys.* **41**, 2629-2657, 2000.
- [2] A.Nersessian and G.S.Pogosyan. On the relation of the oscillator and Coulomb systems on (pseudo)spheres. quant-ph/0006118; *Phys.Rev* **A63**, 20103 (R), 2000.

- [3] E.G.Kalnins, W.Miller Jr. and G.S.Pogosyan. The Coulomb-Oscillator Relation on n-Dimensional Spheres and Hyperboloids. *Physics of Atomic Nuclei*, **65(6)**, 1086-1094, 2002.
- [4] V.M.Ter-Antonyan. Dyon -Oscillator Duality, quant-ph/0003106, 2000.
- [5] S.Flügge, *Problems in Quantum Mechanics*, V.1 Springer - Verlag, Berlin - Heidelberg - New York, (1971).
- [6] A.Erdelyi, W.Magnus, F.Oberhettinger and F.Tricomi. *Tables of Integral Transforms*, (McGraw-Hill, New York, 1954), Vol II.