Z_N Gauge Theories on a Lattice and Quantum Memory

Gaku Arakawa¹ and Ikuo Ichinose²

Department of Applied Physics, Graduate School of Engineering, Nagoya Institute of Technology, Nagoya, 466-8555 Japan

Abstract

In the present paper we shall study (2 + 1) dimensional Z_N gauge theories on a lattice. It is shown that the gauge theories have two phases, one is a Higgs phase and the other is a confinement phase. We investigate low-energy excitation modes in the Higgs phase and clarify relationship between the Z_N gauge theories and Kitaev's model for quantum memory and quantum computations. Then we study effects of random gauge couplings(RGC) which are identified with noise and errors in quantum computations by Kitaev's model. By using a duality transformation, it is shown that timeindependent RGC give no significant effects on the phase structure and the stability of quantum memory and computations. Then by using the replica methods, we study Z_N gauge theories with time-dependent RGC and show that nontrivial phase transitions occur by the RGC.

1 Introduction

In the last few years, discrete gauge theories have got renewed interests as a possible device for the quantum computations, a quantum computer. This idea was first proposed by Kitaev in his seminal paper[1], and after that there appeared interesting works on this idea[2, 3, 4, 5]. One of the most difficult problem of making a quantum computer and performing quantum computations fault-tolerantly is the stability of the quantum states which participate in quantum memory and computations. There must be a (large) energy gap between these states and others in the system and also mixings of these states must be suppressed by certain effects or selection rules. Then one can conceive that topological interactions such as the Aharonov-Bohm(AB) effect may play an important role there. The AB effect in the two spatial dimensions gives nontrivial statistics to particles with gauge interactions, i.e., anyons. The groundstates of the anyons are degenerate if the space is a torus and almost no mixing occurs between them because of the *topological* quantum number. Whereas the gauge symmetry should be descrete in order to avoid long-range interactions

¹e-mail address: e101608@phys.kyy.nitech.ac.jp

²e-mail address: ikuo@ks.kyy.nitech.ac.jp

besides the topological ones. Kitaev's model is based on the anyonic excitations in qubits system. However its detailed relationship to the gauge theory is still missing.

In this paper, we shall study discrete Z_N gauge theories in (2 + 1) dimensions. There are two phases in these model, one is a confinement phase and the other is a Higgs phase. We show that Kitaev's system of qubits corresponds to some limit of the Higgs phase of the Z_2 gauge models. Stability of Kitaev's model is closely related with the confinement-Higgs phase transition of the Z_N gauge models.

This paper is organized as follows. In Sec.2, we study Z_N gauge theories which appear as a result of spontaneous breakdown of U(1) gauge symmetry. We clarify the relationship between the gauge system and Kitaev's model for quantum memory and computations. In Sec.3, low-energy excitations in the Higgs phase are investigated. There appear anyonic excitations, magnetic vortices and dyons in a natural way as in the spontaneously broken gauge systems in the continuum space[6]. In Sec.4, phase structure and effects of the (static) random gauge couplings(RGC) are investigated by using a duality transformation. The Z_N gauge systems are transformed to spin systems which are more tractable than the gauge systems. In Sec.5, effects of the time-dependent RGC are studied by the replica methods. It is found that nontrivial phase transitions occur as the RGC varies. Section 6 is devoted to conclusion.

2 U(1) and Z_N gauge theories

Let us start with the following U(1) Abelian gauge-Higgs model on a 2-dimensional (2D) square lattice. Hamiltonian is given by,

$$H_{U(1)} = g^{2} \sum_{link} E_{xi}^{2} - \frac{1}{g^{2}} \sum_{plaquette} UUUU + \frac{1}{\kappa} \sum_{x} (\Pi_{x}^{\phi})^{2} - \kappa \sum_{link} \phi_{x+i}^{\dagger} U_{xi}^{N} \phi_{x} - \gamma \sum_{link} \psi_{x+i}^{\dagger} U_{xi}^{q} \psi_{x} + M \sum_{x} \psi_{x}^{\dagger} \psi_{x} - \gamma \sum_{link} \varphi_{x+i}^{\dagger} U_{xi}^{-q} \varphi_{x} + M \sum_{x} \varphi_{x}^{\dagger} \varphi_{x} + \text{H.c.},$$
(1)

where U_{xi} is the U(1) gauge field on the link $(x, i)(x = \text{site}, i = \hat{1} \text{ or } \hat{2})$ and E_{xi} is the conjugate electric field. The Higgs field $\phi_x \in U(1)$ carries U(1) charge N whereas the charge of the fermion field¹ $\psi_x(\varphi_x)$ is q(-q) which is an integer. Π_x^{ϕ} is the conjugate field of ϕ_x , the gauge coupling is g and the fermion mass is M. Other notations are standard. We are interested in the case $N \neq 1$. In this case there are two phases in the model, one is the Higgs phase and the other is the confinement phase. In particular in the limit $g^2 \to 0$, the gauge field U_{xi} is restricted to the pure-gauge configuration and the model reduces to a Hamiltonian description of the classical 3D XY spin model plus the free

¹We often call ψ_x and φ_x fermion because they satisfy fermionic anticommutation relations. As a result of the gauge interactions, they obey anyonic statistics in the Higgs phase. See later discussion.

fermion system. The classical 3D XY model exhibits a phase transition from the magnetized phase to the disordered phase at a critical coupling κ_c . On the other hand for large κ , quantum fluctuations of ϕ_x are suppressed and low-energy excitations of the gauge and Higgs fields are restricted as

$$U_{xi}^N \sim 1, \quad \phi_x \sim 1, \tag{2}$$

up to (time-independent) local gauge transfomation. Then we can put

$$U_{xi} \sim Z_{xi},\tag{3}$$

where the Z_N gauge operator Z_{xi} is explicitly given as follows by $(N \times N)$ matrix,

$$Z_{xi} = \begin{pmatrix} 1 & 0 & \cdot & \cdot & 0 \\ 0 & e^{\frac{2\pi}{N}i} & 0 & \cdot & 0 \\ \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & 0 \\ 0 & \cdot & \cdot & \cdot & e^{\frac{2\pi(N-1)}{N}i} \end{pmatrix}.$$
 (4)

Corresponding to the above representation of Z_{xi} , we introduce "conjugate matrix" X_{xi} as follows,

$$X_{xi} = \begin{pmatrix} 0 & 1 & 0 & \cdot & 0 \\ 0 & 0 & 1 & 0 & 0 \\ \cdot & \cdot & \cdot & \cdot & \cdot \\ 0 & 0 & \cdot & 0 & 1 \\ 1 & 0 & \cdot & \cdot & 0 \end{pmatrix}.$$
 (5)

One can easily verify the following commutation relations,

$$X_{xi}Z_{xi} = e^{\frac{2\pi}{N}i}Z_{xi}X_{xi}, \quad X_{xi}Z_{yj} = Z_{yj}X_{xi} \text{ for } (x,i) \neq (y,j).$$
(6)

The electric term in Eq.(1) is reduced to the following term in the reduced Z_N gauge-field space,

$$E_{xi}^2 \sim -(X_{xi} + X_{xi}^\dagger).$$

The above result can be shown by using the eigenstates of the electric fields as basis vectors. Let us define an "empty state" $|0\rangle$ as

$$E|0\rangle = 0,\tag{7}$$

where we have omitted link index for notational simplicity. By using the following commutation relation,

$$[E, U] = U, (8)$$

we can show

$$E^2 U|0\rangle = U|0\rangle \equiv |1\rangle. \tag{9}$$

Then the gauge field U is the rising operator of the electric field.

For the Z_N case, we also define "empty" state for the X operator,

$$X|0\rangle_X = |0\rangle_X, \quad X^{\dagger}|0\rangle_X = |0\rangle_X.$$
 (10)

The state $|0\rangle_X$ can be expressed by the eigenstates of the Z operator $|k\rangle_Z$, $Z|k\rangle_Z = e^{i\frac{2\pi}{N}k}|k\rangle_Z$,

$$|0\rangle_X = \frac{1}{\sqrt{N}} \Big(|1\rangle_Z + |2\rangle_Z + \dots + |N\rangle_Z \Big).$$
(11)

Then one can easily show,

$$Z|0\rangle_X = |1\rangle_Z + e^{\frac{2\pi}{N}i}|2\rangle_Z + \dots + e^{2\pi\frac{N-1}{N}i}|N\rangle_Z,$$
(12)

and therefore

$$XZ|0\rangle_X = e^{\frac{2\pi}{N}i}Z|0\rangle_X, \tag{13}$$

$$(X + X^{\dagger})Z|0\rangle_X = 2\cos\left(\frac{2\pi}{N}\right)Z|0\rangle_X.$$
 (14)

From Eqs.(10) and (14), Z is the lowering operator of $X + X^{\dagger}$ and therefore,

$$E_{xi}^2 \sim -(X_{xi} + X_{xi}^{\dagger}), \tag{15}$$

up to irrelevant additive and multiplicative constants. Then for large κ , the U(1) gauge theory (1) reduces to the following Z_N gauge theory,

$$H_{\mathcal{T}} = H_Z + H_Z^{\psi} + H_Z^{\varphi},$$

$$H_Z = -\lambda_1 \sum X_{xi} - \lambda_2 \sum ZZZZ + \text{H.c.},$$

$$H_Z^{\psi} = -\gamma \sum \psi_{x+i}^{\dagger} Z_{xi}^q \psi_x + M \sum \psi_x^{\dagger} \psi_x + \text{H.c.},$$

$$H_Z^{\varphi} = -\gamma \sum \varphi_{x+i}^{\dagger} Z_{xi}^{-q} \varphi_x + M \sum \varphi_x^{\dagger} \varphi_x + \text{H.c.},$$
(16)

where λ_1 and λ_2 are coupling constants of the Z_N gauge theory and they relate to the U(1) gauge coupling g^2 as $\lambda_1 \sim g^2$ and $\lambda_2 \sim 1/g^2$.

The above "derivation" of the Z_N gauge theory (16) from the U(1) gauge system (1) is rather sketchy but it might be useful for realization of discrete gauge systems in architecture of the quantum computers. For example, spontaneous breaking of U(1) gauge symmetry occurs in the superconductivity. In most of the superconductors including the high-temperature ones, the "Cooper pair" carries electric charge 2e. Then a discrete Z_2 gauge system close to the present one might be realized in some superconductors. The Hamiltonian (16) is directly obtained from the path-integral formalism of the Z_N gauge theory on 3D lattice by taking the continuum limit of the time-like direction. In the $3D Z_N$ gauge theory, there exist two phases, i.e., confinement and Higgs phases as we show later on. Phase transition occurs at a certain critical coupling $(\lambda_1/\lambda_2)_c$. In the original U(1) gauge theory, there exists a *critical line* connecting the XY phase transition at $(g = 0, \kappa = \kappa_c)$ and Z_N gauge phase transition at $(g = g_c, \kappa = \infty)$ (see Fig.1)[7].

Physical state of the system (16) must be gauge-invariant and this condition is given as follows,

$$G_x \equiv \left(\prod_{(y,i)\in x} \tilde{X}_{yi}\right) e^{-\frac{2\pi q}{N}i(\psi_x^{\dagger}\psi_x - \varphi_x^{\dagger}\varphi_x)}, \quad G_x|phys\rangle = |phys\rangle, \tag{17}$$

where $(y,i) \in x$ denotes 4 links emanating from site x and $\tilde{X}_{yi} = X_{yi}$ for y = x whereas $\tilde{X}_{yi} = X_{xi}^{\dagger}$ for y - i = x. It is proved that G_x is the gauge-transformation operator at site x and the Hamiltonian $H_Z^{\psi} + H_Z^{\phi}$ in Eq.(16) commutes with G_x .

Recently Kitaev proposed a 2-dimensional qubits system for fault-tolerant quantum memory and computations[1]. This system is closely related to the Z_2 gauge theory and contains "anyonic excitations". The system is defined on a torus and the Hamiltonian is given as follows in our notation,

$$H_K = -\sum_x \prod_{(y,i)\in x} X_{yi} - \sum_{pl} ZZZZ,$$
(18)

where Z_{xi} and X_{xi} are explicitly given by the Pauli matrices in the Z_2 case, i.e., $Z = \sigma^z$ and $X = \sigma^x$. The groundstates and excited states of the Hamiltonian (18) are easily obtained since the first and second terms of (18) commute with each other. The groundstates satisfy

$$\prod_{(y,i)\in x} X_{yi} |GS\rangle_K = |GS\rangle_K, \quad \prod_{pl} Z |GS\rangle_K = |GS\rangle_K, \tag{19}$$

for all sites and plaquettes. The groundstates are four-fold degenerate on the torus, as we explain in the following section. These degenerate lowest-energy states form basis for quantum memory[1].

The first excited states are explicitly given by

$$\prod_{(y,i)\in x} X_{yi}|1st\rangle_K = -|1st\rangle_K, \text{ or } \prod_{pl} Z|1st\rangle_K = -|1st\rangle_K,$$
(20)

for some specific site or plaquette and otherwise they satisfy Eq.(19). It is not so difficult to see that Kitaev's model is equivalent to the model (16) with N = 2, $\gamma = 0$, M = 2, q = 1 and $\lambda_1 = 0$, $\lambda_2 = 1$. With these parameters and the physical state condition (17), the groundstates of the gauge model are given as

$$\prod_{pl} Z|GS\rangle_Z = |GS\rangle_Z, \quad \psi_x^{\dagger}\psi_x|GS\rangle_Z = 0, \quad \varphi_x^{\dagger}\varphi_x|GS\rangle_Z = 0, \tag{21}$$

for all plaquettes and sites. From (17), the second and third conditions of (21) mean $\prod_{(y,i)\in x} X_{yi}|GS\rangle_Z = |GS\rangle_Z$. On the other hand, the first excited states of the gauge system are given by,

$$\prod_{pl} Z|1st\rangle_Z^V = -|1st\rangle_Z^V, \text{ or } \psi_x^\dagger \psi_x|1st\rangle_Z^\psi = |1st\rangle_Z^\psi, \text{ or } \varphi_x^\dagger \varphi_x|1st\rangle_Z^\varphi = |1st\rangle_Z^\varphi,$$
(22)

for some specific plaquette or site. From Eq.(17), the second condition in (22) is equivalent to $\prod_{(y,i)\in x} X_{yi}|1st\rangle_Z^{\psi} = -|1st\rangle_Z^{\psi} \text{ for } e^{\pi i \psi_x^{\dagger} \psi_x}|1st\rangle_Z^{\psi} = -|1st\rangle_Z^{\psi} \text{ and energy increases by 2 because of the mass term in (16) with } M = 2$. Similarly for the other fermion φ_x . In the original paper by Kitaev[1], relationship between his model and gauge theories was slightly discussed but full relationship was missing. In the following sections we shall study phase structure of the present Z_N gauge model, low-energy excitations, effects of random gauge couplings, etc. All these discussions give an important insight to the stability problem of Kitaev's model.

3 Low-energy excitations in the Higgs phase

As we show in the following section, there are two phases in the Z_N gauge theory in (2+1) dimensions H_Z in (16). For large λ_2/λ_1 , fluctuation of the gauge field Z_{xi} is small and the Higgs phase is realized whereas for small λ_2/λ_1 , the gauge field Z_{xi} fluctuates strongly and the confinement phase is realized. The Higgs phase of the model can be used for a quantum memory. Coupling of the matter fields ψ_x etc. enhances the Higgs phase.

Let us study the model on the *torus* and focus on the Higgs phase for large λ_2/λ_1 . In particular for $\lambda_1 = 0$, the groundstates are given by Eq.(21) and low-energy excited states are *particle states* of ψ_x , φ_x and states of *plaquette magnetic excitation* or *vortex*, i.e., $\prod_{pl} Z|1st\rangle_Z^V = -|1st\rangle_Z^V$ for specific plaquette. As we study the model on the torus, we have the following "trivial" identities

$$\prod_{\text{all sites}} \prod_{(y,i)\in x} \tilde{X}_{yi} = 1, \quad \prod_{\text{all pl's } pl} \prod_{pl} Z = 1,$$
(23)

and therefore the above excitations must appear in pairs. As the groundstates satisfy Eq.(21), there is no mangetic flux in each plaquette. Then one may think that the groundstate is unique. However this is not the case. There are two nontrivial cycles on the torus, and let us call them a-cycle and b-cycle, i.e., noncontractible closed paths. We introduce the dual lattice in the usual way, and choose certain noncontractible closed loops on the original and dual lattices. We use notations such that $C_Z^a(C_Z^b)$ for a suitably chosen closed loop corresponding to the a-cycle(b-cycle) on the *original* lattice and $C_X^a(C_X^b)$ for a loop corresponding to the a-cycle(b-cycle) on the *dual* lattice. Later discussion does not depend on the choice of the loops. Then we define the following operators, Z_a, Z_b, X_a and X_b ,

$$Z_a = \prod_{C_Z^a} Z_{xi}, \quad Z_b = \prod_{C_Z^b} Z_{xi},$$
$$X_a = \prod_{C_X^a} X_{xi}, \quad X_b = \prod_{C_X^b} X_{xi}, \tag{24}$$

where X_{xi} 's in X_a cross C_X^a and similarly for X_b . These operators are obviously invariant under gauge transformation and commute with H_Z when $\lambda_1 = 0$. Furthermore they satisfy the following commutation relations,

$$X_a Z_b = e^{\frac{2\pi}{N}i} Z_b X_a, \quad X_b Z_a = e^{\frac{2\pi}{N}i} Z_a X_b, \tag{25}$$

and otherwise commute. Therefore the groundstates are eigenstate of the the operators, e.g., Z_a and Z_b and they are N^2 -fold degenerate. This result holds even in the presence of the fermions ψ_x and φ_x since Z_a and Z_b commute with $H_{\mathcal{T}}$ for vanishing λ_1 in Eq.(16).

Fermions ψ_x and φ_x move in an unfluctuating "background" field of Z_{xi} 's with vanishing magnetic field. However they distinguish the above N^2 -fold degenerate Z's groundstates. In fact while ψ_x (or φ_x) fermion moves along a closed loop of the a-cycle, it acquires phase factor which is an eigenvalue of $(Z_a)^q$, and similarly for the b-cycle. Then the Higgs phase is a "topologically ordered" phase. The N^2 groundstates work as *qudit* for quantum memory and the quantum states of the qudit are distinguishable by using matter fields like ψ_x .

Let us discuss excitations in detail. As we explained above, the fermions must appear in a pair. Two-fermion state at sites x and y is explicitly given as,

$$|F;C_{xy}\rangle = \psi_y^{\dagger} \Big(\prod_{C_{xy}} Z^q \Big) \varphi_x^{\dagger} |GS\rangle_Z,$$
(26)

where C_{xy} is a certain path on the *original* lattice connecting x and y, and the state (26) obviously satisfies the physical-state condition (17). On the other hand two-vortex state at *dual* sites x^* and y^* is given as,

$$|V; \tilde{C}_{x^*y^*}\rangle = \Big(\prod_{\tilde{C}_{x^*y^*}} X\Big)|GS\rangle_Z,\tag{27}$$

where $\tilde{C}_{x^*y^*}$ is a certain path on the *dual* lattice connecting x^* and y^* and X's in (27) are on the links crossing $\tilde{C}_{x^*y^*}$ (see Fig.2). This state is also a physical state. Other physical excitations are produced by appling the gauge-invariant operators in Eqs.(26) and (27) successively on the groundstates.

Fermionic excitations and magnetic vortices satisfy a nontrivial statistics. This is an Aharonov-Bohm effect of the Z_N gauge theory. To see this, we consider the state like

$$\psi_{y_1}^{\dagger} \Big(\prod_{C_{x_1y_1}} Z^q \Big) \varphi_{x_1}^{\dagger} \cdot \Big(\prod_{\tilde{C}_{x_2^* y_2^*}} X \Big) |GS\rangle_Z, \tag{28}$$

and assume that the paths $C_{x_1y_1}$ and $\tilde{C}_{x_2^*y_2^*}$ do not entangle with each other. Let us move the φ_x fermion at x_1 around the vortex at x_2^* once counterclockwise (and not y_2^*) and then return it to the original position x_1 . The resultant path $C'_{x_1y_1}$ encircles x_2^* once and $C'_{x_1y_1}$ and $\tilde{C}_{x_2^*y_2^*}$ cross with each other. Then the state can be written as

$$\psi_{y_1}^{\dagger} \Big(\prod_{C'_{x_1y_1}} Z^q \Big) \varphi_{x_1}^{\dagger} \cdot \Big(\prod_{\tilde{C}_{x_2^* y_2^*}} X \Big) |GS\rangle_Z = \psi_{y_1}^{\dagger} \Big(\prod_{C_{x_1y_1}} Z^q \cdot \prod_{C_{\text{closed}}} Z^q \Big) \varphi_{x_1}^{\dagger} \cdot \Big(\prod_{\tilde{C}_{x_2^* y_2^*}} X \Big) |GS\rangle_Z, \quad (29)$$

where C_{closed} is the closed path $(C'_{x_1y_1} \cdot C^{-1}_{x_1y_1})$ which encircles x_2^* once and has a single common link (or odd number of links) with $\tilde{C}_{x_2^*y_2^*}$. Because of the nontrivial commutation relation between Z_{xi} and X_{xi} and (21), the resultant state differs from the original one by the phase factor $e^{\frac{2q\pi}{N}i}$,

$$\psi_{y_1}^{\dagger} \Big(\prod_{C_{x_1y_1}} Z^q \cdot \prod_{C_{\text{closed}}} Z^q \Big) \varphi_{x_1}^{\dagger} \cdot \Big(\prod_{\tilde{C}_{x_2^* y_2^*}} X\Big) |GS\rangle_Z = e^{\frac{2q\pi}{N}i} \psi_{y_1}^{\dagger} \Big(\prod_{C_{x_1y_1}} Z^q \Big) \varphi_{x_1}^{\dagger} \cdot \Big(\prod_{\tilde{C}_{x_2^* y_2^*}} X\Big) |GS\rangle_Z. \tag{30}$$

The above anyonic properties of the low-energy excitations are closely related with the groundstate degeneracy. In the continuum spacetime, a Chern-Simons(CS) gauge theory is often employed for describing anyons which are a nontrivial representation of the braid group. In anyon systems on a torus, movement of an anyon along noncontractible loops like the a-cycle and/or b-cycle is a nontrivial element of the braid group. On the torus, the zero modes of the CS gauge field play an important role and the groundstate wave function of anyons becomes multi-component because of the zero modes[8]. Similar phenomenon occurs in the present Z_N -gauge system as we explained above.

One may conceive that the system has dyonic excitations as in the continuum theories[6]. The answer is positive. Dyon d_x with "electric charge" Q_E and "magnetic charge" R is described by the following Hamiltonian,

$$H_D = -\sum d_{x+i}^{\dagger} Z_{xi}^{Q_E} X_{xi}^R d_x + \text{H.c.}, \qquad (31)$$

where we assume that the fields d_x and d_x^{\dagger} themselves satisfy the fermionic commutation relations for simplicity. The link $(\bar{x}i)$ is associated with the link (xi) and defined as follows,

link
$$(\bar{x}i) = \begin{cases} (x+1,2) & \text{for } i=1, \\ (x+2,1) & \text{for } i=2. \end{cases}$$
 (32)

From the above definition (31) and (32), it is obvious that the electric charge Q_E of the dyon d_x is located at the site x whereas its magnetic charge R is located at the nearest-neighbor plaquette (see Fig.3). Regularization is naturally introduced by the spatial lattice. It is not so difficult to show that the above dyon satisfies nontrivial representation of the braid group and there appears the phase factor like $-\exp(\pm \frac{2(Q_E+R)\pi i}{N})$ when two dyons interchange with each other. When we turn on the parameter λ_1 in H_Z (16), the operator Z_a and Z_b do not commute with H_T anymore and therefore degeneracy of the groundstate disappears. This stems from the fact that because of the term $\lambda_1 \sum X_{xi}$, Z_{xi} becomes dynamical and it fluctuates quantum mechanically and then genuine anyonic properties of the low-energy excitations break down. However for small λ_1 , there is still an energy gap between the N^2 "groundstates" with fine structure and the other excited states. Furthermore, these N^2 states are far apart with each other in the quantum-mechanical configuration space and are hardly mixed if the torus is sufficiently large. Therefore the system with small value of λ_1 is still suited for a quantum memory as Kitaev suggested first. However as λ_1 increases, a phase transition occurs as we show in the following section. In the new phase, a confinement phase, the gauge field fluctuates randomly and the system is useless as a quantum memory.²

4 Duality transformation, phase transition and random gauge couplings

In the previous section, we discussed that for the quantum memory and commputations the Higgs phase must be realized in the present system. In this section we shall study the phase structure of the gauge-theory model H_Z in (16). To this end, the system is defined on a large spatial square lattice. We shall perform a duality transformation which transforms the gauge-theory model into a more tractable spin model. For the Z_2 gauge theory, the duality transfomation is discussed in Kogut's review article[10].

Let us consider the pure gauge system H_Z in (16) with the physical state condition,

$$\prod_{(y,i)\in x} \tilde{X}_{yi} = X_{x1}X_{x,-1}X_{x2}X_{x,-2} = X_{x1}X_{x-1,1}^{\dagger}X_{x2}X_{x-2,2}^{\dagger} = 1.$$
(33)

By solving the above condition (33), the operator X_{x2} is given as follows by the remaining operators,

$$X_{x2} = X_{x1}^{\dagger} X_{x,-1}^{\dagger} X_{x-2,1}^{\dagger} X_{x-2,-1}^{\dagger} \cdots .$$
(34)

As the "conjugate" operators Z_{x2} of X_{x2} commute with the Hamiltonian H_Z , we can set it as a constant, $Z_{x2} = 1$.

Then we introduce the following dual operators W_{x^*} and V_{x^*} which reside on sites of the dual $^{-2}$ It is very interesting to see that similar gauge-theory argument can be applied to neural network models for brain[9]. There Higgs phase corresponds to good brains and the confinement phase to dementia.

lattice,

$$W_{x^*} = \prod_{(yi)\in x^*} Z_{yi},$$

$$V_{x^*} = \prod_{l\geq 0} X_{x-2l,1},$$
 (35)

where $(yi) \in x^*$ denotes 4 links to the plaque on the original lattice which is dual to the site x^* of the dual lattice. From the definition (35), one can easily verify relations like,

$$W_{x^*}^N = V_{x^*}^N = 1, \quad V_{x^*} W_{x^*} = e^{i\frac{2\pi}{N}} W_{x^*} V_{x^*},$$

$$V_{x^*} W_{y^*} = W_{y^*} V_{x^*}, \text{ for } x^* \neq y^*,$$

$$V_{x^*} W_{x^*}^{\dagger} = e^{-i\frac{2\pi}{N}} W_{x^*}^{\dagger} V_{x^*},$$

(36)

$$V_{x^*}V_{x^*-2}^{\dagger} = X_{x1}, \quad V_{x^*}^{\dagger}V_{x^*-1} = X_{x2}.$$
(37)

From Eqs.(36) and (37), the Hamiltonian H_Z in (16) can be rewritten in terms of V_{x^*} and W_{x^*} ,

$$H_Z = -\lambda_1 \sum_{x^*, i=1,2} V_{x^*} V_{x^*-i}^{\dagger} - \lambda_2 \sum_{x^*} W_{x^*} + \text{H.c.}$$
(38)

The above quantum Hamiltonian (38) is nothing but that of the 3D classical Z_N Ising model(the clock model) which is obtained by the transfer-matrix methods and taking the continuum limit of one direction.

The Hamiltonian (38) is more tractable than the original one (16). There are two phases, i.e., ordered and disordered phases, and a phase transition occurs as the value λ_1/λ_2 varies. For small λ_1/λ_2 limit, the groundstate is given by

$$W_{x^*}|0\rangle_S = W_{x^*}^{\dagger}|0\rangle_S = |0\rangle_S.$$
(39)

In the representation,

$$W_{x^*} = \begin{pmatrix} 1 & 0 & \cdot & \cdot & 0 \\ 0 & e^{\frac{2\pi}{N}i} & 0 & \cdot & 0 \\ \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & 0 \\ 0 & \cdot & \cdot & \cdot & e^{\frac{2\pi(N-1)}{N}i} \end{pmatrix},$$
(40)

the above groundstate $|0\rangle_S$ is explicitly given as,

$$|0\rangle_{S} = \prod_{x^{*}} |0\rangle_{x^{*}}, \ |0\rangle_{x^{*}} = \begin{pmatrix} 1\\ 0\\ .\\ .\\ 0\\ 0 \end{pmatrix}.$$
 (41)

For small but nonvanishing λ_1/λ_2 , the groundstate is obtained by the usual perturbative calculation, and the term $V_{x^*}V_{x^*-i}^{\dagger}$ tilts nearest-neighbor W_{x^*} and W_{x^*-i} by $\pm \frac{2\pi}{N}$, respectively. In this phase, there is no "magnetization", i.e.,

$${}_{S}\langle 0|V_{x^{*}}|0\rangle_{S} = 0.$$
 (42)

Low-energy excitations are given by,

$$\frac{1}{\sqrt{N_s}} \sum_{x^*} e^{ip \cdot x^*} V_{x^*} |0\rangle_S, \quad \frac{1}{\sqrt{N_s}} \sum_{x^*} e^{ip \cdot x^*} V_{x^*}^{\dagger} |0\rangle_S, \tag{43}$$

where 2-vector p is a momentum and N_s is the number of the sites. Excited energy of the above states (43) can be easily calculated and obtained as follows for small λ_1/λ_2 ,

$$E = 2\lambda_2 \left(1 - \cos(\frac{2\pi}{N}) \right) + \cdots .$$
(44)

From (44), the energy gap is a decreasing function of N.

For large λ_1/λ_2 , on the other hand, the groundstate of the spin system (38) is given by

$$V_{x^*}|\tilde{0}\rangle_S = e^{i\alpha}|\tilde{0}\rangle_S, \quad V_{x^*}^{\dagger}|\tilde{0}\rangle_S = e^{-i\alpha}|\tilde{0}\rangle_S, \tag{45}$$

where $e^{i\alpha} \in Z_N$ and therefore it is N-fold degenerate.³ There is a nonvanishing magnetization for large λ_1/λ_2 ,

$${}_{S}\langle \tilde{0}|V_{x^{*}}|\tilde{0}\rangle_{S} \neq 0.$$

$$\tag{46}$$

From Eqs.(35), (46) and the discussion in the previous section, it is obvious that vortex condensation occurs in the gauge-system state corresponding to $|\tilde{0}\rangle_S$. This means that for large λ_1/λ_2 the confinement phase is realized and therefore the gauge system does not work as a quantum memory. This result is important for the architecture of the quantum computer.

It is interesting and also important to study another type of disturbance for realization of the Higgs phase, or, a good quantum memory, i.e., the effect of random gauge couplings which corresponds to noise and errors in quantum computations. In this section, we consider *static* random

³This result of the spin system does *not* mean that the original gauge system has the degenerate groundstates. Actually from (16), the groundstate satisfies $X_{xi}|GS$; gauge $\rangle = 1$ for all links (xi) in the large λ_1/λ_2 limit.

gauge coupling (RGC) with random variables $\tau_{x^*} \in Z_N$, and the Hamiltonian is given by,

$$H_Z^R = -\lambda_1 \sum_{link} X_{xi} - \lambda_2 \sum_{pl} \tau_{x^*} Z Z Z Z.$$
(47)

We assume a simple local correlation for the random variables,

$$[\tau_{x^*}\tau_{y^*}] \propto \delta_{x^*y^*},\tag{48}$$

where $[\cdots]$ denotes the ensemble average.

It seems rather difficult to study the above random gauge system (47). However by using the duality transformation (35), we can rewrite the Hamiltonian H_Z^R as in the nonrandom case which we studied above,

$$H_Z = -\lambda_1 \sum_{x^*, i=1,2} V_{x^*} V_{x^*-i}^{\dagger} - \lambda_2 \sum_{x^*} \tau_{x^*} W_{x^*} + \text{H.c.}$$
(49)

Then we *redefine* the dual operators W_{x^*} as follows,

$$\tilde{W}_{x^*} \equiv \tau_{x^*} W_{x^*}.\tag{50}$$

One can easily verify that the new operator \tilde{W}_{x^*} and the old one W_{x^*} satisfy exactly the same operator equations in (36), and also there are *no* spatial correlations of \tilde{W}_{x^*} because of (48). Then H_Z^R is equivalent to the original H_Z and the random system has the same phase structure with the nonrandom one. The groundstate, excitated states, etc. are different in the two systems but there exists one-to-one correspondence between them. This result can be partly seen in the original gauge system (47). For *static* RGC τ_{x^*} and $\lambda_1 = 0$, there is a *unique* Z-field configuration of the lowest energy up to local gauge transformations. Vortex excitations are generated by applying the string operator ($\prod X_{xi}$) on this lowest-energy configuration as in the nonrandom case. Then there is one-to-one correspondence. The perturbative term $\lambda_1 X_{xi}$ generates a pair of vortices in a nearest-neighbor plaquettes from the lowest-energy state as in the nonrandom system.

In the following section, we shall study random Z_2 gauge system with full RGC by the replica methods and show that nontrivial phase structure appears.

5 Replica mean-field theory

In this section we shall study the *d*-dimensional random Z_2 gauge theories by the replica mean-field theory(RMFT). RMFT has been often applied to the random spin systems and spin-glass problems. The random gauge theories have been less studied and as far as we know there is no systematic studies on the random gauge theories by the replica methods. Numerical Monte-Carlo simulations are also useful to obtain phase diagram in the p-T plane, where p is the concentration of plaquettes of "wrong sign" and T is the temperature[11, 12]. In the previous sections we used the Hamiltonian formalism, but in this section we employ the path-integral Lagrangian formalism since the pathintegral formalism is more suitable for the MFT.

Let us first study the nonrandom Z_2 gauge theory on a *d*-dimensional lattice by the MFT[13]. The partition function Z is given by,

$$Z = \operatorname{Tr} e^{-\beta S}, \quad S = -\sum_{pl} \sigma \sigma \sigma \sigma - h \sum_{link} \sigma_{xi}, \quad (51)$$

where the Z_2 gauge variables $\sigma_{xi}(i = 1, \dots, d)$ take ± 1 , Tr means $\sum_{\sigma_{xi}=\pm 1}, \beta$ is inverse temperature and h is an external "magnetic field". It is not so difficult to drive MFT action S_M . To this end, let us decompose σ_{xi} as $\sigma_{xi} = U_0 + \delta \sigma_{xi}$ where U_0 is the MF for σ_{xi} and $\delta \sigma_{xi}$ is the fluctuation from it[14]. In terms of the new variables,

$$S = -U_0^4 N_P - U_0^3 \cdot 2(d-1) \sum_{link} \delta \sigma_{xi} - h \sum_{link} \sigma_{xi} + O((\delta \sigma_{xi})^2)$$

= $-U_0^4 N_P - U_0^3 \cdot 2(d-1) \sum_{link} \sigma_{xi} + 2(d-1)U_0^4 N_L - h \sum_{link} \sigma_{xi} + O((\delta \sigma_{xi})^2),$ (52)

where N_P and N_L are the numbers of plaquettes and links of the lattice, respectively and $N_L = \frac{2}{d-1}N_P$. From (52), S_M is obtained as,

$$S_M = 3U_0^4 N_P - \{2(d-1)U_0^3 + h\} \sum_{link} \sigma_{xi}.$$
(53)

Then it is straightforward to calculate the partition function from S_M in (53),

$$Z_{MF} = \operatorname{Tr} e^{-\beta S_M} = e^{-3\beta U_0^4 N_P} \left[2 \cosh \beta \{ 2(d-1)U_0^3 + h \} \right]^{N_L}.$$
(54)

The "magnetization" m per link is calculated from (54) as

$$m = \frac{1}{Z_{MF}N_L} \frac{\partial Z_{MF}}{\partial h}$$

= $\tanh \beta \{2(d-1)U_0^3 + h\}.$ (55)

Similarly the free energy is obtained as,

$$F = -\frac{1}{\beta} \frac{2}{d-1} N_P \log \left[2 \cosh \beta \{ 2(d-1)U_0^3 + h \} \right] + 3U_0^4 N_P.$$
(56)

Numerical calculation of the free energy F in (56) is given in Fig.4 as a function U_0 for vanishing hand at various inverse temperatures β . From Eqs.(55) and (56), it is verified that the magnetization m is equal to the value of U_0 at stationary points of $F = F(U_0)$. Result in Fig.4 shows that there is a first-order phase transition as the temperature varies and at low temperature the magnetiztion m is novanishing.

It is known that there is a *second-order* phase transition in $3D Z_2$ gauge theory which is dual to the 3D Ising model as we showed in the previous section[10]. This means that the MFT gives correct results only at large spatial dimensions as it is well known for the spin systems, etc. However we believe that the MFT is still useful for obtaining rough estimations of the physical quantities, phase stucture of systems, etc.

Let us turn to random gauge theories (RGT). We study the random Z_2 theory with the following action,

$$S_R = -\sum_{pl} J_p \sigma \sigma \sigma \sigma, \tag{57}$$

where we assume that the RGC J_p has the probability distribution like,

$$P(J_p) = \frac{1}{J\sqrt{2\pi}} \exp\left\{-\frac{1}{2J^2}(J_p - J_0)^2\right\},$$
(58)

with positive parameters J and J_0 . We choose (58) in which J_p takes continuous real value instead of the discrete distribution $J_p = \pm J_0$, because it is more tractable.

We apply replica tricks to the above RGT and then the partition function is given as,

$$[Z^n] = \int \left\{ \prod_p dJ_p P(J_p) \right\} \operatorname{Tr} \exp\left(\beta \sum_p J_p \sum_{\alpha=1}^n \prod_p \sigma^\alpha + \beta h \sum_{link} \sigma^\alpha\right),$$
(59)

where α is the replica index which takes $\alpha = 1, \dots, n$, and we shall take the limit $n \to 0$ in the final stage of the calculation. Because of the replica tricks, integration over J_p can be done for each p,

$$\int dJ_p e^{-\frac{1}{2J^2}(J_p - J_0)^2} e^{\beta J_p \sum_{\alpha=1}^n \prod_p \sigma_\alpha} = e^{\frac{1}{2}\beta^2 J^2 \sum_{\alpha,\beta} \prod_p \sigma^\alpha \prod_p \sigma^\beta + J_0 \beta \sum_\alpha \prod_p \sigma^\alpha}$$
(60)

We introduce the MF $U_{0\alpha}$ for σ_{xi}^{α} and the glass MF(GMF) $Q_{\alpha\beta}$ for $\sigma_{xi}^{\alpha}\sigma_{xi}^{\beta}$. Then the terms in the action (60) can be rewritten as follows as in the MFT for the nonrandom case (53),

$$\sum_{pl} \prod_{p} \sigma^{\alpha} \to -3N_{P}U_{0\alpha}^{4} + 4CU_{0\alpha}^{3} \sum_{link} \sigma_{xi}^{\alpha},$$
$$\sum_{pl} \prod_{p} \sigma^{\alpha} \prod_{p} \sigma^{\beta} \to -3N_{P}Q_{\alpha\beta}^{4} + 4CQ_{\alpha\beta}^{3} \sum_{link} \sigma_{xi}^{\alpha} \sigma_{xi}^{\beta},$$
(61)

where we have put $C = \frac{d-1}{2}$.

From Eqs.(59), (60) and (61),

$$[Z^n] = \exp\left(-3\beta^2 J^2 N_P \sum_{\alpha < \beta} Q^4_{\alpha\beta} - 3J_0\beta N_P \sum_{\alpha} U^4_{0\alpha} + N_L \log \operatorname{Tr} e^L\right),\tag{62}$$

where

$$L = 4\beta^2 J^2 C \sum_{\alpha < \beta} Q^3_{\alpha\beta} \sigma^{\alpha}_{xi} \sigma^{\beta}_{xi} + \beta \sum_{\alpha} (4J_0 C U^3_{0\alpha} + h) \sigma^{\alpha}_{xi}.$$
 (63)

We assume a replica symmetric (RS) solution for $U_{0\alpha} = U_0$ and $Q_{\alpha\beta} = Q$. In the RS case, log Tr e^L can be evaluated as follows,

$$\log \operatorname{Tr} e^{L} = \log \operatorname{Tr} \sqrt{\frac{4\beta^{2}J^{2}CQ^{3}}{2\pi}} \int_{-\infty}^{\infty} dz \exp\left(-\frac{4\beta^{2}J^{2}CQ^{3}}{2}z^{2} + 4\beta^{2}J^{2}CQ^{3}z \sum_{\alpha} \sigma_{xi}^{\alpha} -2\beta^{2}J^{2}CnQ^{3} + \beta(4J_{0}CU_{0}^{3} + h) \sum_{\alpha} \sigma_{xi}^{\alpha}\right)$$
$$= \log\left(1 + n \int_{-\infty}^{\infty} Dz \log(2\cosh\beta\tilde{H}(z)) - 2n\beta^{2}J^{2}CQ^{3} + O(n^{2})\right), \quad (64)$$

where

$$Dz = dz \; \frac{e^{-\frac{z^2}{2}}}{\sqrt{2\pi}}, \quad \tilde{H}(z) = 2J\sqrt{CQ^3}z + 4J_0CU_0^3 + h. \tag{65}$$

From (64), the free energy F_R is evaluated as

$$-\beta F_R = \lim_{n \to 0} \frac{[Z^n] - 1}{n} \\
= N_P \left(-\frac{3\beta^2 J^2}{2} (n-1)Q^4 - 3J_0\beta U_0^4 + \frac{1}{Cn}\log \operatorname{Tr} e^L \right) \\
= N_P \left(\frac{3}{2}\beta^2 J^2 Q^4 - 3J_0\beta U_0^4 + \frac{1}{C}\int Dz \,\log(2\cosh\beta\tilde{H}(z)) - 2\beta^2 J^2 Q^3 \right). \quad (66)$$

The values of MF's U_0 and Q are determined by the stationary condition of F_R ,

$$\frac{\partial F_R}{\partial U_0} = 0, \quad \frac{\partial F_R}{\partial Q} = 0.$$
 (67)

Numerical calculation is necessary for solving Eq.(67), and the result is given in Fig.5.

Let us explain physical meanings of the "order parameters" $U_{0\alpha}$ and $Q_{\alpha\beta}$. In the ordinary gauge theories with constant gauge coupling, the confinement and deconfinement phases are distinguished by the expectation value of the Wilson loop operator W(C),

$$W(C) = \prod_{(xi)\in C} \sigma_{xi},\tag{68}$$

where C is a large closed loop on the original lattice. If the system is in the confinement phase, $\langle W(C) \rangle \propto e^{-\operatorname{Area}(C)}$, whereas in the deconfinement phase, $\langle W(C) \rangle \propto e^{-\operatorname{Perimeter}(C)}$.

In the RGT, on the other hand, the ensemble average must be taken in order to obtain physical quantities. Then order parameter is given by $[\langle W(C) \rangle]$. Nonvanishing of the MF $U_{0\alpha}$ means the perimeter law $[\langle W(C) \rangle] \propto e^{-\text{Perimeter}(C)}$, which indicates that the system is in the deconfinement phase, the Higgs phase in the present case. As shown in Fig.5, the RMFT predicts that the Higgs phase exists in the RGT if the fluctuation of the RGC J_p is not so large[15]. Result in Fig.5 also indicates the existence of a "gauge glass" phase[4]. In this phase, $U_{0\alpha} = 0$ whereas $Q_{\alpha\beta} \neq 0$. This

means $[\langle W(C) \rangle] \propto e^{-\operatorname{Area}(C)}$ whereas $[\langle W(C) \rangle^2] \propto e^{-\operatorname{Perimeter}(C)}$. This prediction of the gauge glass is itself interesting but the spatial dimension must be probably large for its realization.⁴

Reliability of the RMFT can be studied as in the usual spin glass models like the Sherrington-Kirkpatrick model. Parisi-type solutions for replica-symmetry breaking are also interesting. These problems are under study and results will be reported in a future publication.

6 Conclusion

In this paper we explicitly showed the relationship between Z_N gauge theories and Kitaev's model for quantum memory and computations. Z_N gauge theories appear as a result of the spontaneous breakdown of the U(1) gauge theory with the "Higgs field" of charge N. The Higgs-phase limit of the Z_N gauge systems corresponds to Kitaev's model. Stability of Kitaev's model was discussed and it was shown that the errors or noise represented by the term like $\sum X_{xi}$ induce the phase transition to the confinement phase in the gauge-theory terminology. In that phase, quantum memory and quantum computations are impossible. Then we studied effects of the RGC which are also regarded as noise and errors in quantum computations. Static RGC gives no significant effect on the phase structure whereas time-dependent RGC induces phase transitions including that to the gauge-glass phase. Application of the present studies to non-Abelian discrete gauge theory is interesting and important for quantum computations.

Acknowledgement

We acknowledge helpful discussions with T.Ohno and T.Matsui.

References

- A.Yu.Kitaev, "Quantum error correction with imperfect gates", in *Proceedings of the Third International Conference on Quantum Communication and Measurement*, ed. O.Hirota, A.S. Holevo, and C.M. Caves (New York, Plenum, 1997); "Fault-tolerant quantum computation by anyons", quant-ph/9707021; Annals Phys.303, 2(2003); M.Freedman, A.Kitaev, M.Larsen, and Z.Wang, "Topological quantum computation", quant-ph/0101025.
- [2] J.Preskill, "Fault-tolerant quantum computation", quant-ph/9712048;

⁴In this paper we are considering systems only on the simple square or cubic lattice. If we consider more complicated lattice or networks, effective spatial dimensions increase and the gauge glass phase may appear.

- [3] E.Dennis, A.Kitaev, A.Landahl, and J.Preskill, "Topological quantum memory", quant-ph/0110143; J.Math.Phys.43, 4452 (2002).
- [4] C.Wang and J.Preskill, "Confinement-Higgs transition in a disordered gauge theory and the accuracy threshold for quantum memory", quant-ph/0207088; Annals Phys. 303, 31 (2003).
- [5] C.Mochon, Phys.Rev.A67, 022315(2003).
- [6] M.W.Propitius and F.A.Bais, hep-th/9511201 and references cited therein.
- [7] Phase structure of the N = 2 case was studied both theoretically and numerically. See for example, E.Fradkin and S.H.Shenker, Phys.Rev.D19, 3682 (1979); G.Bhanot and B.A.Freedman, Nucl.Phys.B190, 2191 (1977); A.Sudbø, E.Smørgav, J.Smiseth, F.S.Nogueira, and J.Hove, Phys.Rev.Lett.89, 226403 (2002); J.Smiseth, E.Smørgav, F.S.Nogueira, J.Hove, and A.Sudbø, "Phase structure of d=2+1 compact lattice gauge theories and the transition from Mott insulator to fractionalized insulator", cond-mat/031297.
- [8] See for example, I.Ichinose and T.Ohbayashi, Nucl.Phys.B419[FS], 529 (1994).
- [9] K.Sakakibara, M.Kemuriyama, and T.Matsui, "Gauged neural network: Phase structure, learning and associative memory", cond-mat/0203136.
- [10] J.B.Kogut, Rev.Mod.Phys.51, 659 (1979).
- [11] T.Ohno, G.Arakawa, I.Ichinose, and T.Matsui, paper in preparation.
- [12] Properties of the three-dimensional random Z_2 gauge theory at zero temperature are numerically investigated in Ref.[4].
- [13] J.M.Drouffe, Nucl.Phys.B170[FS1], 211 (1980).
- [14] As it is well-known, Elitzur's theorem forbids the nonvanishing expectation value of the gaugevariant quantities like σ_{xi} . Discussion on the compatibility of the MFT and Elitzur's theorem is given in Ref.[13].
- [15] Recent numerical studies on $3D Z_2$ random gauge theory in Refs.[4] and [11] indicate that the Higgs phase at low temperature disappears quite rapidly as the concentration p of the "wrong-sign" plaquettes increases.




Figure 1: Phase diagram of the gauge-Higgs model




Figure 2: Path $\widetilde{c}_{x^*y^*}$ connecting dual sites x^* and y^*




Figure 3: The links $(\bar{x}i)$ on which $X_{\bar{x}i}^R$ operaters. Dyon is composed of electric charge on site and magnetic vortex on plaquette




Figure 4: Free energy of the Z_2 gauge system by the MFT




Figure 5: Phase diagram of the RGT obtained by the RMFT. The Higgs phase exists at low temperature(T) and small fluctuation of the RGC. Gauge-glass phase appears at low T and large fluctuation of the RGC. In the Higgs phase, the MF's $U_0 \neq 0$ and also $Q \neq 0$, whereas in the gauge-glass phase $U_0 = 0$ and $Q \neq 0$.