## Correlations of observables in chaotic states of macroscopic quantum systems

Ayumu Sugita\*

Department of Applied Physics, Osaka City University 3-3-138 Sugimoto, Sumiyoshi-ku, Osaka, 558-8585, Japan

## Akira Shimizu<sup>†</sup>

Department of Basic Science, University of Tokyo, 3-8-1 Komaba, Tokyo 153-8902, Japan and PRESTO, Japan Science and Technology Corparation, 4-1-8 Honcho, Kawaguchi, Saitama, Japan (Dated: October 30, 2018)

We study correlations of observables in energy eigenstates of chaotic systems of a large size N. We show that the bipartite entanglement of two subsystems is quite strong, whereas macroscopic entanglement of the total system is absent. It is also found that correlations, either quantum or classical, among less than N/2 points are quite small. These results imply that chaotic states are stable. Invariance of these properties under local operations is also shown.

PACS numbers: 03.67.Mn, 05.45.Mt, 03.65.Yz, 05.45.Jn

It is generally believed that almost all macroscopic systems have chaotic dynamics, because otherwise thermodynamics would not hold. However, properties of chaotic quantum states in macroscopic systems are poorly understood as compared with those in systems of small degrees of freedom [1]. Among such properties, correlations of observables at different points are of particular interest. For example, such correlations are directly related to entanglement [2], which is the central subject of quantum information theory [3]. The correlations also define the 'cluster property', which is one of the most fundamental notions of field theory [4]. Moreover, the correlations determine the magnitudes of fluctuations of 'additive operators' (see below), which are macroscopic observables. Furthermore, Shimizu and Miyadera [5] (Hereafter referred to as SM) showed that two-point correlations determine the stabilities against classical noises, perturbations from environments, and local measurements. In particular, they showed that states with 'macroscopic entanglement' are unstable. Since chaotic dynamics is generally believed to promote entanglement production [6], and since classical chaos is characterized by extreme sensitivity to the initial condition, it might be tempting to conjecture that chaotic states would be unstable. However, as we will show, such a naive expectation is wrong. In this work, we study these points for chaotic quantum states in macroscopic systems, i.e., systems with a large but finite degrees of freedom.

Two-point correlations and fluctuations of additive operators: We consider an energy eigenstate  $|\Psi\rangle$  of a quantum chaotic system which is composed of  $N \gg 1$  sites, where the Hilbert space is the tensor product of local Hilbert spaces. Let  $\{\hat{a}_{\alpha}(l)\}$  be a basis of local observables at a site l. Assuming that  $\hat{a}_{\alpha}(l)$ 's are bounded, we normalize them as  $\text{Tr}[\hat{a}_{\alpha}^{\dagger}(l)\hat{a}_{\beta}(l)] = \text{constant} \times \delta_{\alpha\beta}$ .

Then, all information about the two-point correlations in  $|\Psi\rangle$  are included in the variance-covariance matrix (VCM),  $V_{\alpha l,\beta l'} \equiv \langle \Psi | \delta \hat{a}^{\dagger}_{\alpha}(l) \delta \hat{a}_{\beta}(l') | \Psi \rangle$ , where  $\delta \hat{a}_{\alpha}(l) \equiv$  $\hat{a}_{\alpha}(l) - \langle \Psi | \hat{a}_{\alpha}(l) | \Psi \rangle$  [7]. This matrix also gives us information about fluctuations  $\langle \Psi | \delta \hat{A}^{\dagger} \delta \hat{A} | \Psi \rangle$  of 'additive operators'  $\hat{A}$ , which are defined as the sums of local operators;  $\hat{A} = \sum_{\alpha,l} c_{\alpha l} \hat{a}_{\alpha}(l)$ . Here,  $c_{\alpha l}$ 's are c-numbers independent of N. Without loss of generality, we here normalize them as  $\sum_{\alpha,l} |c_{\alpha l}|^2 = N$ . As shown in Ref. [7], the maximum fluctuation of the additive operators is  $Ne_{\max}$ , where  $e_{\max}$  is the maximum eigenvalue of the VCM. For example,  $e_{\text{max}} = O(1)$  [13] for any product state  $|\Psi\rangle = \bigotimes_l |\psi_l\rangle$ , whereas  $e_{\text{max}} = o(N)$  for 'vacuum states' of many-body physics in accordance with thermodynamics [5, 8]. Interestingly, as special states in large but finite systems, there also exist pure states for which  $e_{\text{max}} = O(N)$  [5, 7]. Such pure states are superpositions of macroscopically distinct states, hence are called macroscopically entangled states [5, 7, 9].

Let us evaluate the VCMs of energy eigenstates in chaotic systems using random matrix theory (RMT). Suppose that an energy eigenstate  $|\Psi_{\lambda}\rangle$ , labeled by a single parameter  $\lambda$ , is represented as  $|\Psi_{\lambda}\rangle = \sum_{i} c_{\lambda i} |i\rangle$ in some basis  $\{|i\rangle\}$ . The ensemble averages, denoted by overline, of products of the coefficients can be calculated easily using RMT [10]. For example, in the  $d \times d$  GUE or GOE,

$$\overline{|c_{\lambda i}|^2} = \frac{1}{d},\tag{1}$$

$$\overline{|c_{\lambda i}|^4} = \frac{2+q}{d(d+1+q)},\tag{2}$$

$$\frac{|c_{\lambda i}|^2}{|c_{\lambda i}|^4} = \frac{1}{d}, \tag{1}$$

$$\frac{|c_{\lambda i}|^4}{|c_{\lambda i}|^2} = \frac{2+q}{d(d+1+q)}, \tag{2}$$

$$\frac{1}{|c_{\lambda i}|^2 |c_{\lambda i'}|^2} = \frac{1}{d(d+1+q)} \text{ for } i \neq i', \tag{3}$$

where q = 0 and 1 for GUE and GOE, respectively. Up to the 4-th order, the other combinations which are not written above vanish identically. By using these formulae, we can evaluate the averages and variances of the matrix elements of VCMs. Since the right-hand sides of

<sup>\*</sup>Electronic address: sugita@a-phys.eng.osaka-cu.ac.jp

<sup>†</sup>Electronic address: shmz@ASone.c.u-tokyo.ac.jp

Eqs. (1)-(3) are independent of  $\lambda$ , we will hereafter drop the label  $\lambda$ . Furthermore, we will denote  $\overline{\langle \Psi_{\lambda}| \cdot |\Psi_{\lambda}\rangle}$  simply by  $\overline{\langle \cdot \rangle}$ .

We here present results for the case where each site is a qubit, which is equivalent to a spin- $\frac{1}{2}$  system. In this case, we can use the Pauli matrices  $\{\sigma_x,\sigma_y,\sigma_z\}$  as a basis of local observables [14]. Therefore, the VCMs of an N-qubit system are  $3N\times 3N$  matrices. It is easy to show that  $\overline{\langle\sigma_\alpha(l)\rangle}=0$  and  $\overline{\langle\sigma_\alpha(l)\rangle\langle\sigma_\beta(l')\rangle}=0$  for  $l\neq l'$   $(\alpha,\beta=x,y,z).$  With the help of these, the average values of the elements of the VCM are calculated as

$$\overline{\langle \delta \sigma_{\alpha}(l) \delta \sigma_{\beta}(l') \rangle} = \frac{2^{N}}{2^{N} + 1 + q} \delta_{\alpha\beta} \delta_{ll'}. \tag{4}$$

This suggests that the VCM converges to the unit matrix I as  $N \to \infty$ , hence  $e_{\max} = O(1)$ . To examine this point, we plot in Fig. 1 the maximum and the minimum eigenvalues  $e_{\max}$  and  $e_{\min}$ , of the VCMs of energy eigenstates of GUE and GOE as functions of N. It is found that both  $e_{\max}$  and  $e_{\min}$  (hence all eigenvalues) concentrate on 1 as N increases, and thus the VCMs approach I. This means that fluctuations of all additive operators are O(N) or less, and that all two-point correlations for  $l \neq l'$  are vanishingly small. In fact, for  $l \neq l'$  we can directly show that

$$\overline{|\langle \sigma_{\alpha}(l)\sigma_{\beta}(l')\rangle|^2} = \frac{1+q}{2^N+1+q},\tag{5}$$

hence  $|\langle \delta \sigma_{\alpha}(l) \delta \sigma_{\beta}(l') \rangle|^2$  is also  $O(2^{-N})$ . Therefore, either entanglement or classical correlation will be found to be exponentially small if one observes arbitrary two points of a chaotic state. Thus, in the sense of SM, chaotic states have the 'cluster property' [15], and are not entangled macroscopically. According to the general results of SM, these imply that chaotic states are stable under local measurements, and that the decoherence rate  $\Gamma$  of chaotic states is O(N) or less under weak perturbations from any noises or environments which have short time correlations. This is in sharp contrast to macroscopically entangled states for which some noise gives  $\Gamma = O(N^2)$  [5]. These points will be discussed later again.

Correlations among many points: We have seen that two-point correlations are infinitesimal. How about m-point correlations for larger m? We can estimate it from bipartite entanglement between two subsystems as follows.

Let us separate the system into two subsystems A and B which contain  $N_{\rm A}$  and  $N_{\rm B}=N-N_{\rm A}$  sites, respectively. The Hilbert space of subsystem A (B) has dimension  $d_{\rm A}=2^{N_{\rm A}}$  ( $d_{\rm B}=2^{N_{\rm B}}$ ). We assume that  $N_{\rm A}\leq N_{\rm B}$ , and evaluate 'purity'  ${\rm Tr}\left(\hat{\rho}_{\rm A}^2\right)$  as a measure of bipartite entanglement, where  $\hat{\rho}_{\rm A}$  is the reduced density operator of A;  $\hat{\rho}_{\rm A}\equiv {\rm Tr}_{\rm B}\left(|\Psi\rangle\langle\Psi|\right)$ . [16] The purity takes the maximum value 1 when the state  $|\Psi\rangle$  is separable (unentangled), and the minimum value  $1/d_{\rm A}$  when  $\hat{\rho}_{\rm A}$  is a scalar matrix  $I/d_{\rm A}$ . For GUE and GOE, Eqs. (1)-(3) yield the




FIG. 1: The maximum and the minimum eigenvalues of the VCMs in spin- $\frac{1}{2}$  systems as functions of the system size N. The solid lines represent the results for eigenstates of GUE. The results from GOE, which are not displayed here, are very similar to those of GUE. The dotted lines are the results for the the central  $(2^{N-1}$ -th) eigenstates of the hamiltonian (8). In both cases the averages have been taken over 100 samples. The error bars show the standard deviation.

average purity as [11]

$$\overline{\text{Tr}(\hat{\rho}_{A}^{2})} = \frac{d_{A} + d_{B} + q}{d_{A}d_{B} + 1 + q} = \frac{1}{d_{A}} \left( 1 + \frac{1}{2^{\Delta N}} + \cdots \right), \quad (6)$$

where  $\Delta N \equiv N_{\rm B} - N_{\rm A}~(\geq 0)$ , and '…' denotes smaller terms. It is found that the average purity is quite small, which means that the bipartite entanglement of two subsystems are strong. With increasing  $\Delta N$ , in particular, the average purity approaches exponentially the minimum value  $1/d_{\rm A}$ . Therefore  $\hat{\rho}_{\rm A}$  converges to  $I/d_{\rm A}$  exponentially with increasing  $\Delta N$ , for almost all states. This is demonstrated for N=12 in Fig. 2, in which  $\hat{\rho}_m=\hat{\rho}_{\rm A}$  and  $m=N_{\rm A}$ . It is noteworthy that the bipartite entanglement is large but not maximum when  $N_{\rm A}=N_{\rm B}$ .

Suppose now that observables  $\hat{a}_1(l_1), \hat{a}_2(l_2), \ldots, \hat{a}_m(l_m)$  at m different points (sites)  $l_1, l_2, \cdots, l_m$  are measured. Since these observables commute with each other, their correlations can be calculated in a manner similar to the case of classical stochastic variables. In particular, the cumulants are given by

$$\langle \hat{a}_1(l_1) \dots \hat{a}_m(l_m) \rangle_c = (-1)^m \frac{\partial^m \ln Z}{\partial J_1 \dots \partial J_m},$$
 (7)

where  $Z(\{J_i\}) \equiv \langle \Psi | \exp\left[-\sum_i J_i \hat{a}_i(l_i)\right] | \Psi \rangle$  is the generator of moments. If we regard the set of m sites  $l_1, l_2, \cdots, l_m$  as subsystem A, we have  $Z(\{J_i\}) = \operatorname{Tr}\left(\hat{\rho}_m \exp\left[-\sum_{i=1}^m J_i \hat{a}_i(l_i)\right]\right)$ , where  $\hat{\rho}_m = \hat{\rho}_{\mathrm{A}}$ . When  $4^{m-N/2} \ll 1$ , i.e., when  $2^m \ll 2^{N-m}$ ,  $\rho_m = I/2^m$ 




FIG. 2:  $-\log_2 \overline{\text{Tr}(\hat{\rho}_m^2)}$  of a subsystem composed of m sites is plotted for N=12. The solid line denotes the upper limit, corresponding to the lower limit of  $\overline{\text{Tr}(\hat{\rho}_m^2)}$ , for each m. The dotted and dashed lines, which look overlapped each other in this figure, represent results for GUE and GOE, respectively. The long dashed line shows the result obtained from the central  $(2^{N-1}$ -th) eigenvector of the hamiltonian (8).

to a good approximation, according to our results above. In this case, we obtain  $\langle \hat{a}_1(l_1) \dots \hat{a}_m(l_m) \rangle_c = 0$ , which means that there is no correlation among  $a_1(l_1), \dots, a_m(l_m)$ . Since they are arbitrary observables at m points, it is concluded that neither entanglement nor classical correlation can be found if  $4^{m-N/2} \ll 1$ , i.e., if the number m of points one performs measurements is somewhat smaller than half the total number of points.

On the other hand, if observables at all points are measured, one can detect all types of entanglement. For example, one can detect the bipartite entanglement by measuring a correlation of the Clauser-Horne-Shimony-Holt (CHSH) type [12]. This correlation becomes almost maximum for chaotic states, because we have already shown that the bipartite entanglement between subsystems A and B is almost maximal (when  $2^{\Delta N} \gg 1$ ). However, the enhancement of the CHSH correlation over that of separable states (or of local classical theories) is at most by a factor of  $\sqrt{2}$ . Recently, another correlation of local observables was found in Ref. [9], which detects macroscopic entanglement in general states (either pure or mixed): It becomes as large as  $O(N^2)$  for macroscopically entangled states, whereas it is at most O(N) for states with  $e_{\text{max}} = O(1)$  (and for any mixtures of such states). For chaotic states, this correlation is at most O(N) because we have shown that  $e_{\text{max}} = O(1)$ .

A model with short-range interactions: We have assumed RMT in the above analysis. One might suspect that RMT would not be fully applicable to macroscopic systems, in which interactions are short-ranged, because

RMT assumes that strengths of interactions are of the same order between any two points. To explore this point, we study eigenstates of the following hamiltonian:

$$H = J \sum_{l=1}^{N} \left\{ \sigma_x(l) \sigma_x(l+1) + \sigma_z(l) \sigma_z(l+1) + \sqrt{2} \cos \phi_l \, \sigma_y(l) \sigma_y(l+1) \right\}$$
$$-h \sum_{l=1}^{N} \left\{ \sin \theta_l \, \sigma_x(l) + \cos \theta_l \, \sigma_z(l) \right\}. \tag{8}$$

Here, J and h are constants,  $\{\phi_l\}$  and  $\{\theta_l\}$  are random variables with  $0 \le \phi_l, \theta_l < 2\pi$ , and  $\sigma_\alpha(N+1) = \sigma_\alpha(1)$ . This hamiltonian describes a one-dimensional spin system in which spins interact only with their nearest neighbors, where the interaction in the y-direction is random. Besides, there is an external magnetic field with strength h, whose direction is random in the x-z plane. When J and h are sufficiently large (say, J=h=1), this system becomes chaotic, except for states around the lower and upper limits of the energy spectrum, in the sense that the level spacing distribution coincides with that of GOE.

The dotted lines in Fig. 1 represent the maximum and the minimum eigenvalues of the VCM for the  $2^{N-1}$ -th eigenstate, which is located at the center of the spectrum, where the chaotic nature appears most clearly. It is seen that the results agree fairly with those of RMT. Furthermore, the long dashed lines in Fig. 2 represent  $-\log_2 \overline{\text{Tr}(\hat{\rho}_m^2)}$  for the same state. The results are close to those for RMT, especially when m is small. Similar results are obtained also for other eigenstates except those near the upper and the lower ends of the spectrum. Since the density of states has a dominant peak at the center of the spectrum, our conclusions from RMT hold for the great majority of the eigenstates of the Hamiltonian (8).[17] We thus see that RMT correctly describes correlations in energy eigenstates of the chaotic system with short-range interactions.

Invariance under local operations: What happens if local operations are performed on a chaotic state? Let  $\{|i_l\rangle\}$  be a basis of the local Hilbert space at site l. An energy eigenstate  $|\Psi^{(N)}\rangle$  of an N-site system can be expanded as

$$|\Psi^{(N)}\rangle = \sum_{i_1} \sum_{i_2} \cdots \sum_{i_N} c_{i_1 i_2 \cdots i_N}^{(N)} |i_1\rangle |i_2\rangle \cdots |i_N\rangle.$$
 (9)

For GUE, the probability density of the  $2^N$  coefficients  $c_{i_1i_2\cdots i_N}^{(N)}$  is given by [10]

$$\frac{(2^N - 1)!}{\pi^{2^N}} \delta \left( \sum_{i_1} \cdots \sum_{i_N} |c_{i_1 \cdots i_N}^{(N)}|^2 - 1 \right). \tag{10}$$

Since this density is invariant under changes of the local basis  $\{|i_l\rangle\}$  for any l, the statistical properties of  $|\Psi^{(N)}\rangle$  are not changed by local unitary transformations. Furthermore, when a local projective measure-

ment that diagonalizes  $\{|i_N\rangle\}$  is performed on the Nth qubit, the post-measurement state (for each measurement) is given by  $|i_N\rangle\sum_{i_1}\cdots\sum_{i_{N'}}c_{i_1\cdots i_{N'}}^{(N')}|i_1\rangle\cdots|i_{N'}\rangle\equiv |i_N\rangle|\Psi^{(N')}\rangle$ , where  $N'\equiv N-1$ , and  $c_{i_1\cdots i_{N'}}^{(N')}\equiv c_{i_1\cdots i_{N'}i_N}^{(N)}/(\sum_{i_1}\cdots\sum_{i_{N'}}|c_{i_1\cdots i_{N'}i_N}^{(N)}|^2)^{1/2}$ . It is easy to show that  $\{c_{i_1\cdots i_{N'}}^{(N')}\}$  also obeys the probability distribution of CULE. bution of GUE, i.e., Eq. (10) with N being replaced with N'. Therefore, all the results for  $|\Psi^{(N)}\rangle$  hold for  $|\Psi^{(N')}\rangle$  as well, with N being replaced with N' [18]. The same can be said when a local projective measurement that diagonalizes another local basis  $\{|j_N\rangle\}$  is performed on the N-th qubit, because the density (10) is invariant under changes of the local basis. The same conclusions can be derived for GOE as well. We therefore conclude that properties of chaotic states which we have found in this work are invariant under 'local operations, 'including local unitary transformations and local projective measurements diagonalizing a local basis. This implies, for example, that a chaotic state cannot be disentangled by local measurements at less than Npoints. This is in sharp contrast to, e.g., a 'cat state,'  $|C\rangle \equiv |\uparrow\uparrow\cdots\uparrow\rangle/\sqrt{2} + |\downarrow\downarrow\cdots\downarrow\rangle/\sqrt{2}$ , for which a single local measurement suffices to disentangle the state [5]. This is consistent with the conclusion, which we have derived above using the theorem by SM, that chaotic states are stable under local measurements because two-point correlations are infinitesimal.

Discussions: If one defines the 'degree' (or 'strength')  $\mathcal{E}$  of entanglement by the minimum number of local operations that are necessary to disentangle it, then, according to our results,  $\mathcal{E}$  of chaotic states in large systems is quite high. However, this does not mean that they are anomalous as macroscopic states. Indeed, we have shown that they are stable against local measurements and that fluctuations of all additive operators are O(N) or less, in accordance with thermodynamics [5, 7, 8]. Furthermore, as we have mentioned above, the correlation of Ref. [9],

which detects macroscopic entanglement, is quite small for chaotic states. Moreover, in the infinite-size limit, all multi-point correlations vanish since the number m of measured points is always finite (hence  $m \ll N/2$  as  $N \to \infty$ ). Although the absence of quantum correlations for finite m at  $N \to \infty$  has been generally proved [9], we have found here that chaotic states do not even have classical correlations in the same limit. Therefore, correlations, either quantum or classical, of chaotic states are neither detectable nor usable in infinite systems.

N-dependences: In this work, we have often discussed N dependences of various quantities. Unlike a uniform state in a uniform system, however, correspondence between states in systems of different sizes is non-trivial in chaotic systems. For completeness, we finally describe what the N dependences mean in this paper.

For each system size N, consider a set S of all energy eigenstates whose energies are in an interval (E - $\Delta/2, E+\Delta/2$ ]. To see the N dependence of some quantity Q, look at its (probability) distribution  $P_N(Q)$  in S. Then, the N dependence of Q can be discussed in terms of the N dependence of  $P_N(Q)$ . For example, we say  $Q \leq Q_0$  for almost all states in S for sufficiently large N' if for arbitrary positive number  $\epsilon$  there exists  $N_0$  such that  $1 - P_N(Q \leq Q_0) \leq \epsilon$  for  $N \geq N_0$ . We can thus discuss N dependences of various quantities for chaotic states. In particular, we can apply the results of SM, in which N dependences play crucial roles. Note that  $P_N(Q)$  is independent of E and  $\Delta$  in the case of RMT. In the short-range interaction model given by Eq. (8), the dependence of  $P_N(Q)$  on E and  $\Delta$  is quite weak if we confine ourselves to states around the center of the energy spectrum. In either case, we can discuss the Ndependence without specifying E and  $\Delta$ .

Acknowledgment We thank H. Matsuful for discussions. A. Sugita was supported by Japan Society for the Promotion of Science for Young Scientists.

M. C. Gutzwiller, Chaos in Classical and Quantum Mechanics (Springer, New York, 1990).

<sup>[2]</sup> A. Sugita, J. Phys. A: Math. Gen. **36**, 9081 (2003).

<sup>[3]</sup> B. M. Terhal, M. M. Wolf and A. C. Doherty, Physics Today, April 2003, 46, and references cited therein.

<sup>[4]</sup> R. Haag, Local Quantum Physics (Springer, Berlin, 1992).

<sup>[5]</sup> A. Shimizu and T. Miyadera, Phys. Rev. Lett. 89, 270403 (2002).

<sup>[6]</sup> See, for example, P. A. Miller and Sarben Sarkar, Phys. Rev. E 60 1542 (1999).

<sup>[7]</sup> T. Morimae, A. Sugita and A. Shimizu, Phys. Rev. A 71 032317 (2005).

<sup>[8]</sup> D. Ruelle, Statistical Mechanics: Rigorous Results (Benjamin, Reading, 1969).

<sup>[9]</sup> A. Shimizu and T. Morimae, preprint.

<sup>[10]</sup> N. Ullah, Nucl. Phys. 58, 65 (1964).

<sup>[11]</sup> The result for GUE, which is a property of random vectors in Hilbert space, has been reported several times in the literature. See, for example, V M. Kendon, K. Życzkowski and W. J. Munro, Phys. Rev. A 66 062310 (2002) and references therein.

<sup>[12]</sup> J. F. Clauser et al., Phys. Rev. Lett. 23, 880 (1969).

<sup>[13]</sup> Throughout this paper, we say that f(N) = O(g(N)) if  $\lim_{N\to\infty} f(N)/g(N) = \text{const } \neq 0$ , and that f(N) = o(g(N)) if  $\lim_{N\to\infty} f(N)/g(N) = 0$ .

<sup>[14]</sup> Actually we need four matrices  $\sigma_x, \sigma_y, \sigma_z$  and I to represent all  $2 \times 2$  matrices as their linear combinations. However, we don't have to consider I because all correlations including it vanish.

<sup>[15]</sup> The cluster property of SM is defined by two-point correlations, whereas the one of Ref. [4] is defined by multipoint correlations. Clearly, a state having the latter cluster property has the former one as well.

- [16] For many-body states, there are many sorts of entanglement and, accordingly, many measures or indices of entanglement. The simplest entanglement is bipartite entanglement, in which a many-body system is regarded as a two-body system composed of two subsystems of the many-body system. For pure states, the von Neumann entropy of the reduced density operator is the unique measure of the bipartite entanglement, whereas the pu-
- rity is a simplified version of it.
- [17] Systematic analysis of eigenstates near the ends of the spectrum will be reported elsewhere.
- [18]  $|\hat{\Psi}^{(N)}\rangle$  is an energy eigenstate of the N-site system, whereas  $|\Psi^{(N')}\rangle$  is not necessarily an energy eigenstate of the N'-site system. This difference does not make any difference in the statistical properties.