## Information Erasure and Recover in Quantum Memory\*

Qing-Yu Cai<sup>†</sup>

Wuhan Institute of physics and mathematics, The chinese Academy of Sciences, Wuhan. 430071. China

We show that information in quantum memory can be erased and recovered perfectly if it is necessary. That the final states of environment are completely determined by the initial states of the system allows that an easure operation can be realized by a swap operation between system and an ancilla. Therefore, the erased information can be recoverd. When there is an irreversible process, e.g. an irreversible operation or a decoherence process, in the erasure process, the information would be erased perpetually. We present that quantum erasure will also give heat dissipation in environment. And a classical limit of quantum erasure is given which coincides with Landauer's erasure principle.

PACS: 03.65.Bz, 03.67.Hk

The formalism of quantum mechanics allows the vast information content of a quantum state to be efficiently processed, at a rate that cannot be matched in real time by any classical means. Recently, a great deal of attention has been focused on quantum computation. Quantum computation is a field of fundamental interest because we believe quantum information processing machines can actually realized in nature. The elementary units of the theory are quantum qubits. When memory in a quantum computer is scarce (since any real physical resource is finite), erasure may plays an important role in its own way since the quantum no-deleting principle states that the linearity of quantum theory does not allow us to delete a copy of an arbitrary quantum state perfectly. As is well known, information of an unknown classical bit can be erased perfectly. In this paper, we will show that information in quantum memory can be erased and recovered perfectly if it is necessary. And the information will be erased perpetually under an irreversible operation. In this case, information can not be recovered. And the heat dissipation is inevitable.

In quantum theory, erasure an quantum state can be described, for an arbitrary state  $|\varphi\rangle$ , [10]

$$\varepsilon |\varphi\rangle \to |0\rangle,$$
 (1)

where  $\varepsilon$  is an erasure operation and  $|0\rangle$  is the standard state. If the state is known, then one can prepare it to a standard state by using a unitary operation. Consider a big number of qubits to be erased of a quantum memory in quantum computer. These qubits should be treated as in the unknown states. Suppose  $\varepsilon$  is unitary and the state is unknown. For two arbitrary states  $|\varphi_0\rangle$  and  $|\varphi_1\rangle$ , we have

$$\varepsilon |\varphi_0\rangle \to |0\rangle,$$
 (2)

$$\varepsilon |\varphi_1\rangle \to |0\rangle$$
. (3)

Since unitary transformations preserve the inner products, it must be that

$$\langle \varphi_0 | \varphi_1 \rangle = \langle 0 | 0 \rangle. \tag{4}$$

Clearly,  $\varepsilon$  is nonunitary. The quantum mechanics postulate states that the evolution of a closed quantum system is described by a unitary transformation. Since  $\varepsilon$  is nonunitary, then an erasure process requires the system open. Without loss of generality, we assume that the erasure process is to unitarily interact the state ( $|\varphi_0\rangle$  or  $|\varphi_1\rangle$ ) with an ancilla  $|e\rangle$ , [11] in the two case one obtaining

$$U|\varphi_0>|e>\to |0>|e_0>, \tag{5}$$

$$U|\varphi_1>|e>\to |0>|e_1>. \tag{6}$$

Taking inner product gives that

$$\langle \varphi_0 | \varphi_1 \rangle = \langle e_0 | e_1 \rangle. \tag{7}$$

<sup>\*</sup>Supported by the National Natural Science Foundation of China under Grant No 10304022.

<sup>†</sup>qycai@wipm.ac.cn

Because  $|\varphi_0\rangle$  and  $|\varphi_1\rangle$  are arbitrary states, then the final states of environment are completely determined by the initial states of system. Since the erasure process can be realized by using a unitary operation U together with ancilla, it seems that an erasure process is reversible. That is the case. But if one performs an irreversible operation, e.g. measurement operation, on the ancilla, the erasure process becomes irreversible immediately.

Since the final states of system is completely determined by the initial states of system, then the operation U can be consider as a swap transformation  $U_S$  if ancilla is in the standard state |0>,

$$U_S|\varphi>|0>\to|0>|\varphi>. \tag{8}$$

When one needs to use the erased information, he can using a swap operation to recover it,

$$U_S|0>|\varphi>\to|\varphi>|0>. \tag{9}$$

However, in a physical erasure process, real environment is a heat reservoir. And the final states of ancilla will be destroyed under the decoherence, which leads this erasure process to irreversibility. That is, a perpetual erasure operation can be realized by a swap transformation together with an irreversible process that may be naturally completed by the decoherence of environment. Consider the case of a single qubit in an unknown state  $|\varphi>=\alpha|0>$  $+\beta|1>$ , where  $\alpha$ ,  $\beta$  are arbitrary complex numbers and  $|\alpha|^2+|\beta|^2=1$ . Suppose that ancilla is in the standard state |0>. Equation (8) tells us that a swap operation

$$U_S = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \tag{10}$$

can help the erasure process to be completed. Equation (8) can be rewritten as

$$\begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} \alpha \\ \beta \end{pmatrix} \otimes \begin{pmatrix} 1 \\ 0 \end{pmatrix} \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \end{pmatrix} \otimes \begin{pmatrix} \alpha \\ \beta \end{pmatrix}. \tag{11}$$

Then the qubit is in a standard state after the swap transformation. The ancilla is in the unknown state  $|\varphi\rangle = \alpha|0\rangle$  $+\beta$ 1 >. Clearly this is an reversible process. However, when we perform a measurement on the ancilla in the basis  $\{|0>, |1>\}$ , the state  $|\varphi>=\alpha|0>+\beta|1>$  immediately collapses to a known state |0> with a probability  $|\alpha|^2$  or |1> with a probability  $|\beta|^2$ . The information of the state  $|\varphi>=\alpha|0>+\beta|1>$  can not be recovered. A perpetual erasure process is realized.

In fact, an erasure operation can be realized by an irreversible operation followed with a unitary operation. One can perform a measurement operation to prepare the memory in a known state. Then the quantum memory can be prepared in the standard state  $|0\rangle$  with a unitary operation. On this occasion, information can not be recovered once it has been erased. Since the quantum memory is open in the erasure process, there is energy exchange between system and environment. In microworld, the minimal energy exchange to perform a logical operation is of order  $\hbar\omega$ . [12] We can reasonably assume that energy exchange between system and environment when erasure one qubit is at least of order  $\hbar\omega$ . Suppose one qubit is a two-level system which has one degree of freedom. Using theorem of equipartition of energy, the average energy exchange when erasure one qubit can be calculated

$$\bar{\varepsilon} = \frac{\sum_{n=0}^{\infty} n\varepsilon_0 e^{-n\varepsilon_0/k_B T}}{\sum_{n=0}^{\infty} e^{-n\varepsilon_0/k_B T}}$$

$$= \frac{\varepsilon_0}{e^{\varepsilon_0/k_B T} - 1},$$
(12)

$$=\frac{\varepsilon_0}{e^{\varepsilon_0/k_BT}-1},\tag{13}$$

where  $\varepsilon_0 = \hbar \omega$ . That is to say, when erasure one qubit in the quantum memory, on average, there is  $\bar{\varepsilon}$  heat dissipation in environment.

Consider the classical limit of the information erasure and recover in quantum memory. Bennett has shown that all the logical irreversible operations can be reconstructed by logical reversible operations. [13] Then the erased information can be recovered in classical computer. From equation (13), when we accept that  $\hbar \to 0$  in classical limit, perpetually erasure one bit in classical computer will give heat dissipation of order

$$\bar{\varepsilon} = k_B T,$$
 (14)

which coincides with Landauer's erasure principle.

In summary, we show that information erasure can be realized by using a unitary operation together with an ancilla. The final states of ancilla is completely determined by the initial states of the quantum memory. The information can be recovered if it is necessary. When an irreversible operation was performed, the erased information can not be recovered forever. And this perpetual erasure operation is corresponding to a heat dissipation in environment.

## I. REFERENCES

- [1] Deutsch D and Jozsa R 1992, Proc. R. Soc. Lond. A 439, 553
- [2] Shor P W 1994, Proceedings of the 35th annual symposium on the foundation of the computer science, IEEE Computer Society Press, Los Alamitos, CA, p124
  - [3] Grover L K 1997, Phys. Rev. Lett. 79 325
  - [4] Pati A. K. and Braunstein 2000 Nature 404 164
  - [5] Feng S, Zheng S and Yim M 2002 Phys. Rev. A 65 042324
  - [6] Qiu D 2002 Phys. Rev A 65 052303
  - [7] Feng J, Gao Y F, Wang J S and Zhan M S 2002 Phys. Rev A 65 052311
  - [8] Landauer R 1961 IBM J Res. Dev. 5 183
  - [9] Bennett C. H. 1982 Int. J. Theor. Phys. 21 905
  - [10] Piechocinsha B 2000 Phys. Rev. A 61 062314
- [11] Nielsen M A and Chuang I L 2000 Quantum Computation and Quantum Information (Cambridge University Press)
  - [12] Gea-Banacloche J 2002 Phys. Rev Lett. 89 217901
  - [13] Bennett C H 1973 IBM J. Res. Dev. 17 525