A New Proof of The Strong Subadditivity Theorem

Yong-Jian Han, Yong-Sheng Zhang*, Guang-Can Guo[†]
Key Laboratory of Quantum Information, University of Science and Technology
of China, CAS, Hefei 230026, People's Republic of China

It is well known that the strong subadditivity theorem is hold for classical system, but it is very difficult to prove that it is hold for quantum system. The first proof of this theorem is due to Lieb by using the Lieb's theorem. Here we use the conditions obtained in our previous work of matrix analysis method to give a new proof of this famous theorem. This new proof is very elementary, it only needs to carefully analyse the minimal value of a function. This proof also shows that the conditions obtained in our previous work are stronger than the strong subadditivity theorem.

PACS number(s): 03.67.-a, 02.10.Yn, 89.70.+c

I. INTRODUCTION

Entropy is an important concept not only for physics but also for information science. From the definition of entropy, we can get some simple properties of it, such as concavity, continuity property, additivity and subadditivity [1]. But some other properties is not so obvious, such as the strong subadditivity theorem (SSA). Among all of the properties of entropy, the most famous one is the SSA, and it is very difficult to prove this theorem for quantum system. The content of this theorem can be expressed as the following: two overlapping subsystem AB and BC, the entropy of their union (ABC) plus the entropy of their intersection (B) does not exceed the sum of the entropies of the subsystems (AB and BC) [2], that is

$$S(\rho_{ABC}) + S(\rho_B) \le S(\rho_{AB}) + S(\rho_{BC}). \tag{1}$$

where $S(\rho) = -Tr(\rho l n \rho)$. It is well known that this theorem is true for classical information theory, but to prove this theorem is true for quantum system is very difficult. This theorem is first conjectured to be true for the quantum system by Lanford and Robibson [3]. The first proof of this conjecture is given by Lieb *et al.* several years later. This proof is based on the concave of the function $S(\rho_{12}) - S(\rho_1)$ in ρ_{12} [4]. Another proof based on the same fact was proposed by Uhlmann [5,6].

Recently, quantum information theory attracts more and more attentions for its misterious properties and its potential applications in science and technology [7,8]. The SSA plays an important role in this new field [9] too. The fundamental scource in quantum information is entanglement between many particles which can be viewed as the relations between the partial particles. So distinguish whether a set of the partial particles come from a single state (N-representability problem) [10,11] and further to obtain its entanglement property are important in quantum information while SSA gives a strong constraint on the partial particles and the whole system. The convenience of the SSA is that it has explicit physical and manipulating meaning. So it is a convenient necessary criterion for the N-representability problem. Recently, we use the matrix analysis method [12] to get some necessary conditions for the N-presentability problem. We find that using these conditions we can get a new proof for the SSA. Our new proof is elementary, we need only to use the Lagrange multiplier method and carefully analyse the minimum of a function.

II. THE THEOREM AND THE PROOF

There is a density matrix ρ_{ABC} , where the particle A,B and C are in L-dimension, M-dimension and N-dimension Hilbert space, respectively. Let $\{\lambda_{AB}^{(1)},\lambda_{AB}^{(2)},\cdots,\lambda_{AB}^{(LM)}\}$, $\{\lambda_{BC}^{(1)},\lambda_{BC}^{(2)},\cdots,\lambda_{BC}^{(MN)}\}$, $\{\lambda_{B}^{(1)},\lambda_{BC}^{(2)},\cdots,\lambda_{ABC}^{(MN)}\}$, are the eigenvalues of the density matrix $\rho_{AB},\rho_{BC},\rho_{B}$ and ρ_{ABC} , respectively (where

^{*}Electronic address: yshzhang@ustc.edu.cn †Electronic address: gcguo@ustc.edu.cn

 ρ_{AB}, ρ_{BC} and ρ_{B} are gotten by tracing the other particles from ρ_{ABC}), and they are arranged in increasing order. We defined vectors $\lambda_{AB} = \{\lambda_{AB}^{(1)}, \lambda_{AB}^{(2)}, \cdots, \lambda_{AB}^{(LM)}\}$, $\lambda_{BC} = \{\lambda_{BC}^{(1)}, \lambda_{BC}^{(2)}, \cdots, \lambda_{BC}^{(MN)}\}$, $\lambda_{B} = \{\lambda_{B}^{(1)}, \lambda_{B}^{(2)}, \cdots, \lambda_{B}^{(M)}\}$, $\lambda_{ABC} = \{\lambda_{ABC}^{(1)}, \lambda_{ABC}^{(2)}, \cdots, \lambda_{ABC}^{(LMN)}\}$ and $\lambda_{AB}^{B} = \{\sum_{i=1}^{L} \lambda_{AB}^{(i)}, \sum_{i=L+1}^{2L} \lambda_{AB}^{(i)}, \cdots, \sum_{i=L(M-1)+1}^{LM} \lambda_{AB}^{(i)}\}$, $\lambda_{BC}^{B} = \{\sum_{i=1}^{N} \lambda_{BC}^{(i)}, \sum_{i=N+1}^{2L} \lambda_{ABC}^{(i)}, \cdots, \sum_{j=(LM-1)N+1}^{LMN} \lambda_{ABC}^{(j)}\}$, and $\lambda_{ABC}^{BC} = \{\sum_{j=1}^{L} \lambda_{ABC}^{(j)}, \sum_{j=L+1}^{2L} \lambda_{ABC}^{(j)}, \cdots, \sum_{j=(LM-1)L+1}^{LMN} \lambda_{ABC}^{(j)}\}$. Using the matrix analysis method, we get the following two lemmas on the eigenvalues [12]. ρ_{AB} , ρ_{BC} and ρ_{B} are gotten by tracing the other particles from ρ_{ABC}), and they are arranged in increasing order. We

Lemma 1. Using the notes defined before, we can get the relations between the eigenvalues of ρ_{BC} , ρ_{AB} , ρ_{B} and ρ_{ABC} as

$$\lambda_{ABC}^{AB} \succ \lambda_{AB} \tag{2.1}$$

$$\lambda_{ABC}^{BC} \succ \lambda_{BC} \tag{2.2}$$

$$\lambda_{BC}^{B} \succ \lambda_{B} \tag{2.3.1}$$

$$\lambda_{AB}^{B} \succ \lambda_{B} \tag{2.3.2}$$

$$\lambda_{AB}^B \succ \lambda_B \tag{2.3.2}$$

Lemma 2. Suppose $rank(\rho_{ABC}) = LMN - Ls$, $rank(\rho_{BC}) = MN - s$, $rank(\rho_{AB}) = LM - r$ and $rank(\rho_{B}) = M - t$, if r and s satisfy the condition $Nr \leq Ls$, there will be

$$t \le \left[\frac{r-1}{L}\right] + 1,\tag{3}$$

where [x] is the maximum integer which is smaller than x.

The notation $y \succ x$ mean that the vector x is majorized by the vector y. The majorization is defined as the following. Let $x = \{x_1, x_2, \dots, x_n\}$ and $y = \{y_1, y_2, \dots, y_n\}$ are n-dimensional vectors and the elements are arranged in increasing order. Then the vector x is majorized by vector y [13], denoted by $y \succ x$, if for each k $(k = 1, 2, \dots, n)$ the following inequality is hold

$$\sum_{i=1}^{k} x_i \ge \sum_{i=1}^{k} y_i$$

and the equality is hold when k=n. Under these two Lemmas, we can find that the SSA is hold in the following. **Theorem** There are four normalized vectors $\lambda^{ABC} = \{\lambda_1^{ABC}, \lambda_2^{ABC}, \cdots, \lambda_{LMN}^{ABC}\}$, $\lambda^{AB} = \{\lambda_1^{AB}, \lambda_2^{AB}, \cdots, \lambda_{LMN}^{AB}\}$, $\lambda^{BC} = \{\lambda_1^{BC}, \lambda_2^{BC}, \cdots, \lambda_{MN}^{BC}\}$ and $\lambda^{B} = \{\lambda_1^{B}, \lambda_2^{B}, \cdots, \lambda_M^{BC}\}$, the elements of these vectors are non-negative and arranged in increasing and define the vectors λ^{ABC}_{AB} , λ^{ABC}_{BC} and λ^{BC}_{BC} , λ^{AB}_{BC} , which are similar to the vectors in lemma 1. If the elements of these vectors satisfy the following conditions

- 1. $\lambda_{AB}^{ABC} \succ \lambda_{B}^{AB}$; 2. $\lambda_{BC}^{ABC} \succ \lambda_{B}^{BC}$; 3. $\lambda_{B}^{BC} \succ \lambda_{B}^{B}$ and $\lambda_{B}^{AB} \succ \lambda_{B}^{B}$; 4. Suppose the vector $\lambda = \lambda_{BC}^{ABC}$ has only Ls zero elements and λ_{B}^{BC} has s zero elements, and if the vector λ_{B}^{AB} has s zero elements, there are at least $\left[\frac{r-1}{L}\right] + 1$ elements of the vector λ_{B}^{AB} are zeroes. If exchange the role of the vector λ_{B}^{AB} are zeroes. λ^{AB} and λ^{BC} , the similar result must be hold also.

Thus the following inequality is hold

$$S(\lambda) + S(\lambda^B) \le S(\lambda^{BC}) + S(\lambda^{AB}),\tag{4}$$

where $S(\lambda) = \sum_{i=1}^{LMN} (-\lambda_i \ln \lambda_i)$.

The proof of the theorem is technical. We use the Lagrange multiplier method to get the minimal value of a function under the conditions 1, 2 and 3. Because there are many possible extreme points, we need to find out the minimal one. We use some facts to find that when the function gets the minimal value, all of the nonzero elements are equal to each other. Then use the condition 4 to get the minimal value of the function.

Proof. At first, we define a function

$$F = \sum_{i=1}^{LM} (-\lambda_i^{AB} \ln \lambda_i^{AB}) + \sum_{i=1}^{MN} (-\lambda_i^{BC} \ln \lambda_i^{BC}) + \sum_{i=1}^{M} (\lambda_i^B \ln \lambda_i^B) + \sum_{i=1}^{LMN} (\lambda_i \ln \lambda_i),$$
 (5)

If we can prove that the minimum of this function is not less than 0 under the conditions 1-4, the theorem is true. So the proof becomes to find the minimal value of a function under some conditions. Obviously, the minimal value of this function exists and is finite. Now we use the Lagrange multiplier method to deal with the conditions 1, 2, 3 and define a new function

$$G = \sum_{i=1}^{LM} (-\lambda_i^{AB} \ln \lambda_i^{AB}) + \sum_{i=1}^{MN} (-\lambda_i^{BC} \ln \lambda_i^{BC}) + \sum_{i=1}^{M} (\lambda_i^{B} \ln \lambda_i^{B}) + \sum_{i=1}^{LMN} (\lambda_i \ln \lambda_i)$$

$$+ \sum_{i=1}^{LM-1} \alpha_i^1 (\sum_{j=1}^{i} \lambda_j^{AB} - \sum_{j=1}^{Ni} \lambda_j - x_{1i}^2) + \sum_{k=1}^{MN-1} \beta_k^1 (\sum_{j=1}^{k} \lambda_j^{BC} - \sum_{j=1}^{Lk} \lambda_j - y_{1k}^2)$$

$$+ \sum_{i=1}^{M-1} \alpha_i^2 (\sum_{j=1}^{Li} \lambda_j^{B} - \sum_{j=1}^{i} \lambda_j^{AB} - x_{2i}^2) + \sum_{k=1}^{M-1} \beta_k^2 (\sum_{j=1}^{Nk} \lambda_j^{BC} - \sum_{j=1}^{k} \lambda_j^{BC} - y_{2k}^2)$$

$$+ \sum_{i=0}^{LMN} \gamma_i (\lambda_{i+1} - \lambda_i - z_i^2) + \sum_{i=1}^{LM} u_i (\lambda_{i+1}^{AB} - \lambda_i^{AB} - r_i^2)$$

$$+ \sum_{i=1}^{MN} v_i (\lambda_{i+1}^{BC} - \lambda_i^{BC} - s_i^2) + \sum_{i=1}^{M} w_i (\lambda_{i+1}^{B} - \lambda_i^{B} - t_i^2)$$

$$+ a_1 (\sum_{i=1}^{LM} \lambda_i^{AB} - 1) + a_2 (\sum_{i=1}^{MN} \lambda_i^{BC} - 1) + a_3 (\sum_{i=1}^{M} \lambda_i^{B} - 1) + a_4 (\sum_{i=1}^{LMN} \lambda_i - 1),$$

where the parameters α_i^j , β_k^j , u_i , v_i and w_i , γ_j , a_i are Lagrange multipliers, x_{1i}^2 , y_{1k}^2 , x_{2i}^2 , y_{2k}^2 , z_i^2 , r_i^2 , s_i^2 and t_i^2 are introduced to make the inequalities to be equations. We have used the conditions that λ_i^{AB} , λ_j^{B} , λ_k^{BC} and λ_i are arranged in increasing order and let $\lambda_0 = 0$.

Then when G get the minimal value, there must be some constraints on the parameters and variables. First, we can get $\alpha_i^j x_{ji} = 0$ $(i = 1, 2, \dots, LM - 1; j = 1, 2)$ and the similar relations between β_k^j and y_{jk} , γ_i and z_i , u_i and r_i , v_i and s_i , w_i and t_i . The second, we get the relations between the elements of the vector λ^{AB} and λ ,

$$\sum_{j=1}^{i} \lambda_j^{AB} - \sum_{j=1}^{Ni} \lambda_j = x_{1i}^2, i = 1, 2, \dots, LM - 1$$
 (7)

and the similar relations between λ and λ^{BC} , λ^{AB} and λ^{B} , λ^{BC} and λ^{B} . Then the relations between the vector λ^{AB} can be gotten

$$\lambda_{i+1}^{AB} - \lambda_i^{AB} = r_i^2, i = 1, \dots, LM - 1;$$

$$\sum_{i=1}^{LM} \lambda_i^{AB} - 1 = 0,$$
(8)

and the similar relations between the vectors of λ^{BC} , λ^{B} and λ .

The most important constraints are the equations between the vectors $\lambda_i^{AB}, \lambda_i^{BC}$, λ_i^{B} and λ_i

$$-\ln \lambda_i^{AB} - 1 + \sum_{j=i}^{LM-1} \alpha_i^1 - \sum_{j=\left[\frac{i-1}{L}\right]+1}^{M-1} \alpha_j^2 + u_{i-1} - u_i + a_1 = 0 \ (i = 1, 2, \dots, LM). \tag{9.1}$$

$$-\ln \lambda_i^{BC} - 1 + \sum_{j=i}^{MN-1} \beta_j^1 - \sum_{j=\left[\frac{i-1}{N}\right]+1}^{M-1} \beta_j^2 + v_{i-1} - v_i + a_2 = 0 \ (i = 1, 2, \dots, MN). \tag{9.2}$$

$$\ln \lambda_i^B + 1 + \sum_{j=i}^{M-1} \alpha_i^2 + \sum_{j=i}^{M-1} \beta_i^2 + w_{i-1} - w_i + a_3 = 0 \ (i = 1, 2, \dots, M). \tag{9.3}$$

$$\ln \lambda_i + 1 - \sum_{j=\left[\frac{i-1}{N}\right]+1}^{LM-1} \alpha_j^1 - \sum_{j=\left[\frac{i-1}{T}\right]+1}^{MN-1} \beta_j^1 + \gamma_{i-1} - \gamma_i + a_4 = 0 \ (i = 1, 2, \dots, LMN).$$
 (9.4)

Since the number of the possible cases are so large, it is very difficult to get the solutions directly. We point out some useful facts to reduce the possible solutions and to find the minimal value of the function G.

Fact 1. When the function G get the minimum, suppose that parameters α_i^1 and α_j^2 are the nearest nonzero parameter act on the elements of vector λ^{AB} , if the parameters u_i and u_{Lj} are zeroes, all the parameter u_k $(i \le k \le Lj)$ are equal to zeroes. This fact is true for the other parameters v_i, w_i, γ_i .

Proof. Without loss of generality, we only consider the parameter u_i . Suppose the fact is not true, there are some parameters u_p ($i < m \le p \le n < Lj$) are not zeroes. For simplicity, we suppose there are no more nonzero parameters α_i^1 and α_i^2 . Then we get the conditions from (9.1)

$$-\ln \lambda_m^{AB} - 1 + \alpha_i^1 - \alpha_j^2 + 0 - u_m + a_1 = 0,$$

$$-\ln \lambda_{m+1}^{AB} - 1 + \alpha_i^1 - \alpha_j^2 + u_m - u_{m+1} + a_1 = 0,$$

$$\vdots$$

$$-\ln \lambda_n^{AB} - 1 + \alpha_i^1 - \alpha_j^2 + u_{n-1} - u_n + a_1 = 0,$$

$$-\ln \lambda_{n+1}^{AB} - 1 + \alpha_i^1 - \alpha_j^2 + u_n - 0 + a_1 = 0,$$

$$(10)$$

where we have used the conditions that the parameters u_{m-1} and u_{n+1} are zeros. Since the parameters u_p ($i < m \le p \le n < Lj$) are nonzero, then we get $\lambda_m^{AB} = \lambda_{m+1}^{AB} = \cdots = \lambda_n^{AB} = \lambda_{n+1}^{AB}$. So we have the relations between these nonzero parameters

$$-u_m = u_m - u_{m+1} = \dots = u_{n-1} - u_n = u_n, \tag{11}$$

that is, $u_n = (n - m + 1)u_m = -u_m$. So $u_m = 0$, then all of the parameters u_p $(i < m \le p \le n < Lj)$ are zeroes, which is inconsistent with our suppose. So this fact is true. QED.

Since the fact 1, the parameters u_k affect the result only when there are some nonzero parameter α_i^1 or α_j^2 make k = i or k = Lj. For this situation, we have the following fact.

Fact 2. When the function G get the minimum, if there are a set of parameters u_k $(m \le k \le n)$ are nonzero and there are some nonzero parameters α_i^1 make $m \le i \le n$, This situation is equal to the situation where the parameters u_k $(m \le k \le n)$ and α_i^1 are zeroes, but two new parameters α_{m-1}^1 and α_{n+1}^1 should be added, and the parameters γ_i $(0 \le i \le LMN)$ should be adjusted.

Proof. Without loss of generality, we suppose only the nonzero parameter α_i^1 satisfy the condition $m \leq i \leq n$. For simplicity, we suppose there is no other nonzero parameters act on the eigenvalues of ρ_{AB} . Since the parameter α_i^1 are nonzero, then

$$\sum_{j=1}^{i} \lambda_j^{AB} - \sum_{j=1}^{Ni} \lambda_j = 0; \tag{12}$$

and the parameters u_k $(m \le k \le n)$ are nonzero, we get

$$\lambda_m^{AB} = \lambda_{m+1}^{AB} = \dots = \lambda_n^{AB} = \lambda_{n+1}^{AB}. \tag{13}$$

Since we have the condition $\sum_{j=1}^{i-1} \lambda_j^{AB} \ge \sum_{j=1}^{N(i-1)} \lambda_j$, together with the equation (12), we get $\lambda_i^{AB} \le \sum_{j=N(i-1)+1}^{Ni} \lambda_j$. On the other hand, $\lambda_{i+1}^{AB} \ge \sum_{j=N(i+1)}^{N(i+1)} \lambda_j$, that is, $\sum_{j=N(i-1)+1}^{Ni} \lambda_j \ge \sum_{j=N(i+1)}^{N(i+1)} \lambda_j$. Because of the condition $\lambda_{N(i-1)+1} \le \lambda_{N(i-1)+2} \le \cdots \le \lambda_{N(i+1)}$, we get the equation $\lambda_{N(i-1)+1} = \lambda_{N(i-1)+1} = \cdots = \lambda_{N(i+1)}$. So we get $\lambda_i^{AB} = \sum_{j=N(i-1)+1}^{Ni} \lambda_j$ and $\lambda_{i+1}^{AB} = \sum_{j=N(i+1)}^{N(i+1)} \lambda_j$, that is

$$\sum_{j=1}^{i-1} \lambda_j^{AB} - \sum_{j=1}^{N(i-1)} \lambda_j = 0, \sum_{j=1}^{i+1} \lambda_j^{AB} - \sum_{j=1}^{N(i+1)} \lambda_j = 0.$$
 (14)

Continue to use this method we can get

$$\sum_{j=1}^{m-1} \lambda_j^{AB} - \sum_{j=1}^{N(m-1)} \lambda_j = 0, \sum_{j=1}^{m-1} \lambda_j^{AB} - \sum_{j=1}^{N(m+1)} \lambda_j = 0.$$
 (15)

From these equations, we can find this is just as there are two nonzero parameters α_{m-1}^1 and α_{n+1}^1 , and the nonzero parameter α_i^1 have no effect in this case. From the constraints on λ_k^{AB} , there will be

$$-\ln \lambda_k^{AB} - 1 + \alpha_i^1 + a_1 = 0 \ (1 \le k \le m - 1), \tag{16.1}$$

$$-\ln \lambda_k^{AB} - 1 + \alpha_i^1 + u_{k-1} - u_k + a_1 = 0 \ (m \le k \le i),$$
(16.2)

$$-\ln \lambda_k^{AB} - 1 + u_{k-1} - u_k + a_1 = 0 \ (i+1 \le k \le n+1), \tag{16.3}$$

$$-\ln \lambda_k^{AB} - 1 + a_1 = 0 \ (n+2 \le k \le LM). \tag{16.4}$$

Since the elements λ_k^{AB} $(m \le k \le n+1)$ are equal to each other, then $u_m = u_m - u_{m+1} = \cdots = u_{i-1} - u_i \equiv \alpha$, $u_i - u_{i+1} = u_{i+1} - u_{i+2} = \cdots = u_n \equiv \beta$ and $\beta = \alpha + \alpha_i^1$. If we let the nonzero parameters $\alpha_{m-1}^1 = -\alpha$ and $\alpha_{n+1}^1 = \beta$, the equations are the same. Now we consider the effect of this substitution on the vector λ . For simplicity, we suppose also that there are only the nonzero parameter α_i^1 act on the vector λ . Then the equations are

$$\ln \lambda_k + 1 - \alpha_i^1 + \gamma_{k-1} - \gamma_k + a_4 = 0 \ (1 \le k \le Ni), \tag{17.1}$$

$$\ln \lambda_k + 1 + \gamma_{k-1} - \gamma_k + a_4 = 0 \ (Ni + 1 \le k \le LMN). \tag{17.2}$$

Insert the parameters α_{m-1}^1 and α_{n+1}^1 into the equations, we can find that

$$\ln \lambda_k + 1 - \alpha_{m-1}^1 - \alpha_{m+1}^1 + \gamma_{k-1} - \gamma_k + a_4 = 0 \ (1 \le k \le Ni),$$
(18.1)

$$\ln \lambda_k + 1 - \alpha_{n+1}^1 + \gamma_{k-1}^{'} - \gamma_k^{'} + a_4 = 0 \ (Ni + 1 \le k \le N(n+1))$$
 (18.2)

$$\ln \lambda_k + 1 + \gamma'_{k-1} - \gamma'_k + a_4 = 0 \ (N(n+1) + 1 \le k \le LMN)$$
 (18.3)

where γ_k' $(Ni+1 \le k \le LMN)$ are the new parameters to make the equations are the same as the equation (17). This is just as the situation that the parameters α_{m-1}^1 and α_{n+1}^1 are nonzero, and the parameter γ_k is adjusted. QED This fact is also true for the parameters β_j^1 . This fact tell us that any solution found in the former situation can be found in the later case. In the following, we always suppose we have already done this change. After making these changes there is no nonzero parameters α_i^1 or β_j^1 make the parameter $u_i(v_j)$ nonzero and the parameters γ_i are

substituted by γ_i' .

Fact 3. When the function G get the minium, if i and j are the nearest indexes to make the equations $\sum_{k=1}^i \lambda_k^{AB(BC)} = \sum_{k=1}^{N(L)i} \lambda_k$ and $\sum_{k=1}^j \lambda_k^{BC(AB)} = \sum_{k=1}^{L(N)j} \lambda_k$ to be hold, the elements of the vector λ between Ni and Lj are equal to each other.

Proof. We suppose this conclusion is not true, without loss of generality, let Lj > Ni. Then there are some elements satisfy the following conditions $\lambda_{Lj} = \lambda_{Lj-1} = \cdots = \lambda_p \equiv \lambda_b > \lambda_a \equiv \lambda_{Ni} = \lambda_{Ni+1} = \cdots = \lambda_q$, for simplification, we suppose that $Lj - p \geq q - Ni$. If we define the following parameters Δ_l and Δ'_m as $\sum_{k=i+1}^l \lambda_k^{AB} - \sum_{k=Ni+1}^{Nl} \lambda_k = \Delta_l$ $(i+1 \leq l \leq \left \lfloor \frac{Lj}{N} \right \rfloor, [x]$ is the maximal integer which is smaller than x) and $\sum_{k=Nm+1}^{Nj} \lambda_k - \sum_{k=m}^j \lambda_k^{BC} = \Delta'_m$ $(\left \lfloor \frac{Ni}{L} \right \rfloor \leq m \leq j)$, we can find that all of these parameters are more than zero. Then we take out the minimal number from $\frac{\Delta_l}{l-i}$ and $\frac{\Delta'_m}{j-m}$, we denote it by Δ , obviously it is more than zero. Now we change the element λ_2 by $\lambda_2 - \frac{\Delta'}{Lj-p}$ and λ_1 by $\lambda_1 + \frac{\Delta'}{q-Ni}$ where the parameter $\Delta' = (q-Ni)\Delta$. After these substitution, the new elements of the vector λ' satisfy all of the conditions. The entropy of the vector λ' is larger than the entropy of the vector λ and the entropy of the other vector is invariable. So the function G for the new vector is smaller than the former which is inconsistent with the suppose. QED.

This fact is also true for the vector λ^B . Since we have the fact 3, then we want to know how many nonzero parameters α^1_i and β^1_j in the section where all of the elements are the same. We have the following fact

Fact 4. When the function G get the minimum, there is no nonzero parameters α_i^1 and β_j^1 in the section where all of the elements of vector λ are the same except for the edge parameters.

Proof. We first point out that there are at most four nonzero parameters α_i^1 or β_j^1 in the section where all of the elements of vector λ are equal to each other if the conclusion is not true. If this assert is not true, there are at least five nonzero parameters act on the section where all of the elements of the vector λ are the same. So at least three of them (such as α_l^1 $(l=i,j,k\cdots)$ or β_l^1 $(l=i,j,k\cdots)$) are act on the same vector. Without loss of generality,

we suppose there are three nonzero parameters α_l^1 (l=i,j,k). Since the elements $\lambda_{Ni+1} = \lambda_{Ni+2} = \cdots = \lambda_{Nj} = \lambda_{Nj+1} = \lambda_{Nj+2} = \cdots = \lambda_{Nk} \equiv \lambda_a$. We have $\lambda_{i+1}^{AB} \geq N\lambda_a$ and $\lambda_k^{AB} \leq N\lambda_a$, since $\lambda_{i+1}^{AB} \leq \lambda_k^{AB}$, then all of the elements λ_l^{AB} $(i+1 \leq l \leq k)$ are equal to each other. Because we have already done the changes in the fact 2, and use the fact 1, we find all of the parameters u_k $(i+1 \leq l \leq k)$ are zeroes. Further more, the parameters α_j^1 are zeroes too. So the number of the nonzero parameters is no more than four, and they divide the section where all the elements are equal into three smaller sections.

Now we only need to prove the case that less than five parameters are also zeroes. If these parameters are nonzero and set on the vectors as figure 1, which makes the function G get the minimum, we take some sufficient small value Δ from the elements of the first section to the third section. At the same time, Δ must be taken from the left side section of the parameters k and j to the right side section. Using the same method of the proof of the fact 3, if the Δ is sufficient small, all the conditions will be satisfied. From the following calculating, we can find that through this manipulation the function G is smaller which is inconsistent with the minimal suppose.

Let the elements of the vectors before the manipulating are $\lambda_{Ni+1} = \lambda_{Ni+2} = \cdots = \lambda_{Lk} \equiv \lambda_a, \lambda_{Nj+1} = \lambda_{Nj+2} = \cdots = \lambda_{Ll} \equiv \lambda_b; \lambda_j^{AB} = \lambda_{j-1}^{AB} = \cdots = \lambda_s^{AB} \equiv \lambda_a^{AB}, \lambda_{j+1}^{AB} = \lambda_{j+2}^{AB} = \cdots = \lambda_t^{AB} \equiv \lambda_b^{AB}; \lambda_k^{BC} = \lambda_{k-1}^{BC} = \cdots = \lambda_u^{BC} \equiv \lambda_a^{BC}, \lambda_{k+1}^{BC} = \lambda_{k+2}^{BC} = \cdots = \lambda_b^{BC} \equiv \lambda_b^{BC}$. After the manipulate, the new elements are $\lambda_a' = \lambda_a - \frac{\Delta}{Lk - Ni}, \lambda_b' = \lambda_a + \frac{\Delta}{Ll - Nj}; \lambda_a^{AB'} = \lambda_a^{AB} - \frac{\Delta}{j - s + 1}, \lambda_b^{AB'} = \lambda_b^{AB} + \frac{\Delta}{t - j}; \lambda_a^{BC'} = \lambda_a^{BC} - \frac{\Delta}{k - u + 1}, \lambda_b^{BC'} = \lambda_b^{BC} - \frac{\Delta}{v - k}$ and the other elements are the same as before. Since Δ is sufficient small, we can expand the function $\ln(\lambda + \frac{\Delta}{K}) = \ln \lambda + \frac{\Delta}{K\lambda}$ in the first order. Using this formula, we can calculate the difference of the function G between these two vectors.

$$G' - G = -(Lk - Ni)\lambda_a \ln \lambda_a - (Ll - Nj)\lambda_b \ln \lambda_b + (Lk - Ni)\lambda_a' \ln \lambda_a' + (Ll - Nj)\lambda_b' \ln \lambda_b'$$

$$+ (j - s + 1)\lambda_a^{AB} \ln \lambda_a^{AB} + (t - j)\lambda_b^{AB} \ln \lambda_b^{AB} - (j - s + 1)\lambda_a^{AB'} \ln \lambda_a^{AB'}$$

$$- (t - j)\lambda_b^{AB'} \ln \lambda_b^{AB'} + (k - u + 1)\lambda_a^{BC} \ln \lambda_a^{BC} + (v - k)\lambda_b^{BC} \ln \lambda_b^{BC}$$

$$- (k - u + 1)\lambda_a^{BC'} \ln \lambda_a^{BC'} - (v - k)\lambda_b^{BC'} \ln \lambda_b^{BC'}$$

$$= \Delta \ln \frac{\lambda_b \lambda_a^{AB} \lambda_b^{BC}}{\lambda_a \lambda_b^{AB} \lambda_b^{BC}}$$
(19)

Since $\lambda_a = \lambda_b$ and $\lambda_a^{AB} < \lambda_b^{AB}, \lambda_a^{BC} < \lambda_b^{BC}$, there will be G' - G < 0. This is inconsistent with the suppose that the function G get the minimum. QED

Fact 5: When the function G get the minimum, there are at most one α_i^1 and one α_j^2 are nonzero and the elements $\lambda_k = 0$ $(k \le Ni \text{ or } k \le Lj), \lambda_k^{AB} = 0$ $(k \le i), \lambda_k^{BC} = 0$ $(k \le j)$.

The proof of this fact is similar to the proof of the second part of the fact 4. If there is another nonzero parameter, we can take some small value from the left of this parameter to the right of it to make the value of the function G smaller, which is inconsistent with the minimal suppose of the function G. This fact means that all of the nonzero elements of the vector λ are equal to each other. If there is no nonzero parameter act on the vector λ , that is, all of the parameters α_i^1 and β_j^1 are zeroes, then all of the elements of the vector λ are $\frac{1}{LMN}$, all of the elements of the vector λ^{BC} are $\frac{1}{MN}$, all of the elements of the vector λ^B are $\frac{1}{M}$. Now the value of the function G is zero. If there is only one parameter (such as α_i^1) is nonzero, we have the following fact.

Fact 6: When the function G get the minimum and there is only one parameter α_i^1 (β_j^1) is nonzero, then all of the nonzero elements of the vector λ^{BC} , λ^{AB} and λ^{B} are equal to each other.

Proof: Without loss of generality, we suppose the nonzero parameter is α_i^1 . The nonzero elements of the vector λ^{AB} is equal to each other. We can get this result by only using the inequality between the elements of the vector λ^{AB} and λ . We focus on the other part of the fact. Since the nonzero elements of the vector λ^{AB} are the same, all of the parameters α_i^2 are zero. Now we only consider the parameters β_j^2 . Suppose the nonzero parameters $\beta_{i_1}^2, \beta_{i_2}^2, \cdots, \beta_{i_k}^2$ are set as the figure II. From the constraints of the elements of the vector λ^B and λ^{BC} in equations (9)

$$-\ln \lambda_i^{BC} - 1 - \sum_{j=\left[\frac{i-1}{N}\right]+1}^{M-1} \beta_j^2 + v_{i-1} - v_i + a_2 = 0 \ (i = 1, 2, \dots, MN), \tag{20.1}$$

$$\ln \lambda_i^B + 1 + \sum_{j=i}^{M-1} \beta_i^2 + w_{i-1} - w_i + a_3 = 0 \ (i = 1, 2, \dots, M).$$
 (20.2)

Then we find the elements of these vectors can be divided into several groups, in each group the elements are equal to each other, that is,

$$\lambda_{Ni_{1}+p_{1}+1}^{BC} = \lambda_{Ni_{1}+p_{1}+2}^{BC} = \cdots = \lambda_{MN}^{BC} \equiv \zeta_{0}^{BC},
\lambda_{Ni_{1}-q_{1}}^{BC} = \lambda_{Ni_{1}-q_{1}+1}^{BC} = \cdots = \lambda_{Ni_{1}+p_{1}}^{BC} \equiv \zeta_{01}^{BC}
\lambda_{Ni_{2}+p_{2}+1}^{BC} = \lambda_{Ni_{2}+p_{2}+2}^{BC} = \cdots = \lambda_{Ni_{1}-q_{1}-1}^{BC} \equiv \zeta_{1}^{BC},
\vdots
\lambda_{1}^{BC} = \lambda_{2}^{BC} = \cdots = \lambda_{Ni_{k}}^{BC} \equiv \zeta_{k}^{BC},$$
(21)

and

$$\lambda_{i_1+1}^B = \lambda_{i_1+2}^B = \dots = \lambda_M^B \equiv \zeta_0^B,$$

$$\lambda_{i_2+1}^B = \lambda_{i_2+2}^B = \dots = \lambda_{i_1}^B \equiv \zeta_1^B,$$

$$\vdots$$

$$\lambda_1^B = \lambda_2^B = \dots = \lambda_{i_k}^B \equiv \zeta_k^B.$$
(22)

We must note that all of the parameters w_i which act on the vector λ^B are zeroes. At first, if all of the indexes i_j satisfy $w_{i_j}=0$, using the fact 1, all of the parameters are zero. The second, if there are some indexes (such as i_j) make the parameter w_{i_j} to be nonzero. Because the elements in the same section are equal to each other for the fact 3, we get $\zeta_{j-i}^B = \zeta_j^B$. Because of $\zeta_{j-1}^B \geq N\zeta_{j-1,j}^{BC}$ and $\zeta_j^B \leq N\zeta_{j-1,j}^{BC}$, then $\zeta_j^B = N\zeta_{j-1,j}^{BC}$. So if we let $l = [\frac{q_j}{N}]$ and $m = [\frac{p_j}{N}]$, we can get the inequality $\zeta_{j-1}^B \geq (p_j - m)\zeta_{j-1,j}^{BC} + (N - p_j + m)\zeta_{j-1}^{BC}$ and $\zeta_j^B \leq (q_j - l)\zeta_{j-1,j}^{BC} + (N - q_j + l)\zeta_j^{BC}$. Since $\zeta_{j-1}^{BC} \geq \zeta_{j-1,j}^{BC} \geq \zeta_j^{BC}$, we can get that $\zeta_{j-1}^{BC} = \zeta_{j-1,j}^{BC} = \zeta_j^{BC}$. Now we can get the conclusion by using the fact 1, that all of the parameters v_k $(i_j - q_{i_j} \leq k \leq i_j - p_{i_j})$ and $\beta_{i_j}^2$ are zeros. So the second situation can be reduced to the first situation. So The constraints of the vectors λ^B and λ^{AB} are reduced to

$$-\ln \lambda_i^{BC} - 1 - \sum_{j=\lfloor \frac{i-1}{N} \rfloor + 1}^{M-1} \beta_j^2 + v_{i-1} - v_i + a_2 = 0 \ (i = 1, 2, \dots, MN), \tag{23.1}$$

$$\ln \lambda_i^B + 1 + \sum_{j=i}^{M-1} \beta_i^2 + a_3 = 0 \ (i = 1, 2, \dots, M).$$
 (23.2)

If let $\frac{\zeta_i^{BC}}{\zeta_0^{BC}} = \chi_i, \frac{\zeta_{i-1,i}^{BC}}{\zeta_0^{BC}} = \chi_{i-1,i}$ and $\frac{\zeta_i^B}{\zeta_0^B} = \eta_i$ $(i=0,2,\cdots,k)$, we can get $\chi_i = \eta_i$ and $\chi_{i-1,i} = \chi_{i-1}^{\omega_i} \chi_i^{\sigma_i}$ where $\omega_i + \sigma_i = 1$ and $\omega_i = \frac{p_i}{p_i + q_i}$, $\sigma_i = \frac{q_i}{p_i + q_i}$. From these definition, we find that all of the parameters χ_i and $\chi_{i-1,i}$ are in the section [0,1]. Since we have the conditions $\sum_{l=Ni_j+1}^{Ni_{j+1}} \lambda_l^{BC} = \sum_{l=i_j+1}^{i_{j+1}} \lambda_l^B$, then we can get the equations $\sum_{l=Ni_j+1}^{Ni_{j+1}} \lambda_l^{BC} = \sum_{l=i_j+1}^{i_{j+1}} \lambda_l^B$. That is

$$\frac{[N(i_j-i_{j+1})-p_{j+1}-q_j]\chi_j+p_{j+1}\chi_{j+1}^{\sigma_{j+1}}\chi_j^{\omega_{j+1}}+q_j\chi_j^{\sigma_j}\chi_{j-1}^{\omega_j}}{MN-Ni_1-p_1+p_1\chi_1^{\sigma_1}}=\frac{(i_j-i_{j+1})\chi_j}{M-i_1}.$$

So we can get the equations

$$(i_j - i_{j+1})p_1(1 - \chi_1^{\sigma_1}) = (M - i_1)[p_{j+1}(1 - (\frac{\chi_{j+1}}{\chi_j})^{\sigma_{j+1}}) + q_j(1 - (\frac{\chi_{j-1}}{\chi_j})^{\omega_j})]. \tag{24}$$

If there is a parameter $w_{Ni_{m+1}}=0$, then the *m*th equations in equations (24) has no item which is including p_{m+1} . According to the number of the parameters which make $w_{Ni_{l+1}}=0$ $(1 \le l \le k)$, we can divide the elements of these vectors into some sections, the last equation of this section has no item which contains p. We can only point out that the parameter v_{Ni_n} must be zero where the parameter n satisfy the condition $\eta_n=0$ and $\eta_{n-1}>0$. Or the condition 4 will not be satisfied. So we always can sum up all of the equations in the same section to get

$$(i_1 - i_{s+1})p_1(1 - \chi_1^{\sigma_1}) = (M - i_1)[q_1(1 - (\frac{1}{\chi_1})^{\omega_1}) + \sum_{l=2}^{s} (p_l + q_l)(1 - \omega_l(\frac{\chi_l}{\chi_{l-1}})^{\sigma_l} - \sigma_l(\frac{\chi_{l-1}}{\chi_l})^{\omega_l})]$$
(25)

where the parameter s means that the parameter $v_{i_{s+1}} = 0$. We first focus on the lhs. of the equation (25), and obviously, it is non-negative. Then we consider the rhs. of this equation, there is a function $f(x) = xa^{1-x} + (1-x)a^{-x}$. The value of this function is not more than 1. Then the rhs. is non-positive. To make the equation to be hold, the two sides of the equation must be zero. That is $\chi_1 = 1$ and $\sigma_i(\omega_i) = 0$ or a = 1. For each section, we can get the same conditions which imply that all of the nonzero elements of the vector λ^B and λ^{BC} are equal to each other. QED

For the case there are two nonzero parameters α_i^1 and β_j^1 , using the similar method before and notice the condition 4, we can get the same result that all of the nonzero elements are equal to each other.

For the facts proved before, we can get the conclusion that when the function G get the minimum, all of the nonzero elements of the vectors λ^{AB} , λ^{BC} , λ^{B} and λ are equal to each other. Using the condition 4, we can calculate that the minimum of the function G is not less than zero. This is the end of the proof of the theorem. QED

Since the Lemma 1 and Lemma 2, the theorem imply that the SSA is hold. This method can be used to prove some other entropy properties between the partial density matrix and the multipartite density matrix, Such as the inequality $S(\rho_{AB}) \leq S(\rho_A) + S(\rho_B)$.

III. CONCLUSION


In this paper we give a new elementary proof of the SSA which is an important property of the entropy for classical information and quantum information. The proof is dependent on the analysis of the minimal value of a function under some conditions. This proof also show that the conditions in our previous work [12] are stronger than the SSA.

IV. ACKNOWLEDGE

This work was funded by the National Fundamental Research Program (2001CB309300), National Natural Science Foundation of China, the Innovation Funds from Chinese Academy of Sciences, and also founded by the outstanding Ph. D thesis award and the CAS's talented scientist award rewarded to Lu-Ming Duan.

- [1] A. Wehrl, Rev. Mod. Phys. **50**, 221 (1978).
- [2] J. Preskill, Quantum Information and Computation, Lecture Notes on physics Vol. 229 (Caltech, Pasadena, 1998).
- [3] D. W. Robinson and D. Ruelle, Commun. Math. Phys. 5, 288 (1967); O. E. Lanford and D. W. Robinson, J. Math. Phys. 9, 1120 (1968).
- [4] E. H. Lieb and M. B. Ruskai, Phys. Rev. Lett. 30, 434 (1973); E. H. Lieb, Bull. Am. Math. Soc. 81, 1 (1975).
- [5] Uhlmann, Rep. Math. Phys. 7, 449 (1975).
- [6] Uhlmann, Commun. Math. Phys. **54**, 21 (1976).
- [7] C. H. Bennett, G. Brassard, C. Crépeau, R. Jozsa, A. Peres, and W. K. Wootters, Phys. Rev. Lett. 70, 1895 (1993).
- [8] A. K. Ekert, Phys. Rev. Lett. 67, 661 (1991).
- [9] M. A. Nielsen and I. L. Chuang, *Quantum Computation and Quantum Information* (Cambridge University Press, Cambridge, England, 2000).
- [10] R. Erdahl and V. H. Smith, Density matrices and Density Functions, (Reidel Publishing Company, 1987).
- [11] A. J. Coleman, J. Math. Phys. 13, 214 (1972); A. J. Coleman, Rev. Mod. Phys., 35, 668 (1963).
- [12] Han et al. to be published.
- [13] A. W. Marshall and I. Olkin, Inequalities: Theory of Majorization and Its Applications, (Academic Press, New York, 1979).
- [14] M. A. Nielsen et al., Phys. Rev. A 67, 052301 (2003).
 Figure caption.

Figure 1. In this figure, the parameters $\alpha_i^1, \alpha_j^1, \beta_k^1, \beta_l^1$ and $\alpha_n^2, \alpha_q^2, \beta_m^2, \beta_p^2$ are nonzero. The eigenvalues $\lambda_v(i < v \le j)$ are equal to each other. Theses nonzero parameters divide the eigenvalues between i and j into three sections. We take sufficient small value Δ from the first section to the third section. And the same time, we must take the same value from the left section of the parameter k and j to the right section in the eigenvalue λ^{BC} and λ^{BC} , respectively. The bold line means that the eigenvalues in the line are the same.






Fig. 1




Fig. 2