LA-UR-02-5568

Gaussian Dynamics is Classical Dynamics

Salman Habib*

MS B285, Theoretical Division, The University of California, Los Alamos National Laboratory, Los Alamos, NM 87545 (Dated: October 29, 2018)

A direct comparison of quantum and classical dynamical systems can be accomplished through the use of distribution functions. This is useful for both fundamental investigations such as the nature of the quantum-classical transition as well as for applications such as quantum feedback control. By affording a clear separation between kinematical and dynamical quantum effects, the Wigner distribution is particularly valuable in this regard. Here we discuss some consequences of the fact that when closed-system classical and quantum dynamics are treated in Gaussian approximation, they are in fact identical. Thus, it follows that several results in the so-called 'semiquantum' chaos actually arise from approximating the classical, and not the quantum dynamics. (Similarly, opposing claims of quantum suppression of chaos are also suspect.) As a simple byproduct of the analysis we are able to show how the Lyapunov exponent appears in the language of phase space distributions in a way that clearly underlines the difference between quantum and classical dynamical situations. We also informally discuss some aspects of approximations that go beyond the Gaussian approximation, such as the issue of when quantum nonlinear dynamical corrections become important compared to nonlinear classical corrections.

PACS numbers: 03.65.-w, 05.45.Mt

I. INTRODUCTION AND DISCUSSION

A formal comparison of closed classical and quantum dynamical systems is conceptually difficult because what is natural in the classical description (trajectories) is not what is natural in the quantum description (amplitudes) and vice versa. While "semiclassical" limits of quantum mechanics are available (WKB, large N, coherent states, etc.) an important obstruction is the singular nature of the limit $\hbar \to 0$, and the fact that this limit does not commute with the limit $t \to \infty$. Nowhere is this problem more important than in the study of quantum chaos.

Here we adopt the viewpoint that the most natural language in which to compare closed classical and quantum systems is the language of phase space distribution functions as first set forth by Wigner [1]. Since quantum mechanics is a probabilistic theory such a description is physically attractive. Moreover, it has also turned out to be fruitful from a calculational perspective: Among several other applications. Wigner functions have been used to compute quantum corrections to classical statistical mechanics [2, 3], to clarify the WKB limit of quantum mechanics for both regular and stochastic orbits [4, 5, 6], to study quantum dynamics [7], and, in particular, dynamical aspects of quantum chaos [8, 9] including the effects of decoherence [10, 11]. However, we do wish to emphasize that results of this paper do not depend in any essential way on using the Wigner function: It is a convenient device to represent the density matrix, but

certainly not necessary. We also emphasize that it is not our purpose here to discuss the quantum-classical transition *per se* as would be the case in a measured (hence, open) quantum dynamical system [12].

In the area of quantum chaos, the well-known work of Berry and Balazs [8] is based on the idea of comparing classical and quantum dynamics in phase space. These authors studied the evolution of a semiclassical (WKB) Wigner function corresponding to a classical curve in phase space, arguing that while at early times the Airy peak of this distribution function tracks the time-dependent classical curve, at later times, once the classical evolution produces structure on sub- \hbar scales in phase space, the quantum distribution is unable to continue this tracking and there is a transition to a qualitatively new regime. By identifying two sorts of nonlinear structures, 'whorls' (generated by stable motion) and 'tendrils,' (generated by unstable motion) Berry and Balazs conjectured that this transition would occur on a timescale $t_c = O(\hbar^{-2/3})$ for whorls and $t_c = O(\ln \hbar^{-1})$ for tendrils, the latter of which is the famous so-called 'log' time. It is important to note that the existence of the log time does not necessarily imply the failure of semiclassical methods beyond this time: later work has demonstrated that suitably improved semiclassical methods can be successful well past the log time [13, 14, 15, 16].

A naive comparison of Wigner functions and classical probability distributions is not possible as Wigner functions are not positive definite in general. This is a consequence of the fact that quantum mechanics admits interference effects, whereas classical mechanics does not. Nevertheless, one can set up a scheme to compare classical and quantum dynamics in the following way: begin

^{*}Electronic address: habib@lanl.gov; URL: http://t8web.lanl.gov/people/salman/

with an initial distribution function that is acceptable both classically and quantum mechanically (this condition may be relaxed to allowing all acceptable quantum states, with or without a positive Wigner distribution) and then, (i) solve the two different classical and quantum Liouville equations, or equivalently, (ii) solve the corresponding infinite hierarchies of dynamical equations for the corresponding moments. The advantage of using the Wigner function is that, by introducing a c-number phase space description of quantum dynamics, one can directly compare the classical and quantum moments.

A key feature of the Wigner formalism that we exploit here is the clear separation of kinematical and dynamical effects. Confusion has arisen in the literature as a conseguence of the failure to recognize that \hbar can occur in some places purely as a consequence of quantum initial conditions and, in other places, as a consequence of quantum dynamical corrections to classical dynamics. To sharpen one's arguments, especially in areas as controversial as quantum chaos, it is important to clearly distinguish between these two situations. (Essentially this point, but phrased a little differently, has also been made by Ballentine, Yang, and Zibin [17].) The importance of this last point is further underscored by the fact that classical and quantum mechanics have different underlying group symmetries (symplectic vs. unitary) which accounts for the singular nature of the limit $\hbar \to 0$ [18, 19].

A clarification of just what is meant by such statements as "the classical limit" is important both conceptually and operationally (as emphasized in Ref. [17]). The usual textbook arguments involving wave packets and Ehrenfest's theorem attempt to compare a classical description in the sense of a *single trajectory* with a quantum analog, usually taken to be a sufficiently narrow wave packet. This is a rather limited point of view, especially in chaotic systems, where, due to their instability, the concept of individual trajectories is not physically useful even classically, and where quantum wave packet dynamics breaks down very quickly. It makes more sense, therefore, to compare classical distribution functions, describing an ensemble of trajectories with their appropriate quantum analogs as discussed above. (Such a viewpoint is in fact implicit in Feynman's discussion of the two-slit experiment [20], a classic example of a highly nonlinear dynamical system.) The above philosophy of quantum-classical comparison has been implemented in some recent work on the quantum-classical correspondence in kicked systems [21], the driven pendulum [22], and the driven anharmonic oscillator [10]. Such an approach [18] is also needed to analyze experiments in quantum chaos utilizing ultracold atoms [23] in order to unambiguously distinguish quantum from classical effects [24].

Turning now to moment expansions [25], the first important point, discussed below, is that quantum dynamical effects do not arise in moments of less than third order. In an approach utilizing moment expansions around a centroid, this means that to first nontrivial order in fluctuations there are *no* dynamical quantum effects: This is the reason why the simple Gaussian approximation fails to sample any aspects of quantum dynamics.

The infinite hierarchy of coupled equations for the moments is almost impossible to deal with analytically; a variational approach is often attempted by assuming a particular functional form for the distribution functions, forcing thereby relations between lower and higher order moments (alternatively, one may utilize a cumulant expansion). A very popular approach is to assume the distribution function to be Gaussian which implies that all odd moments vanish and that all even moments may be expressed in terms of powers of the second order moments. It is useful to distinguish between three different variants of the Gaussian approximation: The 'time dependent variational principle' (TDVP) method [26], the 'truncated Gaussian approximation' (TGA) of which Heller's method [27] is a non-self-consistent variant, and finally the multiple classical trajectory version of the second technique [13], which we will refer to as the 'multiple trajectory Gaussian approximation' (MTGA) (although in principle the MTGA can be applied to both types of single Gaussian approximations). To avoid confusion, the term 'Gaussian approximation' will be used when discussing general properties of all three variants, otherwise they will be referred to specifically.

It is easy to show that in TGA, and for the individual packets in MTGA, no quantum dynamical corrections are included as only quadratic order moments are considered. This is a generic problem with all Gaussian approximations: in TDVP, while one does resum a certain series (which in a naive moment expansion is merely truncated) this resummation still does not incorporate terms arising from quantum dynamics. (As a consequence, it is not obvious whether the TDVP is any better than the truncated approximations.) For chaotic systems, both the TDVP and TGA will lose accuracy once the initial distribution becomes significantly non-Gaussian, which will happen at least on a Lyapunov timescale (typically this breakdown occurs much sooner) [28]. The MTGA is better behaved because a coherent sum of Gaussians can be a much better approximation for the full distribution function than a single Gaussian. The MTGA also keeps track of relative quantum phases between its constituent wave packets and this is the reason it can handle quantum revivals and other coherent effects.

What is true for Gaussian approximations in nonlinear systems is also obviously true for linear systems where the Gaussian approximation is exact. One consequence of this is that the well-known independent oscillator model [29], long considered a paradigm for quantum Brownian motion, in fact contains only classical dynamics. All quantum features in this model reside in the initial conditions. Once this key fact is realized, the derivation of the Master equation can be radically streamlined with no need to use path integral methods. Moreover, new light is shed on the derivation of classical Langevin equations and the interpretation of precisely what is meant by 'noise' when the dynamics is not Markovian, yet the

The main point of the present paper may be stated in the following way. Classically, or quantum mechanically, when one follows the evolution of a 'mean field' there are 'fluctuation' corrections (back-reaction) to the (classical) equations for the mean field. The quantum moment expansion described above allows one to separate the back-reaction into its classical and quantum components. When the exact moment expansion is approximated by either a quadratic truncation of the hierarchy, or by the Gaussian approximation, the quantum back-reaction is completely neglected. The only quantum feature still remaining is the uncertainty principle constraint on the initial condition for the distribution function. Therefore, if one wishes to compare the strengths of the quantum versus the classical back-reaction, the Gaussian approximation is wholly inadequate. These comments apply with equal force to Gaussian (Hartree) approximations which have attained a certain popularity in studying initial value problems in quantum field theory. Off-Gaussian corrections need to be taken into account, as is possible in principle, e.g., in a systematic 1/N expansion [31] (though this approximation has its own problems [32]). We will discuss some of these aspects further below.

The Gaussian approximation has been applied to study chaos in quantum systems [33, 34]. In Ref. [33], the leading order 1/N expansion (equivalent to a quadratic TGA in that particular case) was used to study a quantum mechanical model which is the (0+1)-dimensional limit of QED. Classical chaos was shown to exist in the dynamical equations for the correlation functions. In Ref. [34], the TDVP was applied to a one-dimensional quartic oscillator, and this system which is classically nonchaotic (in contrast to the situation in Ref. [33]), was shown, in the approximation, to display chaos. In interpreting these examples of "semiquantal" chaos, two questions immediately arise. The first is whether the chaos is just an artifact of the Gaussian approximation, and the second, whether it has anything to do with quantum dynamics. The existence of chaos in the general case of dynamical variational approaches had been previously noted by Caurier et al [35] who argued that the onset of chaos signaled the breakdown of the approximation scheme. Motivated by the results of Ref. [34], the question was also tackled by Sundaram and Milonni [36] using operator moments leading to a similar conclusion. In yet another investigation, it was also demonstrated that Gaussian approximations break down before the timescale on which the approximate dynamics becomes chaotic [28]. (For a recent study of semiquantum chaos in a different approximation, see Ref. [37].) The answer to the second question, given here, is that Gaussian semiquantum chaos, even if approximate, has nothing to do with quantum dynamics. Thus the result of Pattanayak and Schieve [34] is not counter-intuitive (how can "adding" quantum mechanics to a nonchaotic classical system produce chaos?): it is a consequence of the approximation, and selective resummation, of only the classical back-reaction terms in the moment expansion. This also rules out the heuristic explanation that their result is due to the inclusion somehow of quantum "noise," which then makes the system chaotic.

The Gaussian approximation has also been applied by Zhang and Feng [38] in an attempt to display an example of quantal suppression of chaos. Just as we reject the reality of semiquantal chaos, so must we reject such an interpretation of the calculation. What has been shown is that the classical back-reaction, in Gaussian approximation, can suppress chaos. This is not surprising in principle as the Gaussian approximation is completely uncontrolled and there is no guarantee that if the original system is nonintegrable, the Gaussian dynamics also will be (the reverse is true in semiquantum chaos where the original system can be integrable, and the Gaussian approximation, not).

The above discussion leads naturally to another set of questions: What are the timescales on which non-Gaussian corrections become important? Are these timescales different for the classical and quantum backreaction terms? As argued in Refs. [17, 36] it is certainly reasonable to expect the TGA and TDVP to break down, for a classically chaotic system, on a timescale set by the largest Lyapunov exponent. Since ensemble averaged quantities do not possess chaotic signatures like a late-time Lyapunov exponent [39], it is also clear that any time the Gaussian approximation is chaotic, the result is simply wrong. The timescale set by this false chaos, in the cases where it occurs, is therefore precisely the timescale over which one expects the classical non-Gaussian corrections to become important. It is not apriori obvious what will happen for the quantum corrections, but given that the classical non-Gaussian corrections must eliminate the chaos on a (relatively) fast timescale, there is no dynamical imperative for these contributions to become dominant at early times.

The concept of a crossover or 'break' time from classical to quantum dynamics in closed systems (chaotic or not) is often invoked. However, the evidence for the utility of such a concept is surprisingly weak: The notion of the crossover time could well depend on the quantities one chooses to compare, and on the dynamical approximations one makes. If, for example, the estimates of this time are based on TGA or TDVP, then all one is computing is when the approximation breaks down. That this may not have anything to do with how well classical systems track quantum evolution (in a suitable coarse-grained sense), both for chaotic and regular systems, has been pointed out in Ref. [17]. Evidence for this point of view is contained, for example, in the work of Fox and Elston [21] on the kicked top and the kicked pendulum. These authors compare the time evolution of classical and quantum distributions, finding excellent agreement at early times (as expected, the Gaussian approximation being accurate in this regime), but also at

late times, when a coarse-grained quantum distribution is compared to the classical distribution. What this says is that the low order moments can be insensitive to the nature of the dynamics, despite the existence of nontrivial quantum dynamical effects. Investigations utilizing high-resolution simulations have shown that in some examples of chaotic dynamical systems, while the classical and quantum moments do track each other very closely, there is a small 'mini-break' which occurs on a timescale longer than the dynamical time [10]. Whether this 'minibreak' has anything to do with log time arguments is not yet clear, though more recent numerical evidence appears to argue against this [40].

The set of equations of the simple, albeit non-selfconsistent, TGA due to Heller [27] provide an instructive link to the computation of the actual classical Lyapunov exponents. Classical mechanics allows arbitrarily strong shrinking of a Gaussian ball in phase space around the fiducial trajectory; this shrinking is what allows the effective Hamiltonian for Gaussian fluctuations to reduce to that for linearized perturbations, which is just what defines the Lyapunov exponents. In this way, even though the classical Liouville equation is linear, the allowance of a singular trajectory limit permits the existence of chaos. However, in the quantum case, because of the finite value of \hbar , the shrinking allowed classically is impossible beyond a certain point, and the effective Hamiltonian for Gaussian fluctuations never becomes that for linearized perturbations.

Finally, it is important to point out that the Gaussian approximation works extremely well in certain situations where localization of the quantum state is guaranteed, as can happen in studying the dynamics of observed quantum systems (quantum trajectories) (Cf. Ref. [12]). In a related context, the use of a Gaussian ansatz for the Wigner function in quantum feedback control has also been shown to be very effective [41, 42, 43], even when the true state can have substantial non-Gaussian features [43]. While this is not the focus of the present paper, some of the general formulae for Gaussian states discussed here are also applicable to these, otherwise quite distinct, situations.

The organization of this paper is as follows. In Section II, we study the classical and quantum Liouville equations, especially in the context of moment expansions. In Section III, we develop Gaussian dynamics using moments and discuss various ramifications of the results. We conclude in Section IV with a short summary.

II. CLASSICAL AND QUANTUM DYNAMICS IN PHASE SPACE

A. Liouville Equations

Throughout this paper we will deal with continuoustime Hamiltonian dynamical systems, with Hamiltonian

$$H = \frac{p^2}{2m} + V(x), \qquad (1)$$

where the potential V(x) is allowed to be explicitly timedependent. We take the system to be one-dimensional: extension to higher dimensions is obvious. The classical equations of motion following from the Hamiltonian (1) are entirely equivalent to the classical Liouville equation

$$\frac{\partial f_c}{\partial t} = L_c f_c \tag{2}$$

for the phase space distribution function $f_c(x, p)$, and where the classical Liouville operator,

$$L_c = -\frac{p}{m}\frac{\partial}{\partial x} + \frac{\partial V}{\partial x}\frac{\partial}{\partial p}.$$
(3)

Point trajectories can be recovered from the Liouville equation (2) if one allows the distribution function to be a delta function over phase space.

An important consequence of the classical evolution equation is the exact conservation of the quantities (when they exist; f is any phase-space distribution function)

$$\sigma_n \equiv \int dx dp \ f^n \tag{4}$$

by the classical evolution equation (2), following from Liouville's theorem. (The trivial case n = 1 arises from the conservation of total probability.) This will acquire more significance when we proceed to the quantum case.

The corresponding quantum Liouville equation is written in terms of the Wigner function [1], which is a 'half-Fourier transform' of the density matrix $\rho(x_1, x_2)$ in the coordinate representation. One introduces the sum and difference variables

$$x = (x_1 + x_2)/2, \quad \Delta = x_1 - x_2$$
 (5)

and then defines the Wigner function as

$$f_W(x,p) = \frac{1}{2\pi\hbar} \int \rho(x,\Delta) \mathrm{e}^{-ip\Delta/\hbar} d\Delta.$$
 (6)

Details on the properties of this function may be found in Refs. [44, 45]. A particularly nice discussion, with special emphasis on conceptual pitfalls and the distinction between kinematical and dynamical aspects, has been given by Tatarskii [46]. (See also, Ref. [47].)

For the purposes of this paper, it is sufficient to note the following: (i) f_W is real and normalized to unity over the (x, p) phase space, (ii) f_W is square integrable, with

$$\sigma_2 = \int dx dp \ f_W^2 \le \frac{1}{2\pi\hbar},\tag{7}$$

the equality holding for pure states $(2\pi\hbar\sigma_2 = Tr\rho^2)$. This is a *partial* statement of the uncertainty principle in the Wigner formalism; localization of the distribution to a delta function on phase space, while classically allowed, is forbidden in quantum mechanics. (iii) The only pure state for which f_W is positive everywhere is one with the Gaussian wave function (Hudson's theorem [48])

$$\psi = e^{-ax^2 + bx + c} \tag{8}$$

with a, b, c complex and $Re \ a > 0$. (iv) Expectation values of operators that are either pure powers in position or pure powers in momentum are directly given by the averages of x^m or p^m taken with respect to f_W . Expectation values of mixed operators like $\hat{x}^2\hat{p}^2$ cannot be obtained directly. The Wigner formalism is associated with Weyl's rule for the ordering of operators [49]: the phase space average of a mixed term like $x^n p^m$ corresponds to the expectation value of a completely symmetrized operator expression.

The dynamics of the Wigner function is easily obtained from the quantum Liouville equation for the density matrix $\rho(x_1, x_2)$:

$$i\hbar \frac{\partial}{\partial t}\rho(x_1, x_2) = [H(x_1) - H(x_2)^*]\rho(x_1, x_2).$$
 (9)

The Wigner transform of this equation gives

$$\frac{\partial f_W}{\partial t} = L_c f_W + \sum_{\lambda \text{ odd}} \frac{1}{\lambda!} \left(\frac{\hbar}{2i}\right)^{\lambda-1} \frac{\partial^\lambda V(x)}{\partial x^\lambda} \frac{\partial^\lambda f_W}{\partial p^\lambda} \quad (10)$$

where $\lambda \neq 1$, and with the assumption that the potential can be Taylor-expanded. Often this is not the case and Eqn. (10) becomes an integro-differential equation; we avoid this added complication in this paper. On comparing Eqns. (10) and (2) we see that quantum and classical dynamics differ due to the last term in the quantum Liouville equation (10). This apparent separation of quantum and classical pieces in the evolution equation is one of the original motivations for the Wigner function approach. However, since the definition of the Wigner function itself involves \hbar , the actual situation is not as straightforward as it appears to be. In particular, the 'quantum correction' in Eqn. (10) is not unitary. (Neither is L_c , but their sum is. Ref. [18] contains a further discussion including the violation of positivity in the classical approximation.)

The quantum evolution equation (10) exactly conserves the uncertainty principle constraint (7). This has a one-to-one correspondence with the classical evolution preserving the corresponding classical quantity σ_2 of Eqn. (4). It is important to note that since the quantum evolution does not obey Liouville's theorem, in this case, $\dot{\sigma}_n \neq 0, \forall n \neq 1, 2$. Thus, since the conservation of σ_1 is trivial in both classical and quantum evolutions, the constancy of only σ_2 in the quantum case acquires added significance.

B. The Moment Hierarchies

In the moment hierarchy approach to dynamics one attempts to replace a knowledge of the distribution function by a knowledge of all-order values of the quantities $\langle x^n \rangle$, $\langle p^n \rangle$, and $\langle x^n p^m \rangle$ where the bracket denotes an average taken over the distribution function. From now on $\langle \rangle_c$ will denote an expectation value taken with respect to a classical distribution, and $\langle \rangle_q$ will denote an expectation to the quantum distribution function. Below we present an elementary derivation of the classical and quantum moment equations using the evolution equations for the corresponding distributions (for alternative approaches, see Ref. [25]).

With the assumption that the distribution function and its derivatives are zero at the phase space boundary, in the classical case [using the classical Liouville equation (2)] it is easy to show that :

$$\frac{d}{dt} \langle x^n \rangle_c = \int dx dp \ x^n \frac{\partial f_c}{\partial t} \\
= \frac{n}{m} \langle x^{n-1} p \rangle_c, \qquad (11) \\
\frac{d}{dt} \langle p^n \rangle_c = \int dx dp \ p^n \frac{\partial f_c}{\partial t} \\
= -n \left\langle p^{n-1} \frac{\partial V}{\partial x} \right\rangle_c, \qquad (12) \\
\frac{d}{dt} \langle x^n p^k \rangle_c = \int dx dp \ x^n p^k \frac{\partial f_c}{\partial t} \\
= \frac{n}{m} \langle x^{n-1} p^{k+1} \rangle_c \\
-k \left\langle x^n p^{k-1} \frac{\partial V}{\partial V} \right\rangle \qquad (13)$$

$$-k\left\langle x^{n}p^{\kappa-1}\frac{\partial x}{\partial x}\right\rangle_{c}.$$
 (13)

Of course, these equations are nothing but the Hamilton equations of motion averaged over the initial conditions.

The quantum hierarchy is equally straightforward to obtain. The evolution of the spatial moments is given by

$$\frac{d}{dt} \langle x^n \rangle_q = \int dx dp \ x^n \frac{\partial f_W}{\partial t} \\
= \frac{n}{m} \langle x^{n-1} p \rangle_q,$$
(14)

formally the same as the classical case. However,

$$\frac{d}{dt} \langle p^{n} \rangle_{q} = \int dx dp \ p^{n} \frac{\partial f_{W}}{\partial t} \\
= -n \left\langle p^{n-1} \frac{\partial V}{\partial x} \right\rangle_{q} \\
+ \sum_{\lambda \ odd}^{n, \ n-1} \Theta(n, \lambda) \left\langle p^{n-\lambda} \frac{\partial^{\lambda} V}{\partial x^{\lambda}} \right\rangle_{q}, \quad (15)$$

where

$$\Theta(n,\lambda) \equiv \frac{(-1)^{\lambda}}{\lambda!} \frac{n!}{(n-\lambda)!} \left(\frac{\hbar}{2i}\right)^{\lambda-1}, \qquad (16)$$

differs from the corresponding classical equation. The upper limit of the sum in (15) is n, if n is odd, and n-1 if n is even. Clearly the first dynamical quantum correction comes in only at third order, where

$$\frac{d}{dt}\left\langle p^{3}\right\rangle_{q} = -3\left\langle p^{2}\frac{\partial V}{\partial x}\right\rangle_{q} + \frac{\hbar^{2}}{4}\left\langle \frac{\partial^{3}V}{\partial x^{3}}\right\rangle_{q}.$$
 (17)

Finally, the cross moments evolve according to

$$\frac{d}{dt} \langle x^n p^k \rangle_q = \int dx dp \ x^n p^k \frac{\partial f_W}{\partial t} \\
= \frac{n}{m} \langle x^{n-1} p^{k+1} \rangle_q - k \left\langle x^n p^{k-1} \frac{\partial V}{\partial x} \right\rangle_q \\
+ \sum_{\lambda \ odd}^{k,k-1} \Theta(k,\lambda) \left\langle x^n p^{k-\lambda} \frac{\partial^{\lambda} V}{\partial x^{\lambda}} \right\rangle_q. (18)$$

As in Eqn. (15), the upper limit of the sum is k if k is odd, and k - 1, if it is even.

A moment expansion around the centroid can now be developed by expanding in 'fluctuations' around 'mean fields,'

$$x = \langle x \rangle_c + \delta, \tag{19}$$

$$p = \langle p \rangle_c + \eta, \tag{20}$$

in the classical case, and

$$x = \langle x \rangle_q + \delta, \tag{21}$$

$$p = \langle p \rangle_q + \eta, \tag{22}$$

in the quantum case. It is clear from the definition of the fluctuations that their expectation values vanish identically.

The strategy to follow now is simple: One begins with the lowest order classical dynamical equations from (11)-(13), or the corresponding quantum equations (15)-(18), and substitutes in the appropriate pair from equations (19)-(22) given above. One then finds terms involving expectation values of higher order fluctuations. Dynamical equations for these quantities can then be found by going to higher order in (11)-(13) or (19)-(22). Care has to be taken in interpreting the moment expansion: the classical and quantum mean field equations are not identical and the fluctuations in the quantum case contain both 'classical' and 'quantum' contributions. An analogous moment expansion was performed by Sundaram and Milonni [36] by working in a Heisenberg operator formalism. The moment hierarchy described below is completely equivalent to the set of equations they derived. The advantage here is that we can compare directly with the corresponding classical hierarchy.

To illustrate the above procedure we now carry out the moment expansion to quadratic order in the fluctuations. As will be seen below, to this order, there is no difference between the classical and quantum hierarchies (this is well-known in quantum optics, see, e.g., Ref. [50]) so it is sufficient merely to present the classical situation. The equations for the average quantities are [set n = 1 in (11) and (12)]:

$$\frac{d}{dt} \langle x \rangle_c = \frac{\langle p \rangle_c}{m}, \tag{23}$$

$$\frac{a}{dt} \langle p \rangle_c = -\left\langle \frac{\partial V}{\partial x} \right\rangle_c$$
$$= -V_c^{(1)} - \frac{1}{2!} \left\langle \delta^2 \right\rangle_c V_c^{(3)} + \cdots, \qquad (24)$$

where

$$V_c^{(n)} \equiv \left[\frac{\partial^n V(x)}{\partial x^n}\right]_{x = \langle x \rangle_c}.$$
 (25)

To obtain a closed set of equations we need to know the equations for the fluctuations correct to quadratic order. We begin by noting that

$$\langle x^2 \rangle_c = \langle x \rangle_c^2 + \langle \delta^2 \rangle_c$$
 (26)

which implies directly in turn

$$\frac{d}{dt}\left\langle x^{2}\right\rangle _{c}=2\left\langle x\right\rangle _{c}\frac{d}{dt}\left\langle x\right\rangle _{c}+\frac{d}{dt}\left\langle \delta^{2}\right\rangle _{c}.$$
(27)

But we know already from the hierarchy that [set n = 2 in (11)],

$$\frac{d}{dt} \langle x^2 \rangle_c = \frac{2}{m} \langle xp \rangle_c$$

$$= \frac{2}{m} (\langle x \rangle_c \langle p \rangle_c + \langle \delta\eta \rangle_c)$$

$$= 2 \langle x \rangle_c \frac{d}{dt} \langle x \rangle_c + \frac{2}{m} \langle \delta\eta \rangle_c. \quad (28)$$

Comparing (27) and (28), we find

$$\frac{d}{dt}\left\langle \delta^{2}\right\rangle _{c}=\frac{2}{m}\left\langle \delta\eta\right\rangle _{c} \tag{29}$$

which is the first of the required equations. Notice that the derivation of this equation is in fact exact as no terms were neglected. This is not true for the other two equations to which we now pay attention. Proceeding as above, we first write

$$\frac{d}{dt}\left\langle p^{2}\right\rangle _{c}=2\left\langle p\right\rangle _{c}\frac{d}{dt}\left\langle p\right\rangle _{c}+\frac{d}{dt}\left\langle \eta^{2}\right\rangle _{c}.$$
(30)

then setting n = 2 in (12), we find

$$\frac{d}{dt} \langle p^2 \rangle_c = -2 \left\langle p \frac{\partial V}{\partial x} \right\rangle_c$$

$$= -2 \left[\langle p \rangle_c V_c^{(1)} + \frac{1}{2!} \langle p \rangle_c \langle \delta^2 \rangle_c V_c^{(3)} + \langle \eta \delta \rangle_c V_c^{(2)} + \cdots \right]$$

$$= 2 \langle p \rangle_c \frac{d}{dt} \langle p \rangle_c - 2 \langle \eta \delta \rangle_c V_c^{(2)} + \cdots \quad (31)$$

Comparing (30) and (31), we obtain

$$\frac{d}{dt}\left\langle \eta^{2}\right\rangle _{c}=-2\left\langle \eta\delta\right\rangle _{c}V_{c}^{(2)}.$$
(32)

It is now apparent how to proceed to get the last equation and we merely write the final result

$$\frac{d}{dt} \left\langle \eta \delta \right\rangle_c = \frac{\left\langle \eta^2 \right\rangle_c}{m} - \left\langle \delta^2 \right\rangle_c V_c^{(2)}. \tag{33}$$

Equations (23), (24), (29), (32), and (33) form a closed set of dynamical equations. Since at most only second order moments were used in the derivation, these five classical equations are formally identical to their quantum counterparts. Had we truncated the moment expansion at one higher order there would have been a difference between the quantum and classical equations since, as we have seen, the first quantum dynamical correction comes in only at third order.

General expressions can be written for the exact classical and quantum moment hierarchies by extending the procedure outlined above. In the classical case, restricting attention only to those moments relevant to future discussion of the Gaussian approximation, one finds,

$$\frac{d}{dt} \langle \delta^n \rangle_c = \frac{n}{m} \langle \delta^{n-1} \eta \rangle_c, \qquad (34)$$

$$\frac{d}{dt} \langle \eta^n \rangle_c = n \left[\langle \eta^{n-1} \rangle_c \left\langle \frac{\partial V}{\partial x} \right\rangle_c - \left\langle \eta^{n-1} \frac{\partial V}{\partial x} \right\rangle_c \right], \qquad (35)$$

$$\frac{d}{dt} \left\langle \delta^n \eta \right\rangle_c = \left\langle \delta^n \right\rangle_c \left\langle \frac{\partial V}{\partial x} \right\rangle_c - \left\langle \delta^n \frac{\partial V}{\partial x} \right\rangle_c + \frac{n}{m} \left\langle \delta^{n-1} \eta^2 \right\rangle_c. \quad (36)$$

The quantum hierarchy is only slightly more complicated. For the same moments considered above,

$$\frac{d}{dt} \langle \delta^n \rangle_q = \frac{n}{m} \langle \delta^{n-1} \eta \rangle_q,$$

$$\frac{d}{dt} \langle \eta^n \rangle_q = n \left[\langle \eta^{n-1} \rangle_q \left\langle \frac{\partial V}{\partial x} \right\rangle \right]$$
(37)

$$-\left\langle \eta^{n-1} \frac{\partial V}{\partial x} \right\rangle_{q} + \Delta^{(n)}, \qquad (38)$$

$$\frac{a}{dt} \left\langle \delta^n \eta \right\rangle_q = \left\langle \delta^n \right\rangle_q \left\langle \frac{\partial V}{\partial x} \right\rangle_q - \left\langle \delta^n \frac{\partial V}{\partial x} \right\rangle_q + \frac{n}{m} \left\langle \delta^{n-1} \eta^2 \right\rangle_q, \qquad (39)$$

where

J

$$\Delta^{(n)} = \sum_{\lambda \text{ odd}}^{n,n-1} \Theta(n,\lambda) \left\langle \left(\langle p \rangle_q + \eta \right)^{n-\lambda} \frac{\partial^{\lambda} V}{\partial x^{\lambda}} \right\rangle_q.$$
(40)

III. THE GAUSSIAN APPROXIMATION

A. Time Dependent Variational Principle

We now proceed to the first variant of the Gaussian approximation. The idea here is to (i) assume that the distribution function is at all times a Gaussian (for more mathematical details on the formal properties of such states, see Refs. [51, 52]) and then, (ii) to write down the corresponding dynamical equations for the moment hierarchy. The approach given here is a generalization of the time dependent variational principle first considered by Dirac [26] by considering mixed as well as pure states. The moment hierarchy approach is far more convenient than a direct application of a variational principle as carried out, for example, in Ref. [53]. As will be apparent in what follows, this approximation only samples the moment hierarchy to quadratic order, and hence contains no dynamical quantum corrections.

We assume the system to be always described by a general Gaussian distribution (here we make no distinction between classical and quantum) of the form

$$f = \frac{1}{N} \exp \left[-a(x - \bar{x}(t))^2 - b(p - \bar{p}(t))^2 + c(x - \bar{x}(t))(p - \bar{p}(t)) \right],$$
(41)

or in a more compact notation:

$$f = \frac{1}{\sqrt{Det(2\pi S)}} \exp\left[-\frac{1}{2} \sum_{j,k} (x_j - \bar{x}_j) S_{jk}^{-1}(x_k - \bar{x}_k)\right],$$
(42)

with $x_1 = x$, $x_2 = p$, and the matrices,

$$S^{-1} = \begin{pmatrix} 2a & -c \\ -c & 2b \end{pmatrix}, \tag{43}$$

$$S = \frac{1}{4ab - c^2} \begin{pmatrix} 2b & c \\ c & 2a \end{pmatrix}.$$
 (44)

The distribution is normalized to unity over phase space, i.e.,

$$N = \sqrt{Det(2\pi S)} = \frac{2\pi}{\sqrt{4ab - c^2}},\tag{45}$$

and satisfies the constraint (7). This extra condition in turn leads to a constraint on the parameters of the distribution:

$$4ab - c^2 = (4\pi\sigma_2)^2. \tag{46}$$

This purely kinematical condition must be satisfied for quantum distributions. In general, classical distributions need not satisfy this condition, but if they do, then, as already discussed, this condition is exactly preserved by the classical dynamics.

Given the distribution function specified above, it is trivial to show that

$$\langle x \rangle = \bar{x}, \quad \langle p \rangle = \bar{p}, \tag{47}$$

and that the quadratic moments are

$$\left< \delta^2 \right> = S_{11} = \frac{1}{2\alpha}, \quad \alpha = a - \frac{c^2}{4b} = \frac{4\pi^2 \sigma_2^2}{b}, \quad (48)$$

$$\langle \eta^2 \rangle = S_{22} = \frac{1}{2\beta}, \quad \beta = b - \frac{c^2}{4a} = \frac{4\pi^2 \sigma_2^2}{a}, \quad (49)$$

$$\langle \eta \delta \rangle = S_{12} = \frac{c}{4a\beta} = \frac{c}{(4\pi\sigma_2)^2}.$$
(50)

The kinematical constraint (46) can now be expressed as

$$\langle \eta^2 \rangle \langle \delta^2 \rangle = \langle \eta \delta \rangle^2 + \frac{1}{(4\pi\sigma_2)^2},$$
 (51)

which is a statement of the uncertainty principle generalized to mixed states. For pure states, $\sigma_2 = 1/2\pi\hbar$, and (51) takes the familiar form (the wave function is now that of some Gaussian state, more general than just a squeezed coherent state)

$$\left\langle \eta^{2} \right\rangle \left\langle \delta^{2} \right\rangle = \left\langle \eta \delta \right\rangle^{2} + \frac{\hbar^{2}}{4}.$$
 (52)

All odd-order moments are zero and the higher order even moments can be computed from the quadratic ones by a simple application of the result, valid for Gaussian distributions,

$$\langle (x_j - \bar{x}_j)g(\mathbf{x}) \rangle = \sum_k S_{jk} \left\langle \frac{\partial g(\mathbf{x})}{\partial x_k} \right\rangle,$$
 (53)

where $\mathbf{x} = (x_1, x_2)$ and $g(\mathbf{x})$ is some arbitrary function of \mathbf{x} . The moments relevant to us are

$$\langle \delta^{2n} \rangle = \frac{(2n-1)!}{2^{n-1}(n-1)!} \langle \delta^2 \rangle^n,$$
 (54)

$$\langle \eta^{2n} \rangle = \frac{(2n-1)!}{2^{n-1}(n-1)!} \langle \eta^2 \rangle^n,$$
 (55)

$$\left\langle \eta \delta^{2n+1} \right\rangle = \frac{(2n+1)!}{2^n n!} \left\langle \eta \delta \right\rangle \left\langle \delta^2 \right\rangle^n,$$
 (56)

where (51) has been used to simplify the last expression. The above expressions are the only ones needed to obtain the moment hierarchy in Gaussian approximation.

The dynamical equations under the Gaussian approximation are easy to obtain. First,

$$\frac{d}{dt}\langle x\rangle = \frac{\langle p\rangle}{m} \tag{57}$$

or

$$\frac{d}{dt}\bar{x} = \frac{\bar{p}}{m} \tag{58}$$

and

$$\frac{d}{dt} \langle p \rangle = \frac{d}{dt} \bar{p} = -\left\langle \frac{\partial V}{\partial x} \right\rangle$$

$$= -V^{(1)} - \frac{\left\langle \delta^2 \right\rangle}{2!} V^{(3)} - \frac{\left\langle \delta^3 \right\rangle}{3!} V^{(4)} - \cdots$$

$$= -\sum_{n=0}^{\infty} \frac{\left\langle \delta^2 \right\rangle^n}{n! 2^n} V^{(2n+1)} \tag{59}$$

where we have used (54) in the last step. The dynamical equation for $\langle \delta^2 \rangle$ is nothing but (29) which is already exact. The equation for $\langle \eta^2 \rangle$ follows from (35) or (38):

$$\frac{d}{dt} \langle \eta^2 \rangle = -2 \left\langle \eta \frac{\partial V}{\partial x} \right\rangle$$

$$= -2 \left[\left\langle \eta \left(V^{(1)} + \delta V^{(2)} + \frac{\delta^2}{2!} V^{(3)} + \frac{\delta^3}{3!} V^{(4)} + \cdots \right) \right\rangle \right]$$

$$= -\sum_{k=0}^{\infty} \frac{V^{(2k+2)}}{2^{k-1}k!} \langle \eta \delta \rangle \left\langle \delta^2 \right\rangle^k.$$
(60)

This result has been written in a simple form by using (56) along with the constraint (51). The last equation we have to deal with is for $\langle \delta \eta \rangle$, this following from (36):

$$\frac{d}{dt} \langle \delta \eta \rangle = \frac{1}{m} \langle \eta^2 \rangle - \left\langle \delta \frac{\partial V}{\partial x} \right\rangle$$

$$= \frac{1}{m} \langle \eta^2 \rangle - \left\langle \delta \left(V^{(1)} + \delta V^{(2)} + \frac{\delta^2}{2!} V^{(3)} + \frac{\delta^3}{3!} V^{(4)} + \cdots \right) \right\rangle$$

$$= \frac{1}{m} \langle \eta^2 \rangle - \sum_{n=0}^{\infty} \frac{\langle \delta^2 \rangle^{n+1}}{2^n n!} V^{(2n+2)}.$$
(61)

Note that we have used (54) in the last step.

We now have five equations, two for the mean fields, and three for the fluctuations. It is easy to check that the constraint (52) is exactly preserved by the dynamical equations. Therefore the constraint may be used to eliminate one of the three dynamical equations for the fluctuations. For example, if we choose to keep $\langle \delta^2 \rangle$ and $\langle \delta \eta \rangle$ as the preferred variables, then the two dynamical equations for the fluctuations are (29) and (61) with $\langle \eta^2 \rangle$ written in terms of $\langle \delta^2 \rangle$ and $\langle \delta \eta \rangle$ using (51). Notice that these equations were obtained without having to go to moments higher than those of order two. Therefore they do not contain any quantum dynamical contributions, just as advertized. Of course, \hbar can still appear in these equations, but in a purely kinematical sense. This will happen if the initial state is pure $(\sigma_2 = 1/2\pi\hbar)$ and we use the constraint to eliminate one of the dynamical equations for the fluctuations.

To conclude our discussion of the TDVP, we note that this approximation is conservative and further, can be mapped to a classical Hamiltonian for a new set of dynamical variables [54]. The first point can be easily checked by writing the expectation value of the Hamiltonian (here assumed explicitly time independent) in Gaussian approximation,

and then verifying, using the dynamical equations given above, that

$$\frac{d}{dt}\left\langle H\right\rangle = 0. \tag{63}$$

It is interesting to note that any consistent truncation of the TDVP is also conservative (by consistent truncation, we mean: keep only the moments up to some finite order, i.e., go up to the *same* finite order in all the equations). As it happens, Heller's approximation [27] is not consistent in this sense because it does not account for the back-reaction on the mean field thereby missing a quadratic term that should have been kept; consequently it fails to be conservative.

It is easy to show that the four dynamical equations for the centroid and two chosen fluctuation variables are in fact Hamiltonian. Eliminating $\langle \eta^2 \rangle$ from (61) using the constraint (51) and introducing the new variables [54][34]

$$\left< \delta^2 \right> \equiv \rho^2 ; \quad \left< \eta \delta \right> \equiv \rho \gamma, \tag{64}$$

in (62), we find that the Hamiltonian

$$H_{G} = \langle H \rangle = \frac{1}{2m} \bar{p}^{2} + \frac{1}{2m} \gamma_{0}^{2} + \frac{1}{2m\rho^{2}(4\pi\sigma_{2})^{2}} + \sum_{n=1}^{\infty} \frac{\rho^{2n}}{2^{n}n!} V^{(2n)}(\bar{x})$$
(65)

generates the correct equations of motion as can be easily verified. It follows trivially from (65) that any consistent TGA is also Hamiltonian.

The expectation value of the Hamiltonian (62) is precisely the quantity that is minimized in the Gaussian effective potential approach [55]. Since \hbar appears only in kinematical guise in (62) it follows that no essential quantum effects (i.e., those without formal classical analogs) can be seen in this approximation. For example, tunneling/overlap corrections in a double-well system are not accessible in simple Gaussian approximation [56].

B. Semiquantum Chaos and Related Issues

In Ref. [34], Pattanayak and Schieve studied a onedimensional anharmonic oscillator in Gaussian approximation. (See also Ref. /citeC2.) Even though the original system is clearly not chaotic, in the approximation it can be, because we now have four coupled nonlinear dynamical equations, rather than the original two. This is the main technical point of their paper and is, of course, correct. The authors stressed the surprising nature of their result, that dynamics of expectation values may be chaotic, "even though the system has regular classical behavior and the quantum behavior has been assumed regular" [34]. Their conclusion has been criticized by Sundaram and Milonni [36] who point out that the Gaussian approximation becomes unreliable very quickly when either the classical system is chaotic, or when the Gaussian dynamical equations are themselves chaotic. The more general theoretical setting is the following: Any variational approximation to the dynamics of a quantum system based on the Dirac action principle leads to a classical Hamiltonian dynamics for the variational parameters [57]. Since this Hamiltonian is generically nonlinear and nonintegrable, the dynamics thus generated can be chaotic, in distinction to the exact quantum evolution [28, 35]; the Gaussian approximation represents a particularly simple special case.

Once we realize that, in fact, the analysis presented in Refs. [33] and [34] is purely classical, the results are easy to understand. First, they are not surprising in the sense that quantum mechanics is *not* somehow catalyzing chaos. Second, we see why the approximation breaks down as it does: if the original system is chaotic then, 'tendrilization,' in the sense of Berry and Balazs [8] will happen on a timescale set by the maximal Lyapunov exponent by which time the Gaussian approximation is definitely wrong. If, on the other hand, the original classical system is not chaotic, but the Gaussian set of equations are, all this tells us is that the non-Gaussian corrections are becoming very important on the timescale set by the maximal Lyapunov exponent of the false chaos, and need to be included to obtain the correct classical dynamics. Third, the explanation that quantum noise is somehow promoting the chaos is not tenable since the Gaussian approximation contains only classical fluctuations.

An ostensibly opposite result was obtained by Zhang and Feng [38]: These authors implemented the TDVP approach for the kicked rotor, finding a suppression of the classical chaos. This, they claimed, was a simple example of a generic suppression of classical chaos by quantum mechanics. Our interpretation of their result is different: since it is a consequence of merely classical dynamics, what they have really displayed is a suppression of chaos by the classical back-reaction, computed in Gaussian approximation. Thus, there really is no information about quantum suppression (or even of a classical 'fluctuation suppression,' since the calculation is only approximate). Real quantum suppression of classical chaos in this model is discussed in Refs. [58, 59].

C. The Truncated Gaussian Approximation

The full Gaussian approximation (TDVP) involves summations to all orders of the fluctuations (though these are somewhat trivial as the higher terms that are summed are disconnected pieces, i.e., no cumulants of order higher than two are encountered). This approximation differs from a truncation approach where one merely truncates the moment hierarchy at some finite order. We can incorporate the truncation idea in Gaussian approximation by requiring the distribution to be Gaussian but still stopping the summations at some finite order. As discussed above, this truncation is conservative and Hamiltonian in the sense that the corresponding equations of motion for dynamical variables can be derived from a Hamiltonian. The TGA may be useful in some situations because of its computational simplicity.

An example of this idea is the approach due to Heller [27]. In our language, Heller's instructions are the following: (i) Drop the fluctuation corrections to the equation for the average momentum (59), and (ii) keep only the leading order (quadratic) corrections in the equations for the moments [Eqns. (29), (60), and (61)]. The motivation for this approach as originally stated is to provide a semiclassical limit that does not quite correspond to the usual $\hbar \to 0$ limit of WKB theory (For example, the quadratic order TGA is exact for the harmonic oscillator, but WKB is not). As will become clear in what follows, dynamically this is still a classical approach and its ostensible quantum features are easily explained: (i) the appearance of \hbar is purely kinematical, and (ii) "penetration" of the Gaussian into classically forbidden territory is simply a classical consequence of the distribution function having support at high momenta. It is no surprise that this approximation has trouble with tunneling (which is accessible in a WKB approach) since it does not incorporate quantum dynamical corrections.

By carrying through with the prescription given above, we obtain the dynamical equations:

$$\frac{d}{dt}\bar{x} = \frac{\bar{p}}{m},\tag{66}$$

$$\frac{d}{dt}\bar{p} = -V^{(1)},\tag{67}$$

$$\frac{d}{dt}\left\langle \delta^{2}\right\rangle = \frac{2}{m}\left\langle \delta\eta\right\rangle,\tag{68}$$

$$\frac{d}{dt}\langle\delta\eta\rangle = \frac{1}{m}\langle\eta^2\rangle - \langle\delta^2\rangle V^{(2)},\tag{69}$$

$$\frac{d}{dt} \langle \eta^2 \rangle = -2 \langle \delta \eta \rangle V^{(2)}. \tag{70}$$

The first two equations (62) and (63) are just the classical equations of motion for a particle in the potential V(x). This approximation thus ignores the back-reaction of fluctuations on the mean field. As for the fluctuations

themselves, they are sensitive only to the second derivative of the potential, as in the quadratic approximation of Section IIB. Heller's method is not equivalent to a quadratic truncation because there is no back-reaction on the mean fields and because it still enforces the constraint (51), reducing the number of dynamical equations from five to four. However, it is not a consistent quadratic TGA because of the failure to include the back-reaction. Just as in the TDVP and quadratic approximations, this approximation only includes classical dynamics.

As a consequence of the lack of self-consistency at quadratic order, the expectation value of the energy is not conserved:

$$\frac{d}{dt}\langle H\rangle = \frac{\langle p\rangle}{2m} V^{(3)} \left\langle \delta^2 \right\rangle. \tag{71}$$

It is a trivial matter to improve this approximation by keeping the back-reaction consistently to quadratic order. (Note, however, that this approximation is conservative for linear systems.) Even though this method is not unitary, it has been applied very successfully, especially in its multiple trajectory variant, to chaotic systems [13, 14].

The implementation of the TGA in the Schrödinger picture also involves taking into account a time varying global (Maslov) phase for the wave function. Since the Wigner function is a transform of the density matrix, it does not contain this phase information. If there was more than one packet (as in MTGA), however, the Wigner function would keep relative phase information for which there is no classical analog.

D. Connection with Lyapunov Exponents

A straightforward observation is that the set of equations (66) - (70) are exactly those needed to determine the classical Lyapunov exponent, with (\bar{x}, \bar{p}) specifying a fiducial classical trajectory, and the fluctuation variables specifying the linearized evolution of a Gaussian ball of initial conditions around the fiducial trajectory. As before, imposition of the constraint arising from the constancy of σ_2 leaves only two equations for the fluctuations, and these are Hamiltonian separately, with

$$H_F = \frac{1}{2m}\gamma^2 + \frac{1}{2m\rho^2(4\pi\sigma_2)^2} + \frac{1}{2}V^{(2)}(\bar{x})\rho^2.$$
 (72)

This would be precisely the linearized Hamiltonian for the system were it not for the second term which enforces the constraint. The definition of the Lyapunov exponent requires the double limit of first squeezing the initial Gaussian ball onto the fiducial trajectory (in *all* phase space directions) and then taking the limit $t \to \infty$. The first limit corresponds to taking $\sigma_2 \to \infty$ (clearly not allowed quantum mechanically for \hbar finite), which is the same as dropping the second term in H_F . Incidentally, this is also a good way to see how chaos can occur in the Liouville theory despite the fact that the classical transport equation is linear (like the Schrödinger equation or the quantum Liouville equation): it arises simply because the delta function or trajectory limit is allowed classically, but not quantum mechanically.

E. The Multiple Trajectory Gaussian Approximation

The idea behind the MTGA is to improve the TGA by including multiple classical trajectories. One does this by propagating more than one Gaussian, using the TGA for each individual packet. There are two reasons why this is a good idea. First, a sum of Gaussians can track the classical distribution even at times when the distribution is highly non-Gaussian (although this may involve a high computational cost as very many Gaussians might be needed in the sum). The second issue involves quantum coherence in phase space. The classical version of this approximation simply sums over the individual Gaussians to give the composite distribution function. This is not true in the quantum case: a coherent sum of Gaussian wave functions, it turns out, is not a sum of the associated Wigner functions, but rather a sum augmented by "off-diagonal" quantum coherence terms [6]. (These off-diagonal terms account for the Airy fringes in the WKB approximation for the Wigner function discussed by Berry [4].) Thus the MTGA contains nontrivial quantum information, and is in this sense, unlike TDVP and TGA, not a classical approximation.

IV. SUMMARY

The comparative evolution of c-number classical and quantum moments yields an obvious method to compare classical and quantum mechanics as dynamical theories. In doing so here, we have restricted attention largely to the case of the Gaussian approximation. The timedependent Gaussian approximation in quantum mechanics has a direct, completely equivalent, classical analog in terms of an ensemble of classical particles also treated in the same approximation. Thus every 'quantum' result in this approximation has an exact classical counterpart. Part of the reason for the misinterpretation of 'semiquantum' chaos as a quantum dynamical effect was simply due to a too naive comparison with classical mechanics: a comparison against trajectories, rather than against swarms of trajectories. Using the more general approach presented here, it becomes clear that, in this context, \hbar plays no fundamental role. The key point is that a dimensionful constant *also* appears in classical transport theory: σ_2 , as defined in Eqn. (4), which plays the role of an effective \hbar there – such a constant is missing from the equations of motion of classical trajectories but arises once classical distributions have to be evolved by the Liouville equation.

V. ACKNOWLEDGEMENTS

It is a pleasure to acknowledge helpful discussions (over a rather extended period of time!) with Tanmoy Bhattacharya, Fred Cooper, Tim Elston, Ronald Fox, Kurt Jacobs, Ronnie Mainieri, Peter Milonni, Arjendu Pattanayak, Daniel Steck, Kosuke Shizume, Bala Sundaram, and George Zaslavsky.

- [1] E.P. Wigner, Phys. Rev. 40, 749 (1932).
- [2] J.H. Irving and R.W. Zwanzig, J. Chem. Phys. 19, 1173 (1951).
- [3] I. Oppenheim and J. Ross, Phys. Rev. 107, 28 (1957).
- [4] M.V. Berry, Philos. Trans. R. Soc. 287, 237 (1977); J. Phys. A 10, 2083 (1977).
- [5] A. Voros, Ann. Inst. Henri Poincaré 24, 31 (1976); *ibid* 26, 343 (1977).
- [6] E.J. Heller, J. Chem. Phys. **67**, 3339 (1977).
- [7] E.J. Heller, J. Chem. Phys. 65, 1289 (1976).
- [8] M.V. Berry and N.L. Balazs, J. Phys. A 12,625 (1979).
- [9] R.F. Fox, Phys. Rev. A 41, 2969 (1990).
- [10] S. Habib, K. Shizume, and W.H. Zurek, Phys. Rev. Lett. 80, 4361 (1998).
- [11] B.D. Greenbaum, S. Habib, K. Shizume, and B. Sundaram, quant-ph/0401174.
- [12] See, e.g., T. Bhattacharya, S. Habib, and K. Jacobs, Phys. Rev. Lett. 85, 4852 (2000); Phys. Rev. A 67, 042103 (2003); and references therein.
- [13] E.J. Heller, J. Chem. Phys. 94, 2723 (1991).

- [14] S. Tomsovic and E.J. Heller, Phys. Rev. Lett. 67, 664 (1991).
- [15] P.W. O'Connor, S. Tomsovic, and E.J. Heller, Physica D 55, 340 (1992).
- [16] D. Provost and P. Brumer, Phys. Rev. Lett. 74, 250 (1995).
- [17] L.E. Ballentine, Y. Yang, and J.P. Zibin, Phys. Rev. A 50, 2854 (1994).
- [18] S. Habib, K. Jacobs, H. Mabuchi, R.D. Ryne, K. Shizume, and B. Sundaram, Phys. Rev. Lett. 88, 040402 (2002).
- [19] A.J. Dragt and S. Habib, in *Quantum Aspects of Beam Physics*, Proceedings of the 1998 Advanced ICFA Beam Dynamics Workshop (World Scientific, New York, 1999); quant-ph/9806056.
- [20] R.P. Feynman, The Feynman Lectures on Physics, Vol. III (Addison-Wesley, New York, 1965).
- [21] R.F. Fox and T.C. Elston, Phys. Rev. E 49, 3683 (1994); *ibid* 50, 2553 (1994).
- [22] B.S. Helmkamp and D.A. Browne, Phys. Rev. E 49, 1831

(1994).

- [23] F.L. Moore, J.C. Robinson, C. Bharucha, P.E. Williams, and M.G. Raizen, Phys. Rev. Lett. **73**, 2974 (1994);
 D.A. Steck, W.H. Oskay, and M.G. Raizen, Science **293**, 274 (2001).
- [24] T. Bhattacharya, S. Habib, K. Jacobs, and K. Shizume, Phys. Rev. A 65, 032115 (2002).
- [25] For previous derivations of the moment hierarchies, see, M. Andrews and M. Hall, J. Phys. A 18, 37 (1985); L.E. Ballentine and S.M. McRae, Phys. Rev. A 58, 1799 (1998).
- [26] P.A.M. Dirac, Proc. Cambridge Philos. Soc. 26, 376 (1930). See also, F. Cooper, S.-Y. Pi, and P.N. Stancioff, Phys. Rev. D 34, 3831 (1986), and references therein.
- [27] E.J. Heller, J. Chem. Phys. **62**, 1544 (1975).
- [28] F. Cooper, J. Dawson, S. Habib, and R.D. Ryne, Phys. Rev. E, 57, 1489 (1998).
- [29] G.W. Ford, M. Kac, and P. Mazur, J. Math. Phys. 6, 504 (1965); R. Zwanzig, J. Stat. Phys. 9, 215 (1973);
 A.O. Caldeira and A.J. Leggett, Physica 121A, 587 (1983); Ann. Phys. (N.Y.) 149, 374 (1983); B.L. Hu, J.P. Paz, and Y. Zhang, Phys. Rev. D 45, 2843 (1992).
- [30] J. Anglin and S. Habib, Mod. Phys. Lett. A 11, 2655 (1996).
- [31] F. Cooper, S. Habib, Y. Kluger, E. Mottola, J.P. Paz, and P.R. Anderson, Phys. Rev. D 50, 2848 (1994);
 F. Cooper, J. Dawson, S. Habib, Y. Kluger, D. Meredith, and H. Shepard, Physica D 83, 74 (1995); D. Boyanovsky,
 H.J. de Vega, R. Holman, D.S. Lee, and A. Singh, Phys. Rev. D 51, 4419 (1995); J. Baacke and K. Heitmann,
 Phys. Rev. D 62, 105022 (2000).
- [32] B. Mihaila, T. Athan, F. Cooper, J. Dawson, and S. Habib, Phys. Rev. D 62, 125015 (2000).
- [33] F. Cooper, J. Dawson, D. Meredith, and H. Shepard, Phys. Rev. Lett. **72**, 1337 (1994).
- [34] A.K. Pattanayak and W.C. Schieve, Phys. Rev. Lett. 72, 2855 (1994).
- [35] E. Caurier, S. Drożdż, J. Okolowicz, and M. Ploszajczak, Acta Phys. Pol. B22, 389 (1991).
- [36] B. Sundaram and P. Milonni, Phys. Rev. E 51, 1971 (1995).
- [37] L.E. Ballentine, Phys. Rev. E 63, 056204 (2001).
- [38] W.M. Zhang and D.H. Feng, Mod. Phys. Lett. A 8, 1417 (1993).

- [39] S. Habib, K. Jacobs, and K. Shizume, in preparation.
- [40] S. Habib and K. Shizume (unpublished).
- [41] A.C. Doherty and K. Jacobs, Phys. Rev. A **60**, 2700 (1999); A.C. Doherty, S. Habib, K. Jacobs, H. Mabuchi, and S.M. Tan, Phys. Rev. A **62**, 012105 (2000).
- [42] A. Hopkins, K. Jacobs, S. Habib, and K. Schwab, Phys. Rev. B 68, 235328 (2003).
- [43] D.A. Steck, K. Jacobs, H. Mabuchi, T. Bhattacharya, and S. Habib, Phys. Rev. Lett. (to appear); quant-ph/0310153
- [44] M. Hillery, R. O'Connell, M.O. Scully, and E.P. Wigner, Phys. Rep. 106, 121 (1984).
- [45] N.L. Balazs and B.K. Jennings, Phys. Rep. 104, 347 (1984).
- [46] V.I. Tatarskii, Usp. Fiz. Nauk **139**, 587 (1983) (Sov. Phys. Usp. **26**, 311 (1983)).
- [47] S. Habib, Phys. Rev. D 42, 2566 (1990).
- [48] R.L. Hudson, Rep. Math. Phys. 6, 249 (1974).
- [49] F.J. Narcowich, in Seminars in Mathematical Physics (Texas A&M Univesity, College Station, 1986). For a more general discussion of operator ordering issues, see J.R. Shewell, Am. J. Phys. 87, 16 (1959).
- [50] C. Fabre and S. Reynaud, in Fundamental Systems in Quantum Optics, Lectures at the Les Houches Summer School, NATO Advanced Study Institute, 25 June - 27 July, 1990, edited by J. Dalibard, J.-M. Raimond, and J. Zinn-Justin (North-Holland, Amsterdam, 1992).
- [51] R.G. Littlejohn, Phys. Rep. **138**, 193 (1986).
- [52] B.L. Schumaker, Phys. Rep. **135**, 317 (1986).
- [53] O. Éboli, R. Jackiw, and S.-Y. Pi, Phys. Rev. D 37, 3557 (1988).
- [54] A.K. Rajagopal and J.T. Marshall, Phys. Rev. A 26, 2977 (1982) where only the pure state case was considered.
- [55] See, e.g., P.M. Stevenson, Phys. Rev. D 30, 1712 (1984).
- [56] See, e.g., F.J. Alexander, S. Habib, and A. Kovner, Phys. Rev. D 48, 4284 (1993).
- [57] K.-K. Kan, Phys. Rev. A 24, 2831 (1981).
- [58] B.V. Chirikov, F.D. Izrailev, and D.L. Shepelyansky, Sov. Sci. Rev. Sect. C 2, 209 (1981).
- [59] S. Fishman, D.R. Grempel, and R.E. Prange, Phys. Rev. Lett. 49, 509 (1982).