Spectra and symmetric eigentensors of the Lichnerowicz Laplacian on S^n

M. Boucetta
Faculté des Sciences et Techniques
BP 549 Marrakech
Morocco

Email: boucetta@fstg-marrakech.ac.ma *

Abstract. We compute the eigenvalues with multiplicities of the Lichnerowicz Laplacian acting on the space of symmetric covariant tensor fields on the Euclidian sphere S^n . The spaces of symmetric eigentensors are explicitly given.

Mathematical Subject Classification (2000):53B21, 53B50, 58C40 Key words: Lichnerowicz Laplacian

1 Introduction

Let (M, g) be a Riemannian n-manifold. For any $p \in \mathbb{N}$, we shall denote by $\Gamma(\otimes^p T^*M)$, $\Omega^p(M)$ and $\mathcal{S}^p M$ the space of covariant p-tensor fields on M, the space of differential p-forms on M and the space of symmetric covariant p-tensor fields on M, respectively. Note that $\Gamma(\otimes^0 T^*M) = \Omega^0(M) = \mathcal{S}^0 M = \mathcal{S}^0 M$

$$C^{\infty}(M, \mathbb{R}), \ \Omega(M) = \sum_{p=0}^{n} \Omega^{p}(M) \text{ and } \mathcal{S}(M) = \sum_{p\geq 0} \mathcal{S}^{p}(M).$$

Let D be the Levi-Civita connection associated to g; its curvature tensor field R is given by

$$R(X,Y)Z = D_{[X,Y]}Z - (D_XD_YZ - D_YD_XZ),$$

^{*}Recherche menée dans le cadre du Programme Thématique d'Appui à la Recherche Scientifique PROTARS III.

and the Ricci endomorphism field $r:TM\longrightarrow TM$ is given by

$$g(r(X), Y) = \sum_{i=1}^{n} g(R(X, E_i)Y, E_i),$$

where (E_1, \ldots, E_n) is any local orthonormal frame.

For any $p \in \mathbb{N}$, the connection D induces a differential operator $D: \Gamma(\otimes^p T^*M) \longrightarrow \Gamma(\otimes^{p+1} T^*M)$ given by

$$DT(X, Y_1, \dots, Y_p) = D_X T(Y_1, \dots, Y_p) = X.T(Y_1, \dots, Y_p) - \sum_{j=1}^p T(Y_1, \dots, D_X Y_j, \dots, Y_p).$$

Its formal adjoint $D^*: \Gamma(\otimes^{p+1}T^*M) \longrightarrow \Gamma(\otimes^pT^*M)$ is given by

$$D^*T(Y_1,\ldots,Y_p) = -\sum_{i=1}^n D_{E_i}T(E_i,Y_1,\ldots,Y_p),$$

where (E_1, \ldots, E_n) is any local orthonormal frame. Recall that, for any differential p-form α , we have

$$d\alpha(X_1, \dots, X_{p+1}) = \sum_{j=1}^{p+1} (-1)^{j+1} D_{X_j} \alpha(X_1, \dots, \hat{X}_j, \dots, X_{p+1}).$$
 (1)

We denote by δ the restriction of D^* to $\Omega(M) \oplus \mathcal{S}(M)$ and we define δ^* : $\mathcal{S}^p(M) \longrightarrow \mathcal{S}^{p+1}(M)$ by

$$\delta^*T(X_1,\ldots,X_{p+1}) = \sum_{j=1}^{p+1} D_{X_j}T(X_1,\ldots,\hat{X}_j,\ldots,X_{p+1}).$$

Recall that the operator trace $Tr: \mathcal{S}^p(M) \longrightarrow \mathcal{S}^{p-2}(M)$ is given by

$$TrT(X_1, \dots, X_{p-2}) = \sum_{j=1}^{n} T(E_j, E_j, X_1, \dots, X_{p-2}),$$

where (E_1, \ldots, E_n) is any local orthonormal frame.

The Lichnerowicz Laplacian is the second order differential operator

$$\Delta_M: \Gamma(\otimes^p T^*M) \longrightarrow \Gamma(\otimes^p T^*M)$$

given by

$$\Delta_M(T) = D^*D(T) + R(T),$$

where R(T) is the curvature operator given by

$$R(T)(Y_1, \dots, Y_p) = \sum_{j=1}^p T(Y_1, \dots, r(Y_j), \dots, Y_p)$$

$$- \sum_{i < j} \sum_{l=1}^n \left\{ T(Y_1, \dots, E_l, \dots, R(Y_i, E_l) Y_j, \dots, Y_p) + T(Y_1, \dots, R(Y_j, E_l) Y_i, \dots, E_l, \dots, Y_p) \right\},$$

where (E_1, \ldots, E_n) is any local orthonormal frame and, in

$$T(Y_1,\ldots,E_l,\ldots,R(Y_i,E_l)Y_j,\ldots,Y_p),$$

 E_l takes the place of Y_i and $R(Y_i, E_l)Y_j$ takes the place of Y_j .

This differential operator, introduced by Lichnerowicz in [15] pp. 26, is self-adjoint, elliptic and respects the symmetries of tensor fields. In particular, Δ_M leaves invariant $\mathcal{S}(M)$ and the restriction of Δ_M to $\Omega(M)$ coincides with the Hodge-de Rham Laplacian, i.e., for any differential p-form α ,

$$\Delta_M \alpha = (d\delta + \delta d)(\alpha). \tag{2}$$

We have shown in [6] that, for any symmetric covariant tensor field T,

$$\Delta_M(T) = (\delta \circ \delta^* - \delta^* \delta)(T) + 2R(T). \tag{3}$$

Note that if $T \in \mathcal{S}(M)$ and g^l denotes the symmetric product of l copies of the Riemannian metric g, we have

$$(Tr \circ \Delta_M)T = (\Delta_M \circ Tr)T, \tag{4}$$

$$\Delta_M(T \odot g^l) = (\Delta_M T) \odot g^l, \tag{5}$$

where \odot is the symmetric product.

The Lichnerowicz Laplacian acting on symmetric covariant tensor fields is of fundamental importance in mathematical physics (see for instance [9], [20] and [22]). Note also that the Lichenrowicz Laplacian acting on symmetric covariant 2-tensor fields appears in many problems in Riemannian geometry (see [3], [5], [19]...).

On a Riemannian compact manifold, the Lichnerowicz Laplacian Δ_M has discrete eigenvalues with finite multiplicities. For a given Riemannian compact manifold, it may be an interesting problem to determine explicitly the eigenvalues and the eigentensors of Δ_M on M.

Let us enumerate the cases where the spectra of Δ_M was computed:

- 1. Δ_M acting on $C^{\infty}(M, \mathbb{C})$: M is either flat toris or Klein bottles [4], M is a Hopf manifolds [1];
- 2. Δ_M acting on $\Omega(M)$: $M = S^n$ or $P^n(\mathbb{C})$ [10] and [11], $M = \mathbb{C}aP^2$ or $G_2/SO(4)$ [16] and [18], $M = SO(n+1)/SO(2) \times SO(n)$ or $M = Sp(n+1)/Sp(1) \times Sp(n)$ [21];
- 3. Δ_M acting on $\mathcal{S}^2(M)$ and M is the complex projective space $P^2(\mathbb{C})$ [22];
- 4. Δ_M acting on $S^2(M)$ and M is either S^n or $P^n(\mathbb{C})$ [6] and [7];
- 5. Brian and Richard Millman give in [2] a theoretical method for computing the spectra of Lichnerowicz Laplacian acting on $\Omega(G)$ where G is a compact semisimple Lie group endowed with the Killing form;
- 6. Some partial results where given in [12]-[14].

In this paper, we compute the eigenvalues and we determine the spaces of eigentensors of Δ_M acting on $\mathcal{S}(M)$ in the case where M is the Euclidian sphere S^n .

Let us describe our method briefly. We consider the (n+1)-Euclidian space \mathbb{R}^{n+1} with its canonical coordinates (x_1, \ldots, x_{n+1}) . For any $k, p \in \mathbb{N}$, we denote by $\mathcal{S}^p H_k^{\delta}$ the space of symmetric covariant p-tensor fields T on \mathbb{R}^{n+1} satisfying:

- 1. $T = \sum_{1 \leq i_1 \leq ... \leq i_p \leq n+1} T_{i_1,...,i_p} dx_{i_1} \odot ... \odot dx_{i_p}$ where $T_{i_1,...,i_p}$ are homogeneous polynomials of degree k;
- $2. \ \delta(T) = \Delta_{\mathbb{R}^{n+1}}(T) = 0.$

The *n*-dimensional sphere S^n is the space of unitary vectors in \mathbb{R}^{n+1} and the Euclidian metric on \mathbb{R}^{n+1} induces a Riemannian metric on S^n . We denote by $i: S^n \hookrightarrow \mathbb{R}^{n+1}$ the canonical inclusion.

For any tensor field $T \in \Gamma(\otimes^p T^*\mathbb{R}^{n+1})$, we compute $i^*(\Delta_{\mathbb{R}^{n+1}}T) - \Delta_{S^n}(i^*T)$ and get a formula (see Theorem 2.1). Inspired by this formula and having in mind the fact that $i^*: \sum_{k>0} \mathcal{S}^p H_k^{\delta} \longrightarrow \mathcal{S}^p S^n$ is injective and its image is

dense in S^pS^n (see [10]), we give, for any k, a direct sum decomposition of $S^pH_k^{\delta}$ composed by eigenspaces of Δ_{S^n} . Thus we obtain the eigenvalues and the spaces of eigentensors with its multiplicities of Δ_{S^n} acting on $S(S^n)$ (see Section 4).

Note that the eigenvalues and the eigenspaces of Δ_{S^n} acting on $\Omega(S^n)$ was computed in [10] by using the representation theory. In [11], I. Iwasaki and K. Katase recover the result by a method using the restriction of harmonic tensor fields and a result in [8]. The formula obtained in Theorem 2.1 combined with the methods developed in [10] and [11] permit to present those results in a more precise form (see Section 3).

2 A relation between $\Delta_{\mathbb{R}^{n+1}}$ and Δ_{S^n}

We consider the Euclidian space \mathbb{R}^{n+1} endowed with its canonical coordinates (x_1, \ldots, x_{n+1}) and its canonical Euclidian flat Riemannian metric <, >. We denote by D be the Levi-Civita covariant derivative associated to <, >. We consider the radial vector field given by

$$\overrightarrow{r} = \sum_{i=1}^{n+1} x_i \frac{\partial}{\partial x_i}.$$

For any p-tensor field $T \in \Gamma(\otimes^p T^*\mathbb{R}^{n+1})$ and for any $1 \leq i < j \leq p$, we denote by $i_{\overrightarrow{T},j}T$ the (p-1)-tensor field given by

$$i_{\overrightarrow{r},j}T(X_1,\ldots,X_{p-1})=T(X_1,\ldots,X_{j-1},\overrightarrow{r},X_j,\ldots,X_{p-1}),$$

and by $Tr_{i,j}T$ the (p-2)-tensor field given by

$$Tr_{i,j}T(X_1,\ldots,X_{p-2})=\sum_{l=1}^{n+1}T(X_1,\ldots,X_{i-1},E_l,X_i,\ldots,X_{j-2},E_l,X_{j-1},\ldots,X_{p-2}),$$

where (E_1, \ldots, E_{n+1}) is any orthonormal basis of \mathbb{R}^{n+1} . Note that $Tr_{i,j}T = 0$ if T is a differential form and $Tr_{i,j}T = TrT$ if T is symmetric.

For any permutation σ of $\{1,\ldots,p\}$, we denote by T^{σ} the p-tensor field

$$T^{\sigma}(X_1,\ldots,X_p)=T(X_{\sigma(1)},\ldots,X_{\sigma(p)}).$$

For $1 \leq i < j \leq p$, the transposition of (i, j) is the permutation $\sigma_{i,j}$ of $\{1, \ldots, p\}$ such that $\sigma_{i,j}(i) = j$, $\sigma_{i,j}(j) = i$ and $\sigma_{i,j}(k) = k$ for $k \neq i, j$. Let \mathcal{T} denote the set of the transpositions of $\{1, \ldots, p\}$.

The sphere $i:S^n\hookrightarrow {\rm I\!R}^{n+1}$ is endowed with the Euclidian metric.

Theorem 2.1 Let T be a covariant p-tensor field on \mathbb{R}^{n+1} . Then,

$$i^{*}(\Delta_{\mathbb{R}^{n+1}}T) = \Delta_{S^{n}}i^{*}T + i^{*}\left(p(1-p)T + (2p-n+1)L_{\overrightarrow{r}}T - L_{\overrightarrow{r}} \circ L_{\overrightarrow{r}}T\right)$$
$$-2\sum_{\sigma \in \mathcal{T}}T^{\sigma} + O(T),$$

where O(T) is given by

$$O(T)(X_{1},...,X_{p}) = 2\sum_{i< j} \langle X_{i}, X_{j} \rangle Tr_{i,j}(X_{1},...,\hat{X}_{i},...,\hat{X}_{j},...,X_{p})$$
$$-2\sum_{j=1}^{p} D_{X_{j}}(i_{\overrightarrow{r},j}T)(X_{1},...,\hat{X}_{j},...,X_{p}),$$

where \hat{X} designs that X is deleted.

Proof. The proof is a massive computation in a local orthonormal frame using the properties of the Riemannian embedding of the sphere in the Euclidian space.

We choose a local orthonormal frame of \mathbb{R}^{n+1} of the form (E_1, \ldots, E_n, N) such that E_i is tangent to S^n for $1 \leq i \leq n$ and $N = \frac{1}{r} \overrightarrow{r}$ where $r = \sqrt{x_1^2 + \ldots + x_{n+1}^2}$.

For any vector field X on \mathbb{R}^{n+1} , we have

$$D_X N = \frac{1}{r} (X - \langle X, N \rangle N), \qquad (6)$$

$$D_N X = [N, X] + \frac{1}{r}(X - \langle X, N \rangle N).$$
 (7)

Let ∇ be the Levi-Civita connexion of the Riemannian metric on S^n . We have, for any vector fields X, Y tangent to S^n ,

$$D_X Y = \nabla_X Y - \langle X, Y \rangle N. \tag{8}$$

Let T be a covariant p-tensor field on \mathbb{R}^{n+1} and (X_1, \ldots, X_p) a family of vector fields on \mathbb{R}^{n+1} which are tangent to S^n . A direct calculation using the definition of the Lichnerowicz Laplacian gives

$$\Delta_{\mathbb{R}^{n+1}}(T)(X_1, \dots, X_p) = D^*D(T)(X_1, \dots, X_p)
= \sum_{i=1}^n \left(-E_i E_i . T(X_1, \dots, X_p) + 2 \sum_{j=1}^p E_i . T(X_1, \dots, D_{E_i} X_j, \dots, X_p) \right)
+ D_{E_i} E_i . T(X_1, \dots, X_p) - \sum_{j=1}^p T(X_1, \dots, D_{D_{E_i} E_i} X_j, \dots, X_p)
- \sum_{j=1}^p T(X_1, \dots, D_{E_i} D_{E_i} X_j, \dots, X_p) - 2 \sum_{l < j} T(X_1, \dots, D_{E_i} X_l, \dots, D_{E_i} X_j, \dots, X_p)
- N.N.T(X_1, \dots, X_p) + 2 \sum_{j=1}^p N.T(X_1, \dots, D_N X_j, \dots, X_p)
+ D_N N.T(X_1, \dots, X_p) - \sum_{j=1}^p T(X_1, \dots, D_N X_j, \dots, X_p)
- \sum_{j=1}^p T(X_1, \dots, D_N D_N X_j, \dots, X_p) - 2 \sum_{l < j} T(X_1, \dots, D_N X_l, \dots, D_N X_j, \dots, X_p).$$

(6)-(8) make it obvious that

$$D_{D_{E_{i}}E_{i}}X_{j} = \nabla_{\nabla_{E_{i}}E_{i}}X_{j} - \langle \nabla_{E_{i}}E_{i}, X_{j} > N - [N, X_{j}]$$

$$-\frac{1}{r}(X_{j} - \langle X_{j}, N > N),$$

$$D_{E_{i}}D_{E_{i}}X_{j} = \nabla_{E_{i}}\nabla_{E_{i}}X_{j} - (\langle E_{i}, \nabla_{E_{i}}X_{j} > +E_{i}. \langle E_{i}, X_{j} >)N$$

$$-\frac{1}{r} \langle E_{i}, X_{j} > E_{i},$$

$$D_{N}D_{N}X = [N, [N, X]] + \frac{2}{r}[N, X] + (\frac{1}{r^{2}} - \frac{1}{r})(X - \langle X, N > N)$$

$$-\frac{2}{r}N. \langle X, N > N.$$

$$(11)$$

By (8)-(10), we get easily, in restriction to S^n ,

$$\sum_{i=1}^{n} \left(2 \sum_{j=1}^{p} E_{i}.T(X_{1}, \dots, D_{E_{i}}X_{j}, \dots, X_{p}) + D_{E_{i}}E_{i}.T(X_{1}, \dots, X_{p}) \right)$$

$$- \sum_{j=1}^{p} T(X_{1}, \dots, D_{D_{E_{i}}E_{i}}X_{j}, \dots, X_{p}) - \sum_{j=1}^{p} T(X_{1}, \dots, D_{E_{i}}D_{E_{i}}X_{j}, \dots, X_{p}) \right) =$$

$$\sum_{i=1}^{n} \left(2 \sum_{j=1}^{p} E_{i}.T(X_{1}, \dots, \nabla_{E_{i}}X_{j}, \dots, X_{p}) + \nabla_{E_{i}}E_{i}.T(X_{1}, \dots, X_{p}) \right)$$

$$- \sum_{j=1}^{p} T(X_{1}, \dots, \nabla_{\nabla_{E_{i}}E_{i}}X_{j}, \dots, X_{p}) - \sum_{j=1}^{p} T(X_{1}, \dots, \nabla_{E_{i}}\nabla_{E_{i}}X_{j}, \dots, X_{p}) \right)$$

$$-2 \sum_{j=1}^{p} X_{j}.T(X_{1}, \dots, \bigcap_{i} \bigcap_{j} (1, \dots, X_{p}) + p(n+1)T(X_{1}, \dots, X_{p}) - nL_{N}T(X_{1}, \dots, X_{p}).$$

On other hand, also by using (8), we have

$$\sum_{l < j} \sum_{i=1}^{n} T(X_{1}, \dots, D_{E_{i}}X_{l}, \dots, D_{E_{i}}X_{j}, \dots, X_{p}) =$$

$$\sum_{l < j} \sum_{i=1}^{n} T(X_{1}, \dots, D_{E_{i}}X_{l}, \dots, \nabla_{E_{i}}X_{j}, \dots, X_{p}) - \sum_{l < j} T(X_{1}, \dots, D_{X_{j}}X_{l}, \dots, \overset{j}{N}, \dots, X_{p}) =$$

$$\sum_{l < j} \sum_{i=1}^{n} T(X_{1}, \dots, \nabla_{E_{i}}X_{l}, \dots, \nabla_{E_{i}}X_{j}, \dots, X_{p}) - \sum_{l < j} T(X_{1}, \dots, \overset{l}{N}, \dots, \nabla_{X_{l}}X_{j}, \dots, X_{p})$$

$$- \sum_{l < j} T(X_{1}, \dots, D_{X_{j}}X_{l}, \dots, \overset{j}{N}, \dots, X_{p}) = \sum_{l < j} \sum_{i=1}^{n} T(X_{1}, \dots, \nabla_{E_{i}}X_{l}, \dots, \nabla_{E_{i}}X_{j}, \dots, X_{p})$$

$$- \sum_{l < j} T(X_{1}, \dots, D_{X_{j}}X_{l}, \dots, \overset{j}{N}, \dots, X_{p}) - \sum_{l < j} T(X_{1}, \dots, \overset{l}{N}, \dots, D_{X_{l}}X_{j}, \dots, X_{p})$$

$$- \sum_{l < j} < X_{l}, X_{j} > T(X_{1}, \dots, \overset{j}{N}, \dots, \overset{j}{N}, \dots, X_{p}).$$

So we get, in restriction to S^n , since $D_N N = 0$

$$\Delta_{\mathbb{R}^{n+1}}(X_{1},\ldots,X_{p}) - \nabla^{*}\nabla T(X_{1},\ldots,X_{p}) =
p(n+1)T(X_{1},\ldots,X_{p}) - nL_{N}T(X_{1},\ldots,X_{p}) - 2\sum_{j=1}^{p} D_{X_{j}}(i_{N,j}T)(X_{1},\ldots,\hat{X}_{j},\ldots,X_{p})
+2\sum_{l< j} < X_{l}, X_{j} > T(X_{1},\ldots,\hat{N},\ldots,\hat{N},\ldots,X_{p}) - N.N.T(X_{1},\ldots,X_{p})
+2\sum_{j=1}^{p} N.T(X_{1},\ldots,D_{N}X_{j},\ldots,X_{p}) - \sum_{j=1}^{p} T(X_{1},\ldots,D_{N}D_{N}X_{j},\ldots,X_{p})
-2\sum_{i< j} T(X_{1},\ldots,D_{N}X_{i},\ldots,D_{N}X_{j},\ldots,X_{p}).$$

Remark that, in restriction to S^n , the following equality holds

$$\sum_{j=1}^{p} D_{X_j}(i_{N,j}T)(X_1, \dots, \hat{X}_j, \dots, X_p) = \sum_{j=1}^{p} D_{X_j}(i_{\overrightarrow{r'},j}T)(X_1, \dots, \hat{X}_j, \dots, X_p).$$

Now by using (7) and (11) and by taking the restriction to S^n , we have

$$2\sum_{j=1}^{p} N.T(X_{1}, \dots, D_{N}X_{j}, \dots, X_{p}) =$$

$$2\sum_{j=1}^{p} N.T(X_{1}, \dots, [N, X_{j}], \dots, X_{p}) + 2\sum_{j=1}^{p} N(\frac{1}{r})T(X_{1}, \dots, X_{j}, \dots, X_{p})$$

$$+2\sum_{j=1}^{p} N.T(X_{1}, \dots, X_{j}, \dots, X_{p}) - 2\sum_{j=1}^{p} N(\langle X_{j}, N \rangle)T(X_{1}, \dots, X_{p}) =$$

$$2\sum_{j=1}^{p} N.T(X_{1}, \dots, [N, X_{j}], \dots, X_{p}) - 2pT(X_{1}, \dots, X_{p}) + 2pN.T(X_{1}, \dots, X_{j}, \dots, X_{p})$$

$$-2\sum_{j=1}^{p} N(\langle X_{j}, N \rangle)T(X_{1}, \dots, X_{p}) - 2\sum_{j=1}^{p} N(\langle X_{j}, N \rangle)T(X_{1}, \dots, X_{p}).$$

$$\sum_{j=1}^{p} T(X_{1}, \dots, D_{N}D_{N}X_{j}, \dots, X_{p}) =$$

$$\sum_{j=1}^{p} T(X_{1}, \dots, [N, [N, X_{j}], \dots, X_{p}) - 2\sum_{j=1}^{p} N(\langle X_{j}, N \rangle)T(X_{1}, \dots, X_{p}).$$

$$\sum_{i < j} T(X_1, \dots, D_N X_i, \dots, D_N X_j, \dots, X_p) =
\sum_{i < j} T(X_1, \dots, [N, X_i], \dots, [N, X_j], \dots, X_p) + \frac{p(p-1)}{2} T(X_1, \dots, X_p)
+ \sum_{i < j} T(X_1, \dots, X_i, \dots, [N, X_j], \dots, X_p) + \sum_{i < j} T(X_1, \dots, [N, X_i], \dots, X_j, \dots, X_p).$$

So we get, in restriction to S^n

$$-N.N.T(X_1, ..., X_p) + 2\sum_{j=1}^p N.T(X_1, ..., D_N X_j, ..., X_p)$$

$$-\sum_{j=1}^p T(X_1, ..., D_N D_N X_j, ..., X_p) - 2\sum_{i < j} T(X_1, ..., D_N X_i, ..., D_N X_j, ..., X_p) =$$

$$-L_N \circ L_N T(X_1, ..., X_p) + 2pL_N T(X_1, ..., X_p) - p(1+p)T(X_1, ..., X_p).$$

The curvature of S^n is given by

$$R(X,Y)Z = \langle X, Y \rangle Z - \langle Y, Z \rangle X$$
 and $r(X) = (n-1)X$.

Hence, a direct computation gives that the curvature operator is given by

$$R(T)(X_1, \dots, X_p) = p(n-1)T(X_1, \dots, X_p) + 2\sum_{\sigma \in \mathcal{T}} T^{\sigma}(X_1, \dots, X_p)$$
$$-2\sum_{i < j} \sum_{l=1}^{n} \langle X_i, X_j \rangle T(X_1, \dots, E_l, \dots, E_l, \dots, X_p).$$

Finally, we get

$$i^*(\Delta_{\mathbb{R}^{n+1}}T) = \Delta_{S^n}i^*T + i^*(p(1-p)T + (2p-n)L_NT - L_N \circ L_NT - 2\sum_{\sigma \in \mathcal{T}}T^{\sigma} + O(T)),$$

One can conclude the proof by remarking that

$$i^*(L_NT) = i^*(L_{\overrightarrow{r}}T)$$
 and $i^*(L_N \circ L_NT) = -i^*(L_{\overrightarrow{r}}T) + i^*(L_{\overrightarrow{r}} \circ L_{\overrightarrow{r}}T)$. Q.E.D.

Corollary 2.1 Let α be a differential p-form on \mathbb{R}^{n+1} . Then

$$i^*(\Delta_{\mathbb{R}^{n+1}}\alpha) = \Delta_{S^n}i^*\alpha + i^*\left((2p-n+1)L_{\overrightarrow{r}}\alpha - L_{\overrightarrow{r}}\circ L_{\overrightarrow{r}}\alpha - 2di_{\overrightarrow{r}}\alpha\right)$$

Corollary 2.2 Let T be a symmetric p-tensor field on \mathbb{R}^{n+1} . Then

$$i^*(\Delta_{\mathbb{R}^{n+1}}T) = \Delta_{S^n}i^*T + i^*\left(2p(1-p)T + (2p-n+1)L_{\overrightarrow{r}}T - L_{\overrightarrow{r}} \circ L_{\overrightarrow{r}}T\right) - 2\delta^*(i_{\overrightarrow{r}}T) + 2Tr(T)\odot <, >\right),$$

where \odot is the symmetric product.

3 Eigenvalues and eigenforms of Δ_{S^n} acting on $\Omega(S^n)$

In this section, we will use corollary 2.1 and the results developed in [10] to deduce the eigenvalues and the spaces of eigenforms of Δ_{S^n} acting on $\Omega^*(S^n)$. We recover the results of [10] and [11] in a more precise form.

Let $\wedge^p H_k$ be the space of all coclosed harmonic homogeneous p-forms of degree k on \mathbb{R}^{n+1} . A differential form α belongs to $\wedge^p H_k$ if $\delta(\alpha) = 0$ and α can be written

$$\alpha = \sum_{1 \le i_1 < \dots < i_p \le n+1} \alpha_{i_1 \dots i_p} dx_{i_1} \wedge \dots \wedge dx_{i_p},$$

where $\alpha_{i_1...i_p}$ are harmonic polynomial functions on \mathbb{R}^{n+1} of degree k. For any $\alpha \in \wedge^p H_k$, we have

$$L_{\overrightarrow{r}}\alpha = di_{\overrightarrow{r}}\alpha + i_{\overrightarrow{r}}d\alpha = (k+p)\alpha. \tag{12}$$

We have (see [10]),

$$i^*: \sum_{k\geq 0} \wedge^p H_k \longrightarrow \Omega^p(S^n)$$

is injective and its image is dense.

For any $\alpha \in \wedge^p H_k$, we put

$$\omega(\alpha) = \alpha - \frac{1}{p+k} di \overrightarrow{r} \alpha. \tag{13}$$

Lemma 3.1 We get a linear map $\omega : \wedge^p H_k \longrightarrow \wedge^p H_k$ which is a projector, i.e., $\omega \circ \omega = \omega$. Moreover,

$$Ker\omega = d(\wedge^{p-1}H_{k+1}), \quad Im\omega = \wedge^p H_k \cap Keri_{\overrightarrow{R}},$$

and hence

$$\wedge^p H_k = \wedge^p H_k \cap Keri_{\overrightarrow{r}} \oplus d(\wedge^{p-1} H_{k+1}).$$

The following lemma is an immediate consequence of Corollary 2.1 and (12).

Lemma 3.2 1. For any $\alpha \in \wedge^p H_k \cap Keri_{\overrightarrow{r}}$, we have

$$\Delta_{S^n} i^* \alpha = (k+p)(k+n-p-1)i^* \alpha.$$

2. For any $\alpha \in d(\wedge^{p-1}H_{k+1})$, we have

$$\Delta_{S^n} i^* \alpha = (k+p)(k+n-p+1)i^* \alpha.$$

Remark 3.1 We have

$$(k+p)(k+n-p-1) = (k'+p)(k'+n-p+1) \Leftrightarrow k = k'+1 \quad and \quad n = 2p.$$

The following Table gives explicitly the spectra of Δ_{S^n} and the spaces of eigenforms with its multiplicities. The multiplicity was computed in [11].

Table I

p	The eigenvalues	The space of eigenforms	Multiplicity
p = 0	$k(k+n-1), k \in \mathbb{N}$	$\wedge^0 H_k$	$\frac{(n+k-2)!(n+2k-1)}{k!(n-1)!}$
$1 \le p \le n, \\ n \ne 2p$	$(k+p)(k+n-p-1),$ $k \in \mathbb{N}^*$ $(k+p)(k+n-p+1),$ $k \in \mathbb{N}$		$\frac{(n+k-1)!(n+2k-1)}{p!(k-1)!(n-p-1)!(n+k-p-1)(k+p)}$ $\frac{(n+k)!(n+2k+1)}{(p-1)!k!(n-p)!(n+k-p+1)(k+p)}$
$1 \le p \le n,$ $n = 2p$	$(k+p)(k+p+1)$ $k \in \mathbb{N}$	$\omega(\wedge^p H_{k+1}) \oplus d(\wedge^{p-1} H_{k+1})$	$\frac{2(2p+k)!(2p+2k+1)}{p!(p-1)!k!(k+p+1)(k+p)}$

4 Eigenvalues and eigentensors of Δ_{S^n} acting on $\mathcal{S}(S^n)$

This section is devoted to the determination of the eigenvalues and the spaces of eigentensors of Δ_{S^n} acting on $\mathcal{S}(S^n)$.

Let $S^p P_k$ be the space of $T \in S^p(\mathbb{R}^{n+1})$ of the form

$$T = \sum_{1 \le i_1 \le \dots \le i_p \le n+1} T_{i_1 \dots i_p} dx_{i_1} \odot \dots \odot dx_{i_p},$$

where $T_{i_1...i_p}$ are homogeneous polynomials of degree k. We put

$$\mathcal{S}^p H_k^{\delta} = \mathcal{S}^p P_k \cap Ker \Delta_{\mathbb{R}^{n+1}} \cap Ker \delta$$
 and $\mathcal{S}^p H_k^{\delta 0} = \mathcal{S}^p H_k^{\delta} \cap Ker Tr$.

In a similar manner as in [10] Lemma 6.4 and Corollary 6.6, we have

$$S^p P_k = S^p H_k^{\delta} \oplus (r^2 S^p P_{k-2} + dr^2 \odot S^{p-1} P_{k-1}), \tag{14}$$

and

$$i^*: \sum_{k\geq 0} \mathcal{S}^p H_k^{\delta} \longrightarrow \mathcal{S}^p S^n$$

is injective and its image is dense in S^pS^n .

Now, for any $k \geq 0$, we proceed to give a direct sum decomposition of $\mathcal{S}^p H_k^{\delta}$ consisting of eigenspaces of Δ_{S^n} and, hence, we determine completely the eigenvalues of Δ_{S^n} acting on $\mathcal{S}^p(S^n)$. This will be done in several steps.

At first, we have the following direct sum decomposition:

$$S^{p}H_{k}^{\delta} = S^{p}H_{k}^{\delta 0} \oplus \bigoplus_{l=1}^{\left[\frac{p}{2}\right]} S^{p-2l}H_{k}^{\delta 0} \odot <, >^{l}, \tag{15}$$

where $<,>^l$ is the symmetric product of l copies of <,>.

The task is now to decompose $S^pH_k^{\delta 0}$ as a sum of eigenspaces of Δ_{S^n} and get, according to (5), all the eigenvalues. This decomposition needs some preparation.

Lemma 4.1 Let $T \in S^p P_k$ and $h \in \mathbb{N}^*$. Then we have the following formulas:

1.
$$\delta^*(i \rightarrow T) - i \rightarrow \delta^*(T) = (p - k)T;$$

2.
$$\delta^{*(h)}(i \rightarrow T) - i \rightarrow \delta^{*(h)}(T) = h(p - k + h - 1)\delta^{*(h-1)}(T);$$

3.
$$\delta^*(i_{\overrightarrow{r}}hT) - i_{\overrightarrow{r}}h\delta^*(T) = h(p-k-h+1)i_{\overrightarrow{r}}h-1T$$

where
$$i_{\overrightarrow{r}^{h}} = \overbrace{i_{\overrightarrow{r}} \circ \ldots \circ i_{\overrightarrow{r}}}^{h}$$
 and $\delta^{*(h)} = \overbrace{\delta^{*} \circ \ldots \circ \delta^{*}}^{h}$.

Proof. The first formula is easily verified and the others follow by induction on h. Q.E.D.

Now, we will construct two linear maps $p_{\delta^*}: \mathcal{S}^p P_k \longrightarrow \mathcal{S}^p P_k$ for $k \leq p$, and $p_{\overrightarrow{r}}: \mathcal{S}^p P_k \longrightarrow \mathcal{S}^p P_k$ for $k \geq p$ satisfying:

1.
$$p_{\delta^*} \circ p_{\delta^*} = p_{\delta^*}, Kerp_{\delta^*} = i \xrightarrow{\sim} (\mathcal{S}^{p+1}P_{k-1}), Imp_{\delta^*} = Ker\delta^* \cap \mathcal{S}^p P_k;$$

2.
$$p_{\overrightarrow{r}} \circ p_{\overrightarrow{r}} = p_{\overrightarrow{r}}, Kerp_{\overrightarrow{r}} = \delta^*(\mathcal{S}^{p-1}P_{k+1}), Imp_{\overrightarrow{r}} = Keri_{\overrightarrow{r}} \cap \mathcal{S}^p P_k$$

The procedure is to put, for $T \in \mathcal{S}^p P_k$,

$$p_{\delta^*}(T) = \sum_{s=0}^k \alpha_s i \overrightarrow{r}^s \delta^{*(s)}(T), \text{ and } p\overrightarrow{r}(T) = \sum_{s=0}^p \beta_s \delta^{*(s)}(i \overrightarrow{r}^s T),$$

and find $(\alpha_0, \ldots, \alpha_k)$ and $(\beta_0, \ldots, \beta_p)$ such that the required properties are satisfied.

A straightforward computation using Lemma 4.1 gives

$$\delta^{*}(p_{\delta^{*}}(T)) = \sum_{s=0}^{k-1} (\alpha_{s} - (s+1)(k-p-s-2)\alpha_{s+1})i_{\overrightarrow{r}^{s}}\delta^{*(s+1)}(T),$$

$$i_{\overrightarrow{r}^{s}}(p_{\overrightarrow{r}^{s}}(T)) = \sum_{s=0}^{p-1} (\beta_{s} - (s+1)(p-k-s-2)\beta_{s+1})\delta^{*(s)}(i_{\overrightarrow{r}^{s+1}}T).$$

Hence, we define p_{δ^*} and p_{\rightarrow} as follows:

$$\begin{cases} p_{\delta^*}(T) = \sum_{s=0}^k \alpha_s i_{\overrightarrow{r}^s} \delta^{*(s)}(T) \\ \alpha_0 = 1 & \text{and } \alpha_s - (s+1)(k-p-s-2)\alpha_{s+1} = 0 & \text{for } 1 \le s \le k-1; \end{cases}$$

$$\begin{cases} p_{\overrightarrow{r'}}(T) = \sum_{s=0}^{p} \beta_s \delta^{*(s)}(i_{\overrightarrow{r'}} T) \\ \beta_0 = 1 & \text{and } \beta_s - (s+1)(p-k-s-2)\beta_{s+1} = 0 & \text{for } 1 \le s \le p-1. \end{cases}$$

From this definition and by using Lemma 4.1, one can check easily that p_{δ^*} and $p_{\overrightarrow{\sigma}}$ satisfy the required properties.

On other hand, it is easy to check that we have, for any symmetric tensor field T on \mathbb{R}^{n+1} ,

$$\Delta_{\mathbb{R}^{n+1}}(i \overrightarrow{r} T) = i \overrightarrow{r} \Delta_{\mathbb{R}^{n+1}}(T) + 2\delta T, \tag{16}$$

$$\delta(i \overrightarrow{r} T) = i \overrightarrow{r} \delta(T) - Tr(T), \tag{17}$$

$$Tr(\delta^*(T)) = -2\delta(T) + \delta^*(Tr(T)), \tag{18}$$

$$Tr(i_{\overrightarrow{r}}T) = i_{\overrightarrow{r}}Tr(T).$$
 (19)

From these formulas and from (3), one deduce easily that $p_{\delta^*}(\mathcal{S}^p H_k^{\delta 0}) \subset \mathcal{S}^p H_k^{\delta 0}$ and $p_{\overrightarrow{r'}}(\mathcal{S}^p H_k^{\delta 0}) \subset \mathcal{S}^p H_k^{\delta 0}$ and thus one get the following direct sum decompositions:

$$S^{p}H_{k}^{\delta 0} = S^{p}H_{k}^{\delta 0} \cap Ker\delta^{*} \oplus i_{\overrightarrow{r}}\left(S^{p+1}H_{k-1}^{\delta 0}\right), \quad \text{if } k \leq p, \tag{20}$$

$$\mathcal{S}^{p}H_{k}^{\delta 0} = \mathcal{S}^{p}H_{k}^{\delta 0} \cap Keri_{\overrightarrow{r}} \oplus \delta^{*}\left(\mathcal{S}^{p-1}H_{k+1}^{\delta 0}\right), \quad \text{if } k \geq p.$$
 (21)

These decompositions are far for being sufficient and, in order to obtain a more sharp direct sum decompositions of $S^pH_k^{\delta 0}$, we need the following lemma.

Lemma 4.2 1. For k < p, $i \rightarrow \mathcal{S}^p P_k \longrightarrow \mathcal{S}^{p-1} P_{k+1}$ is injective.

- 2. For k > p, $\delta^* : \mathcal{S}^p P_k \longrightarrow \mathcal{S}^{p+1} P_{k-1}$ is injective.
- 3. For k = p, $Ker\delta^* = Keri \rightarrow$.

Proof.

1. Let $T \in \mathcal{S}^p P_k$ such that $i \xrightarrow{r} T = 0$. The second formula in Lemma 4.1 gives, for any $h \ge 1$,

$$i \overrightarrow{r} \delta^{*(h)}(T) = -h(p-k+h-1)\delta^{*(h-1)}(T).$$

Since $\delta^{*(h)}(T) = 0$ for $h \ge k+1$ and $h(p-k+h-1) \ne 0$ for any $h \ge 1$, we get form this relation that $\delta^{*(h-1)}(T) = 0$ for any $h \ge 1$, in particular for h = 1, we get T = 0.

2. The same argument as 1. using the third formula in Lemma 4.1.

3. Let $T \in \mathcal{S}^p P_p$ such that $i_{\overrightarrow{r}}T = 0$. From Lemma 4.1, we get $i_{\overrightarrow{r}}\delta^*(T) = 0$. Since $\delta^*(T) \in \mathcal{S}^{p+1}P_{p-1}$ and from 1. we deduce that $\delta^*(T) = 0$ and hence $Keri_{\overrightarrow{r}} \subset Ker\delta^*$. The same argument using Lemma 4.1 and 2. will give the other inclusion. Q.E.D.

By combining (20) and (21) with Lemma 4.2, we obtain the following lemma.

Lemma 4.3 We have:

1. if k < p

$$\mathcal{S}^{p}H_{k}^{\delta 0} = \bigoplus_{l=0}^{k} i_{\overrightarrow{r}^{l}} \left(\mathcal{S}^{p+l}H_{k-l}^{\delta 0} \cap Ker\delta^{*} \right);$$

2. if k > p

$$\mathcal{S}^{p}H_{k}^{\delta 0} = \bigoplus_{l=0}^{p} \delta^{*l} \left(\mathcal{S}^{p-l} H_{k+l}^{\delta 0} \cap Keri \overrightarrow{r} \right);$$

3. If k = p, for any $0 \le l \le p$,

$$i_{\overrightarrow{r}^{l}}\left(\mathcal{S}^{p+l}H_{p-l}^{\delta 0}\cap Ker\delta^{*}\right)=\delta^{*l}\left(\mathcal{S}^{p-l}H_{p+l}^{\delta 0}\cap Keri_{\overrightarrow{r}^{\prime}}\right),$$

and

$$\mathcal{S}^{p}H_{p}^{\delta 0} = \bigoplus_{l=0}^{p} i_{\overrightarrow{r}^{l}} \left(\mathcal{S}^{p+l}H_{p-l}^{\delta 0} \cap Ker\delta^{*} \right) = \bigoplus_{l=0}^{p} \delta^{*l} \left(\mathcal{S}^{p-l}H_{p+l}^{\delta 0} \cap Keri_{\overrightarrow{r}^{l}} \right).$$

Now, we use Corollary 2.2 to show that the decompositions of $S^p H_k^{\delta 0}$ given in Lemma 4.3 are composed by eigenspaces of Δ_{S^n} .

Theorem 4.1 We have:

1. If
$$k \leq p$$
, for any $0 \leq q \leq k$ and any $T \in i_{\overrightarrow{r}^{(k-q)}} \left(\mathcal{S}^{p+k-q} H_q^{\delta 0} \cap Ker \delta^* \right)$,
$$\Delta_{S^n} i^* T = \left((k+p)(n+p+k-2q-1) + 2q(q-1) \right) i^* T;$$

2. If
$$k \geq p$$
, for any $0 \leq q \leq p$ and for any $T \in \delta^{*(p-q)} \left(\mathcal{S}^q H_{k+p-q}^{\delta 0} \cap Keri \overrightarrow{r} \right)$,
$$\Delta_{S^n} i^* T = \left((k+p)(n+p+k-2q-1) + 2q(q-1) \right) i^* T.$$

Proof.

1. Let $T = i_{\overrightarrow{r}^{(k-q)}}(T_0)$ with $T_0 \in \mathcal{S}^{p+k-q}H_q^{\delta 0} \cap Ker\delta^*$. We have from Corollary 2.2

$$\Delta_{S^n} i^* T = i^* \left(2p(p-1)T + (n-2p-1)L_{\overrightarrow{r}}T + L_{\overrightarrow{r}} \circ L_{\overrightarrow{r}}T + 2\delta^*(i_{\overrightarrow{r}}T) - 2Tr(T)\odot <, > \right).$$

We have

$$TrT = 0$$
, $L_{\overrightarrow{r}} = (k+p)T$ and $L_{\overrightarrow{r}} \circ L_{\overrightarrow{r}}T = (k+p)^2T$.

Moreover, by using Lemma 4.1, we have

$$2\delta^*(i\overrightarrow{r}T) = 2\delta^*(i\overrightarrow{r}^{(k-q+1)}T_0)$$

$$\stackrel{\delta^*(T_0)=0}{=} 2(k-q+1)(p+k-q-q-k+q-1+1)i\overrightarrow{r}^{(k-q)}T_0$$

$$= 2(k-q+1)(p-q)T.$$

Hence

$$\Delta_{S^n}i^*T = (2p(p-1) + (n-2p-1)(k+p) + (k+p)^2 + 2(p-q)(k-q+1))i^*T.$$

One can deduce the desired relation by remarking that

$$2p(p-1) + 2(p-q)(k-q+1) = 2(k+p)(p-q) + 2q(q-1).$$

2. This follows by the same calculation as 1. Q.E.D.

From the fact that

$$i^*: \sum_{k>0} \mathcal{S}^p H_k^{\delta} \longrightarrow \mathcal{S}^p S^n$$

is injective and its image is dense in S^pS^n , from (15), and from Lemma 4.3 and Theorem 4.1, note that we have actually proved that the eigenvalues of Δ_{S^n} acting on S^pS^n belongs to

$$\left\{ (k+p-2l)(n+p+k-2l-2q-1) + 2q(q-1), \\ k \in \mathbb{N}, 0 \le l \le [\frac{p}{2}], 0 \le q \le \min(k,p-2l) \right\}.$$

Our next goal is to sharpen this result by computing $\dim \mathcal{S}^p H_k^{\delta 0} \cap Ker \delta^*$ if $k \leq p$ and $\dim \mathcal{S}^p H_k^{\delta 0} \cap Ker i \xrightarrow{}$ if $k \geq p$.

Lemma 4.4 We have the following formulas:

- 1. $\dim \mathcal{S}^p H_k^{\delta} = \dim \mathcal{S}^p P_k \dim \mathcal{S}^p P_{k-2} \dim \mathcal{S}^{p-1} P_{k-1} + \dim \mathcal{S}^{p-1} P_{k-3}$
- 2. $\dim \mathcal{S}^p H_k^{\delta 0} = \dim \mathcal{S}^p H_k^{\delta} \dim \mathcal{S}^{p-2} H_k^{\delta}$
- 3. $\dim(\mathcal{S}^p H_k^{\delta 0} \cap Ker\delta^*) = \dim \mathcal{S}^p H_k^{\delta 0} \dim \mathcal{S}^{p+1} H_{k-1}^{\delta 0} (k \le p),$
- 4. $\dim(\mathcal{S}^p H_k^{\delta 0} \cap Keri_{\overrightarrow{r}}) = \dim \mathcal{S}^p H_k^{\delta 0} \dim \mathcal{S}^{p-1} H_{k+1}^{\delta 0} (k \ge p).$

Note that we use the convention that $S^p P_k = S^p H_k^{\delta} = S^p H_k^{\delta 0} = 0$ if k < 0 or p < 0.

Proof.

1. The formula is a consequence of (14), the relation

$$(r^2 \mathcal{S}^p P_{k-2}) \cap (dr^2 \odot \mathcal{S}^{p-1} P_{k-1}) = r^2 (dr^2 \odot \mathcal{S}^{p-1} P_{k-3})$$

and the fact that $dr^2\odot.:\mathcal{S}^pP_k\longrightarrow\mathcal{S}^{p+1}P_{k+1}$ is injective.

- 2. The formula is a consequence of (15).
- 3. The formula is a consequence of (20) and Lemma 4.2.
- 4. The formula is a consequence of (21) and Lemma 4.2. Q.E.D.

A straightforward calculation using Lemma 4.4 and the formula

$$\dim \mathcal{S}^p P_k = \frac{(n+p)!}{n!p!} \frac{(n+k)!}{n!k!}$$

gives $\dim \mathcal{S}^p H_k^{\delta 0} \cap Ker \delta^*$ if $k \leq p$ and $\dim \mathcal{S}^p H_k^{\delta 0} \cap Ker i \rightarrow if k \geq p$. We summarize the results on the following Table.

Table II

Space	Dimension	Conditions on k and p
$\mathcal{S}^0 H_k^{\delta 0} \cap Keri \xrightarrow{r}$	$\frac{(n+k-2)!(n+2k-1)}{k!(n-1)!}$	$k \ge 0$
$\mathcal{S}^p H_0^{\delta 0} \cap Ker \delta^*$	$\frac{(n+p-2)!(n+2p-1)}{p!(n-1)!}$	$p \ge 0$
$\mathcal{S}^1 H_k^{\delta 0} \cap Keri \xrightarrow{r}$	$\frac{(n+k-3)!k(n+2k-1)(n+k-1)}{(n-2)!(k+1)!}$	$k \ge 1$
$\mathcal{S}^p H_1^{\delta 0} \cap Ker \delta^*$	$\frac{(n+p-3)!p(n+2p-1)(n+p-1)}{(n-2)!(p+1)!}$	$p \ge 1$
$\mathcal{S}^p H_k^{\delta 0} \cap Ker \delta^*$	$\frac{(n+k-4)!(n+p-3)!(n+p+k-2)}{k!(p+1)!(n-1)!(n-2)!} \times (n-2)(n+2k-3)(n+2p-1)(p-k+1)$	2 < k < n
$\mathcal{S}^p H_k^{\delta 0} \cap Keri_{\overrightarrow{r}}$	$\frac{(n+k-3)!(n+p-4)!(n+p+k-2)}{(k+1)!p!(n-1)!(n-2)!} \times$	$2 \le k \le p$
	(n-2)(n+2k-1)(n+2p-3)(k-p+1)	$k \ge p \ge 2$

Remark 4.1 Note that, for n = 2, we have

$$\dim(\mathcal{S}^p H_k^{\delta 0} \cap Ker \delta^*) = 0 \quad \text{for} \quad 2 \le k \le p,$$

$$\dim(\mathcal{S}^p H_k^{\delta 0} \cap Ker i_{\overrightarrow{P}}) = 0 \quad \text{for} \quad k \ge p \ge 2.$$

For simplicity we introduce the following notations.

$$S_{0} = \left\{ (k, l, q) \in \mathbb{N}^{3}, 0 \leq l \leq \left[\frac{p}{2}\right], 0 \leq k \leq p - 2l, 0 \leq q \leq k \right\},$$

$$S_{1} = \left\{ (k, l, q) \in \mathbb{N}^{3}, 0 \leq l \leq \left[\frac{p}{2}\right], k > p - 2l, 0 \leq q \leq p - 2l \right\},$$

$$V_{q, l}^{k} = i_{\overrightarrow{r}^{k-q}} \left(\mathcal{S}^{p-2l+k-q} H_{q}^{\delta 0} \cap Ker\delta^{*} \right) \odot <, >^{l} \text{ for } (k, l, q) \in S_{0},$$

$$W_{q, l}^{k} = \delta^{*(p-2l-q)} \left(\mathcal{S}^{q} H_{p-2l+k-q}^{\delta 0} \cap Keri_{\overrightarrow{r}} \right) \odot <, >^{l} \text{ for } (k, l, q) \in S_{1}.$$

Let us summarize all the results above.

Theorem 4.2 1. For n = 2, we have:

(a) The set of the eigenvalues of Δ_{S^2} acting on S^pS^2 is

$$\left\{ (k+p-2l)(p+k-2l+1), \ k \in \mathbb{N}, 0 \le l \le [\frac{p}{2}] \right\};$$

(b) The eigenspace associated to the eigenvalue $\lambda(k,l) = (k+p-2l)(k+p-2l+1)$ is given by

$$V_{\lambda(k,l)} = \begin{cases} \bigoplus_{\substack{a=0 \\ \min(l, [\frac{k}{2}])}}^{\min(l, [\frac{k}{2}])} \left(V_{0,l-a}^{k-2a} \oplus V_{1,l-a}^{k+1-2a} \right) & if \quad 0 \le k \le p-2l \\ \bigoplus_{\substack{a=0 \\ \min(l, [\frac{k}{2}])}}^{\min(l, [\frac{k}{2}])} \left(W_{0,l-a}^{k-2a} \oplus W_{1,l-a}^{k+1-2a} \right) & if \quad k > p-2l; \end{cases}$$

(c) The multiplicity of $\lambda(k, l)$ is given by

$$m(\lambda(k,l)) = 2(\min(l, \lfloor \frac{k}{2} \rfloor) + 1)(1 + 2p + 2k - 4l).$$

- 2. For $n \geq 3$, we have:
 - (a) The set of the eigenvalues of Δ_{S^n} acting on S^pS^n is

$$\left\{ (k+p-2l)(n+p+k-2l-2q-1) + 2q(q-1), \\ k \in \mathbb{N}, 0 \le l \le [\frac{p}{2}], 0 \le q \le \min(k,p-2l) \right\};$$

(b) The space

$$\mathcal{P} = \sum_{k \geq 0} \mathcal{S}^p H_k^{\delta} = (\bigoplus_{(k,l,q) \in S_0} V_{q,l}^k) \oplus (\bigoplus_{(k,l,q) \in S_1} W_{q,l}^k)$$

is dense in S^pS^n and, for any $(k,q,l) \in S_0$ (resp. $(k,q,l) \in S_1$), $V_{q,l}^k$ (resp. $W_{q,l}^k$) is a subspace of the eigenspace associated to the eigenvalue (k+p-2l)(n+p+k-2l-2q-1)+2q(q-1);

(c) The dimensions of $V_{q,l}^k$ and $W_{q,l}^k$ are given in Table II since

$$\dim V_{q,l}^k = \dim \left(\mathcal{S}^{p-2l+k-q} H_q^{\delta 0} \cap Ker \delta^* \right) \quad \text{for} \quad (k,l,q) \in S_0,$$

$$\dim W_{q,l}^k = \dim \left(\mathcal{S}^q H_{p-2l+k-q}^{\delta 0} \cap Ker i_{\overrightarrow{r}} \right) \quad \text{for} \quad (k,l,q) \in S_1.$$

References

[1] E. Bedford and T. Suwa, Eigenvalues of Hopf manifolds, American Mathemaical Society, Vol. 60 (1976), 259-264.

- [2] B. L. Beers and R. S. Millman, The spectra of the Laplace-Beltrami operator on compact, semisimple Lie groups, Amer. J. Math., **99** (4) (1975), 801-807.
- [3] M. Berger and D. Ebin, Some decompositions of the space of symmetric tensors on Riemannian manifolds, J. Diff. Geom., 3 (1969), 379-392.
- [4] M. Berger, P. Gauduchon and E. Mazet, Le spectre d'une variété riemannienne, Lecture Notes in Math., Vol 194, Springer Verlag (1971).
- [5] **A. Besse**, Einstein manifolds, Springer-Verlag, Berlin-Hiedelberg-New York (1987).
- [6] M. Boucetta, Spectre des Laplaciens de Lichnerowicz sur les sphères et les projectifs réels, Publicacions Matemàtiques, Vol. 43 (1999), 451-483.
- [7] M. Boucetta, Spectre du Laplacien de Lichnerowicz sur les projectifs complexes, C. R. Acad. Sci. Paris, t. 333, Série I, (2001), 571-576.
- [8] S. Gallot and D. Meyer, Opérateur de courbure et laplacien des formes différentielles d'une variété riemannienne, J. Math. Pures Appl., **54** (1975), 259-289.
- [9] G. W. Gibbons and M. J. Perry, Quantizing gravitational instantons, Nuclear Physics B, Vol. 146, Issue I (1978), 90-108.
- [10] **A. Ikeda and Y. Taniguchi,** Spectra and eigenforms of the Laplacian on S^n and $P^n(\mathbb{C})$, Osaka J. Math., **15** (3) (1978), 515-546.
- [11] I. Iwasaki and K. Katase, On the spectra of Laplace operator on $\wedge^*(S^n)$, Proc. Japan Acad., **55**, Ser. A (1979), 141-145.
- [12] **E. Kaneda,** The spectra of 1-forms on simply connected compact irreducible Riemannian symmetric spaces, J. Math. Kyoto Univ., **23** (1983), 369-395 and **24** (1984), 141-162.
- [13] **A. Lévy-Bruhl-Laperrière,** Spectre de de Rham-Hodge sur les formes de degré 1 des sphères de \mathbb{R}^n $(n \ge 6)$, Bull. Sc. Math., 2^e série, **99** (1975), 213-240.
- [14] A. Lévy-Bruhl-Laperrière, Spectre de de Rham-Hodge sur l'espace projectif complexe, C. R. Acad. Sc. Paris 284 Série A (1977), 1265-1267.
- [15] **A. Lichnerowicz,** Propagateurs et commutateurs en relativité générale, Inst. Hautes Etude Sci. Publ. Math., **10** (1961).
- [16] **K. Mashimo,** Spectra of Laplacian on $G_2/SO(4)$, Bull. Fac. Gen. Ed. Tokyo Univ. of Agr. and Tech. **26** (1989), 85-92.
- [17] **K. Mashimo,** On branching theorem of the pair $(G_2, SU(3))$, Nihonkai Math. J., Vol. 8 No. 2 (1997), 101-107.

- [18] **K. Mashimo**, Spectra of the Laplacian on the Cayley projective plane, Tsukuba J. Math., Vol. **21** No. 2 (1997), 367-396.
- [19] **R. Michel,** Problème d'analyse géométrique liés à la conjecture de Blaschke, Bull. Soc. Math. France, **101** (1973), 17-69.
- [20] K. Pilch and N. Schellekens, Formulas of the eigenvalues of the Laplacian on tensor harmonics on symmetric coset spaces, J. Math. Phys., **25** (12) (1984), 3455-3459.
- [21] **C. Tsukamoto,** The sepctra of the Laplace-Beltrami operator on $SO(n+2)/SO(2) \times SO(n)$ and $Sp(n+1)/Sp(1) \times Sp(n)$, Osaka J. Math. **18** (1981), 407-226.
- [22] **N. P. Warner,** The spectra of operators on $\mathbb{C}P^n$, Proc. R. Soc. Lond. A **383** (1982), 217-230.