A simple test of quantumness for a single system

Robert Ali
ki

Institute of Theoretical Physics and Astrophysics, University of Gdańsk, Wita Stwosza 57, PL 80-952 Gdańsk, Poland Ni
holas Van Ryn

S
hool Of Physi
s, Quantum Resear
h Group, University of KwaZulu-Natal,

Westville Campus, Private Bag x54001, Durban, South Afri
a.

November 17, 2018

Abstract

We propose a simple test of quantumness which can decide whether for the given set of accessible experimental data the classical model is insufficient. Take two observables A, B such that for any state ψ their mean values satisfy $0 \le \langle \psi | A | \psi \rangle \le \langle \psi | B | \psi \rangle \le 1$. If there exists a state ϕ such that the second moments fulfill the inequality $\langle \phi | A^2 | \phi \rangle$ > $\langle \phi | B^2 | \phi \rangle$ then the system cannot be described by the classical probabilistic scheme. An example of an optimal triple (A, B, ϕ) in the case of a qubit is given.

Although we are confident that the proper theory describing all physical phenomena is the quantum theory, there are many situations where the classical description in terms of functions and probability distributions over a suitable "phase-space" is sufficient. In particular, the systems consisting of a large number of particles and/or emerging in quantum states characterized by large quantum numbers are supposed to behave classically. The standard explanation of this fact refers to the stability properties of quantum states with respect to the interaction with an environment. For large quantum systems the interaction with the environment is so strong that most quantum states rapidly decay (decohere) and the remaining manifold of experimentally accessible states can be described by classical models. However, the actual border between quantum and classical worlds is still a topic of theoretical debate and experimental efforts [1]. This question is particularly important in the field of quantum information. Useful large s
ale quantum omputations would demand preservation of some quantum properties for rather large physical systems, say at least for 10³ qubits. Moreover, some promising implementations of a qubit are based on mesoscopic systems which lie on the aforementioned border. In particular, the so-called superconducting qubits are systems composed of $10^8 - 10^9$ particles (Cooper pairs). Therefore, it is important to find a simple operational test of quantumness whi
h ould be applied to a single system implementing a qubit. It is generally believed that the simplest operational and model independent test of quantumness (or strictly speaking non-classicality) is based on Bell inequalities which involve systems composed of at least two parts [2]. However, this test is problematic for systems which cannot be spatially well-separated ("locality loophole").

We propose here a much simpler operational test which can be applied to a single system and formulated in terms of inequalities between mean values of properly hosen observables. In ontrast to Bell inequalities this test does not refer to the notion of locality, associated with the space-time, but involves only the most fundamental differences between classical and quantum observables. It is based on the observation that for any two real functions f, g satisfying

$$
0 \le f(x) \le g(x) \tag{1}
$$

and any probability distribution $\rho(x)$ the following inequality holds

$$
\langle f^2 \rangle_{\rho} \equiv \int f^2(x) \rho(x) dx \le \int g^2(x) \rho(x) dx \equiv \langle g^2 \rangle_{\rho} . \tag{2}
$$

On the other hand, as illustrated below for the case of a single qubit, for quantum systems we can always find a pair of observables A, B such that for all states ψ their mean values satisfy $0 \le \langle \psi | A | \psi \rangle \le \langle \psi | B | \psi \rangle$ but there exists a state ϕ such that for the second moments $\langle \phi | A^2 | \phi \rangle > \langle \phi | B^2 | \phi \rangle$.

The above statement follows from the interesting and nontrivial mathematical result in the theory of C^* -algebras [3], which is presented below for completeness.

Consider an abstract formalism where (bounded) observables are elements of a certain C^* -algebra A , and states are positive normalized functionals $A \ni A \mapsto \langle A \rangle_\rho$ with $\langle A \rangle_\rho$ denoting the mean value of an observable A in a state ρ . For all practical purposes we can restrict ourselves to two extreme cases: the first being a classical model where A is an algebra of functions on a certain "phase-space" with $\langle A \rangle_{\rho} = \int A(x) \rho(x) dx$, where $\rho(x)$ is some probability distribution, and the second being a finite quantum model in which A is an algebra of matrices and $\langle A \rangle_\rho = \text{Tr}(\rho A)$, where ρ is some density matrix. For any pair of observables $A, B \in \mathcal{A}$, the order relation $A \leq B$ means that $\langle A \rangle_\rho \leq \langle B \rangle_\rho$ for all states ρ (in fact it is enough to take all pure states). Now we can formulate the following:

Theorem. If the following implication

$$
0 \le A \le B \Longrightarrow A^2 \le B^2 \tag{3}
$$

always holds then the algebra A is commutative, *i.e.* isomorphic to the algebra of continuous functions on a certain compact space.

As a consequence of the above theorem, for any quantum system there exists a pair of observables (identified with matrices) (A, B) such that the eigenvalues of A, B and B – A are nonnegative but the matrix $B^2 - A^2$ possesses at least one negative eigenvalue.

In order to apply our test of quantumness in an experiment, one should first guess a pair of observables A, B with nonnegative values of possible outcomes and perform a statistical test of the inequality $\langle A \rangle_\rho \leq \langle B \rangle_\rho$ with as large as possible number of different, generally mixed, initial states ρ . These states should be as pure as possible, otherwise the quantumness could be not detected. Then one should search amongst these for any states σ satisfying the relation for the second moments $\langle A^2 \rangle_{\sigma} > \langle B^2 \rangle_{\sigma}$. If the violation of the classical relation [\(2\)](#page-0-0) holds, it means that the system exhibits some quantum characteristics. Although there are always infinitely many triples (A, B, σ) , the effects of external noise acting on the system and measuring apparatus can easily wash out the deviations from "classicality". Therefore, instead of a random guess it is useful to find the examples of such triples which maximally violate classicality and can be used to optimally design the experimental setting. This can easily be achieved in the ase of a qubit whi
h is the most important example for quantum information.

We search for a pair of 2×2 matrices A, B and a pure state ϕ which satisfy

$$
0 \le A \le B \le I \ , \langle A^2 \rangle_{\phi} \equiv \langle \phi | A^2 | \phi \rangle > \langle B^2 \rangle_{\phi} \equiv \langle \phi | B^2 | \phi \rangle \ , \tag{4}
$$

where the observables A and B are normalized in such a way that their upper bound is the identity. Such a triple is given as an example as

$$
A = \begin{pmatrix} a_1 & \xi \\ \xi^* & a_2 \end{pmatrix}, \ B = \begin{pmatrix} 1 & 0 \\ 0 & b \end{pmatrix}, \ \phi = \begin{pmatrix} \alpha \\ \beta \end{pmatrix}.
$$
\n⁽⁵⁾

The matrix B is chosen to be diagonal, with the identity as its upper bound, and this choice of basis can be made since the solution to this problem is unique up to unitary equivalence. From the upper bound it can be seen that both eigenvalues of B should be at most 1, and to maximize the violation of (3) one of the eigenvalues is chosen to be fixed at 1. The usual condition $|\alpha|^2 + |\beta|^2 = 1$ applies for the parameters of the state ϕ . The positivity of the observables A, B and $(B - A)$ is ensured by the requirement that both their diagonal elements and determinants are positive. These onditions are expressed below as;

$$
0 \le b \le 1,\tag{6}
$$

$$
0 \le a_1 \le 1,\tag{7}
$$

$$
0 \le a_1 a_2 - |\xi|^2,\tag{8}
$$

and

$$
0 \le (1 - a_1)(b - a_2) - |\xi|^2. \tag{9}
$$

In order for this triple to satisfy equation [\(4\)](#page-1-1), it is sufficient that one of the eigenvalues of $(B^2 - A^2)$ is found to be negative while remaining within the onstraints listed above.

The eigenvalues of the matrix $(B^2 - A^2)$ are found to be

$$
\frac{1}{2}\left(b^2+1-a_1^2-a_2^2-2|\xi|^2\pm\sqrt{(b^2-1+a_1^2-a_2^2)^2+4(a_1+a_2)^2|\xi|^2}\right).
$$
\n(10)

In order to find the most negative value which one of the eigenvalues can attain, one need only consider one of the two. Since the square root is always positive, one eigenvalue always remains greater than the other. A numerical technique is used to calculate the maximal violation of the inequality [\(3\)](#page-1-0) since we are unable to find an exact solution to this optimisation problem. In finding the optimal set of parameters a_1, a_2, b and real ξ , it can be seen

that the maximal violation of the inequality [\(3\)](#page-1-0) arises when the onditions [\(8\)](#page-1-2) and [\(9\)](#page-1-3) are equalities rather than inequalities. Using

$$
a_1 a_2 = |\xi|^2,\t\t(11)
$$

and

$$
(1 - a_1)(b - a_2) = |\xi|^2 \tag{12}
$$

reduces the problem from four unknowns to two, and the triplet can then be written as

$$
A = \begin{pmatrix} a_1 & \sqrt{a_1 a_2} \\ \sqrt{a_1 a_2} & a_2 \end{pmatrix}, \ B = \begin{pmatrix} 1 & 0 \\ 0 & \frac{a_2}{1 - a_1} \end{pmatrix}, \ \phi = \begin{pmatrix} \alpha \\ \beta \end{pmatrix}.
$$
 (13)

The parameters which result in one of the eigenvalues of $(B^2 - A^2)$ attaining it's most negative value, and thus maximally violating inequality [\(3\)](#page-1-0), while still remaining within the onstraints give us the triplet

$$
A = \begin{pmatrix} 0.724 & 0.249 \\ 0.249 & 0.0854 \end{pmatrix}, \ B = \begin{pmatrix} 1 & 0 \\ 0 & 0.309 \end{pmatrix}, \ \phi = \begin{pmatrix} 0.391 \\ 0.920 \end{pmatrix}.
$$
 (14)

In this example, the value of $\langle \phi | B | \phi \rangle - \langle \phi | A | \phi \rangle$ is 0.0528, a positive value, whereas it can be seen that $\langle \phi | B^2 | \phi \rangle \langle \phi | A^2 | \phi \rangle = -0.0590$ which clearly demonstrates the quantum nature of this example. The eigenvectors and corresponding eigenvalues of A using these parameters is calculated to be

$$
A \begin{pmatrix} 0.946 \\ 0.325 \end{pmatrix} = 0.809 \begin{pmatrix} 0.946 \\ 0.325 \end{pmatrix}
$$
 (15)

and

$$
A\begin{pmatrix} -0.325\\ 0.946 \end{pmatrix} = \begin{pmatrix} 0\\ 0 \end{pmatrix}.
$$
 (16)

To give a concrete example, one can apply these results to the polarization of a single photon. Choosing a polarization basis as $|H\rangle, |V\rangle$ and attributing the values 1 to $|H\rangle$ and 0.309 to $|V\rangle$ we obtain the observable B. The observable A corresponds to a rotated polarization basis $|H'\rangle = \cos(19°)|H\rangle + \sin(19°)|V\rangle$, $|V'\rangle =$ $-\sin(19°)|H\rangle + \cos(19°)|V\rangle$ with the eigenvalues 0.809 and 0, respectively. The maximal violation of classicality should be observed in the neighborhood of the state $|\phi\rangle = \cos(67^\circ)|H\rangle + \sin(67^\circ)|V\rangle$.

In principle, the proposed test with the parameters obtained above could be used to support the quantum picture for different implementations of a qubit including, for example, "superconducting qubits" with a still questionable quantum character [4].

Acknowledgements. The authors thank P. Badziag, M. Horodecki, R. Horodecki, W.A. Majewski and M. Zukowski for discussions. Financial support by the POLAND/SA COLLABORATION PROGRAMME of the National Research Foundation of South Africa and the Polish Ministry of Science and Higher Education and by the European Union through the Integrated Project SCALA is acknowledged.

Referen
es

- [1] E. Joos, H.D. Zeh, C. Kiefer, D. Giulini, J. Kupsch, and I,-O. Stamatescu, Decoherence and the Appearance of a Classi
al World in Quantum Theory, 2nd ed. Springer,Berlin 2003.
- [2] J.S. Bell, Rev. Mod. Phys. $38, 447$ (1966)
- [3] R.V. Kadison and J.R. Ringrose, Fundamentals of the Theory of Operator Algebras: Elementary Theory, A
ademi Press, New York ,1983
- [4] R. Alicki, [quant-ph/0610008](http://arxiv.org/abs/quant-ph/0610008)