

SYMMETRIES IN THE SYSTEM OF TYPE $D_4^{(1)}$

YUSUKE SASANO

ABSTRACT. In this paper, we propose a 4-parameter family of coupled Painlevé III systems in dimension four with affine Weyl group symmetry of type $D_4^{(1)}$. We also propose its symmetric form in which the $D_4^{(1)}$ -symmetries become clearly visible.

0. STATEMENT OF MAIN RESULTS

In [9, 10, 11], we presented some types of coupled Painlevé systems with various affine Weyl group symmetries. In this paper, we present a 4-parameter family of coupled Painlevé III systems with affine Weyl group symmetry of type $D_4^{(1)}$. Since these universal Bäcklund transformations have Lie theoretic origin, similarity reduction of a Drinfeld-Sokolov hierarchy admits such a Bäcklund symmetry. The aim of this paper is to introduce the system of type $D_4^{(1)}$. After our discovery of this system, they were studied from the viewpoint of Drinfeld-Sokolov hierarchy by K. Fuji independently (cf. [1]), and he succeeded to obtain our system by similarity reduction of the Drinfeld-Sokolov hierarchy of type $D_4^{(1)}$.

At first, we propose a 4-parameter family of autonomous ordinary differential systems with the invariant divisors f_i as variables:

$$(1) \quad \left\{ \begin{array}{l} \frac{df_0}{dt} = -(2f_1g_1 + \alpha_1)f_0 - \alpha_0f_1, \\ \frac{df_1}{dt} = -(2f_0g_1 + \alpha_0)f_1 - \alpha_1f_0, \\ \frac{df_2}{dt} = \{(f_0 + f_1)g_1 + (f_3 + f_4)g_2 + 1\}f_2 - 2\alpha_2g_1g_2, \\ \frac{df_3}{dt} = -(2f_4g_2 + \alpha_4)f_3 - \alpha_3f_4, \\ \frac{df_4}{dt} = -(2f_3g_2 + \alpha_3)f_4 - \alpha_4f_3, \\ \frac{dg_1}{dt} = (f_0 + f_1)g_1^2 - \{(f_3 + f_4)g_2 - \alpha_0 - \alpha_1\}g_1 + (f_3 + f_4)f_2, \\ \frac{dg_2}{dt} = (f_3 + f_4)g_2^2 - \{(f_0 + f_1)g_1 - \alpha_3 - \alpha_4\}g_2 + (f_0 + f_1)f_2. \end{array} \right.$$

Here $f_0, f_1, \dots, f_4$ and g_1, g_2 denote unknown complex variables and $\alpha_0, \dots, \alpha_4$ are the parameters satisfying the condition:

$$\alpha_0 + \alpha_1 + 2\alpha_2 + \alpha_3 + \alpha_4 = 1.$$

Proposition 0.1. *This system has the following invariant divisors:*

invariant divisors	parameter's relation
$f_0 := 0$	$\alpha_0 = 0$
$f_1 := 0$	$\alpha_1 = 0$
$f_2 := 0$	$\alpha_2 = 0$
$f_3 := 0$	$\alpha_3 = 0$
$f_4 := 0$	$\alpha_4 = 0$

Theorem 0.1. *This system is invariant under the transformations $s_0, \dots, s_4$ defined as follows: with the notation $(*) := (f_0, f_1, \dots, f_4, g_1, g_2; \alpha_0, \alpha_1, \dots, \alpha_4)$,*

$$\begin{aligned}
s_0 : (*) &\rightarrow (f_0, f_1, f_2 + \frac{\alpha_0 g_2}{f_0}, f_3, f_4, g_1 + \frac{\alpha_0}{f_0}, g_2; -\alpha_0, \alpha_1, \alpha_2 + \alpha_0, \alpha_3, \alpha_4), \\
s_1 : (*) &\rightarrow (f_0, f_1, f_2 + \frac{\alpha_1 g_2}{f_1}, f_3, f_4, g_1 + \frac{\alpha_1}{f_1}, g_2; \alpha_0, -\alpha_1, \alpha_2 + \alpha_1, \alpha_3, \alpha_4), \\
s_2 : (*) &\rightarrow (f_0 - \frac{\alpha_2 g_2}{f_2}, f_1 - \frac{\alpha_2 g_2}{f_2}, f_2, f_3 - \frac{\alpha_2 g_1}{f_2}, f_4 - \frac{\alpha_2 g_1}{f_2}, g_1, g_2; \\
&\quad \alpha_0 + \alpha_2, \alpha_1 + \alpha_2, -\alpha_2, \alpha_3 + \alpha_2, \alpha_4 + \alpha_2), \\
s_3 : (*) &\rightarrow (f_0, f_1, f_2 + \frac{\alpha_3 g_1}{f_3}, f_3, f_4, g_1, g_2 + \frac{\alpha_3}{f_3}; \alpha_0, \alpha_1, \alpha_2 + \alpha_3, -\alpha_3, \alpha_4), \\
s_4 : (*) &\rightarrow (f_0, f_1, f_2 + \frac{\alpha_4 g_1}{f_4}, f_3, f_4, g_1, g_2 + \frac{\alpha_4}{f_4}; \alpha_0, \alpha_1, \alpha_2 + \alpha_4, \alpha_3, -\alpha_4).
\end{aligned}
\tag{2}$$

Theorem 0.2. *This system has two first integrals:*

$$\frac{d(f_0 - f_1)}{dt} = \frac{d(f_3 - f_4)}{dt} = 0, \quad \frac{d(f_2 - g_1 g_2)}{dt} = f_2 - g_1 g_2.$$

From this, we have

$$f_0 = f_1 - 1, \quad f_3 = f_4 - 1, \quad f_2 - g_1 g_2 = e^{(t+c)}.$$

Here we set

$$t + c = \log T, \quad x := g_1, \quad y := f_1, \quad z := g_2, \quad w := f_4,$$

then we obtain a 4-parameter family of coupled Painlevé III systems in dimension four with affine Weyl group symmetry of type $D_4^{(1)}$ explicitly given by

$$\begin{cases}
\frac{dx}{dT} = \frac{2x^2 y - x^2 + (\alpha_0 + \alpha_1)x}{T} - 1 + 2w, \\
\frac{dy}{dT} = \frac{-2xy^2 + 2xy - (\alpha_0 + \alpha_1)y + \alpha_1}{T}, \\
\frac{dz}{dT} = \frac{2z^2 w - z^2 + (\alpha_3 + \alpha_4)z}{T} - 1 + 2y, \\
\frac{dw}{dT} = \frac{-2zw^2 + 2zw - (\alpha_3 + \alpha_4)w + \alpha_4}{T}
\end{cases}
\tag{3}$$

with the Hamiltonian

$$\begin{aligned}
H &= \frac{x^2 y^2 - x^2 y + (\alpha_0 + \alpha_1)xy - \alpha_1 x}{T} - y \\
&+ \frac{z^2 w^2 - z^2 w + (\alpha_3 + \alpha_4)zw - \alpha_4 z}{T} - w + 2yw.
\end{aligned}
\tag{4}$$




FIGURE 1. The transformations s_i satisfy the relations: $s_i^2 = 1$ ($i = 0, 1, 2, 3, 4$), $(s_0 s_1)^2 = (s_0 s_3)^2 = (s_0 s_4)^2 = (s_1 s_3)^2 = (s_1 s_4)^2 = (s_3 s_4)^2 = 1$, $(s_0 s_2)^3 = (s_1 s_2)^3 = (s_3 s_2)^3 = (s_4 s_2)^3 = 1$.

Theorem 0.3. *This system is invariant under the transformations $s_0, \dots, s_4, \pi_1, \pi_2, \pi_3$ defined as follows: with the notation $(*) := (x, y, z, w, T; \alpha_0, \alpha_1, \alpha_2, \alpha_3, \alpha_4)$,*

$$\begin{aligned}
 s_0 : (*) &\rightarrow (x + \frac{\alpha_0}{y-1}, y, z, w, T; -\alpha_0, \alpha_1, \alpha_2 + \alpha_0, \alpha_3, \alpha_4), \\
 s_1 : (*) &\rightarrow (x + \frac{\alpha_1}{y}, y, z, w, T; \alpha_0, -\alpha_1, \alpha_2 + \alpha_1, \alpha_3, \alpha_4), \\
 s_2 : (*) &\rightarrow (x, y - \frac{\alpha_2 z}{xz+T}, z, w - \frac{\alpha_2 x}{xz+T}, T; \\
 &\quad \alpha_0 + \alpha_2, \alpha_1 + \alpha_2, -\alpha_2, \alpha_3 + \alpha_2, \alpha_4 + \alpha_2), \\
 s_3 : (*) &\rightarrow (x, y, z + \frac{\alpha_3}{w-1}, w, T; \alpha_0, \alpha_1, \alpha_2 + \alpha_3, -\alpha_3, \alpha_4), \\
 s_4 : (*) &\rightarrow (x, y, z + \frac{\alpha_4}{w}, w, T; \alpha_0, \alpha_1, \alpha_2 + \alpha_4, \alpha_3, -\alpha_4), \\
 \pi_1 : (*) &\rightarrow (-x, 1 - y, z, w, -T; \alpha_1, \alpha_0, \alpha_2, \alpha_3, \alpha_4), \\
 \pi_2 : (*) &\rightarrow (x, y, -z, 1 - w, -T; \alpha_0, \alpha_1, \alpha_2, \alpha_4, \alpha_3), \\
 \pi_3 : (*) &\rightarrow (z, w, x, y, T; \alpha_3, \alpha_4, \alpha_2, \alpha_0, \alpha_1).
 \end{aligned}
 \tag{5}$$

REFERENCES

- [1] K. Fuji and T. Suzuki, *The sixth Painlevé equation arising from $D_4^{(1)}$ hierarchy*, J. Phys. A: Math. Gen. **39** (2006), 12073–12082.
- [2] T. Matano, A. Matsumiya and K. Takano, *On some Hamiltonian structures of Painlevé systems, II*, J. Math. Soc. Japan **51** (1999), 843–866.
- [3] M. Noumi and Y. Yamada, *Affine Weyl Groups, Discrete Dynamical Systems and Painlevé Equations*, Comm Math Phys. **199** (1998), 281–295.
- [4] M. Noumi, K. Takano and Y. Yamada, *Bäcklund transformations and the manifolds of Painlevé systems*, Funkcial. Ekvac. **45** (2002), 237–258.
- [5] M. Noumi, *Painlevé Equations Through Symmetry*, American Mathematical Society, **223** (2004).
- [6] Y. Sasano, *Coupled Painlevé V systems in dimension 4*, Funkcial. Ekvac. **49** (2006), 133–161.
- [7] Y. Sasano, *The phase space of coupled Painlevé III system in dimension four*, to appear in Kyoto Journal.
- [8] Y. Sasano, *Coupled Painlevé II systems in dimension four and the systems of type $A_4^{(1)}$* , Tohoku. Math. J. **58** (2006), 529–548.

- [9] Y. Sasano, *Higher order Painlevé equations of type $D_l^{(1)}$* , RIMS Kokyuroku **1473** (2006), 143–163.
- [10] Y. Sasano, *Coupled Painlevé VI systems in dimension four with affine Weyl group symmetry of types $B_6^{(1)}$, $D_6^{(1)}$ and $D_7^{(2)}$* , submitted to Nagoya Journal.
- [11] Y. Sasano, *Four-dimensional Painlevé systems of types $D_5^{(1)}$ and $B_4^{(1)}$* , submitted to Nagoya Journal.
- [12] Y. Sasano and Y. Yamada, *Symmetry and holomorphy of Painlevé type systems*, RIMS Kokyuroku **B2** (2007), 215–225.
- [13] T. Shioda and K. Takano, *On some Hamiltonian structures of Painlevé systems I*, Funkcial. Ekvac. **40** (1997), 271–291.
- [14] M. Suzuki, N. Tahara and K. Takano, *Hierarchy of Bäcklund transformation groups of the Painlevé equations*, J. Math. Soc. Japan **56** (2004), 1221–1232.
- [15] N. Tahara, *An augmentation of the phase space of the system of type $A_4^{(1)}$* , Kyushu J. Math. **58** (2004), 393–425.

GRADUATE SCHOOL OF MATHEMATICAL SCIENCES, THE UNIVERSITY OF TOKYO, TOKYO 153-8914, JAPAN.

E-mail address: sasano@ms.u-tokyo.ac.jp