The Dark Matter Puzzle And Other Issues

B.G. Sidharth

International Institute for Applicable Mathematics & Information Sciences Hyderabad (India) & Udine (Italy)

B.M. Birla Science Centre, Adarsh Nagar, Hyderabad - 500 063 (India)

Abstract

We consider the problem of the flattening of the velocity curves in galactic discs and the consequent postulation of dark matter from three different but converging perspectives– a change in the large scale dimensionality of space, a variation of G and the MOND approach. We also discuss the paradigm of the universe itself being a Black Hole.

1 Introduction

It is many decades now since the existence of dark, that is non luminous matter was postulated, though the identity of this dark matter has only been a matter of guess work. The reason for invoking the hypothetical dark matter is well known– the velocities of stars in galaxies should tend to zero using usual dynamics, as we approach the edge of the disc. Instead astrophysical observation has consistently shown that the velocity curves flatten out, that is the velocities tend a constant rather than zero. So Zwicky and others postulated that there was matter other than the visible matter which gave a greater mass to the galaxies, and this in turn would explain the velocity discrepancy [\[1\]](#page-17-0).

The question then arose, "What exactly is this dark matter?". Over the years several hypotheses have been put forth– it could be hot dark matter or it could be cold dark matter. These could range from weakly interacting massive particles (WIMPS) to cold neutrinos. Or there could be the missing monopoles, or undetectable brown dwarf stars or even black holes and so on. To this day the question has remained unresolved.

There have however been alternative suggestions to explain the flat velocity curves. We will discuss three of these, two put forward by the author and the other by Milgrom, and try to find an interrelationship.

2 Less Than Three Dimensional Space?

The author (with A.D. Popova) [\[2\]](#page-17-1) suggested that the dimensionality of space falls off asymptotically, and this would explain astrophysical observations including the dark matter problem. Indeed it had already been argued that the dimensionality of space could be expressed by a non integer number that is less than three on large scales [\[3,](#page-17-2) [4\]](#page-17-3). Indeed the three dimensionality of our immediate space may be necessary, for the very existence of atoms, as was pointed by Ehrenfest long ago [\[5\]](#page-17-4). Similarly this dimensionality may also be required for usual wave propagation $[6, 7, 8]$ $[6, 7, 8]$ $[6, 7, 8]$. All this is at what we may call intermediate scales. At different scales of measurement, the dimensionality could be different [\[9\]](#page-17-8). This fact could explain the dark matter problem, as we will now argue.

More generally, the dependence of matter on distance $M(r)$, obtained from observing 21cm neutral hydrogen emission of gas clouds moving around a galaxy far from its visual bounds (the continuation of a rotation curve) [\[10,](#page-17-9) [11\]](#page-17-10) is

$$
M \propto r^{1.2 \div 1.3} \tag{1}
$$

This conclusion reflects the fact that the observed rotation velocity slightly increases at outer parts of galaxies, so the growth of M is interpreted as the presence of some dark halo besides luminous matter. Moreover, the amount of dark matter grows relatively to luminous matter when coming to larger and larger scales [\[12\]](#page-17-11).

However, even the nonrelativistic (Newtonian) consideration of gravitational forces in spaces with lesser than three dimensions, enables us in principle to bring in correlation dynamics and "the shortage" of luminous matter. It would be very difficult to take into account the smooth fall of dimensionality because we do not know the law of such a fall. In order to make some estimates we roughly assume that on some relative distance R_0 the dimensionality changes by a leap from 3 to $n < 3$. We consider the rotation curves of disk galaxies –similar considerations apply for the dynamics of double galaxies and the dispersion of velocities in elliptical (spheroidal) galaxies. We show how one can lower the estimates of masses of these systems under

the assumption that the dimensionality is lesser than three starting even at scales of the order of a typical galaxy's size. We also discuss the possible hierarchical change of dimensionality. We only demonstrate the principal possibility to lower dynamical mass and refrain from making any numerical estimates as yet because of too many uncertain parameters.

Certainly, we know now of a constructive physical model which can describe noninteger and nonconstant dimensionality. We can only outline some suggestive arguments for it. The first of possible suggestions comes from the fractal theory [\[13\]](#page-17-12); the space itself may have a fractal-like structure. The second suggestion is that effectively, if we consider individually each object in the Universe leaving aside other objects, then we can perceive a space between us and this object as 2-dimensional because one spatial direction is fixed as a line from us to the object, and the other direction can be fixed by a vector of relative velocity of the object with respect to us. Thus, may be our space filled by distant separated objects consists of a set of (perhaps non-connected) 2-dimensional subspaces for which effectively $2 \leq n \leq 3$. The third suggestion comes from the existence of the large-scale structure of the Universe in distribution of galaxies, their groups, clusters, voids and superclusters. There is the tendency for matter to perform oblate structures at each hierarchical level. Possibly, the structure and dimensionality of our space itself might reflect the distribution of (luminous) cosmic matter.

3 Newtonian Consideration

In 3 and n dimensions, the expressions for the gravitational forces acting between the mass M and a unit mass separated by the distance r are

$$
F^{(3)} = -\frac{G^{(3)}M}{r^2} \tag{2}
$$

and

$$
F^{(n)} = -\frac{G^{(n)}M}{r^{n-1}},
$$
\n(3)

respectively, where $G^{(3)}$ and $G^{(n)}$ are relevant gravitational constants. The corresponding potentials ($\vec{F} = -\vec{\nabla}\Phi$ by definition) up to arbitrary constants $C^{(3)}$ and $C^{(n)}$ are $\gamma(2)$

$$
\Phi^{(3)} = -\frac{G^{(3)}M}{r} + C^{(3)},\tag{4}
$$

and for $n \neq 2$

$$
\Phi^{(n)} = -\frac{1}{n-2} \frac{G^{(n)}M}{r^{n-2}} + C^{(n)} \tag{5}
$$

In this case $n = 2$,

$$
\Phi^{(2)} = G^{(2)}Mlnr + C^{(2)}.
$$

Now, let us assume that at some relative distance R_0 from a body the dimensionality changes by leap from 3 to n . The condition of matching the forces [\(2\)](#page-2-0) and [\(3\)](#page-2-1) at R_0 gives the connection between the gravitational constants

$$
G^{(n)} = G^{(3)} R_0^{n-3} \tag{6}
$$

Thus, the improved force in accordance with our conception is [\(2\)](#page-2-0) for $r \leq R_0$ and [\(3\)](#page-2-1) for $r > R_0$ with [\(4\)](#page-2-2):

$$
F^{imp} = \begin{cases} F^{(3)}, & r \le R_0, \\ F^{(n)}, & r > R_0. \end{cases}
$$
 (7)

The condition of matching the potentials (4) and (5) at R_0 is also required, and leads to the following expression for the "n-dimensional" constant $C^{(3)}$ = 0 is chosen in (4) ,

$$
C^{(n)} = \frac{3 - n G^{(3)}M}{n - 2 R_0}
$$

for $n \neq 2$, and

$$
C^{(2)} = -\frac{G^{(3)}M}{R_0}(1 + \ln R_0)
$$

for $n = 2$.

Thus, the improved potential in accordance with our conception is [\(4\)](#page-2-2) for $r \leq R_0$ and [\(5\)](#page-3-0) for $r > R_0$:

$$
\Phi^{imp} = \begin{cases} \Phi^{(3)}, & r \le R_0, \\ \Phi^{(n)}, & r > R_0. \end{cases}
$$
 (8)

Let us stress that we can only think of R_0 as of a relative distance between any bodies. Otherwise, first the conception of relativity of space which is an achievement of Einstein's physics, would be violated. Second, there would be troubles with the universality of a gravitational attraction.

Indeed, let a sphere with the radius R_0 be "rigidly fixed" in space, and let bodies freely come in it and come out of it. Let a body of the mass M

be placed inside the sphere at some point O which does not coincide with the center of the sphere O , and let other bodies of the equal small mass m be outside of this sphere at the points A_1 and A_2 in such a way that $O'A_1 = O'A_2 = r$. Then, we have for the gravitational forces between each body of the mass m and that of the mass M

$$
F_1^{(n)} = -G_1^{(n)} \frac{mM}{r^{n-1}}, F_2^{(n)} \frac{mM}{r^{n-1}}
$$

where $G_1^{(n)} = G^{(3)} R_1^{n-3}$ and $G_2^{(n)} = G^{(3)} R_2^{n-3}$ are the gravitational constants found from the continuity of forces at the sphere, and R_1 and R_2 are the distances from the point O' to the points of intersections of the rays $O'A_1$ and $O'A_2$ with the sphere. Thus, if the above consideration were true, then the "gravitational constant" would not be a constant, and the law of gravitation attraction would be anisotropic and in-homogeneous. Moreover, in the case $n < 3$, the body of the mass M at O' which tends to the sphere on the insider $(R_1 \rightarrow 0 \text{ or } R_2 \rightarrow 0)$ would produce an infinitely divergent "gravitational" constant".

That is why we admit R_0 as only relative distance. Perhaps, our consideration would be less rough, if we consider the change of dimensionality which occurs by leaps several times from 3 to n_1 at R_0 , from n_1 to n_2 at R_1 , and so on, and from n_j to $n_{j+1} > 2$ at R_j . Then, we have the chain of relations between the gravitational constants

$$
G^{(n_1)} = G^{(3)} R_0^{n_1 - 3}
$$

$$
G^{(n_2)} = G^{(n_1)} R_1^{n_2 - n_1} = G^{(3)} R_0^{n_1 - 3} R_1^{n_2 - n_1}
$$
(9)
$$
G^{(n_j + 1)} = G^{(n_j)} R_j^{n_{j+1}^{1 - n_j}} = \dots = G^{(3)} R_0^{n_1 - 3} R_1^{n_2 - n_1} \dots R_j^{n_{j+1}^{-n_j}}
$$

The chain of relations between the constants in potentials is rather cumbersome; however the recurrence relation is

$$
C^{(n_{j+1})}=G^{(n_j)}R_j^{n_{j+1}^{-n_j}}=\cdots=G^{(3)}R_0^{n_1-3}R_1^{n_2-n_1}\cdots R_j^{n_{j+1}-n_j}.
$$

The chain of relations between the constants in potentials is rather cumbersome; however the recurrence relation is

$$
C^{(n_{j+1})} = C^{(n_j)} + \frac{n_j - n_{j+1}}{(n_j - 2)(n_{j+1} - 2)} \frac{G^{(n_j)}M}{R_j^{n_j - 2}}
$$

The three subsections below present the application of the force [\(7\)](#page-3-1) (the potential of [\(8\)](#page-3-2) to determinations of galactic masses. We shall return to discussing the problem of the dimensionality change after considering the three examples below.

4 Rotation Curves of Disk Galaxies

A rough calculation of rotation velocity v at the distance r far from the center of a galaxy (i.e., when all its mass is effectively concentrated near the center or spherically distributed around it) is based on the equality of the centrifugal force and gravitational force.

In the 3-dimensional space, using [\(2\)](#page-2-0)

$$
\frac{v^2}{r} = \frac{G^{(3)}M_g^{(3)}}{r^2} \tag{10}
$$

In accordance with our conception, we can write

$$
M_g^{dyn} \equiv M_g^{(3)} = \frac{v^2 r}{G^{(3)}},
$$

i.e., we call the dynamical galactic mass, M_g^{dyn} , a mass calculated as if our space were 3-dimensional.

In the *n*-dimensional space, if $r \gg R_0$ then equality [\(10\)](#page-5-0) should be replaced by the following (with the use of [\(3\)](#page-2-1):

$$
\frac{v^2}{r} = \frac{G^{(n)}M_g^{(n)}}{r^{n-1}},
$$

so that we call a mass calculated in the n -dimensional space the true galactic mass, M_g^{true} :

$$
M_g^{true} \equiv M_g^{(n)} = \frac{v^2 r^{n-2}}{G^{(n)}} = \left(\frac{R_0}{r}\right)^{3-n} M_g^{dyn}
$$
 (11)

where in the last equality [\(6\)](#page-3-3) is used. When $2 < n < 3$ the factor at M_g^{dyn} is lesser than unity, therefore $M_g^{true} < M_g^{dyn}$.

The more accurate calculation of the rotation curve can be done for the case when the disk of a galaxy lies in the 2-dimensional space (or plane), and there

exist no other spatial dimensions. Let the distribution of the 2-dimensional matter density in the disk satisfy the law

$$
\rho = \rho_0 exp\left(\frac{r}{R_d}\right) \tag{12}
$$

where ρ_0 is the 2-density in the disk center, and R_d is some characteristic radius. Let $R_0 \ll R_d$. The distribution [\(12\)](#page-6-0) corresponds to the observed distribution of luminous matter in [\[13\]](#page-17-12). The velocity square in this case is given as follows

$$
v^{2}(r) = 2\pi\rho_{0}G^{(3)}\frac{R_{d}^{2}}{R_{d}}\left[1 - \left(1 + \frac{r}{R_{d}}\right)exp\left(-\frac{r}{R_{d}}\right)\right]
$$
(13)

The function [\(13\)](#page-6-1) monotonically increases from zero and tends to a constant value at $r \to \infty$. That is why the dynamical mass calculated with the aid of [\(13\)](#page-6-1) tends to grow linearly at larger $r : M^{dyn} \propto r$. However, at any finite r we can effectively write $M_g^{dyn} \propto r^{\beta}$ where always $\beta > 1$. Probably, this fact could explain the dependence [\(1\)](#page-1-0), meaning that our real space has the dimensionality which is very near to two at the scales of the outer parts of galaxies. This also explains the flattening of the rotational curves, without invoking dark matter.

Alternatively, we note that from the above, for $w = 2$, we get

$$
v^2 = GMlnr,
$$

or

$$
v\frac{dv}{dr} \propto \frac{1}{r} \to 0
$$

as r becomes large, so that $\frac{dv}{dr} \to 0$, because v does not $\to 0$. So, $v \to a$ constant value.

5 The Time Variation of the Gravitational Constant

We now come to the author's cosmological model which in 1997 predicted a dark energy driven accelerating universe with a small cosmological constant. It may be recalled that at that time the ruling paradigm embodied in the hot big bang standard cosmological model was exactly the opposite. This

model has been discussed in detail (Cf.ref.[\[7,](#page-17-6) [8,](#page-17-7) [14,](#page-17-13) [15\]](#page-17-14)). In this model it turns out that the gravitational constant, rather as in Dirac cosmology, has the following time dependence

$$
G = \frac{\beta}{T} \tag{14}
$$

where in our cosmology β is given in terms of the constant microphysical parameters.

We next observe that from [\(14\)](#page-7-0) it follows that

$$
G = G_0 \left(1 - \frac{t}{t_0} \right) \tag{15}
$$

where G_0 is the present value of G and t_0 is the present age of the universe and t the time elapsed from the present epoch. Similarly one could deduce that $(Cf.ref.[1]),$ $(Cf.ref.[1]),$ $(Cf.ref.[1]),$

$$
r = r_0 \left(\frac{t_0}{t_0 + t}\right) \tag{16}
$$

We next use Kepler's Third law [\[16\]](#page-17-15):

$$
\tau = \frac{2\pi a^{3/2}}{\sqrt{GM}}\tag{17}
$$

 τ is the period of revolution, a is the orbit's semi major axis, and M is the mass of the sun. Denoting the average angular velocity of the planet by

$$
\dot{\Theta} \equiv \frac{2\pi}{\tau},
$$

it follows from (15) , (16) and (17) that

$$
\dot{\Theta} - \dot{\Theta}_o = \dot{\Theta}_0 \frac{t}{t_o},
$$

where the subscript *o* refers to the present epoch, Whence,

$$
\omega(t) \equiv \Theta - \Theta_o = \frac{\pi}{\tau_o t_o} t^2 \tag{18}
$$

Equation [\(18\)](#page-7-4) gives the average perhelion precession at time $'t'$. Specializing to the case of Mercury, where $\tau_o = \frac{1}{4}$ $\frac{1}{4}$ year, it follows from [\(18\)](#page-7-4) that the average precession per year at time $'t'$ is given by

$$
\omega(t) = \frac{4\pi t^2}{t_0} \tag{19}
$$

Whence, considering $\omega(t)$ for years $t = 1, 2, \dots, 100$, we can obtain from [\(19\)](#page-7-5), the usual total perhelion precession per century as,

$$
\omega = \sum_{n=1}^{100} \omega(n) \approx 43'',
$$

if the age of the universe is taken to be $\approx 2 \times 10^{10}$ years.

Conversely, if we use the observed value of the precession in [\(19\)](#page-7-5), we can get back the above age of the universe.

Interestingly it can be seen from [\(19\)](#page-7-5), that the precession depends on the epoch.

We next demonstrate that orbiting objects will have an anamolous inward radial acceleration.

Using the well known equation for Keplarian orbits (cf.ref.[\[16\]](#page-17-15)),

$$
\frac{1}{r} = \frac{GMm^2}{l^2}(1 + e\cos\Theta)
$$
\n(20)

$$
\dot{r}^2 \approx \frac{GM}{r} - \frac{l^2}{m^2 r^2} \tag{21}
$$

 l being the orbital angular momentum constant and e the eccentricity of the orbit, we can deduce such an extra inward radial acceleration, on differentiation of (21) and using (15) and (16) ,

$$
a_r = \frac{GM}{2t_o r \dot{r}}\tag{22}
$$

It can be easily shown from [\(20\)](#page-8-1) that (on the average),

$$
\dot{r} \approx \frac{eGM}{rv} \tag{23}
$$

For a nearly circular orbit $rv^2 \approx GM$, whence use of [\(23\)](#page-8-2) in [\(22\)](#page-8-3) gives,

$$
a_r \approx v/2t_oe \tag{24}
$$

For the earth, [\(24\)](#page-8-4) gives an anomalous inward radial acceleration $\sim 10^{-9}$ cm/sec², which is known to be the case [\[17\]](#page-17-16).

We could also deduce a progressive decrease in the eccentricity of orbits. Indeed, e in (20) is given by

$$
e^{2} = 1 + \frac{2El^{2}}{G^{2}m^{3}M^{2}} \equiv 1 + \gamma, \gamma < 0.
$$

Use of [\(15\)](#page-7-1) in the above and differenciation, leads to,

$$
\dot{e} = \frac{\gamma}{et_o} \approx -\frac{1}{et_o} \approx -\frac{10^{-10}}{e} \text{per year},
$$

if the orbit is nearly circular. (Variations of eccentricity in the usual theory have been extensively studied (cf.ref.[\[18\]](#page-17-17) for a review). On the other hand, for open orbits, $\gamma > 0$, the eccentricity would progressively increase.

We finally consider the anomalous accelerations given in [\(22\)](#page-8-3) and [\(24\)](#page-8-4) in the context of space crafts leaving the solar system.

If in [\(22\)](#page-8-3) we use the fact that $\dot{r} \leq v$ and approximate

$$
v \approx \sqrt{\frac{GM}{r}},
$$

we get,

$$
a_r \ge \frac{1}{et_o} \sqrt{\frac{GM}{r}}
$$

For $r \sim 10^{14}$ cm, as is the case of Pioneer 10 or Pioneer 11, this gives, $a_r > 10^{-11}$ cm/sec²

Interestingly Anderson et al., [\[19\]](#page-18-0) claim to have observed an anomalous inward acceleration of $\sim 10^{-9} cm/sec^2$.

6 Other Consequences

We could also explain the correct gravitational bending of light. Infact in Newtonian theory also we obtain the bending of light, though the amount is half that predicted by General Relativity $[20, 21, 22]$ $[20, 21, 22]$ $[20, 21, 22]$. In the Newtonian theory we can obtain the bending from the well known orbital equations $(Cf. also (20)),$ $(Cf. also (20)),$ $(Cf. also (20)),$

$$
\frac{1}{r} = \frac{GM}{L^2}(1 + e\cos\Theta)
$$
\n(25)

where M is the mass of the central object, L is the angular momentum per unit mass, which in our case is bc, b being the impact parameter or minimum approach distance of light to the object, and e the eccentricity of the trajectory is given by

$$
e^2 = 1 + \frac{c^2 L^2}{G^2 M^2} \tag{26}
$$

For the deflection of light α , if we substitute $r = \pm \infty$, and then use [\(26\)](#page-9-0) we get

$$
\alpha = \frac{2GM}{bc^2} \tag{27}
$$

This is half the General Relativistic value.

We next note that the effect of time variation of r is given by equation $(16)(\text{cf.}rel. [23])$ $(16)(\text{cf.}rel. [23])$ $(16)(\text{cf.}rel. [23])$ $(16)(\text{cf.}rel. [23])$. Using [\(16\)](#page-7-2) the well known equation for the trajectory is given by (Cf.[\[16,](#page-17-15) [24,](#page-18-5) [25\]](#page-18-6))

$$
u'' + u = \frac{GM}{L^2} + u\frac{t}{t_0} + 0\left(\frac{t}{t_0}\right)^2\tag{28}
$$

where $u=\frac{1}{r}$ $\frac{1}{r}$ and primes denote differenciation with respect to Θ . The first term on the right hand side represents the Newtonian contribution while the remaining terms are the contributions due to (16) . The solution of [\(28\)](#page-10-0) is given by

$$
u = \frac{GM}{L^2} \left[1 + e\cos\left\{ \left(1 - \frac{t}{2t_0} \right) \Theta + \omega \right\} \right]
$$
 (29)

where ω is a constant of integration. Corresponding to $-\infty < r < \infty$ in the Newtonian case we have in the present case, $-t_0 < t < t_0$, where t_0 is large and infinite for practical purposes. Accordingly the analogue of the reception of light for the observer, viz., $r = +\infty$ in the Newtonian case is obtained by taking $t = t_0$ in [\(29\)](#page-10-1) which gives

$$
u = \frac{GM}{L^2} + e\cos\left(\frac{\Theta}{2} + \omega\right) \tag{30}
$$

Comparison of [\(30\)](#page-10-2) with the Newtonian solution obtained by neglecting terms $\sim t/t_0$ in equations [\(16\)](#page-7-2),[\(28\)](#page-10-0) and [\(29\)](#page-10-1) shows that the Newtonian Θ is replaced by $\frac{\Theta}{2}$, whence the deflection obtained by equating the left side of [\(30\)](#page-10-2) to zero, is

$$
\cos\Theta\left(1 - \frac{t}{2t_0}\right) = -\frac{1}{e} \tag{31}
$$

where e is given by [\(26\)](#page-9-0). The value of the deflection from [\(31\)](#page-10-3) is twice the Newtonian deflection given by [\(27\)](#page-10-4). That is the deflection α is now given not by [\(28\)](#page-10-0) but by the formula,

$$
\alpha = \frac{4GM}{bc^2},\tag{32}
$$

The relation [\(32\)](#page-10-5) is the correct observed value and is the same as the General Relativistic formula.

We now come to the problem of galactic rotational curves $(cf.ref.|1|)$. We would expect, on the basis of straightforward dynamics that the rotational velocities at the edges of galaxies would fall off according to

$$
v^2 \approx \frac{GM}{r} \tag{33}
$$

However it is found that the velocities tend to a constant value,

$$
v \sim 300 \, \text{km/sec} \tag{34}
$$

This has lead to the postulation of as yet undetected additional matter, the so called dark matter.(However for an alternative view point Cf.[\[26\]](#page-18-7). We observe that from (16) it can be easily deduced that $[27]$

$$
a \equiv (\ddot{r}_o - \ddot{r}) \approx \frac{1}{t_o} (t\ddot{r}_o + 2\dot{r}_o) \approx -2\frac{r_o}{t_o^2}
$$
 (35)

as we are considering infinitesimal intervals t and nearly circular orbits. Equation (35) shows $(Cf.\text{ref}[23]$ $(Cf.\text{ref}[23]$ also) that there is an anomalous inward acceleration, as if there is an extra attractive force, or an additional central mass, as indeed we saw a little earlier. So,

$$
\frac{GMm}{r^2} + \frac{2mr}{t_o^2} \approx \frac{mv^2}{r}
$$
\n(36)

From [\(36\)](#page-11-1) it follows that

$$
v \approx \left(\frac{2r^2}{t_o^2} + \frac{GM}{r}\right)^{1/2} \tag{37}
$$

From [\(37\)](#page-11-2) it is easily seen that at distances within the edge of a typical galaxy, that is $r < 10^{23}$ cms the equation [\(33\)](#page-11-3) holds but as we reach the edge and beyond, that is for $r \geq 10^{24} \text{cm} s$ we have $v \sim 10^{7} \text{cm} s$ per second, in agreement with [\(34\)](#page-11-4).

Thus the time variation of G explains observation without invoking dark matter.

7 The MOND Approach

Milgrom [\[28\]](#page-18-9) approached the problem by modifying Newtonian dynamics at large distances. This approach is purely phenomenological. The idea was that perhaps standard Newtonian dynamics works at the scale of the solar system but at galactic scales involving much larger distances perhaps the situation is difference. However a simple modification of the distance dependence in the gravitation law, as pointed by Milgrom would not do, even if it produced the asymptotically flat rotation curves of galaxies. Such a law would predict the wrong form of the mass velocity relation. So Milgrom suggested the following modification to Newtonian dynamics: A test particle at a distance r from a large mass M is subject to the acceleration a given by

$$
a^2/a_0 = M G r^{-2},
$$
\n(38)

where a_0 is an acceleration such that standard Newtonian dynamics is a good approximation only for accelerations much larger than a_0 . The above equation however would be true when a is much less than a_0 . Both the statements can be combined in the heuristic relation

$$
\mu(a/a_0)a = MGr^{-2} \tag{39}
$$

In [\(39\)](#page-12-0) $\mu(x) \approx 1$ when $x >> 1$, and $\mu(x) \approx x$ when $x << 1$. It must be stressed that [\(38\)](#page-12-1) or [\(39\)](#page-12-0) are not deduced from theory, but rather are an ad hoc fit to explain observations. Interestingly it must be mentioned that most of the implications of MOND do not depend strongly on the exact form of μ .

It can then be shown that the problem of galactic velocities is solved [\[29,](#page-18-10) [30,](#page-18-11) [31,](#page-18-12) [32,](#page-18-13) [33\]](#page-18-14).

8 Interrelationship

It is interesting to note that there is interesting relationship between the varying G approach, which has a theoretical base and the purely phenomenological MOND approach. Let us write

$$
\beta \frac{GM}{r} = \frac{r^2}{t_0^2} \text{ or } \beta = \frac{r^3}{GMt_0^2}
$$

Whence

$$
\alpha_0 = v^2/r = \frac{GM}{r^2}\,\alpha = \frac{r}{t_0^2}
$$

So that

$$
\frac{\alpha}{\alpha_0} = \frac{r^3}{GMt_0^2} = \beta
$$

At this stage we can see a similarity with MOND. For if $\beta \ll 1$ we are with the usual Newtonian dynamics and if $\beta > 1$ then we get back to the varying G case exactly as with MOND.

Furthermore, as can be seen from (6) , when the dimensionality n gets smaller than 3, effectively G starts falling off as in the time varying case seen in sections 5 and 6.

9 Discussion

It is interesting to note that the varying G approach leads to several observations that have been carried out including the precession of the perihelion of the planets, in particular Mercury, the Pioneer anomaly, the shortening of the orbital periods of binary pulsars and so on [\[23,](#page-18-4) [34,](#page-18-15) [7,](#page-17-6) [8\]](#page-17-7).

A further interesting observation is the fact noted by Milgrom that there is a curious coincidence in MOND viz.,

$$
a_0 \sim H(\sim 10^{-7} \, \text{cm} \, \text{sec}^{-2}) \tag{40}
$$

where H is the Hubble constant. In fact this follows from the varying G theory. For, we have in this case from (22) , (23) , (24) and (37) ,

$$
a_0 \sim r/t_0^2
$$

Feeding the values of r, the radius of the universe $= ct_0$ and the fact that $H \sim \frac{1}{t_0}$ $\frac{1}{t_0}$, we get [\(40\)](#page-13-0), which now shows up no longer as a coincidence.

Finally, it may be noted that the Boomerang results are in tune with MOND rather than the Dark Matter scenario, the WMAP model notwithstanding [\[35\]](#page-18-16).

It is also interesting to note that for large r , (37) gives the Hubble law, here deduced from the G variation.

10 Higher Dimensionality

The fact that asymptotically the universe appears to be two dimensional is interesting from a totally different point of view. This is that a collection of ultra relativistic masses would appear as two dimensional as discussed in detail in ref.[\[36\]](#page-18-17). This is what may be called an inside view of the universe. We now come to what may be called the outside view. The starting point is the observation that the universe shows up as a black hole. This is because we have,

$$
R \sim \frac{GM}{c^2} \tag{41}
$$

Equation [\(41\)](#page-14-0) is not phenomenological– indeed it can be deduced from theory as discussed in detail (Cf.ref.[\[8\]](#page-17-7)). The equation describes a Schwarzchild black hole. In [\(41\)](#page-14-0), dark matter if any is not included. If however dark matter were included in the mass then the case for a black hole becomes even stronger, as the left scales becomes smaller than the right side.

We now argue that this three dimensional black hole, which describes our universe is imbedded in a four dimensional manifold as in Wheeler's Superspace or in the author's multiple universe model [\[37\]](#page-18-18). We now extend the Lorentz transformation to a four dimensional space. In this case the angular momentum

$$
\int \vec{r} \times \vec{p} dV
$$

is replaced by

$$
\vec{\omega}=\int\vec{r}\Lambda\vec{p}dV
$$

where Λ represents the usual anti-symmetrical product. Whence we have

$$
\vec{r} \cdot \vec{\omega} = 0 \tag{42}
$$

The significance of [\(42\)](#page-14-1) is that it represents a three dimensional hyper surface in four dimensional space, remembering that \vec{r} has four space components. This apart the two dimensional surface in the ultra relativistic case referred to above follows precisely from [\(42\)](#page-14-1), but this time in three dimensional space. To sum up what this shows is that indeed the three dimensional universe can be considered to be a black hole imbedded in four dimensional space. There is a totally different route to the above consideration. As shown by the author and Popova [\[2\]](#page-17-1), (Cf. also sections 3 and 4), we have

$$
\dot{\epsilon} + n \frac{\dot{a}}{a} \epsilon = 0 \tag{43}
$$

where a is the size of the universe and ϵ the energy, and n the number of dimensions. An integration gives

$$
\epsilon \sim Mc^2 a^{-n} \tag{44}
$$

From here it follows that

$$
R \equiv a = \sqrt{\frac{(n-1)c^2}{K^{(n)}\epsilon}}\tag{45}
$$

where

$$
K^{(n)} = \frac{n-1}{n-2} \cdot \frac{2\pi^{n/2}}{\Gamma(n/2)} \cdot \frac{G^{(n)}}{c^2}
$$
 (46)

where $G^{(n)}$ has been defined earlier. Substitution of [\(46\)](#page-15-0) in [\(45\)](#page-15-1) for the case $n = 4$ gives us back [\(41\)](#page-14-0) as required for a black hole. The same conclusion can be drawn alternatively from similar reasoning [\[38\]](#page-19-0) in which case we get

$$
a^{(n)}_{max}=[0(1)\cdot \frac{G^nM}{c^2}]^{\frac{1}{n-2}}
$$

which gives back

$$
R(\sim 10^{28} cm) \sim \frac{GM}{c^2}
$$

for the case $n = 4$.

Interestingly if the universe were treated as a black hole, we could associate with it a spin given by

$$
\bar{h} = MRc \sim 10^{93}
$$

As pointed out elsewhere [\[7\]](#page-17-6) this value seems to be vindicated by COBE observations and of the order of the spin in Godel's solution of Einstein's equations. Equally interesting is the fact that with the scaled up value of the spin h , the universe could be considered to be a wave packet with the "Compton wavelength"

$$
R=\frac{\bar{h}}{Mc}
$$

To put it another way (Cf.[\[8\]](#page-17-7)) the universe could be a wave packet of a scaled up Schrodinger equation.

Inany case, we had argued that dimensionality increases with scale $([8])$ $([8])$ $([8])$ – from one at the Planck scale to two at the Compton scale, through three at our scale and now four and beyond at scales large than R above (Cf. also [\[37\]](#page-18-18)).

We conclude with the following remarks, in the context of some of the above considerations. As discussed in detail elsewhere (Cf. for example [\[8\]](#page-17-7)) the Virial Theorem gives in the astrophysical context

$$
\frac{GM}{R} \sim v^2
$$

where v is the dispersion in velocity, M is the mass and R the extent of a relatively isolated gravitational system. On the other hand as we saw above we have

$$
\frac{R^2}{M^2 R^2} \sim v^2
$$

Whence we have

$$
G \sim \frac{\bar{h}^2}{M^3 R}
$$

which gives the correct value of the gravitational constant, if we feed in the mass of the universe and its radius and also the above value of the scaled Planck constant h . What we have done is, we have obtained the expression for v from two totally different considerations. The first was, based on the gravitation of the matter in this mass collection. The second was based on non dynamical considerations, namely a generalized Schrodinger equation and Gaussian wave packets. So the latter considerations show up gravitation as being non fundamental and being more of an effective force. Indeed the non fundamental nature of gravitation has been commented upon in detail in the above reference.

The other aspect is that the Hubble law is a consequence of the spread of the above Gaussian wave packet of the scaled up Schrodinger equation. Indeed all this including the varying G was related to a dark energy driven universe in the author's work.

The interesting point is that the above considerations apply at other scales also as discussed in the reference. For example at the scale of galaxies we have a scaled up Planck constant $\sim 10^{74}$. If we now carry over the Gauss packet considerations to this scale, the decay of the Gauss packet which is the counterpart of the Hubble expansion of the universe now comes up as the gradual increase in size of the galaxies. (Of course we can also deduce the correct value of the universal constant of gravitation in this case too by similar considerations).

References

- [1] Narlikar, J.V., "Introduction to Cosmology", Cambridge University Press, Cambridge, 1993, p.57.
- [2] Sidharth, B.G., and A.D. Popova, Nonlinear World, 1997, 4.
- [3] Popova, A.D., Astron. and Astroph. Trans. 5 (1994), 31.
- [4] Popova, A.D., Astron. and Astroph. Trans. 6 (1995), 165.
- [5] Ehrenfest, P., Proc. Amsterdam Acad. 20 (1917), 200; Ann. Phys. 61 (1920), 440.
- [6] Barrow, J.D., and Parsons, P., Phys.Rev.D., Vol.55, No.4, 15 February 1997, p.1906ff.
- [7] Sidharth, B.G., "Chaotic Universe: From the Planck to the Hubble Scale", Nova Science Publishers, Inc., New York, 2001.
- [8] Sidharth, B.G., "The Universe of Fluctuations", Springer, Dordrecht, 2005.
- [9] Sidharth, B.G., Chaos, Solitons and Fractals, 12, 2001, 1369-1370.
- [10] Burstein, D., and Rubin, V.S., Astroph. J., 297 (1985), 423.
- [11] Kormendy, J., and Knapp, G. (eds.), "Dark Matter in the Universe", Reidel, Dordrecht, 1987.
- [12] Davies, M., et al., Astroph. J. 238 (1980), L113.
- [13] Turner, S., Astroph. J. 208 (1976), 304.
- [14] Sidharth, B.G., Int.J.Mod.Phys.A, 13 (15), 1998, p.2599ff.
- [15] Sidharth, B.G., Int.J.Th.Phys., 37 (4), 1998, p.1307ff.
- [16] Goldstein, H., "Classical Mechanics", Addison-Wesley, Reading, Mass., 1966, pp.76ff.
- [17] Kuhne, R.W., xxx.lanl.gov/gr-qc/9809075.
- [18] Berger, A.L., Astronomy and Astrophysics, 51, 1976, p.127-135.
- [19] Anderson, J.L. et al., xxx.lanl.gov/gr-qc/9808081.
- [20] Denman, H.H., Am.J.Phys. 51(1), 1983, 71.
- [21] Silverman, M.P., Am.J.Phys. 48, 1980, 72.
- [22] Brill, D.R. and Goel, D., Am.J.Phys. 67(4), 1999, 317.
- [23] Sidharth, B.G., Nuovo Cimento, 115B (12) (2), 2000, pp.151ff.
- [24] Bergman, P.G., "Introduction to the Theory of Relativity", Prentice-Hall (New Delhi), 1969, p248ff.
- [25] Lass, H., "Vector and Tensor Analysis", McGraw-Hill Book Co., Tokyo, 1950, p295 ff.
- [26] Sivaram, C., Sivaram and V. de Sabbata, Foundations of Physics Letters, 6 (6), 1993.
- [27] Sidharth, B.G., Chaos, Solitons and Fractals, 12, 2001, 1101-1104.
- [28] Milgrom, M., "MOND A Pedagogical Review" Presented at the XXV International School of Theoretical Physics "Particles and Astrophysics-Standard Models and Beyond", Ustron, Poland, September 2001.
- [29] Milgrom, M., APJ 270 (1983), 371.
- [30] Milgrom, M., APJ 302 (1986), 617.
- [31] Milgrom, M., Comm. Astrophys. 13:4 (1989), 215.
- [32] Milgrom, M., Ann.Phys. 229 (1994), 384.
- [33] Milgrom, M., Phys.Rev. E 56 (1997), 1148.
- [34] Sidharth, B.G., Found.Phys.Lett., 19 (6), 2006, 611-617.
- [35] McGauge, S.S., APJ Lett. 523 (1999), L99.
- [36] Moller, C., "The Theory of Relativity", Clarendon Press, Oxford, 1952, pp.171ff.
- [37] Sidharth, B.G., "Multiply Connected Universe", to appear in New Advances in Physics.

[38] Popova, A.D., Differential Equations and Dynamical Systems, Vol.6, 165-194.