

Reciprocal Symmetric Boltzmann Function and Unified Boson-Fermion Statistics

Mushfiq Ahmad

Department of Physics, Rajshahi University, Rajshahi, Bangladesh

E-mail: mushfiqahmad@ru.ac.bd

Muhammad O. G. Talukder

Department of ETE and CSE, University of Development Alternative, Dhaka,
Bangladesh

Abstract

The differential equation for Boltzmann's function is replaced by the corresponding discrete finite difference equation. The difference equation is, then, symmetrized so that the equation remains invariant when step d is replaced by $-d$. The solutions of this equation come in Boson-Fermion pairs. Reciprocal symmetric Boltzmann's function, thus, unifies both Bosonic and Fermionic distributions.

1. Introduction

We have seen that there exists a relation between reciprocal symmetry and spin¹. It is, then, reasonable to expect that there exists a relation between reciprocal symmetry and quantum statistics. We shall study this below.

If the differential equation for the exponential function is replaced by the corresponding discrete symmetric difference equation, solutions come in pairs². One of them corresponds to classical exponential function and the other is an oscillating function and has no classical analogue. We have seen that the other solution corresponds to Fermi-Dirac statistics³. We want to study in this paper how to exploit this pair solution to obtain a Boson-Fermion unified statistical relation.

2. Boltzmann's Finite Difference Equation

Classical Boltzmann's function g satisfies the differential equation⁴

$$\frac{dg}{dH} = -wg = -w \cdot \exp(-w.H) \quad (2.1)$$

The number N^H of particles in energy level H is given by Boltzmann's probability distribution relation⁵

$$N^H = N^0 \exp(-H.w) \quad (2.2)$$

Where N^0 is the number of particles when $H=0$

3. Planck's Radiation Formula

The formula for the distribution of light in a black body is given by⁶

$$I(\omega)d\omega = \langle H \rangle \frac{\omega^2 d\omega}{\pi^2 c^2} \quad (3.1)$$

(2.2) together with Planck's hypothesis have been invoked to find $\langle H \rangle$ ⁷.

$$\langle H \rangle = \frac{\eta\omega}{\exp(\eta\omega/kT) - 1} \quad (3.2)$$

(2.2) will lead to (3.2) if

$$H = n\eta\omega \quad (3.3a)$$

$$w = 1/kT \quad (3.3b)$$

4. Boltzmann's Symmetric Equation

To exploit its symmetry properties we replace differential equation (2.1) by the corresponding symmetric finite difference equation⁸

$$\frac{Dg_{\pm}^H}{D(H, \delta)} = -W \cdot g_{\pm}^H \quad (4.1)$$

where

$$\frac{Dg_{\pm}^H}{D(H, \delta)} = \frac{g_{\pm}^{H+\delta} - g_{\pm}^{H-\delta}}{2\delta} \quad (4.2)$$

By replacing (2.1) by (4.1) we have tacitly assumed $H = n\delta$ where n is an integer. The difference quotient (4.1) has the following symmetry under the change $\delta \rightarrow -\delta$

$$\frac{Dg_{\pm}^H}{D(H, -\delta)} = \frac{Dg_{\pm}^H}{D(H, \delta)} \quad (4.3)$$

We require that (4.1) should go over to (2.1) in the limit $\delta \rightarrow 0$

$$\frac{Dg_{\pm}^H}{D(H, \delta)} = -Wg_{\pm}^H \xrightarrow{\delta \rightarrow 0} \frac{dg}{dH} = -wg \quad (4.4)$$

With

$$g_{\pm}^H \xrightarrow{\delta \rightarrow 0} g \text{ and } W \xrightarrow{\delta \rightarrow 0} w \quad (4.4)$$

5. Boltzmann's Symmetric Function

(4.1) has solutions in pairs. One pair is

$$g_{\pm}^H = \exp\left(\frac{2\pi i}{\delta} s_{\pm} \mu w\right) H \quad (5.1)$$

Where s_{+} is an integer and s_{-} is a half-integer.

Both g_{+}^H and g_{-}^H satisfy (4.1) with

$$W = \frac{\sinh(w\delta)}{\delta} \quad (5.2)$$

g_{+}^H and g_{-}^H are related through reciprocity relation

$$g_{+}^H \cdot g_{-}^H = (-1)^{H/\delta} \quad (5.3)$$

6. Normalization

g_{+}^H of (3.1) corresponds to classical Boltzmann's function. It is the classical function if we set $s_{+} = 0$.

g_{-}^H diverges for positive H as $H \rightarrow \infty$. Therefore, normalization requirement⁹ imposes the condition that H must be negative.

7. Symmetric Number of Particles

If classical g is replaced by symmetrized g_+^H or g_-^H , the number of particles in energy level H becomes

$$N_{\pm}^H = N^0 g_{\pm}^H \quad (7.1)$$

Where N^0 is the number of particles when $H=0$. If we set $N^0 = 1$ and $H = n.\delta$ (See comment following (4.2))

$$N_{\pm}^H = g_{\pm}^H = g_{\pm}^{n.\delta} \quad (7.2)$$

When $n=1$

$$N_{\pm}^{\delta} = g_{\pm}^{\delta} = \exp\left(\frac{2\pi i}{\delta} s_{\pm} \pm w\right) \delta \quad (7.3)$$

8. Symmetric Radiation Formula

We shall call $\langle H_{\pm} \rangle$ the average energy corresponding to $\langle H \rangle$ when the classical number of particles N^H of (2.2) is replaced by the corresponding symmetric number N_{\pm}^H as given by (4.2). Following the procedure¹⁰ for finding (3.2), we find for $\langle H_{\pm} \rangle$

$$\langle H_{\pm} \rangle = \frac{\delta}{g_{\pm}^{\delta} - 1} \quad (8.1)$$

Using (5.1) and (3.3b), (8.1) becomes

$$\langle H_{\pm} \rangle = \frac{\delta}{\pm \exp(\pm \delta / kT) - 1} = \frac{\pm \delta}{\exp(\pm \delta / kT) \mu 1} \quad (8.2)$$

9. Planck's Hypothesis

Planck's hypothesis has two parts: (a) Energy is discrete $H = n.\delta$ and that (b) $\delta = \eta\omega$.

Part (a) has already been taken care of in sections 2 and 3. If we include part (b), (8.2) becomes

$$\langle H_{\pm} \rangle = \frac{\pm \eta \omega}{\exp(\pm \eta \omega / kT) \pm 1} \quad (9.1)$$

10. Unified Boson-Fermion Statistics

The upper sign in (9.1) gives Planck's relation

$$\langle H_{+} \rangle = \frac{\eta \omega}{\exp(\eta \omega / kT) - 1} \quad (10.1)$$

The lower sign in (9.1) gives the corresponding Fermi-Dirac relation¹¹

$$\langle H_{-} \rangle = \frac{-\eta \omega}{\exp(-\eta \omega / kT) + 1} = \frac{\eta \omega'}{\exp(\eta \omega' / kT) + 1} \quad (10.2)$$

Where $\omega' = -\omega$

11. Conclusion

Symmetric Boltzmann's function is enough to get both Bose-Einstein (10.1) and Fermi-Dirac (10.2) radiation relations from the corresponding classical relation¹² just as Planck's hypothesis is enough to obtain Planck's relation (3.2). To derive (10.2) we have not invoked Pauli Exclusion Principle.

¹Mushfiq Ahmad. Reciprocal Symmetry and Origin of Spin. [arXiv:math-ph/0702043v1](https://arxiv.org/abs/math-ph/0702043v1)

²Mushfiq Ahmad. Reciprocal Symmetry and Equivalence between Relativistic and Quantum Mechanical Concepts [arXiv:math-ph/0611024v1](https://arxiv.org/abs/math-ph/0611024v1)

³Mushfiq Ahmad. Reciprocal Symmetric and Origin of Quantum Statistics. [arXiv:physics/0703194v1](https://arxiv.org/abs/physics/0703194v1)

⁴Rudolf Kurth. Axiomatics of Classical Statistical Mechanics. Pergamon Press. 1960.

⁵<http://www.answers.com/topic/boltzmann-distribution>

⁶Feynman, Leighton, Sands. Feynman Lectures on Physics. Addison-Wesley Pub. Co.

⁷Feynman, Leighton, Sands. Feynman Lectures on Physics. Addison-Wesley Pub. Co

⁸Mushfiq Ahmad Reciprocal Symmetric and Origin of Quantum Statistics.

<http://www.arxiv.org/abs/physics/0703194>

⁹Rudolf Kurth. Axiomatics of Classical Statistical Mechanics. Pergamon Press. 1960.

¹⁰Feynman, Leighton, Sands. Feynman Lectures on Physics. Addison-Wesley Pub. Co

¹¹ http://en.wikipedia.org/wiki/Fermi-Dirac_statistics

¹²Feynman, Leighton, Sands. Feynman Lectures on Physics. Addison-Wesley Pub. Co