## The Orthopositronium Problem and $e-\mu-\tau$ -Universality

B.M.Levin\*

A.F.Ioffe Physical Technical Institute, 194021 St. Petersburg, Russia

August 10, 2021

## Abstract

The quantitative description of the orthopositronium anomalies ("isotope anomaly" in a gaseous neon for the "resonance conditions" and " $\lambda_T$ -anomaly" in non-resonance conditions) is possible on the basis of a hypothesis about restoration of spontaneously broken *complete relativity* (A.F.Andreev, 1982) of the limited space-time "volume" ("defect" of the space-time) in a final state of  $\beta^+$ -decay ( $\Delta J^{\pi} = 1^+$ ) of nucleus such as <sup>22</sup>Na, <sup>68</sup>Ga, etc. This addition of the Standard Model in a final state of the topological quantum transition (non-steady-state, a generalized "displacement current") supposes the description on the common basis timelike fundamental particles (locality) and spacelike fundamental structure (*non-locality without causal paradoxes*). One of achievements of expansion of Standard Model can be a substantiation of topological connection *e-µ-τ*-Universality with discrete structure (quantization) "defect" of the space-time.

Consideration by analogy with orthopositronium  $[o - Ps \equiv {}^{3}(e_{\beta}^{+} e^{-})_{1}]$  of the atoms  ${}^{3}(\mu^{+}\mu^{-})_{1}$  and  ${}^{3}(\tau^{+}\tau^{-})_{1}$  finds out basic difference of the top generations of leptons: impossibility of realization for  ${}^{3}(\mu^{+}\mu^{-})_{1}$ - and  ${}^{3}(\tau^{+}\tau^{-})_{1}$ -atoms, even in some Gedanken experiment, a conditions similar to the "resonance conditions" at supervision of the "isotopic anomaly" in samples of a gaseous neon of various isotope abundance [1,2]. If to treat it as spontaneous breaking of the horizontal symmetry of the leptons generations, then their participation in formation of the fundamental space-like structure within the framework of "additional  $G\hbar/c$ -physicis" would be natural consequence of restoration of the mentioned symmetry in a final state of  $\beta^{+}$ -decay for a time  $\tau_{\mu} \sim R_{\mu}/c \sim 10^{-6}$ s [3-5].

Let's present discrete, space-like and two-sign (±, on mass and all charges, including baryon charge) structure as a tachyonic medium [6], formed in a final state of the  $\beta^+$ -decay of nucleus  ${}^{22}Na$ , etc ( $\Delta J^{\pi} = 1^+$ ). The linear sizes of the fundamental space-like structure presented in [4] – "virtual fundamental length"  $\Delta \simeq 5.5 \cdot 10^{-2} \ cm$  and the size of the "long-range atom"  $2R_{\mu} \simeq 1.1 \cdot 10^5 \ cm$  –

we shall compare with a the characteristic length of the tachyonic instability [6]

$$L > C / \Gamma, \tag{1}$$

where C – the characteristic rate agreed in this specific case with velocity of light c, and  $\Gamma$  is an inverse time of development of instability  $(s^{-1})$ .

For revealing the physical content of the assumed comparison is important presented in [6] isomorphism of the equations of the tachyonic field

$$\left(\omega^2 + C^2 \cdot \nabla^2 + \Gamma^2\right) \cdot \Psi = 0 \qquad (\hbar = 1)$$

and of the Schrödinger equation for the stationary state of the "particle" with mass  $m \ (m = 1/2)$ 

$$\left[\nabla^2 + E - V(x)\right] \cdot \Psi = 0,$$

i.e. a conformity takes place

$$\omega^2/C^2 \iff E, \quad \Gamma^2/C^2 \iff -V(x),$$

if to consider value  $\Gamma$  depending from the spatial coordinate x.

It follows that tachyonic instability ( $\omega^2 < 0$ ,  $\omega = i\Gamma$ ,  $\Psi \sim (exp |\Omega| t)$  is appropriate to the bound state  $-\Gamma^2/C^2$  in an attraction field, and a negative energy of a compensating field (the "mirror Universe" [3-5]) can be presented as the binding energy  $E_b = |M_{Pl}| \cdot c^2$  of the vacuumlike substance component with mass  $M_{Pl} > 0$  of the whole fundamental space-like structure. We receive the increment increase of the tachionic field [6]

$$\Psi \sim exp(\Omega t), \quad \Omega = C \cdot (-E_b)^{1/2},$$

and  $\Omega$  in a considered context accepts value

$$\Omega = \frac{c \cdot [2M_{Pl}(-M_{Pl} \cdot c^2)]^{1/2}}{\hbar} \sim i t_{Pl}^{-1} \sim i \cdot 10^{43} \ s^{-1}.$$
 (2)

All "sites" of the crossing of "spaces" X (in the observable world) and X' (an inverse "space" – the "mirror Universe") "stick together" in "point" in the space 3+N'-dimensions:  $\{X\} = \{X'\} = 0$ . In the space of the observer it allocates the "central" site of "lattice" of the fundamental macroscopical space-like structure with which in a final state of the  $\beta^+$ -decay <sup>22</sup>Na  $(e^+, \nu)^{22*}Ne$  the daughter nucleus <sup>22\*</sup>Ne has been linked. The four-dimensional fundamental space-like structure of the size  $2R_{\mu}$  in the 3 + N'-dim space is captured by Planckian length  $l_{Pl}$ . Process of the tachyonic instability by the conserve of the angular momentum is limited. Specifically, a "self-unwinding" of the fundamental space-like structure in the  $\beta^+$ -decay final state by a difference of the angular moments of the final and initial states (<sup>22</sup>Na, <sup>68</sup>Ga, etc) is limited

$$\Delta J = \hbar > I_{\mu} \cdot c/R_{\mu} , \qquad (3)$$

as velocity of the fundamental space-like structure on its "surface" cannot exceed the velocity of light. In (3)  $I_{\mu} \sim m_{\mu} \cdot R_{\mu}^2$  is the moment inertia of a

spherical huming-top,  $m_{\mu} \sim \hbar/R_{\mu}c$  is its mass. Hence, practically instant "selfunwinding" of the fundamental space-like structure ( $|\Omega| \sim 10^{43} s^{-1}$ ) periodically undergoes a failure and again renews with a casual directions of the angular moment. This result give proof to the postulate about the casual rotation of the "mirror Universe" with a velocity  $V \sim -c$  in respect to the ground-based laboratory (to an *observer*) [7].

The observer perceives additional discrete structure of the "mirror Universe" as discrete scalar *C-field* [8] with a negative sign on all physical "charges" (including mass of "holes" – "proton" and "electron" in the structure of "lattices") [4]. *C*-field compensates of the limited "volume" of the space-time (~ 1 km<sup>3</sup>, during ~  $2 \cdot 10^{-6} s$ ) – the vacuumlike state of matter (VSM) [9] ( the "long-range atom" with Planckian mass also perceived by the observer as microstructure of the VSM and 1-dim discrete space (N' = 1).

As a whole the final state of the  $\beta^+$ -decay ( $\Delta J^{\pi} = 1^+$ ) represents 5-dim (3+1', t) with discrete structure of the spatial dimensions in the limited "volume" of the space-time.

Inequalities (comp. with (1))

$$\Delta > \frac{c}{\Gamma_{\tau}} \ cm \quad \text{and} \quad 2R_{\mu} > \frac{c}{\Gamma_{\mu}} \ cm$$

$$\tag{4}$$

where  $\Gamma_{\tau}$  (~ 3.3 · 10<sup>12</sup> s<sup>-1</sup>) and  $\Gamma_{\mu}$  (~ 4.5 · 10<sup>5</sup> s<sup>-1</sup>), accordingly, are the decay width of the  $\tau$ - and  $\mu$ -leptons, show that in structure of the value  $\Gamma(x)$  is available the contribution of a topological component of the increase of the tachyonic field – width of decay of leptons  $\Gamma_{\tau}$ ,  $\Gamma_{\mu}$  ( $\Gamma_{\tau / \mu}$ ), defining the rate development of an instability.

The topological component limits the rate of increase of the instability as a time of the consecutive transitions is defined by sum of characteristic times

$$\frac{1}{\Gamma\left(x \sim \Delta \ / \ x \sim R_{\mu}\right)} = \frac{1}{\Gamma_{\tau \ / \ \mu}} + \frac{1}{\mid \Omega \mid}$$

and

$$\Gamma\left(x \sim \Delta \mid x \sim R_{\mu}\right) = \frac{\Gamma_{\tau \mid \mu} \cdot \mid \Omega \mid}{\Gamma_{\tau \mid \mu} + \mid \Omega \mid} \cong \Gamma_{\tau \mid \mu} \quad (\mid \Omega \mid >> \Gamma_{\tau/\mu}).$$
(5)

The width of the leptons decay is defined by the the weak interaction constant. Thus, unification of the all interactions on the basis of antipodic symmetry [10] in the "additional  $G\hbar/c$ -physics" (superantipodic symmetry [3]) on the basis of their own coupling constants take place, instead of a uniform constant as it is supposed in the standard "Theory of Everything" at ultrahigh energy. It means, that the impossibility to fix a coupling constant for the Goldstone field in the theory with spontaneously broken complete relativity [11] is not lack of the theory.

In the offered concept "additional  $G\hbar/c$ -physics" the top generations of the leptons adjust the bifurcation attitude mutually supplementing aspects of strong interaction: one – with participation of color (quantum cromodynamics/aromadynamics) [12], and another – a non-stationary long-range for a baryon (and lepton) charge [3-5].

Let's note, that the estimation of the mass presented here fundamental spacelike structures  $m_{\mu} \sim \hbar/R_{\mu}c \sim 3 \cdot 10^{-10} eV$  is kept at extremely wide limits admissible by Standard Model for mass invisible acsion, is closer to its lower limit  $\sim 10^{-12} eV$ . If the "long-range atom" at the moment of a birth in final state of  $\beta^+$ -decay or as result of the its subsequent diffusion is brought into gravitational field with the critical value of acceleration of free falling

$$\gamma_{cr} = \frac{\hbar \cdot c}{m_p \cdot R_{\mu}^2} \geqslant 0.01 \ cm/s^2,$$

there is its "splitting" on plus/minus Planck masses (non-steady-state, the generalized displacement current), i.e. in one-stage (from "nothing"!) "elementary" macroscopical "domain" of a dark matter with mass  $2|M_{Pl}|$  is born [13].

\*E-mail: bormikhlev@mail.ioffe.ru, bormikhlev@mail.ru

## References

- B.M.Levin, L.M.Kochenda, A.A.Markov, and V.P.Shantarovich. Time spectra of annihilation of positrons (<sup>22</sup>Na) in gaseous neon of various isotopic compositions. Sov. J. Nucl. Phys., 1987, v. 45(6), p. 1119.
- B.M.Levin. Orthopositronium: a program for critical experiments. Sov. J. Nucl. Phys., 1990, v. 52(2), p.342.
- B.M.Levin. Orthopositronium: 'Annihilation of positron in gaseous neon'. http://arXiv.org/abs/quant-ph/0303166
- B.A.Kotov, B.M.Levin, V.I.Sokolov. Orthopositronium: "on the possible relation of gravity to electricity". http://arXiv.org/abs/quant-ph/0602063
- [5] B.M.Levin, V.I.Sokolov. On an additional realization of supersymmetry in orthopositronium lifetime anomalies. http://arXiv.org/abs/quant-ph/0704117
- [6] A.Yu.Andreev, D.A.Kirzhnits. Tachyons and the instability of physical systems. Usp. Fiz. Nauk, 1996, v. 166(10), p. 1135.
- [7] B.M.Levin. On the Kinematics of One-Photon Annihilation of Orthopositronium. Phys. At. Nucl., 1995, v. 58(2), p. 332.
- [8] F.Hoyle, and J.V.Narlikar. C-field as direct field of particles. Proc. Roy. Soc., 1964, v. A282, p.178.
- [9] E.B.Gliner. Algebraic properties of energy-momentum tensor and vacuumlike states of matter. Sov. Phys. JETP – USSR, 1966, v. 22, p.378.

- [10] A.D.Linde. The multiplication of the Universe and problem of cosmological constant. Phys. Lett., 1988, v. B200(3), p. 272.
- [11] A.F.Andreev. Spontaneously broken complete relativity. JETP Lett., 1982, v. 36(3), p.100.
- [12] A.S.Sakharov, M.Yu.Khlopov. Horizontal unification as the phenomenology of the theory of everything. Phys. At. Nucl., 1994, v. 57(4), p.651.
- [13] B.M.Levin, V.I.Sokolov. Whether can the decision of the orthopositronium problem to stimulate studying the problem of dark matter in the Universe? http://arXiv.org/abs/astro-ph/0610063