

Boundary behaviour of Loewner Chains

Alexander Kuznetsov

November 13, 2018

Abstract

In paper found conditions that guarantee that solution of Loewner-Kufarev equation maps unit disc onto domain with quasiconformal rectifiable boundary, or it has continuation on closed unit disc, or it's inverse function has continuation on closure of domain.

1 Introduction

In recent time there was a significant interest to evolution processes of domains in complex plane. This phenomenas are well described by equations of Loewner-Kufarev type.

Let $D(t)$ be a simple connected domain in the complex plane, changing in time. Without loss of generality we can assume that $D(t)$ contains origin. Let a function $f(z, t)$ ($f'(0, t) > 0, f(0, t) = 0$) for each fixed t map unit disc $\mathbb{D} = \{z : |z| < 1\}$ onto domain $D(t)$. If $D(t_1) \subset D(t_2), t_1 > t_2$, then function $f(z, t)$ satisfies the Loewner-Kufarev equation

$$\frac{\partial f}{\partial t} = -\frac{\partial f}{\partial t} z p(z, t), z \in \mathbb{D}, \quad (1)$$

where $p(z, t) : \mathbb{D} \times [0, +\infty) \rightarrow \mathbb{C}$ is such that for each fixed $t \in [0, +\infty)$, $\Re p(z, t) > 0, p(0, t) > 0$ and for each fixed $z \in \mathbb{D}$, $p(z, t)$ is mesurable on interval $[0, +\infty)$. Making change of variable (1) we can assume, that for each fixed t function $p(z, t)$ belongs to class \mathcal{P} , consisting of functions $p(0) = 1, \Re p(z) > 0$.

Such type equations arise in description of Hele-Shaw flow [6, 15, 16, 22]. In paper [5] L. Carleson and N. Makarov used this equations for the study of DLA processes. Parametric method of representations of univalent functions [1, 17] is based on this approach. The importance of this method and approach is shown by L. de Branges proof of famous Bieberbach hypothesis [4], that was the central problem of functions theory for long time.

If $p(z, t) = \frac{e^{iu(t)} + z}{e^{iu(t)} - z}$, where $u(t)$ is real function, then equation will be Loewner equation [11]. Recently there was a great activity in the study of this equation [9, 10, 19, 20], there $u(t)$ describes one-dimensional Brownian motion. In this case equation (1) becomes Stochastic Loewner Equation (SLE). The study

of geometrical characteristics of solutions of SLE, by various parameters of Brownian motion is of a great interest.

In paper [13] it was proved, that solution of Loewner equation maps unit disc on quasilit-disc if and only if function $u(t)$ is Hoelder continuous with exponent $1/2$. Also in it was shown if equation

$$\inf_{\varepsilon > 0} \sup_{|t-s| < \varepsilon} \frac{|u(t) - u(s)|}{\sqrt{|t-s|}} < c,$$

holds for some small $c > 0$, (in paper of Lind [12] was proved, that $c = 4$), when solution of Loewner equation maps unit-disc on quasilit-disc.

Let \mathcal{P}_s be some subclass of class \mathcal{P} and \mathcal{PT}_s be a class of functions $p(z, t) : \mathbb{D} \times [0, +\infty) \rightarrow \mathbb{C}$ such that for each fixed $t \in [0, +\infty)$, $p(z, t) \in \mathcal{P}_s$ and, for each fixed $z \in \mathbb{D}$, $p(z, t)$ is mesurable on interval $[0, +\infty)$. The main goal of this paper is to find conditions of class \mathcal{P}_s , under that the solution of equation (1) with $p(z, t) \in \mathcal{PT}_s$ will have certain boundary behaviour.

Similar question was studied by J. Becker. In papers [2, 3] it was proved, if

$$\left| \frac{p(z, t) - 1}{p(z, t) + 1} \right| \leq k < 1,$$

then solution of equation (1) has quasiconformal extension. Recently A. Vasil'ev [21] gave a description of $p(z, t)$ under conditions that, solution $f(z, t)$ of (1) has quasiconformal extension for each t .

In the papers [7, 8] were discussed problems of a similar nature, but from the point of view of covering and distortion theorems for various classes of univalent functions, corresponding to various classes \mathcal{P}_s .

2 Behaviour of inverse function

Let us show that the study of boundary behaviour of solution of equation (1) for various classes \mathcal{PT}_s , can be reduced to the study of Loewner-Kufarev ODE.

Let us consider a function $w^*(z, t, s) = f^{-1}(f(z, t_2), t_1)$, that satisfies the following equation

$$\frac{dw^*}{dt_1} = w^* p(w^*, t_1), \quad w(z, s, s) = z, \quad z \in \mathbb{D}.$$

Let t_1 change from 0 to t , and $t_2 = t$. When $w(z, 0, t) = f(z, t)$. Assuming that $\tau = 1 - t_1$, we have, that $w(z, t) = f(z, t)$, where $w(z, \tau)$ is solution of equation

$$\frac{dw}{d\tau} = -w p(w, \tau), \quad w(z, 0) = z, \quad z \in \mathbb{D}. \quad (2)$$

We have the necessary result.

Theorem 1 *If there are constants $0 < \alpha < 1$ and $C_1, C_2 > 0$ such that*

$$\frac{|p(z, t)|}{\Re p(z, t)} \leq \frac{C_1}{(1-r)^\alpha}, \quad r = |z|, \quad (3)$$

$$\Re p(z, t) \geq C_2(1-r)^\alpha, \quad r = |z|. \quad (4)$$

Then inverse function to solution $w(z, t)$ of equation (2) has a continuation on boundary of domain $w(\mathbb{D}, t)$.

Let us consider a integral curve $s_{(z_0, t_0)}$, passing through point (z_0, t_0) , and find estimate on it's part, laying in cylinder $C_r = \{z : 1/2 < r < |z| < 1\}$.

Separating the real and the imaginary part in equation (2), we have

$$\begin{cases} \frac{d\rho}{dt} = -\rho \Re p(w, t), \\ \frac{d \arg w}{dt} = -\Im p(w, t), \end{cases} \quad (5)$$

where $\rho = |w|$.

From system (5) follows that function $\rho(t) = |w(z, t)|$ is monotonic on t . Thus, we can make a change of variable in equation (2) from t on ρ . Making it, we have that length of part of integral curve $s_{(z_0, t_0)}$, that laying in cylinder C_r , equals to

$$l_s(r) = \int_r^1 \sqrt{\left(\frac{dw}{d\rho}\right)^2 + \left(\frac{dt}{d\rho}\right)^2} d\rho.$$

Using inequalities (3), (4) and $|w(z, t)| \leq z$ we obtain

$$l_s(r) \leq \int_r^1 \sqrt{\frac{C_1^2}{(1-\rho)^{2\alpha}} + \frac{1}{C_2^2 \rho^2 (1-\rho)^{2\alpha}}} d\rho \leq C_3(1-r)^{1-\alpha}.$$

Where C_3 is some constant, that depends only on C_1, C_2 . Thus,

$$\lim_{r \rightarrow 1} l_s(r) = 0. \quad (6)$$

Let $w(z, t)$ be a solution of equation (2), passing through (z_0, t_0) and τ be minimal number, for that $w(z, t)$ exits on interval (τ, t_0) . Using monotony of $|w(z, t)|$ on t and (6), we have that there is $\lim_{t \rightarrow \tau} w(z, t) = \gamma(z_0, t_0)$ and $|\gamma(z_0, t_0)| = 1$

Let define a function $s_{t_1}(z_0, t_0), |z| \leq 1, 0 \leq t_1 \leq t_0$ in a following way. If $w(z, t)$ solution of equation (2) with initial condition (z_0, t_0) exists for t_1 , we assume that $s_{t_1}(z_0, t_0) = w(z, t_1)$. Let Q_0 be a set of pairs (z, t_0) for that this is true. Let, $s_{t_1}(z_0, t_0) = \gamma(z_0, t_0)$, if $z \in Q_1 = \{(z_0, t_0) : |z_0| < 1, (z_0, t_0) \notin Q_0\}$ and if $z \in Q_2 = \{(z_0, t_0) : |z_0| = 1\}$, then $s_{t_1}(z_0, t_0) = (z_0, t_0)$.

Let us show that function $s_{t_1}(z_0, t_0)$ is a continuous one.

By theorem of continuous depending of solution form initial conditions for differential equations in integral form (see for e.g. [18] p.182), set Q_0 is open and function $s_{t_1}(z_0, t_0)$ is continuous on it.

Let $(z_0, t_0) \in Q_2$ and point (z', t') satisfy the following equation $|z_0 - z'| + |t_0 - t'| < \varepsilon$. The length of part of curve $w(z, t)$, passing through point (z', t') and laying in cylinder C_ε , is not greater then $C_3\varepsilon^\alpha$. So we have following inequality

$$|s_{t_1}(z_0, t_0) - s_{t_1}(z', t')| \leq C_3\varepsilon^{1-\alpha} + \varepsilon.$$

This inequality means that function $s_{t_1}(z, t)$ is continuous on set Q_2 .

Let $(z_0, t_0) \in Q_1$, t^* , such that $1 - |w(z, t^*)| = 2\varepsilon$ (for enough small ε such point exists) and O be neighbourhood of point $(t^*, w(z_0, t^*))$, that is defined by the following inequality

$$|t^* - t'| + |w(z, t^*) - z'| < \varepsilon. \quad (7)$$

Using the theorem of continuous depending of solution form initial conditions for differential equations we have, that there is $\delta > 0$ so that, form inequality $|t_0 - t'_0| + |z_0 - z'_0| < \delta$ follows that $w^*(z', t') \in O$, where $w^*(z, t)$ integral curve passing through point (z'_0, t'_0) . Taking into account, that the length of parts of integral curves $w^*(z, t)$ and $w(z, t)$, laying in cylinder $C_{2\varepsilon}$, is not greater then $2C_3(2\varepsilon)^{1-\alpha}$, we have

$$|s_{t_1}(z_0, t_0) - s_{t_1}(z', t')| < 2C_3(2\varepsilon)^{1-\alpha} + \varepsilon.$$

This inequality means that function $s_{t_1}(z, t)$ is continuous on set Q_1 . Noting that $f^{-1}(z, t) = s_0(z, t)$ we have the necessary result.

3 Boundary behaviour

Theorem 2 *Following statmets are true:*

1. *If inequality*

$$\frac{\Re zp_z(z, t)}{|z|\Re p(z, t)} \leq \frac{k}{1-r} + O\left(\frac{1}{(1-r)^\alpha}\right), \quad k < 1, r = |z| \quad (8)$$

holds, then function $w(z, t)$ is Hoelder continuous with exponent $1 - k$ in closed unit disc.

2. *If inequality*

$$\frac{\Re zp_z(z, t)}{|w|\Re p(z, t)} = O\left(\frac{1}{(1-r)^\alpha}\right), \quad r = |z| \quad (9)$$

holds, then function $w(z, t)$ maps unit disc on domain with rectifiable and quasiconformal boundary.

In the proof we are using ideas similar to ideas in papers [7, 8]. Differentiating equation (2) on z , we obtain

$$\frac{dw_z}{dt} = -w_z p(w, z) - ww_z p_z(w, t), \quad |z| < 1, w_z(z, 0) = 1.$$

Making change of variable from t to ρ , we have

$$\frac{dw_z}{d\rho} = -w_z \frac{p(w, t) + wp_z(w, t)}{\rho \Re p(w, t)}.$$

Integrating of this equations gives us

$$w_z(z, t) = e^{\int_{r_1}^{|z|} \frac{p+wp_z}{\rho \Re p} d\rho}.$$

From that we have

$$|w_z(z, t)| \leq e^{\int_{r_1}^{|z|} \frac{1}{\rho} + \frac{\Re w p_z}{\rho \Re p} d\rho}. \quad (10)$$

Taking into account what $r_1 = O(t)$ and inequality (8), we have

$$C_1(1 - |z|)^k \leq |w_z(z, t)| \leq C_2 \frac{1}{(1 - |z|)^k}.$$

From this follows that, function $w(z, t)$ is Hoelder continuous with exponent $1 - k$ in closed unit disc (see for e.g. [17] p.300). First part of the theorem is proved.

Taking into account (9) and (10), we have that, there is t_0 such that for all $t < t_0$ we have $|w_z(z, t) - 1| \leq 1/4$. Let us consider three points on circle z_1, z_2, z_3 , so that point z_2 lays between points z_1, z_3 . From equality

$$|w(z_1, t) - w(z_2, t)| = \left| \int_{z_1}^{z_2} w_z(z, t) dz \right|,$$

we obtain

$$\frac{3}{4}|z_1 - z_2| \leq |w(z_1, t) - w(z_2, t)| \leq \frac{5}{4}|z_1 - z_2|.$$

So we have

$$\frac{|w(z_1, t) - w(z_2, t)|}{|w(z_1, t) - w(z_3, t)|} \leq \frac{5}{3} \frac{|z_1 - z_2|}{|z_1 - z_3|}.$$

Therefore, boundary of domain $w(\mathbb{D}, t)$ is quasiconformal curve and function $w(z, t)$ have quasiconformal extension on hole complex plane, for $t < t_0$.

Let $f_i(z) = w_i(z, t^*)$, where $t^* = t/n, i = 1, 2, \dots, n$ and w_i is solution of equation

$$\frac{dw_i}{dt} = w_i p(w_i, t - it^*), w_i(z, 0) = z.$$

When

$$w(z, t) = f_1 \circ f_2 \circ \dots \circ f_n.$$

Thus choosing n so that $t^* = t/n < t_0$ we have that all $f_i, i = 1, \dots, n$ have quasiconformal extension. Therefore $w(z, t)$ as well. Also we have $|w_z(z, t)| \leq (3/2)^n$ so $w(z, t)$ maps unit disc on domain with rectifiable boundary. The second part of theorem is proved.

4 Examples

In this section we consider various subclasses of \mathcal{P} , that have a simple geometrical description and satisfy conditions of theorems 1 and 2. Let Ω be convex subdomain of the right half-plane and \mathcal{P}_Ω be class of functions $p : \mathbb{D} \rightarrow \Omega$. Through p_Ω we define a map from \mathbb{D} onto Ω . Any function $p \in \mathcal{P}_\Omega$ can be represented as $p_\Omega(\varphi(z))$, where $\varphi : \mathbb{D} \rightarrow \mathbb{D}$ and $\varphi(0) = 0$. From this and Schwarz lemma follows that if function p_Ω satisfies conditions of theorem 1, then all functions from class $p \in \mathcal{P}_\Omega$ also satisfy the conditions.

Let us consider a characteristic of function p :

$$\mathcal{H}(p) = \sup_{|z| < 1} \frac{(1 - |z|^2)|p'(z)|}{\Re p(z)},$$

and show that $\mathcal{H}(p(\varphi(z))) \leq \mathcal{H}(p(z))$ for all $\varphi : \mathbb{D} \rightarrow \mathbb{D}$.

$$\begin{aligned} \mathcal{H}(p(\varphi(z))) &= \sup_{|z| < 1} \frac{(1 - |z|^2)|\varphi'(z)p'(\varphi(z))|}{\Re p(\varphi(z))} = \\ &= \frac{(1 - |z|^2)(1 - |\varphi(z)|^2)|\varphi'(z)||p'(\varphi(z))|}{(1 - |\varphi(z)|^2)\Re p(\varphi(z))}. \end{aligned}$$

Taking into account that

$$\frac{(1 - |z|^2)|\varphi'(z)|}{(1 - |\varphi(z)|^2)} \leq 1,$$

we obtain the necessary result.

We have

$$\frac{\Re zp'(z)}{|z|\Re p(z)} \leq \left| \frac{p'(z)}{\Re p(z)} \right| \leq \frac{\mathcal{H}(p)}{(1 - r)(1 + r)}.$$

If $\frac{1}{1+r} \leq 1 + \varepsilon$, then

$$\frac{\Re zp'(z)}{|z|\Re p(z)} \leq \frac{\mathcal{H}(p)(1 + \varepsilon)}{2(1 - r)},$$

for $|z| > r$. Taking into account that $\left| \frac{p'(z)}{\Re p(z)} \right|$ is bound in disc $|z| \leq r$, for all $\varepsilon > 0$ we have that

$$\frac{\Re zp'(z)}{|z|\Re p(z)} \leq \frac{\mathcal{H}(p)(1 + \varepsilon)}{2(1 - r)} + O\left(\frac{1}{(1 - r)^\alpha}\right).$$

Thus to find some result for class \mathcal{P}_Ω , it is sufficient to study only function p_Ω .

Theorem 3 *If $\Re p(z, t) \geq k > 0$, then function $p(z, t)$ satisfies conditions of theorem 1.*

In this case $\Omega = \{z : \Re z > k\}$ and function $p_\Omega = (1-k)\frac{1+z}{1-z} + k$. We only need to show that equation (4) is true. Assuming $z = re^{i\phi}$, thus we have

$$\frac{|p_\Omega|}{\Re p_\Omega} = \frac{\left| \frac{1+(1-2k)re^{i\phi}}{1-re^{i\phi}} \right|}{\Re \left(\frac{1+(1-2k)re^{i\phi}}{1-re^{i\phi}} \right)}, 0 < r < 1, \phi \in \mathbb{R}.$$

From this

$$\frac{|p_\Omega|}{\Re p_\Omega} = \frac{\sqrt{1+2(1-2k)r \cos \phi + (1-2k)^2 r^2} \sqrt{1-2r \cos \phi + r^2}}{1-2kr \cos \phi - (1-2k)r^2}.$$

Calculation of derivative on ψ gives us that maximum of this function can be only in points $\cos \phi = \pm 1$, $\cos \phi = \frac{2kr}{1-r^2+2kr^2}$. Thus

$$\frac{|p_\Omega|}{\Re p_\Omega} = \frac{|p_\Omega|}{\Re p_\Omega} \leq \max \left\{ \sqrt{\frac{(1+(2k-1)r^2)^2}{(1-r^2)(1-(1-2k)r^2)}}, 1 \right\}.$$

Taking into account that $1 > k > 0$ we have that there is a constant $C_1 > 0$, such that

$$\frac{|p_\Omega|}{\Re p_\Omega} \leq C \frac{1}{(1-r)^{1/2}}.$$

Theorem 4 *If $0 < a < \Re p(z, t) < b$ then the solution $w(z, t)$ of equation (4) is Hoelder function in the closed unit disk and the boundary of domain the $w(\mathbb{D}, t)$ is a Jordan curve.*

In this case $\Omega = \{z : a < \Im m z < b\}$, $0 < a < 1 < b < \infty$ and $p_\Omega(w) = \frac{b-a}{2} \frac{i}{\pi} \log \frac{1+z}{1-z} + \frac{a+b}{2}$.

$$\mathcal{H}(p_\Omega) = \sup_{|z|<1} \frac{(1-|z|^2)^{\frac{b-a}{2\pi}} \frac{1}{|1-z^2|}}{\frac{b-a}{2\pi} \arg \frac{1+z}{1-z} + \frac{a+b}{2}}.$$

Assuming $z = re^{i\pi}$ we have

$$\mathcal{H}(p_\Omega) = \frac{(1-r^2)}{|1-r^2 e^{i2\psi}| (\arg \frac{1+re^{i\psi}}{1-re^{i\psi}} + \delta)},$$

where $\delta = \pi \frac{b+a}{b-a}$. Thus

$$\mathcal{H}(p_\Omega) = \frac{(1-r^2)}{|1-r^2 e^{i2\psi}| (\arg \frac{1+re^{i\psi}}{1-re^{i\psi}} + \delta)}.$$

Taking into account that

$$\arg \frac{1+re^{i\psi}}{1-re^{i\psi}} = \arctan \frac{2r \sin \psi}{1-r^2}$$

and

$$|1 - r^2 e^{i2\psi}| = \sqrt{1 - 2r^2 \cos 2\psi + r^4} = \sqrt{(1 - r^2)^2 + 4r^2 \sin^2 \psi},$$

we obtain

$$\mathcal{H}(p_\Omega) = \frac{2}{\sqrt{1+x^2}(\arctan x + \delta)},$$

where $x = \frac{2r \sin \psi}{1-r^2}$. Using $\sqrt{1+x^2} \arctan x \geq 1$ we have

$$\mathcal{H}(p_\Omega) < \frac{1}{1+\delta}.$$

Therefore function $w(z, t)$ can be extended to a Hoelder function in the closed unit disk with exponent $1 - \frac{1+\varepsilon}{1+\pi \frac{b+a}{b-a}}$ for each $\varepsilon > 0$. Thus boundary L of domain $w(\mathbb{D}, t)$ is curve. From previous theorem we have that inverse function of $w(z, t)$ is continuous on closer of domain $w(\mathbb{D}, t)$. So each point from L has only one preimage. This means that L is Jordan curve.

Theorem 5 *If*

$$\frac{\Im p(z, t)}{\Re p(z, t)} < C, \quad (11)$$

then the solution $w(z, t)$ of equation (4) is a Hoelder function in the closed unit disk and the boundary of domain the $w(\mathbb{D}, t)$ is a Jordan curve.

In this case Ω is a sector, symmetrical with respect to real axis, with angle $2 \arctan C$. Thus $p_\Omega = \left(\frac{1+z}{1-z}\right)^\alpha$, where $\alpha = \frac{2}{\pi} \arctan C$.

$$\mathcal{H}(p_\Omega) = \frac{\alpha \left|\frac{1+z}{1-z}\right|^{\alpha-1} \frac{2}{|1-z|^2}}{\Re \left(\frac{1+z}{1-z}\right)^\alpha}.$$

Assuming $z = re^{i\phi}$, we have

$$\mathcal{H}(p_\Omega) = \frac{4\alpha \frac{|1+re^{i\phi}|^{\alpha-1}}{|1-re^{i\phi}|^{\alpha+1}}}{\Re \left(\alpha \frac{1+re^{i\phi}}{1-re^{i\phi}}\right)^\alpha + \left(\frac{1+re^{-i\phi}}{1-re^{-i\phi}}\right)^\alpha}.$$

Thus

$$\mathcal{H}(p_\Omega) = \frac{4\alpha(1+2r \cos \phi + r^2)^{\frac{\alpha-1}{2}}(1-2r \cos \phi + r^2)^{\frac{\alpha-1}{2}}}{\Re(1+i2r \sin \phi - r^2)^\alpha + (1-i2r \sin \phi - r^2)^\alpha}.$$

Note that

$$\Re(1 \pm i2r \sin \phi - r^2)^\alpha \geq (1-r^2)^\alpha,$$

and equality achieves only for $\sin \phi = 0$. Note that

$$4(1 + 2r \cos \phi + r^2)^{\frac{\alpha-1}{2}}(1 - 2r \cos \phi + r^2)^{\frac{\alpha-1}{2}} \leq (1 - r^2)^{\frac{\alpha-1}{2}}$$

and equality achieves only for $\cos \phi = \pm 1$. Thus $\mathcal{H}(p_\Omega) = 2\alpha$. So, function $w(z, t)$ is Hoelder continuous in closed disc with exponent $1 - \alpha - \varepsilon$.

From inequality (11) follows that $|p_\Omega(z)| \leq \sqrt{1 + C^2} \Re p_\Omega(z)$ or conditions (3) in theorem 1 Taking into account previous argumentation we have

$$\Re p_\Omega(z) \geq \frac{(1 - r^2)^\alpha}{(1 + r)^{2\alpha}} = \frac{(1 - r)^\alpha}{(1 + r)^\alpha},$$

this means inequality (4). Repeat argumentation from proof of theorem 5 we have the necessary result.

5 Application

Often in many phenomenas, that are described by Loewner-Kufarev equation, function $p(z, t)$ is defined by integral representation. It is well known, what any function $p(z)$, $\Re p(z) > 0$ can be represented as

$$\int_0^{2\pi} \frac{e^{i\theta} + z}{e^{i\theta} - z} d\mu(\theta),$$

where $\mu(\theta)$ is a non-decreasing function.

Let $\xi(\theta) = \mu'(\theta)$. Thus if

$$\xi(\theta) = \frac{1}{|f'(e^{i\theta})|^2},$$

then equation (1) will describe the Hele-Show flow. L. Carleson and N. Makarov [5] used

$$\xi(\theta) = \inf\{\epsilon : \text{dist}(f(e^{i\theta}(1 - \epsilon)), t), \partial f(\mathbb{D}, t) = \delta\}, \delta > 0$$

for the description of processes similar to DLA. In this part we connect behaviour of function $\xi(\theta)$ with boundary behaviour of solution of Loewner-Kufarev equation.

First let us assume that $0 < a < \xi(\theta) < b$. Then taking into account that, $p(0, t) > 0$, we can make a change of time from t on τ in equation 1 so, that $p^*(0, \tau) = 1$ and $p^*(z, t) = \frac{dt}{d\tau} p(z, t)$. We obtain that $\frac{a}{b} < p^*(z, \tau) < \frac{b}{a}$. Thus function $p^*(z, \tau)$ satisfies conditions of theorem and we have that solution of Loewner-Kufarev equation maps unit disc onto domain with Hoelder boundary that is Jordan curve.

Let us add condition on $\xi(\theta)$, that it is Hoelder continuous this exponent $0 < k < 1$. Taking into account that (see for e.g. [14] p.69-79)

$$|p_z^*(z, t)| = O\left(\frac{1}{(1 - r)^{1-k}}\right).$$

Note that $\Re p(z, t) > \frac{a}{b}$ we have, that the function satisfies conditions of item 2 in theorem 2. So we have that the solution $w(z, t)$ maps unit disc onto domain with rectifiable and quasiconformal boundary.

References

- [1] I. A. Aleksandrov, *Parametric continuations in the theory of univalent functions*, Nauka, Moscow, 1976. (in Russian)
- [2] J. Becker, *Loewnersche Differentialgleichung und quasikonform fortsetzbare schlichte Funktionen*, J. Reine Angew. Math. **255** (1972), 23-43.
- [3] J. Becker, *Loewnersche Differentialgleichung und Schlichtheitskriterien*, Math. Ann. **202** (1973), 321-335.
- [4] L. de Branges, *A proof of the Bieberbach conjecture*, Acta Math. **154** (1985) no. 1-2, 137-152.
- [5] L. Carleson, N. Makarov, *Aggregation in the plane and Loewner's equation*, Comm. Math. Phys. **216** (2001), 583-607.
- [6] L. A. Galin, *Unsteady filtration with a free surface*, Dokl. Akad. Nauk USSR, **47** (1945), 246-249. (in Russian)
- [7] V. V. Goryainov, *Distortion theorems for some class of univalent function*. Theory of functions and mappings, Naukova Dumka, Kiev, 1979, 75-85. (in Russian)
- [8] V. Ya. Gutlyanski, *On some classes of univalent functions*. Theory of functions and mappings, Naukova Dumka, Kiev, 1979, 85-97. (in Russian)
- [9] G. Lawler, O. Schramm, W. Werner, *Values of Brownian intersection exponents I: Half-plane exponents*, Acta Math. **187** (2001), 237-273.
- [10] G. Lawler, O. Schramm, W. Werner *The dimension of the planar Brownian frontier is 4/3*, Math. Res. Lett. **8** (2001), 401-411.
- [11] Ch. Loewner, *Untersuchungen über schlichte konforme Abbildungen des Einheitskreises*, Math. Ann. **89** (1923), 103-121.
- [12] J. Lind, *A sharp condition for the Loewner equation to generate slits*, Ann. Acad. Sci. Fenn. **30** (2005), 143-158.
- [13] D.E. Marshall and S. Rohde *The Loewner differential equation and slit mappings* J. Amer. Math. Soc. **18** (2005), 763-778.
- [14] N. I. Muskhelishvili, *Singular Integral Equations*, Nauka, Moscow, 3ed, 1968. (in Russian)

- [15] P. Ya. Polubarinova-Kochina, *On a problem of the motion of the contour of a petroleum shell*, Dokl.Akad. Nauk USSR, **47** (1945), no. 4, 254-257. (in Russian)
- [16] P. Ya. Polubarinova-Kochina, *Concerning unsteady motions in the theory of filtration*, Prikl. Matem. Mech., **9** (1945), no. 1, 79-90. (in Russian)
- [17] Ch. Pommerenke, *Univalent functions, with a chapter on quadratic differentials by G. Jensen*, Vandenhoeck & Ruprecht, Goettingen, 1975.
- [18] L. S. Pontryagin, *Ordinary differential equations*, Nauka, Moscow, 4ed, 1974. (in Russian)
- [19] S. Rohde, O. Schramm, *Basic properties of SLE*, arXiv:math.PR/0106036
- [20] O. Schramm, *Scaling limits of loop-erased random walks and uniform spanning trees*, Israel Jour. Math. **118** (2000), 221-288.
- [21] A. Vasil'ev, *Evolution dynamics of conformal maps with quasiconformal extension* Bull. Sci. Math. **129** (2005), no. 10, 831-859.
- [22] A. Vasil'ev, *Univalent functions in two-dimensional free boundary problems*, Acta Applic. Math. **79** (2003), no. 3, 249-280

Alexander Kuznetsov:

Alexander.A.Kuznetsov@gmail.com

Saratov State University, Department of Mathematics and Mechanics, Astrakhan-
skaya Str. 83, 410012 Saratov, Russia