

Hierarchical Spatial Organization of Geographical Networks

Bruno A. N. Travençolo and Luciano da F. Costa

Cybernetic Vision Research Group, GII-IFSC, Universidade de São Paulo, São Carlos, SP, Caixa Postal 369, 13560-970, Brazil, luciano@if.sc.usp.br.

(Dated: November 20, 2018)

In this work we propose the use of a hierarchical extension of the polygonality index as a means to characterize and model geographical networks: each node is associated with the spatial position of the nodes, while the edges of the network are defined by progressive connectivity adjacencies. Through the analysis of such networks, while relating its topological and geometrical properties, it is possible to obtain important indications about the development dynamics of the networks under analysis. The potential of the methodology is illustrated with respect to synthetic geographical networks.

PACS numbers:

Geographical complex networks have been used to model biological systems - e.g. bone structure [1] and mammalian cortical areas [2]. In these systems, the physical proximity and communication between neighboring elements are critical to guarantee proper development and biological function. Another important property of these systems is its spatial organization. It is known that many biological systems depend on the proper adjacency of cells for correct development - e.g. cell communication, distribution of cells in retina, kidney structures, among many others.

Despite the large set of tools and measurements used to characterize the networks that underly these systems [3], relatively little attention has been given to the spatial organization of the nodes. In order to address this issue, we propose the use of the polygonality index [4] - a robust measurement able to quantify the spatial order of system of points - over the nodes of a geographical network.

The polygonality index assigns to each node of the network a value that indicates the amount of the organization around the node with respect to its neighbors. This value is computed based on the angle α_i formed between successive neighbors of a node, as exemplified in Fig. 1 and expressed by the following equation:

$$h_k = \frac{1}{\sum_{i=1}^{N_k} |\alpha_i - \beta| + 1} \quad (1)$$

where k is the node under analysis, N_k is the number of neighbors of the node k and β_k is a parameter whose value can be fixed - to characterize specific spatial arrangements - or can vary accordingly to the number of the neighbors of the nodes, i.e, $\beta = 2\pi/N_k$. A fixed value of β allows one to identify whether the spatial position of the adjacent nodes obeys a specific arrangement (hexagonal, orthogonal, etc.). For example, if $\beta = \pi/3$, it is possible to quantify how much the nodes of network differ from an hexagonal arrangement. When the value of β depends on the number of the neighbors, the polygonality index indicates if the angles between neighbors are equally distributed. In both cases, the polygonality index varies between 0 (total lack of spatial order) and 1




FIG. 1: **Quantifying the spatial organization around a node.** All the angles α_i formed between successive neighbors (q_i) of the node p_k are used to compute the polygonality index (see Eq. 1).

(fully organized system).

In this article we extend the concept of polygonality in order to allow the quantification of the spatial order in a complex network including several hierarchical levels, allowing the quantification of the spatial organization along a wider neighborhood around each node. Figure 2 illustrates the computation of such a polygonality index for three different hierarchical levels of a node. Note that *virtual edges*[5] are established between the central node and its hierarchical neighbors. The angles between successive virtual edges are taken into account to compute the polygonality index, as expressed in Eq. 1.

Figure 3 shows six networks and respective polygonality indices. In (a-c) the networks are perfectly arranged in hexagonal lattices, while in (d-f) the networks originated from a hexagonal lattice, but the positions of their nodes were perturbed in order to reduce the overall order. In (a) and (d) the first hierarchical level was considered to estimate the polygonality index, while in (b) and (e) the second level was used. The results in (c) and (f) consider the third level. In all these images the nodes were colored accordingly to their polygonality index, as expressed in the color-bar.

Analyzing the values of the polygonality index in Figure 3, it is possible to note that in (a), as the nodes and




FIG. 2: **Hierarchical neighbors used to estimate the spatial organization.** Hierarchical neighbors 1 (a), 2 (b) and 3 (c). The virtual edges (in blue) between the central node and its hierarchical neighbors are used to estimate the angles α_i for polygonality index computation (see Eq. 1) .

its neighbors are arranged in a hexagonal way, the polygonality index for all nodes is the maximum possible. In (b), the angles defined between successive neighbors of the nodes are constant and equal $\beta = \pi/6$. As consequence, the polygonality index is also at its maximum value. On the other hand, in (c) the angles between successive neighbors of the third hierarchy are no longer constant, but the polygonality index is the same for all nodes. In the case of the perturbed lattices (Figure 3(d-f)), the polygonality index for all hierarchies are, as expected, lower than the regular lattices.

All in all, the presented measurements allowed a comprehensive characterization of the spatial organization of

geographical complex networks. They can also be compared with several other measurements related to the dynamics and connectivity of the network, paving the way to obtaining new insights about the topological and geometrical properties of these networks.

Acknowledgments

Luciano da F. Costa is grateful to CNPq (301303/06-1) and FAPESP (05/00587-5). Bruno A. N. Travençolo thanks to FAPESP for financial support (03/13072-8).

-
- [1] L. da F. Costa, M. Viana, and M. Beletti, *Applied Physics Letters* **88**, 033903 (2006).
 - [2] L. da F. Costa, M. Kaiser, and C. C. Hilgetag, *BMC Syst Biol* **1**, 16 (2007), URL <http://dx.doi.org/10.1186/1752-0509-1-16>.
 - [3] L. da F. Costa, F. A. Rodrigues, G. Travieso, and P. R. Villas Boas, *Advances in Physics* **56**, 167 (2007).
 - [4] L. da F. Costa, F. Rocha, and S. M. A. de Lima,

- Physical Review E (Statistical, Nonlinear, and Soft Matter Physics) **73**, 011913 (pages 10) (2006), URL <http://link.aps.org/abstract/PRE/v73/e011913>.
- [5] L. Fontoura Costa and F. Silva, *Journal of Statistical Physics* **125**, 841 (November 2006), URL <http://www.ingentaconnect.com/content/klu/joss/2006/00000125>




FIG. 3: Hierarchical polygonality index for a perfect hexagonal lattice (a-c) and for a perturbed hexagonal lattice (d-e). The hierarchical level (R), the mean (μ) and standard deviation (σ) values of the polygonality index are: (a) $R = 1$, $\mu = 1$, $\sigma = 0$; (b) $R = 2$, $\mu = 1$, $\sigma = 0$; (c) $R = 3$, $\mu = 0.73$, $\sigma = 0$; (d) $R = 1$, $\mu = 0.63$, $\sigma = 0.07$; (e) $R = 2$, $\mu = 0.64$, $\sigma = 0.05$; (f) $R = 3$, $\mu = 0.61$, $\sigma = 0.05$.