

Comment to the paper "Radiation induced by relativistic electrons propagating through random layered stacks: Numerical simulation results" by A.A.Varfolomeev and et al NIM B 256, 705 (2007)

Zh.S. Gevorkian^{1,2,3,*}, J.Verhoeven⁴

¹ *Institute of Physics, Academia of Sinica, Nankang, Taipei 11529, Taiwan*

² *Institute of Radiophysics and Electronics, Ashtarak-2, 378410, Armenia.*

³ *Yerevan Physics Institute, Alikhanian Brothers St. 2, Yerevan 375036, Armenia.*

⁴ *FOM-Instituut voor Atoom-en Molecuulfysica, Kruislaan 407, 1098SJ Amsterdam, The Netherlands.*

* *Corresponding author: gevorkia@phys.sinica.edu.tw*

Abstract

We show that the numerical code used in the above mentioned paper does not take into account the multiple scattering effects of electromagnetic field properly and is therefore incorrect.

PACS numbers: 41.60.-m, 07.85.Fv, 41.75.Fr, 87.59.-e

Radiation originating from relativistic charged particle passing through a random stack of foils is considered by Varfolomeev and et al [1] using some numerical code. We treated the same problem analytically [2, 3] before. The results of this numerical simulation qualitatively coincide with the part of our analytical consideration representing the contribution of single scattering to the radiation intensity[2].

One of the main assumptions of the code used in [1] is the constancy of the transversal to foils component of the photon momentum. Writing this condition explicitly for two consecutive foils one finds $\sqrt{\varepsilon_i} \sin \theta_i = \sqrt{\varepsilon_{i+1}} \sin \theta_{i+1}$, where ε_i and θ_i are the dielectric constant and the angle of the photon momentum with the normal to foils in the i -th foil. One immediately recognizes in this condition the Fresnel law of refraction. As it is well known Fresnel laws of reflection and refraction are correct in the geometrical optics limit when photons are treated as rays. Therefore the claim of paper [1] that their code is universal and applicable for all photon wavelengths is incorrect.

The condition for applicability of geometrical optics approximation is that the photon wavelength is smaller than all characteristic sizes of the system. In our case the characteristic size is the foil thickness d . The corrections to geometrical optics when considering scattering on a single foil are of order λ/d . However for a multiple foil system these corrections become of order $N\lambda/d$ and can be large even for $\lambda \ll d$ if number of foils $N \gg 1$. Therefore using geometrical optics approximation for solving Maxwell equations in the multiple stack even for $\lambda \ll d$, as in [1], is incorrect.

As we mentioned above the numerical results of [1] qualitatively reproduce the results of single scattering contribution to radiation intensity of our analytical consideration [2]. This means that the contribution of multiple scattering effects of electromagnetic field to the spectral-angular radiation intensity within the geometrical optics approximation is negligible compared to the single scattering contribution which is simply the incoherent sum of transition radiation photons from random interfaces. However if one properly takes into account the multiple scattering effects their contribution to radiation intensity can exceed the single scattering contribution as it is shown in [2].

Note that we also in our analytical consideration [2, 3] use the geometrical optics approximation. However we only use this to consider of single scattering by a foil in order to find the transmission coefficient through a foil. But we do not use it for solving whole Maxwell equations for a random stack as in [1]. Note also that in our analytical consideration [2] the transversal component of photon momentum in contrary to [1] is not a constant but is a variable integrated quantity.

It is obvious that the angular distribution of emitted photons in the weak absorbing multiply-scattering random layered stack should be the same in the forward and backward directions far away from the radiating system no matter what is the origin of appearing photons: Cherenkov radiation, transition radiation or other mechanisms. Even if the photons initially appear on the forward to particle velocity direction as, for example in the Cherenkov radiation case, after several scattering events on the random spaced interfaces the number of backward and forward photons will become approximately equal. Note that if this was not so we could not observe such a fundamental phenomena as diffuse scattering of light. Instead the numerical simulation shows exceeding of forward intensity up to four order even in the absence of absorption. Of course it is very difficult to achieve the photon multiple scattering regime $\lambda \ll l \ll l_{in}, L$, where l, l_{in} are elastic and inelastic mean free paths of photon in the medium, respectively and L is the size of the system, in the X-ray region. However we think that in some special cases it is still possible for UV and soft X-rays [3]. At the same time it is much more easy to achieve photon multiple scattering regime in the optical region that we have demonstrated experimentally [4]. Summarizing, the code used in [1] to consider the radiation of charged particles in random stack reproduces qualitatively only single scattering contribution of our analytical

consideration and does not take into account multiple scattering effects properly. Therefore in all those cases where the photon multiple scattering effects are important it is incorrect.

The problem of numerical simulation of radiation of a relativistic particle passing through a random stack remains a difficult and open problem. The paper of Varfolomeev and et al [1] contributes to a better understanding of problem. We hope that authors will take into account these comments and will come up with the revised simulation.

-
- [1] A.A.Varfolomeev, M.J.van der Wiel, T.V.Yarovoy and D.A.Ovchinnikov,NIM B **256**,705,(2007).
 - [2] Zh.S.Gevorkian, Phys.Rev.E **57**, 2338 (1998); Sov.Phys.JETP **114**,91, (1998).
 - [3] Zh.S.Gevorkian and J.Verhoeven, NIM B **252**,57,(2006).
 - [4] Zh.S.Gevorkian and et al,Phys.Rev.Lett.,**97**,044801,(2006).