## LIFSHITZ ASYMPTOTICS FOR HAMILTONIANS MONOTONE IN THE RANDOMNESS

IVAN VESELIĆ

ABSTRACT. This is a note for the report on the Oberwolfach Mini-Workshop: Multiscale and Variational Methods in Material Science and Quantum Theory of Solids, published in [MW].

In various aspects of the spectral analysis of random Schrödinger operators monotonicity with respect to the randomness plays a key role. In particular, both the continuity properties and the low energy behaviour of the *integrated density of states* (IDS) are much better understood if such a monotonicity is present in the model than if not.

In this note we present Lifshitz-type bounds on the IDS for two classes of random potentials. One of them is a slight generalisation of a model for which a Lifshitz bound was derived in a recent joint paper with Werner Kirsch [KV]. The second one is a breather type potential which is a sum of characteristic functions of intervals. Although the second model is very simple, it seems that it cannot be treated by the methods of [KV]. The models and the proofs are motivated by well-established methods developed for so called alloy type potentials. The basic notions of random Schrödinger operators and the IDS can be inferred e.g. from [CL90, PF92, Sto01, KM07, Ves06].

§1 Random Schrödinger operators and the IDS. We consider Schrödinger operators on  $L^2(\mathbb{R}^d)$  with a random,  $\mathbb{Z}^d$ -ergodic potential. More precisely, the random potential  $W_\omega\colon\mathbb{R}^d\to\mathbb{R}$  is determined by an i.i.d. family of non-trivial, bounded random variables  $\lambda_k\colon\Omega\to[\lambda_-,\lambda_+]=:J$  indexed by  $k\in\mathbb{Z}^d$  and distributed according to the measure  $\mu$ , and a jointly measurable single site potential  $u\colon J\times\mathbb{R}^d\to\mathbb{R}$ . We assume that  $\lambda_-\in\sup\mu$  and that  $\sup_{\lambda\in J}|u(\lambda,\cdot)|\in\ell^1(L^p),\ p>\max(2,d/2)$ . Under these assumptions the random potential

(1) 
$$W_{\omega}(x) = \sum_{k \in \mathbb{Z}^d} u(\lambda_k(\omega), x - k)$$

is relatively bounded with respect to the Laplacian with relative bound zero, uniformly in  $\omega$ . Consequently, for a bounded  $\mathbb{Z}^d$ -periodic potential  $W_{\mathrm{per}}$  the operators  $H_{\mathrm{per}} := -\Delta + W_{\mathrm{per}}$  and  $H_{\omega} := H_{\mathrm{per}} + W_{\omega}$  are selfadjoint on the domain of  $\Delta$  and lower bounded uniformly in  $\omega$ . Moreover,  $(H_{\omega})_{\omega}$  forms an ergodic family of operators. Hence there exist a closed  $\Sigma \subset \mathbb{R}$  and an  $\Omega' \subset \Omega$  of full measure, such that for all  $\omega \in \Omega'$  the spectrum of  $H_{\omega}$  coincides with  $\Sigma$ . For  $\Lambda_L := [-L/2, L/2]^d, L \in \mathbb{N}$  define the distribution function  $N(E) := L^{-d} \mathbb{E} \left\{ \mathrm{Tr}[\chi_{]-\infty,E]}(H_{\omega}) \chi_{\Lambda_L} \right\}$ . This function is independent of L and is called IDS or spectral distribution function. The support of the associated measure coincides with  $\Sigma$ . The IDS can be approximated in the sense of distribution functions by its finite volume analogs  $N_{\omega}(E) := L^{-d} \sharp \{ \text{eigenvalues of } H_{\omega}^L \leq E \}$  almost surely. Here  $H_{\omega}^L$  denotes the restriction of  $H_{\omega}$ 

to  $\Lambda_L$  with Neumann boundary conditions. For many types of random Hamiltonians the IDS is expected to be very "thin" near the spectral minimum  $E_0 := \min \Sigma$ . More precisely I. M. Lifšic conjectured in [Lif63, Lif64] an asymptotic behaviour of the form  $N(E) \sim ce^{-\tilde{c}(E-E_0)^{-d/2}}$  for  $E-E_0$  small and positive, where  $c, \tilde{c}$  denote some positive constants. The spectrum near  $E_0$  corresponds to very rare configurations of the randomness and  $E_0$  is consequently called a fluctuation boundary.

§2 A class of potentials monotone in the randomness. Here we present a slight extension of the main result in [KV]. Assume that the potentials u and  $W_{\rm per}$  satisfy the following

HYPOTHESIS A. For any  $\lambda \in J$  we have supp  $u(\lambda, \cdot) \subset \Lambda_1$  as well as  $u(\lambda, x) \ge u(\lambda_-, x)$  for all  $x \in \mathbb{R}^d$ . There exist  $\epsilon_1, \epsilon_2 > 0$  such that for all  $\lambda \in [\lambda_-, \lambda_- + \epsilon_2]$ 

$$\int_{\mathbb{R}^d} dx \, u(\lambda, x) \ge \epsilon_1 \, (\lambda - \lambda_-) + \int_{\mathbb{R}^d} dx \, u(\lambda_-, x)$$

and for all  $\lambda \in [\lambda_- + \epsilon_2, \lambda_+]$ 

$$\int_{\mathbb{R}^d} dx \, u(\lambda, x) \ge \int_{\mathbb{R}^d} dx \, u(\lambda_- + \epsilon_2, x)$$

hold. The function  $\lambda \mapsto u(\lambda, x)$  is Lipschitz continuous at  $\lambda_-$ . More precisely, for some  $\kappa$ , all  $x \in \Lambda_1$  and all  $\lambda \in [\lambda_-, \lambda_- + \epsilon_2]$  we have  $u(\lambda, x) - u(\lambda_-, x) \le \kappa(\lambda - \lambda_-)$ . If  $d \ge 2$ , then for any  $\lambda \in J$  the functions  $u(\lambda, \cdot)$  and  $W_{\rm per}$  are reflection symmetric with respect to all d coordinate axes

Typical examples of potentials u satisfing Hypothesis A are: an alloy type potential, i.e.  $u(\lambda, x) = \lambda f(x)$  with  $L_c^{\infty}(\Lambda_1) \ni f \geq 0$ , and a breather type potential, i.e.  $u(\lambda, x) = f(x/\lambda)$  with supp  $f \subset \Lambda_{\lambda_-}, \lambda_- > 0, f \in C^1(\mathbb{R}^d \setminus \{0\})$  and  $L^{\infty}(\mathbb{R}^d) \ni g(x) := -x \cdot (\nabla f)(x) \geq 0$ .

THEOREM B. (LIFSHITZ BOUND) Under the Hypothesis A the IDS of the Schrödinger operator  $H_{\omega} := -\Delta + W_{\rm per} + W_{\omega}$  satisfies

(2) 
$$\lim_{E \searrow E_0} \frac{\log |\log N(E)|}{\log (E - E_0)} \le -\frac{d}{2}$$

Thus for  $E-E_0$  small and positive, asymptotically the bound  $0 < N(E) \le e^{-\tilde{c}(E-E_0)^{-d/2}}$  holds. The proof is essentially the same as in [KV].

§3 Breather potentials with characteristic functions of intervals. We consider a very explicite class of random potentials on  $\mathbb{R}$ . Let  $(\lambda_k)_{k\in\mathbb{Z}}$  be as before with  $\lambda_-=0$ ,  $\lambda_+=1$ . The breather type potential

(3) 
$$W_{\omega}(x) = \sum_{k \in \mathbb{Z}} u(\lambda_k(\omega), x - k), \text{ where } u(\lambda, x) = \chi_{]0,\lambda]}(x)$$

does not satisfy the Lipschitz condition in Hypothesis A. Nevertheless we have

THEOREM C. The IDS of the Schrödinger operator  $H_{\omega} := -\Delta + W_{\omega}$ , where  $W_{\omega}$  is as in (3), satisfies the Lifshitz bound (2). Note that  $E_0 = 0$  for this model.

It seems that the reason why the method of [KV] is not applicable to the potential (3) is the use of Temple's inequality [Tem28]. For Temple's inequality to yield an efficient estimate, the second moment  $\langle H_{\omega}^L \psi, H_{\omega}^L \psi \rangle$  in an well chosen state  $\psi$  has to be much smaller than the first moment  $\langle \psi, H_{\omega}^L \psi \rangle$ . For the current application the best choice of  $\psi$  seems to be the periodic, positive ground state of  $H_{\rm per}$ . However for such  $\psi$  and for the potential (3), the first and second moment coincide! It turns out that Thirring's inequality [Thi94, 3.5.32] is better adapted to the model under consideration. It was used before in [KM83] in a similar context.

Sketch of Proof: As before the superscript L denotes the Neumann b. c. restriction to [-L/2, L/2]. Since  $N(E) \leq L^{-1} \text{Tr}[\chi_{]-\infty, E]}(-\Delta^L)] \mathbb{P}\{\omega \mid E_1(H_\omega^L) \leq E\}$ for any  $L \in \mathbb{N}$ , it is sufficient to derive an exponential bound on the probability that the first eigenvalue  $E_1$  of  $H^L_{\omega}$  does not exceed E.

We set  $I_L := \Lambda_L \cap \mathbb{Z}, H_0 := -\Delta - \alpha/4L^2, \ \psi = L^{-1/2}\chi_{\Lambda_L} \text{ and } V_{\omega}(x) = \alpha/4L^2 + W_{\omega}(x)$ . Then  $E_1(H_0^L) = -\alpha/4L^2$  and  $E_2(H_0^L) \geq 3\alpha/4L^2$ , cf. [KS87]. Since  $V_{\omega}$  does not vanish,  $V_{\omega}^{-1}$  is well-defined and we calculate  $L(\int_{\Lambda_L} V_{\omega}(x)^{-1} dx)^{-1} = 0$  $\frac{\alpha}{4L^2} \frac{4L^2 + \alpha}{4L^2 - 4L^2 S_L + \alpha}$ . We use the notation  $S_L := L^{-1} \sum_{k \in I_L} \lambda_k$  for averages,  $\tilde{\lambda}_k := L^{-1} \sum_{k \in I_L} \lambda_k$  $\min(\lambda_k, 1/2)$  for cut-off random variables and similarly  $\tilde{V}_{\omega}$ ,  $\tilde{S}_L$  for the potential and the averages. Then  $E_1(H_0^L) + \langle \psi, \tilde{V}_{\omega}^{-1} \psi \rangle^{-1} \leq \alpha/4L^2 < E_2(H_0^L)$ , thus Thirring's inequality is applicable and yields

$$E_1(H_{\omega}^L) \ge E_1(\tilde{H}_{\omega}^L) \ge E_1(H_0^L) + \langle \psi, \tilde{V}_{\omega}^{-1} \psi \rangle^{-1} \ge \frac{\alpha \tilde{S}_L}{5L^2}$$

as soon as  $L^2 \geq \alpha$ . For a given E > 0 chose  $L := \lfloor \beta E^{-1/2} \rfloor$ , then  $\mathbb{P}\{E_1(H_\omega^L) \leq E\} \leq \mathbb{P}\{\alpha \tilde{S}_L/5L^2 \leq E\} \leq \mathbb{P}\{\alpha \tilde{S}_L/5 \leq \beta^2\}$ . Since  $0 < \mathbb{E}\{\tilde{S}_L\} = \mathbb{E}\{\tilde{\lambda}_k\} \leq 1/2$ it is possible to choose  $0 < \beta \le \sqrt{\alpha \mathbb{E}\{\tilde{\lambda}_k\}/10}$ . With this choice we have  $\mathbb{P}\{\tilde{S}_L \le$  $5\beta^2/\alpha\} \leq \mathbb{P}\{\tilde{S}_L \leq \mathbb{E}\{\tilde{S}_L\}/2\}$ . A large deviation estimate bounds this probability by  $e^{-cL^d} = e^{-\tilde{c}E^{-d/2}}$  for some positive constants  $c, \tilde{c}$ . This completes the proof.

The higher dimensional analog of this model is currently under study.

## References

- [CL90] R. Carmona and J. Lacroix. Spectral Theory of Random Schrödinger Operators. Birkhäuser, Boston, 1990.
- [KM07] W. Kirsch and B. Metzger. The integrated density of states for random Schrödinger operators. Proceedings of Symposia in Pure Mathematics, 76:649-698. AMS, 2007.
- W. Kirsch and F. Martinelli. Large deviations and Lifshitz singularity of the integrated density of states of random Hamitonians. Commun. Math. Phys., 89:27-40, 1983.
- W. Kirsch and B. Simon. Comparison theorems for the gap of Schrödinger operators. J. [KS87] Funct. Anal., 75:396-410, 1987.
- [KV]W. Kirsch and I. Veselić. Lifschitz tails and localisation for Schrödinger operators with random breather-type potential. http://arxiv.org/math-ph/0609014.
- I. M. Lifshitz. Structure of the energy spectrum of the impurity bands in disordered solid [Lif63] solutions. Sov. Phys. JETP, 17:1159-1170, 1963.
- [Lif64] I.M. Lifshitz. The energy spectrum of disordered systems. Adv. Phys., 13:483-536, 1964.
- [MW]Mini-Workshop: Multiscale and Variational Methods in Material Science and Quantum Theory of Solids organized by: I. Catto, I. Chenchiah, I. Veselić, and J. Zimmer. Oberwolfach Reports, Volume 4(1):371-416, 2007.
- [PF92] L. A. Pastur and A. L. Figotin. Spectra of Random and Almost-Periodic Operators. Springer-Verlag, Berlin, 1992.
- P. Stollmann. Caught by disorder: A Course on Bound States in Random Media, vol-[Sto01] ume 20 of Progress in Mathematical Physics. Birkhäuser, 2001.
- [Tem28] G. Temple. The theory of Rayleigh's principle as applied to continuous systems. Proc. Roy. Soc. London A, 119:276-293, 1028.
- W. Thirring. Lehrbuch der mathematischen Physik. Band 3. Springer-Verlag, Wien, sec-[Thi94] ond edition, 1994. Quantenmechanik von Atomen und Molekülen.
- I. Veselić. Existence and regularity properties of the integrated density of states of random [Ves06] Schrödinger Operators, 2006. Habilitation thesis. TU Chemnitz. To be published in the Springer-Verlag LNM series.

TU CHEMNITZ, FAKULTÄT FÜR MATHEMATIK, 09107 CHEMNITZ, GERMANY & EMMY-NOETHER-Programme of the Deutsche Forschungsgemeinschaft