

An Equilibrium Balance of the Universe

Ernst Fischer

e-mail: e.fischer.stolberg@t-online.de

Contents

1	Introduction	3
2	Back to Einstein	5
3	Red shift	11
4	Black holes - the source of matter	15
5	Ingredients of the intergalactic medium	19
5.1	The intergalactic plasma	20
5.2	The radiation field	21
5.3	Cosmic dust	23
6	Structure formation	26
6.1	Gravitational instability	26
6.2	Thermal instability	29
6.3	Formation of galaxies	30
6.4	The life of galaxies	33
6.5	Galaxy clusters	33
6.6	Spiral galaxies	37
6.7	Active galaxies and quasars	38
7	The universe - a zero-sum game?	41
A	The definition of density and potential energy	45
B	Basic equations for a homogeneous universe	48
C	Red shift - energy loss by gravitational radiation	50
D	X-ray emission of the intergalactic plasma	54
E	Intergalactic dust	56
E.1	Formation of dust	56
E.2	Size selection	58

E.3	Global distribution and stability	59
E.4	Detection of intergalactic dust	60
F	Jeans-instability	62
G	The mass balance of galaxy clusters	64
H	Idea of a symmetric model of elementary particles	68

Chapter 1

Introduction

During the last decades our knowledge of the structure of the universe has taken an enormous step forward due to improved methods of observation and the development of new devices. At the same time the theoretical models to describe the structure and history of the universe have been developed further. Today there exists a so called 'concordance model' that can describe the development of the cosmos, beginning from a 'big bang' up to the present and even forecasting its future.

But in spite of the ability of the mathematical model, to describe many of the observed phenomena rather accurately, some people feel uncomfortable with this model, as it requires rigorous changes of our accustomed thinking. Already Einstein's idea, to abandon space and time as an absolute reference system, was difficult to accept. But what is still worse with the concordance model is, that it shifts the conditions, which led to the present day state of the universe, into some initial conditions and physical principles, which can neither be proved nor disproved observationally, as they only prevailed at some instant of time, when our known physical laws were not valid. In addition the postulated existence of matter or energy fields with rather strange properties, which should dominate the development of the universe, but which cannot be observed in earthbound experiments, makes the model questionable.

But it is the initial singularity, the 'big bang', which appears scarcely acceptable from the physical point of view. Of course, mathematics can easily handle singularities, and in some cases it may be convenient to use singular distributions to describe some effect mathematically, but physically a state of infinite density of some field appears impossible. In physics all our experience says that every model theory, which leads to singularities, is wrong or at least inaccurate under the corresponding conditions. With the big bang theory also the excuse does not help much that near the singularity the known laws of physics are no longer valid. As long as we know no alternative laws, the value of such a statement is zero. Besides, singularities are not restricted to the initial condition, but they are the inevitable final state of matter, when a star or galaxy collapses into a 'black hole' under the influence of gravitation.

According to the generally discussed model of today the size of the universe is infinite, but at the same time it is expanding as a whole. While in the static model, which was originally proposed by Einstein, the curvature of space could explain in an elegant way, why the universe can be of finite size, but at the same time has no limits, this is not the case with the 'concordance model'.

Besides of that, to remain in agreement with observations, the 'concordance model' requires the existence of completely new ingredients to the universe with properties, which are different from all, what was known by now. 95% of gravitational interaction should be dominated by dark matter and dark energy, some stuff, which interacts with conventional matter only by gravity. But the formation of this stuff in the big bang remains a riddle, just as the force or interaction which finally drives the expansion of the universe.

But nevertheless with a suitable adjustment of parameters the 'concordance model' has succeeded to create a system, which is able to describe a large number of observations. As well the formation and abundance of chemical elements as the cosmological microwave background and its inhomogeneities, the red shift of spectral lines and the formation of observed structures can be explained. But it is just this explanation of cosmic structures, which runs into difficulties with increasing improvement of observations. It is scarcely possible to explain, how shortly after the big bang galaxies and galaxy clusters should have been formed, which have similar appearance and metal abundance as those of today.

The concordance model in its present form gives an acceptable description of the observational material, but on the expense of many assumptions, which cannot be verified or falsified and which are scarcely acceptable by our accustomed physical understanding. We must put the question, if there is no alternative to this model, which does not need so many unprovable assumptions and singularities. Hasn't an error crept already into the basic assumptions, which finally leads us into a blind alley? Aren't we in a state like Ptolemaeus, who tried to keep his geocentric universe in accordance with observations by adding more and more epicycles?

In this paper we will try to find possible causes for such a erroneous development and to propose a concept, by which an alternative to the big bang model is possible, which avoids singularity problems and does not need the existence of the mysterious dark matter or dark energy, though explaining all the observations. Of course, it is scarcely possible within the scope of this paper to examine all observations, which have been made over many years, with respect to their compatibility with the model presented here. Instead we will propose a new way of thinking and a new interpretation of observed facts, which may finally lead to an improved understanding of the universe.

We will not concentrate on detailed observations and dispense most references to literature. Literature in the field of astrophysics and cosmology is so copious that a nearly complete reference list would be out of the scope of this paper. Only where we directly refer to data or results from individual authors, references are given.

Chapter 2

Back to Einstein

The probably most important step towards our present day cosmological models was the development of the General Theory of Relativity (GRT) by Albert Einstein. His starting point, to use the equivalence of gravitational mass and inertial mass to describe the complete gravitational interaction by the geometrical properties of the space-time continuum, led to completely new understanding of space and time.

Already with the Special Theory of Relativity and its various experimental confirmations it had become clear that space and time coordinates do not have an absolute meaning, but only the relation of bodies with respect to each other. With his General Theory of Relativity Einstein succeeded in describing the dynamics matter and energy under the influence of gravitation in a way, independent of the reference system, which is only determined by the metric of the space-time continuum.

By the methods of GRT it was not only possible to explain some phenomena like the shift of perihelion of Mercury or the aberration of starlight passing the sun, the new theory offered also the opportunity to make statements on the universe as a whole. With the assumption that space is curved it appeared possible that the universe has no limit, but in spite of that its size is finite. The view of an infinite size of space, which appeared uncomfortable for most physicist of that time, disappeared with Einstein's model. Of course, this model demanded the introduction of a new fundamental quantity, the so called 'cosmological constant', which led to a negative pressure and thus to a repulsive force, counteracting the attractive gravitational force of matter.

A few years after publication of Einstein's theory the mathematician Friedmann showed that there existed a time-dependent solution of Einstein's field equations without the necessity of a cosmological constant under the proposition that space is continuously expanding. As just at the same time the red shift of spectral lines of distant galaxies was discovered, it appeared obvious to take this as a prove that the universe is continuously expanding. Also Einstein himself changed his mind to this interpretation, especially as his original static model proved unstable within the framework of time-dependent solutions of the

basic equations. With a cosmological constant it was possible to establish an equilibrium to the gravitational attraction at a certain radius of curvature, but any small deviation of the curvature would lead to an irreversible expansion or to collapse. Later Einstein defined the introduction of the cosmological constant as his 'greatest blunder'.

With the acceptance of the paradigm of the expanding universe practically the path to the modern 'concordance model' was traced out. The starting point as a singularity and the limited time, available for the development of structures from the big bang to the present, was fixed. All the further corrections, as for instance the addition of an inflationary phase of expansion, or in the last years the claim for the existence of dark energy, were only necessary to keep the model in accordance with the enhanced accuracy of observations.

Inflation, the exponential expansion in the beginning, was necessary to explain, why all parts of the universe have developed in a similar way. That means that they must have had a causal connection, though this was impossible in later times due to the expansion of space. But on the other side introduction of the inflation model required that the total density of all matter, which contributes to gravitation, should equal a critical density, which, however, cannot be explained by the observed matter nor by the chemical element abundances resulting from the big bang nucleosynthesis.

To save the matter balance then the existence of some 'dark matter' was claimed, which should contribute to gravitation, but not interact with normal matter or radiation in any other way. But even this correction was not sufficient, as such matter would cluster in the same way as visible matter due to gravitational interaction. But as from new observations the existence of this dark matter in quantities, necessary to reach the critical density of the universe, must be excluded, finally 'dark energy' was created, some stuff which brings the matter balance to the critical value, but does not conglomerate, as it possesses a negative pressure and thus a repulsive gravitation. With this new form of matter we are practically back with Einstein's cosmological constant without having a better explanation than Einstein had ninety years ago.

If we want to find out the origin of any failure in the development, we must go back further than to the starting point of the theory of an expanding universe. The basic assumptions of GRT should be scrutinised. Of course, since the time of Einstein until today this theory has been confirmed with steadily increasing accuracy, but all these confirmations are related to parameter regions, where small deviations from Newtonian theory of gravitation are discussed. The basics, which were confirmed, were changes of distances or running times of signals, caused by the curvature of space-time, and the existence of gravitational radiation resulting from the fact that changes of the metric of space-time are limited by the speed of light.

There is however one conceptual problem in GRT, to which we should direct our attention: the definition of density. In pre-relativistic times we were accustomed to define the density of a quantity by the ratio of the amount of this quantity and of the volume element in which it is contained. This ratio or, for spatially varying quantities its limiting value, when the size of the volume

element approaches zero, could be considered as the local density value of the corresponding quantity.

According to the concept of GRT, however, the size of a volume element depends on the metric and thus on the distribution of matter or energy in the surrounding space. Thus there is no longer a unique definition of density in the conventional sense. But Einstein easily found a way out of this situation. As a local quantity, which should enter the field equations, he defined the density in the 'tangential space', that means the density value one would measure, if all the surrounding masses were removed to infinity.

In this way the problem of a unique definition is solved, but at the cost of another one. Calculation of the integral of the so defined density over the region of an extended mass distribution, the result is different from what we would get by counting the number of atoms and multiplying it with their characteristic mass. For the example of a spherically symmetric mass distribution, for which the exact solution of the Einstein equations is known, to the first approximation the difference is just the value of potential energy, that means the binding energy, which according to Newtonian theory has to be supplied, to distribute the mass into infinite space against the action of gravitation (see appendix for mathematical details).

But which of the two different values of the mass determines the gravitational action to the outside? Is it, as Einstein said, that 'the inertia of mass is enhanced, when ponderable matter is accumulated in its surrounding' (and according to the principle of equivalence also the gravitational mass), remains it unaffected or does it eventually decrease?

The easiest way to answer this question from our experience is, to consider the limiting case of the collapse of a matter distribution under the influence of gravitation, which finally ends up in an extremely compact state, according to present day concepts, even in a singularity. If the gravitational interaction with the outside world would increase due to the enhanced binding energy, the dynamics of the universe would be governed completely by such singular systems, a fact, which contradicts all observations. But a decreasing gravitational interaction appears improbable as well. In this case it should approach zero in a theoretical model which contains no characteristic scale length. Observations show, however, that there exist very compact objects in space with gravitational masses, corresponding to those of stars or even millions of stars.

Thus we must conclude that gravitational interaction of matter to the outside remains unchanged, when the internal structure is changed. It cannot be true that it is the density, defined in the Euclidean tangent space and integrated over the volume of the curved space, which determines the gravitation to the outside. We have to subtract the binding energy from the matter density to obtain the correct gravitating mass after integration.

The energy tensor, which was used by Einstein to describe the universe as a whole, does not contain the binding energy or, to express it more generally, the (negative) potential energy in any form. Instead he takes over the energy tensor for homogeneously distributed matter from special theory of relativity, in which only the density of matter and the kinetic energy of the particles,

that means the pressure, contribute to the total energy. Of course, this can be regarded as an allowed approximation, as long as the metric and, resulting from this, the dynamics of test particles is dominated by the matter distribution. For the aberration of light, passing the sun, the contribution of potential energy compared to the mass-energy is less then 10^{-6} of the total action at maximum. Of similar size are the corrections also with other observations which have been used as confirmations of the general theory of relativity.

But neglecting potential energy in the energy tensor, which appears as the tiny correction in this case, has strong consequences for the general concept. According to the general theory of relativity, as it is conventionally used, conservation laws of energy and momentum are valid only locally, while changes of the metric can generate or destroy energy. The fact that the dynamical behaviour of test particles within the gravitational field of surrounding matter is described correctly is no indication, however, that the energy tensor, the source term of the field equations, is fully correct. Only by adding potential energy it is possible to develop a model, in which, as in Newtonian theory, the conservation laws are valid locally and in integral form as well.

But the question is: how can we include potential energy within the scope of a covariant theory, that means, in a form which is independent from the reference system? To fulfil this condition any formula for the potential energy may contain only locally defined variables of state beside of the metric tensor and quantities derived from it without differentiation. In addition potential energy should vanish in empty space, that means, it should be proportional to the density of matter. It should vanish in the limiting case of Euclidean geometry, where the special theory of relativity is valid, and in the limiting case of weak curvature potential energy must result in analogy to Newtonian theory (for the mathematical formulation see appendix).

Formally the only sensible possibility to fulfil these conditions is an expression, which for a spatially homogeneous solution looks like the 'cosmological term' in Einstein's static universe, that means, a product of a scalar quantity and the metric tensor. But now the scalar factor is no longer a constant, but it is a quantity, which is proportional to the local energy or matter density and to the inverse of the radius of curvature. The essential difference to the 'cosmological constant' is the fact that an expansion of the universe, that means, a change of curvature, affects potential energy more than mass density. As a consequence a static universe is stable, contrary to the unstable Einstein universe, where the 'cosmological constant' can balance gravitational attraction only at one fixed value of curvature.

There is, however on more serious difference. The field equations of general relativity are relations between tensors. Every tensor component has to fulfil the corresponding equality conditions. But the assumption of a homogeneous medium reduces the number of degrees of freedom to such an extent that it poses additional conditions to the possible forms of the energy tensor, which must be satisfied to allow time dependent solutions. The energy fields have to obey certain equations of state. Only a pressureless mass distribution can change proportional to the volume. Radiation fields, that means, fields with

pressure proportional to $1/3$ of the energy density, must vary with the fourth power of the scale variable. The cosmological term with its negative pressure must be constant, independent of any expansion.

The expression, which we follow from our considerations as a reasonable description of potential energy, does not fulfil this condition. With other words: all time dependent solutions of the field equations for a homogeneous universe lead to contradiction. There is no solution but the trivial one, in which the potential energy of matter is in equilibrium with matter itself. The decisive fact is that potential energy due to its negative pressure acts as a repulsive force counteracting gravitating attraction of matter.

The conception of negative pressure is not as strange as one would think. Also in Newtons theory of gravitation the attraction of masses generates a negative pressure. What is different in general relativity is the fact that this pressure contributes to the gravitational force.

We can imagine the action of negative pressure in Newtonian physics from a simple thought experiment. If we think of space, homogeneously filled with particles at rest, every particle is in equilibrium with its neighbouring masses, as the gravitational forces compensate each other. If, however, the homogeneous distribution contains a little cavity which does not contain matter, a 'vacuum bubble', equilibrium is distorted for the particles near the boundary of the bubble. There are less particles pulling inwards than outwards, so that there is a resulting force tending to enhance the volume of the bubble, as if the bubble had a positive pressure compared to the mass filled space. If we assign zero pressure to the vacuum, the pressure in the surrounding medium must be considered negative.

That we never observe this negative pressure in practice is caused by the fact that under all terrestrial conditions the positive kinetic pressure outbalances it by many orders of magnitude. Thus any 'vacuum bubble' would be immediately filled up by the motion of particles. Generally speaking, positive pressure leads to a homogeneous distribution of matter, while negative pressure leads to clumping and formation of structures. We shall come back to this point later.

An essential difference to the 'big bang' model should be mentioned here. In its presently preferred form this model contains a term, which formally acts like Einstein's cosmological constant. Most people ascribe this term to the zero point energy of quantum fields. But there exists no reasonable answer to the question, why this term should be different from zero, but smaller by many orders of magnitude than estimated by any quantum mechanical model.

In the equilibrium model considered here this problem does not exist. Any energy term, which has the form of a cosmological constant, contributes exactly equal amounts to the density of positive energy and to the negative potential energy. Thus it is without any effect in the field equations (see appendix).

As an important result of the considerations in this chapter we have to state that a universe, in which conservation of mass and energy is valid not only locally, but also globally, has to be generally static, if it is homogeneous except for local fluctuations. Thus there cannot be a 'big bang' and all the structures which we observe must be regarded as local distortions of homogeneity, as 'cos-

mic weather'. If this view of the universe shall be regarded as correct, it must also give explanations for all the observations, which have been explained by the 'concordance model' more or less successfully. What is the reason for the cosmological red shift, if not expansion of space? How can we explain the abundance of elements? What is the source of the microwave background radiation, why is it so homogeneous and what causes the slight inhomogeneities, and how have the matter structures been formed?

In the next chapters we will try to answer these questions within the scope of known and proved physical laws.

Chapter 3

Red shift

The strongest argument, which has led to the development of the 'big bang' model, was the discovery of the cosmological red shift, which was interpreted as change of the wavelength of photons due to the general temporal change of all length scales in the universe. On the other side this concept is based on the assumption that with the propagation of light in a static universe the wavelength of photons should remain unchanged. But this is an assumption, which is not contained in the basic equations of general relativity, but postulated in analogy to the force-free motion in Euclidean space.

As in Einstein's theory the motion of particles or quanta should completely determined by the geometry of space-time, it appeared reasonable to assume that the motion of a test particle should be analogous to the force-free motion in Euclidean space. That means that analogous to the motion with constant energy along a straight line, motion under the influence of gravitation should follow a 'geodesic line', the straightest line in curved space-time. That the energy of motion should be constant along this line does not follow from the field equations. The derivation of the geodesic equation from the principle of minimal action does not require that the action integral is zero, but only that it is minimal.

Does a force-free motion actually exist in curved space? Isn't every motion along a curved path an accelerated motion? In Newtonian physics we have learned that the action of compulsive forces is zero, as they act perpendicular to the direction of motion. But this is true only, if the action is immediate. One characteristic difference between Newtonian gravity and general relativity is the fact that gravitational interaction is limited to the speed of light, similar to electromagnetic interaction. As a consequence the compulsive force has also a component in the direction of motion. There exists a retardation of the interaction, as we know it from electromagnetic theory.

Anyway the principle of geodesic motion can be regarded only as an approximation. It is based on the assumption that a test mass is moving inside a metric, which is determined by the distribution of other matter fields. But at the same time every test mass is part of the matter distribution and thus

contributes to the metric. When in a initially completely homogeneous universe some mass moves from A to B, the universe is no longer homogeneous. It causes a change of the metric which propagates into space at the speed of light.

In most cases this propagating 'gravitational wave' is so weak that to measure it appears scarcely possible. Only with extremely strong changes of the matter distribution, as they occur with the explosion of stars, we can hope to register such waves with very sophisticated detectors.

The only observation which has by now confirmed the existence of gravitational waves, as they are predicted by general relativity, and the loss of energy associated with them, is the decrease of the rotation frequency of the 'double pulsars'. The properties of these systems of collapsed double stars can be accurately measured by their emission of quasi-periodic electromagnetic signals. The observed frequency shift is exactly as expected from general relativity. In this case the energy loss by gravitational waves corresponds to the quadrupole radiation in electromagnetism. Dipole radiation, which dominates in electromagnetic radiation, cannot occur as gravitational radiation, as there are no negative masses, which could form a dipole with the positive ones.

But there should be a 'monopole radiation', the equivalent to electromagnetic bremsstrahlung. Every accelerated electric charge radiates energy, be it bremsstrahlung, which we observe from interactions between free electrons and ions in plasma, or the cyclotron radiation, which is generated by the transversal acceleration, which occurs when electrons are fixed to a curved path by magnetic fields.

Today there exists no exact derivation of this gravitational bremsstrahlung from the field equations of general relativity. But with a simple model it is possible to understand the general properties of this radiation. To do this we start from Newtonian gravity, but regard the fact that the action of forces has to be retarded due to the finite speed of interaction, like it is done to calculate bremsstrahlung in electromagnetic interaction. To avoid the singularities, which occur in Newtonian theory by integration over an infinite volume, we assume that there exists a homogeneous curvature of space so that the volume remains finite. Correspondingly the gravitational force between two points should be assumed to be directed along the geodesic connecting them instead of a straight line. The strength of force should decrease with the square of the distance as measured along this line.

From theoretical point of view this mixture of Newtonian gravity with general relativity may appear somewhat inaccurate, but it gives us the opportunity to understand the mechanism of gravitational bremsstrahlung in a simple way.

Let us first consider a test mass in the middle between two other masses. The gravitational forces of the two outer masses exerted on the test mass are in equilibrium. This holds as well for Newtonian gravity as in general relativity as long as the test mass is at rest. The retardation of forces is equal for both outer masses. But things are different, if the test mass is in motion. According to Newtonian gravity the equilibrium conditions do not change, as the forces act instantaneously. Due to finite speed of signals in general relativity, however, we have to consider the running time of the signal. That means to calculate the

forces we have to use the distances at an earlier point of time. At that time the distance in the direction of motion was larger and thus the attractive force was weaker and against the direction of motion the effect is reversed. Thus there exists a resulting force opposite to the direction of motion, which gets stronger with increasing velocity of the test mass (for the mathematical details see appendix).

This principle can easily applied also to a test mass in the gravitational field of an homogeneous matter distribution, if we include the modifications of Newtonian theory, mentioned above, to consider the curvature of space. Integration of the contributions of all volume elements to the force on a moving test mass leads to loss of momentum, which is the higher the stronger the curvature of space (see appendix). Apparently the principle of momentum loss is valid for every motion with respect to some matter background which can be regarded at rest. That means, it is valid not only in the non-relativistic limit, but also for relativistic particles or energy quanta, if inertia is expressed by its connection to momentum or energy.

This shows that in the theory of general relativity there exists a simple explanation of the cosmological red shift, which does not need any expansion of the universe. The momentum of photons and correspondingly also their energy decreases in proportion to their actual momentum resp. energy. That means: the frequency decreases exponentially with running time or distance from the source, the wavelength increases correspondingly. As a first approximation this is a linear increase of wavelength with distance, just the same dependence as given from the big bang model and which is found from observations.

In our picture the red shift is a consequence of the general principle of momentum loss, which affects every motion, and which will prove necessary for understanding the concept of the dynamical equilibrium model of the universe presented here. One might argue that the loss of momentum or kinetic energy does not agree with the existence of global energy conservation, called for in the beginning. But here again we have the fact that motion leads to equalisation of all contrasts of the matter distribution and thus to diminishing of the negative potential energy, so that the global balance remains unchanged. In the same way as during gravitational collapse kinetic energy is produced on cost of potential energy, motion with respect to the background leads to a more homogeneous distribution of matter and thus to a diminished average of the negative gravitational potential.

Within a model, which takes a homogeneous universe as the starting point, such an effect can scarcely be described, of course. But the fact that the total energy of the universe remains constant, when the radiation field changes, becomes clear from the following consideration:

We regard the universe as an homogeneous matter filled space. The curvature of space is determined by the density of matter. If we add to this space an additional radiation field or a field of homogeneously distributed kinetic energy, without taking into account the potential energy the Einstein field equations would lead to an enhanced curvature of space. But this increase of curvature is equivalent to a stronger negative potential and thus to an increase of the nega-

tive potential energy of matter. This change of the potential energy of matter just compensates the virtual energy supply by the radiation field.

Of course, such an homogeneous creation of kinetic energy is purely hypothetical, but it illustrates, how the global energy balance remains unchanged, if a kinetic energy or radiation field is changing. In the real world such exchange between potential and kinetic energy takes place as localised fluctuations. Only when we discuss the effects which, within the global equilibrium, lead to local instabilities and thus are responsible for the formation or disintegration of structures, we have to consider this exchange of kinetic and potential energy more in detail.

For the moment it is sufficient to state that within general theory of relativity there is a clear explanation of the cosmological red shift which does not require any expansion of the universe. Though an exact theoretical analysis of the effect was not given here, it should be clear that momentum loss for every motion in curved space is a necessary result of Einstein's theory of gravitation.

Chapter 4

Black holes - the source of matter

In the last chapters we have shown that the universe as a whole must be static, if we assume that it is homogeneous on large scale, that means that no region is preferred to others by special properties of the energy tensor. The variety of cosmic phenomena, which we observe, results from local fluctuations or perturbations of homogeneity. The interplay of structure formation by attraction of masses, the tendency of kinetic energy to establish a homogeneous distribution, and finally the repulsive action of the gravitational potential lead to a continuous change of the local equilibria.

In this context it should be mentioned that according to the general theory of relativity it is not the energy or matter density alone, which is responsible for the attractive or repulsive action, but what counts is the invariant trace of the energy tensor. In the case of matter at rest this is only the energy equivalent of mass, that means, according to Einstein's renowned formula $E = mc^2$. For moving masses or for a massless radiation field we have to add the momentum part to the energy density, that means, the pressure of a gas or the radiation pressure, respectively. This amounts to one third of the energy density per degree of freedom, so that such fields count double in the balance, compared to matter at rest. Just the other way things are for potential energy. In this case the pressure is negative, so that also the trace of the energy tensor is negative. Thus the action is repulsive, and here again with twice the value compared to the energy density.

When some matter field collapses, for instance by the formation of a star or a galaxy, the process starts with the generation of kinetic energy of inward motion, which is then transformed into thermal energy by collisions and thus into gas pressure. At the same time with increasing pressure the potential energy of matter is changed, so that the balance, as defined by general relativity, remains unchanged.

The balance of a homogeneous universe is defined by the fact that the energy

equivalent of matter is in equilibrium with its own potential energy. Thus the total energy content is equivalent to zero, if we regard total energy as the trace of the energy tensor in the sense of general relativity. Locally this equilibrium is unstable, however. Any local perturbation generates energy of motion, which remains local, while the change of the gravitational potential, the change of the metric, expands into space with the speed of light and remains scarcely detectable locally.

Only in the limiting case of extremely high matter concentrations, when the density of potential energy is comparable to the density of matter energy, the repulsive action of the potential can stop the attractive force of matter. According to the prevailing concepts of gravity contraction of matter fields would continue into a singularity. A 'black hole' would form, as there exists no force which could halt the collapse. The kinetic energy, which matter takes up from the gravitational potential, would even enforce collapse. But if the repulsive action of the potential energy is taken into account, instead of a singularity under ideal conditions the final state is a stable configuration, in which potential energy and mass energy are in equilibrium. There are no 'black holes' in the sense of conventional theory, regions, in which mass is inevitably lost from the active parts of the universe.

'Black holes' must not be regarded as the end of cosmic evolution, but they are the source, from which matter is continuously regenerated in the universe. To understand this, we must look somewhat closer to their properties. Most important for the understanding of the global mass balance of the universe are the 'supermassive black holes', which are found in the centre of most galaxies. From the dynamical behaviour of the surrounding stars these appear to be very compact aggregations with million or even billion times the solar mass.

According to Newtonian theory of gravitation galaxies are stable configurations, in which angular momentum of matter together with the pressure of gas and radiation are in equilibrium with the gravitational force. Only by radiative and dissipative losses this equilibrium may be disturbed, so that finally the system collapses and more and more matter is accreted onto the central matter concentration.

According to general relativity there exists an additional loss mechanism: gravitational radiation. As has been mentioned in the last chapter, for every motion with respect to background of the universe gravitational radiation reduces momentum and, of course, also angular momentum. This leads to a continuous reduction of the rotational energy, so that on the time scale of the Hubble constant every rotational motion is changed from a circular to a spiral path, and finally all matter is collected in the centre of motion.

We must expect, however, that in the final stage of accretion the time scale becomes so short that angular momentum is conserved approximately, so that the 'black hole', which forms at the centre, will rotate rapidly. Spectroscopic observations of the accretion disks of spiral galaxies have strongly confirmed this picture of a rotating 'black hole'. Even details like the broadening of spectral lines can well be explained by the accretion model. But a fact, which was not well understood from conventional models, is the emission of intense, highly

relativistic matter jets, which are emitted in the direction of the axis of rotation of many galaxies and which can be observed along distances of many Mpc.

From dynamics of the accretion disk this mechanism cannot be explained, if we assume that the inflowing matter inevitably vanishes into a black hole. But if, as we have considered here, the final state of collapse is a configuration, in which rest energy and kinetic energy form an equilibrium with potential energy, the formation of these jets can easily be explained, and we can predict some of their properties.

In the idealised case of an exactly spherical collapse a stable final equilibrium configuration would exist, but by the continuous inflow of matter this equilibrium will be disturbed. Increase of the total amount of matter leads to an overshoot of the repulsive force: matter must be expelled again. The matter flowing inwards consists not only of distributed gas but complete stars may be sucked in, bringing with them their angular momentum and the surrounding magnetic fields. Even if the angular momenta and magnetic fields of individual stars will compensate each other by part, one must assume that after accretion of millions of stars and after the reduction to a very small volume the resulting black hole will rotate rapidly and will be surrounded by a strong magnetic field.

Charged matter particles can escape from such a configuration only from the polar region, as otherwise their motion is bound to the closed magnetic field lines. This effect gives us a simple explanation for the emission of strongly collimated matter jets.

From the mechanism of formation we can deduce some of the properties of these jets. We know that the matter is expelled from a region, in which potential energy is of the same order as the rest energy of particles. As the final stage of accretion is nearly adiabatic, the kinetic energy must be of the same order, too. The energy per particle is so high that in this region only the elementary building blocks of matter can exist, that means, protons, electrons, neutrons and neutrinos in equilibrium.

Correspondingly matter in the jet will first be composed from these particles, too. The jet starts as an equilibrium plasma with the mean energy per particle, corresponding to about the rest energy of protons, that means, an equilibrium temperature of about $10^{13}K$. During expansion first the plasma components can react with each other, but due to expansion inside the jet the plasma becomes more and more diluted, so that the equilibrium is frozen out with a composition which has been achieved by then. The jet expands into space with this composition, but keeping its high temperature.

The formation mechanism of the composition equilibrium is practically the same as adopted in big bang models to explain the synthesis of light elements. Also there one assumes that the primordial composition of light elements is fixed by freezing out from a hot, but expanding equilibrium plasma. But whereas in big bang models one tries to draw conclusions on the actual state of the universe from the singular event of elementary synthesis in the beginning, this is not possible as easily, if formation of light elements takes place continuously. They are formed at different locations from black holes with different histories, which may have different masses and may vary with respect to angular momentum

and magnetic field strength.

But there remains the fact that black holes and their jets constitute a sink of matter that has been processed in stars, and at the same time they are the source of matter of primordial composition. Thus the formation of galaxies and stars and the chemical processes during their evolution are no one way street, but there exists also an opposite effect, which closes the matter cycle of the universe.

Chapter 5

Ingredients of the intergalactic medium

In the last chapter we have shown that gravitational collapse of galaxies does not lead to singularities and to the final vanishing of matter from the balance of change in the universe, but that collapse leads to restoration of the initial state of matter, from which new cosmic structures can be formed. There exists a continuous cycle of matter from the hot plasma in the jets of collapsed systems through the formation of large structures, the fragmentation into individual galaxies or clusters of galaxies, the formation of stars with their complex chemical and physical dynamics, to the renewed gravitational collapse of stars or of complete galaxies, from which matter is recovered in its initial state.

Before we start discussing details of this cycle, first we will consider the properties and composition of the intergalactic medium, the properties of the substrate, from which new structures can emerge. It is not only the matter, expelled from black holes, which is distributed homogeneously into space. Also radiation is emitted by various processes and fills the universe. In addition large amounts of dust are produced by the explosion of stars, which by part possess so much momentum that they can leave the galaxy, in which they have been produced, and which are distributed into free space.

If the universe regarded as static on large scale there must be an equilibrium of production and annihilation, not only for the hot plasma, but also for the radiation field and the dust component. We will try to estimate, how this equilibrium will look like, and show up, how these estimations are confirmed by observations.

As a first step we will consider only those processes which determine the global properties of the intergalactic medium, leaving out all detailed structures. Only those processes are regarded, which can take place in a homogeneously distributed medium. As the discussion below will show, this is justified by the fact that the overwhelming part of matter is in such a state of nearly homogeneous distribution.

5.1 The intergalactic plasma

Let us first look at the hot plasma component. As was discussed in the last chapter, matter of primordial composition is continuously introduced by the matter jets of quasars (or 'black holes'), mainly hydrogen and helium nuclei and electrons, all with mean particle energies, which are in the order of the proton rest energy. Though these particles carry electrical charge, due to the high energy their interaction with the existing plasma is so weak that the mean free path is several times the radius of the universe. The only effective energy loss mechanism at that time is by gravitational radiation. Only during several Hubble times the energy is diminished so far that the mean free path, which decreases with the square of energy, becomes so short that a considerable energy exchange with the existing plasma takes over. This process then leads to a fast thermalisation of the plasma.

Thus we expect that the intergalactic plasma consists of a thermal component and an additional component of very high energy. The equilibrium temperature of the thermal component results from the balance of the rates, by which energy is introduced from the matter jets and the loss rate by gravitational radiation. Also after thermalisation gravitational radiation remains the dominant loss mechanism.

Only when, caused by local fluctuations of density or temperature, losses by electromagnetic bremsstrahlung become important, formation of structures will start, which finally leads to galaxies and stars. Questions of structure formation will be considered more in detail in the next chapter.

What should be stated here is that the period from the formation of matter in the hot jets of quasars to the onset of structures takes many Hubble times, while according to conventional view the lifetime of galaxies, the time from formation of a galaxy to collapse into a black hole, is in the order of one Hubble time. Thus we must expect that the dominant fraction of matter in the universe is contained in the intergalactic plasma and only a small part is in form of galaxies.

The model developed by now was only based on the theoretical considerations without regarding the question, if these ideas are confirmed by observational facts. By principle the existence of a hot plasma can be proved by its optical properties, that means, by emission, absorption, and scattering of electromagnetic radiation. But these measurements are very difficult, because there exist no characteristic structures in the plasma, so that it is scarcely possible to separate the effects from backgrounds of the measuring device.

The only extinction effect, which might be measured in a fully ionised plasma, is Compton scattering from free electrons. By this effect light from distant galaxies or quasars should appear weakened, but the effect is so small that it cannot be used to confirm or discard our model. Even for the most distant quasars, observed by now, the extinction would just be in the order of 10%, if we assume that the total density of the universe is close to that, derived from the model presented here.

A more promising method is, to detect the emission from the hot plasma, the electromagnetic bremsstrahlung. Even though this radiation does only play

a minor role in the energy balance, it is detectable reasonably well. That such a diffuse x-ray radiation background exists is well known from several measurements. But by now the proponents of the big bang cosmology interpret it as a superposition of emission from discrete sources, which cannot be spatially resolved with present day instruments. The reason is that in this model there is simply no room for emission from a continuously distributed hot plasma. But by now, in spite of large effort in this field, no discrete sources could be detected, which can contribute considerable amounts to the observed radiation in the spectral range beyond 10 keV.

From the equilibrium model presented here not only an accurate description of the observed spectrum can be derived, but also the particle density coincides well with the expected value from the calculations, applying the modified formulas of general relativity (see appendix). The thermalised fraction corresponds to a plasma with a mean energy of 100 keV and a density of $4 \cdot 10^{-30} \text{g/cm}^3$. The contribution of the high energy electrons is much lower due to the much lower collision probability, so that it is difficult to estimate from the data. It is in the order of $1 \cdot 10^{-30} \text{g/cm}^3$ to $2 \cdot 10^{-30} \text{g/cm}^3$. This estimation agrees well with the data derived from the equations of general relativity for the total matter density. With a value of 70km/sec/Mpc we get a value of $6.4 \cdot 10^{-30} \text{g/cm}^3$, that means, a similar value as that, derived from spectroscopy. This can be regarded as a strong confirmation of the model and as an hint that really the overwhelming fraction of matter is contained in the intergalactic plasma, as was expected from our considerations with respect to energy loss by gravitational radiation.

5.2 The radiation field

In the last section we have discussed production and development of material particles of the intergalactic medium. But the mechanism of the origin is quite different from all ideas which are discussed in big bang theory. For the radiation the situation is somewhat different. We know a large number of processes, which all lead to the production of photons in various energy ranges, beginning from gamma- and x-ray emission, as it is observed in stellar explosions, the UV, visible, and IR radiation from stars, and finally the radio emission in the long wavelength range of the spectrum.

All these photons expand nearly unhindered into space. If the universe has no beginning, the essential question which must be answered is: Why does the number of photons not grow to infinity? Why is the night sky dark? Of course, one part of the answer is already contained in the general principle of momentum loss by gravitational radiation. By this process all photons, which are emitted, lose energy, they become 'red shifted' more and more. From the viewpoint of energy balance this leads to a limiting range of the total energy of the radiation field. The equilibrium level is fixed by the balance between the loss by gravitational radiation and the emission processes.

But this balance, which appears evident from the viewpoint of energy, is not trivial in the world of photons. Red shift changes the energy of the photons, but

not their number. Simply looking this would lead to an 'infrared catastrophe': With decreasing energy the number of photons would exceed every limit. But it is just this 'infrared catastrophe', which already led Max Planck to his famous law of radiation and to the beginning of quantum mechanics.

Basically it is the simple reasoning that photons are wave packets. They can be regarded only as individual entities as long as the wave packets do not overlap. Otherwise interference or possible extinction has to be considered. The photon gas can no longer be regarded as a system of independent particles, but must be considered as a quantum mechanical ensemble, in which the filling of the possible energy levels is governed by their quantum mechanical probability. This considerations are valid, of course, also for the photons of the cosmic radiation field, when the wave packets of the individual photons begin to interfere.

Thus we expect that similar as for the intergalactic matter there exists a distribution of high energy photons and in addition a low energy background, to which the emitted photons deliver their energy, but with an energy distribution, which corresponds to a quantum mechanical equilibrium distribution, that means, a spectrum of black body radiation. That this background radiation exist, is known for more than 40 years, only the explanation within the big bang model is completely different from the interpretation given here.

Also in big bang theory people assume that in the beginning, prior to the formation of material particles, a radiation field existed with a Planckian energy distribution. Still after matter began to form in the expanding and cooling space the radiation field was the dominating constituent and the material particles remained in equilibrium with this field. Only with further cooling and recombination of electrons and ions into neutral atoms interactions were no longer sufficient to keep matter and radiation in equilibrium.

It remains vague, however, why the mechanism, which was responsible for keeping the photon spectrum in equilibrium prior to formation of matter, should stop to work thereafter. Why should it stop just at the recombination era, so that the inhomogeneities from this time are frozen out and can be observed still today. In our understanding of black body radiation there is no hint to an effect, which restricts the formation of equilibrium to some special region of the spectrum or energy range. If there is a mechanism which, independent from the existence and the special properties of material walls, leads to an equilibrium distribution of photons in every closed space, this mechanism should work, no matter if the cavity is the interior of the sun, an oven or a freezer, or eventually the whole universe. The only essential is the available energy density, which defines the temperature of the radiation field.

It appears scarcely reasonable that, corresponding to the big bang model, after formation of neutral matter the processes, which by then had guaranteed the radiation equilibrium, suddenly should work no longer. Especially strange is the argument that the fluctuations, which had been present at that time, should be present still today, though the universe has expanded to one thousand times its size and various structures have been formed during this time. An equilibrium without interactions must be unstable. Every perturbation will lead to irrecoverable deviations from equilibrium. Thus it is scarcely intelligible, how

one should be able to draw conclusions from the deviations from equilibrium, observed today, to the state of the universe at a much earlier time.

As has been discussed above, in our considerations we assume that the process of equilibration works continuously, but also perturbations by interaction with developing matter structures go on. Deviations from equilibrium may give us information about these structures, on their present form or due to running time of signals also about their past. But they represent no indication of any development from some big bang to the present. To the imprint, which is expected from interaction of the radiation field with the matter distribution, we will come back with the discussion of structure formation.

Here we have to discuss the question, if the mechanism of continuous red shift and the thermalisation into a nearly homogeneous radiation background of 2.73 K is reasonable. That means: Is the rate of production of photons by the various emission processes compatible with the loss rate by quantum interference?

The number of photons, which are lost from the background field can be computed rather accurately. With a temperature of 2.73 K the energy density of the radiation field is $6.3 \cdot 10^{-13}$ erg/cm³, which corresponds to a number of about 1700 photons per cm³. Taking a value of 70 km/sec/Mpc for the Hubble constant, the loss rate is $3.8 \cdot 10^{-15}$ /cm³/sec.

Estimation of the production rate is much more difficult. If all matter in the universe would radiate with the intensity of the sun, a matter density of $3.8 \cdot 10^{-15}$ /cm³/sec would lead only to a production rate $1.6 \cdot 10^{-17}$ /cm³/sec. But we must keep in mind that there are many sources of radiation, which are less spectacular than stars, but which radiate much more effectively in relation to their mass. A dust grain of 1 μ m radius in relation to its mass emits 50 times more energy than the sun, even if its temperature is only 3 K, if both are regarded as black body radiators. The number of emitted photons is higher even by a factor of 10^5 due to the lower photon energy.

This demonstrates the difficulty to get a reasonable estimation of the production rate of photons, as we have no exact knowledge of these low energy processes like the emission of radiation from dust grains. It is just the existence and distribution of dust in the intergalactic medium, of which we have only a very vague knowledge. By principle we know several mechanisms, which lead to the production of dust and discuss this topic more in detail in the next section. But with respect to the radiation balance we must restrict our considerations to the fact that at least from the order of magnitude it appears reasonable that the diffuse background radiation field is determined by the mechanism described here.

5.3 Cosmic dust

That space between the stars of galaxies is not empty, but filled with gas and dust particles, is well known since long ago, and various mechanisms of dust production appear well confirmed. Apparently the main source of dust are stellar

explosions, by which atoms of heavy elements, which have been formed by nuclear fusion in the interior of stars or during explosion, are expelled with the blast wave. During cooling on their way out they first form stable molecules, which then coagulate into larger clusters and finally into macroscopic dust grains.

The composition of the dust grains may be rather different depending on the history of the explosion. It depends on the ratio of carbon and oxygen in the expanding cloud and on the range, to which the formation of heavy elements like silicon, magnesium or iron has proceeded by fusion reactions. Also the size distribution of dust grains may differ dependent on the development of density and temperature within the blast wave. Besides the dust content of the medium, into which the cloud expands, may be different, supplying different amounts of nucleation seed. Homogeneous nucleation requires much higher supersaturation than attachment of molecules to an existing seed.

The expansion velocity of the clouds may be very different. The highest speeds are observed in supernova explosions of type Ia. They are in the range of 10^4 km/sec, that means well beyond the local escape velocity of their galaxy. Dust grains which are formed within a cloud at such high velocity, would thus generally leave their parent galaxy, if they would move unhindered into vacuum. But as the interstellar space is not empty, the dust grains will be slowed down by collisions with gas atoms. The relative loss of momentum is the higher, the smaller the grains. Only the largest grains can really leave their galaxy. An estimation of the minimum size (see appendix) shows that for a supernova, which explodes in an environment similar as that of the sun, the limiting size of the dust grains is in the order of $1\mu\text{m}$. That dust grains of this size can be formed in the blast wave of a supernova also results from simple estimations of growth rates under the conditions inside the cloud (see appendix).

Supernova explosions represent a selective production mechanism, by which large dust grains of the order $1\mu\text{m}$ are emitted into the intergalactic medium, while smaller grains remain within the galaxy, serving there as nucleation seed for later generations of explosions. The existence of these small dust particles in galaxies can easily be proved by their selective absorption properties. The light of stars, which transverses a layer of dust, appears reddened compared to that of stars without absorption.

The existence of dust grains of the order $1\mu\text{m}$, as we expect them in the intergalactic medium, is more difficult to establish, as there is practically no selective absorption of visible light. Only a general lowering of the intensity of light from distant sources can be expected. To measure this absorption, however, it is necessary to know the absolute intensity of the source. Today the favoured standard candles for this purpose are supernovas of type Ia. On one hand they have extremely high intensity, so that they can be observed over very long distances, on the other hand observations of supernovas in nearby galaxies have shown that their light curve follows a universal pattern within close limits.

As has been confirmed by measurements of several research groups, indeed the light of very distant supernova explosions appears considerably dimmer than would be expected from their distance, determined by red shift. The adherents of the big bang theory, however, ascribe this to an accelerated expansion of the

universe which, according to their view, has to be interpreted as the action of some energy field with repulsive gravitation, the so called 'dark energy', but without any physical explanation of its origin. The fraction of this dark energy should amount to about 70% of the total energy.

Interpreting the attenuation of the light from distant supernovas as absorption or scattering by dust grains, the observed facts can be explained without any unknown energy component. Sometimes it has been argued against this interpretation that for very distant supernovas (red shift $z > 1$) the extinction is less than would be expected from a pure absorption model. But this is true only, if space is considered as flat. In curved space the effective angle of observation of very distant objects appears enlarged, compared to Euclidean space. Considering both effects, absorption and curvature of space, the observed data can be explained very accurately (see appendix).

To estimate the amount of dust in the intergalactic medium, one had to know the size distribution of the grains. If we assume that the dominant species are spherical silicate grains of about $1\mu\text{m}$, the observed absorption would correspond to a mass density of about $2 \cdot 10^{-32}\text{g/cm}^3$, that means, 0.3% of the total mass density, as it was estimated at the beginning of this chapter.

This value appears reasonable according to the observed number of supernova explosions. As estimated from observations in neighbouring galaxies, in a galaxy like ours there should be one explosion every thirty years. During the life time of a galaxy, which we estimate in the order of one Hubble time, a few percent of matter are processed by supernova explosions, while the main fraction ends up in a central black hole.

If we further assume that a few percent of the stellar mass in an explosion is transformed into large grains, which can leave their parent galaxy, a dust fraction of 0.3% in the intergalactic medium appears quite reasonable.

Due to the high initial velocity of the explosion dust cloud in the order of 10^4 km/sec, after leaving the galaxy, the dust grains will be slowed down by collisions with the intergalactic plasma only slightly, so that they can spread uniformly into space. Even the largest voids with extensions of 100Mpc can be traversed within less than the Hubble time.

Thus it is reasonable to assume that the intergalactic medium, which is the substratum for the formation of all structures, consists, apart from a radiation field, not only of a hot plasma, but of a mixture of fully ionised gas and of a dust component. Only by the combined action of these components the observed structures can be understood.

Chapter 6

Structure formation

In the last chapter we have discussed the composition of the intergalactic medium under the proposition that the medium is distributed uniformly in space. Certainly this appears satisfied for the high energy plasma component, as the interaction probability with other components is so weak that the mean free path is several times the radius of the universe.

But for the thermalised component of the plasma this assumption is no longer valid. The mean free path is still in the range of 100 Mpc, but it is small compared to the radius of the universe. Thus we must expect that local perturbations of homogeneity occur. This can lead to instabilities, by which instead of a uniform distribution some other more stable configuration develops.

6.1 Gravitational instability

Already more than one hundred years ago James Jeans has shown that perturbations of the equilibrium between pressure and gravitational forces can lead to instability, if the length scale of the perturbations is larger than some critical value, which results from the balance equations (see appendix). It should be mentioned here that the assumptions, which Jeans used to derive his stability criterion, were - long before Einstein developed his general theory of relativity - in contradiction to the accepted theory of that time. Jeans assumed that in a homogeneous mass filled space, in contradiction to the Poisson equation of Newtonian theory, there exists a constant gravitational potential, and that perturbations of the homogeneous matter distribution lead to perturbations of this potential and thus to motion in the matter field. Perturbations of small wavelength lead to periodical oscillations, but if the wavelength exceeds a certain limit, they grow exponentially. The minimum wavelength which leads to instability, the so called 'Jeans length' appears as a useful resource in the discussion of gravitative instabilities.

The assumption of a constant potential in matter filled space, which appeared as a deficit of the Jeans theory and was denoted as 'Jeans swindle'

frequently at his time, appears quite natural within the scope of general relativity and as an anticipation of the correct description. According to general relativity a homogeneous matter filled universe has to be considered as a space of constant curvature. The amount of curvature is a measure of the gravitational potential. Local perturbations of the matter distribution lead to distortions of curvature, which are regarded as perturbations of the gravitational potential. Thus the Jeans theory is just the correct description by general relativity in the quasi-Newtonian approximation.

But in spite of that, the Jeans model is only of limited value for the description of cosmic structures. It gives some hints to the conditions, under which instabilities can occur. But as it is only a linear description of small perturbations, it cannot give answers to the question, which structures are formed, when the instabilities grow and if there exists some new equilibrium state, into which they develop.

When we pose the question, which equilibrium distribution develops, when the homogeneous distribution becomes unstable, it appears more reasonable to presuppose the existence of instabilities and consider directly the properties of the final new equilibrium state.

In the second chapter we have discussed the thought experiment of the formation of a 'vacuum bubble', the formation of a void within a otherwise homogeneous medium. We have stated there that the formation of such vacuum bubbles is not observed under terrestrial conditions, as pressure forces due to thermal motion of atoms exceed gravitational forces by many orders of magnitude. Under cosmic conditions this is no longer true, however. The Jeans criterion shows that the balance can be in favour of gravitational forces.

Within the linear ansatz of the Jeans model positive and negative deviations from equilibrium and thus enhancement and reduction of matter density have equal probability. But a more detailed consideration shows that the formation of voids is much more probable. With every enhancement of density the potential energy, which is set free, is applied to the interior in form of kinetic energy leading to an increase of pressure, which counteracts any further enhancement. Density reduction, on contrary, leads to transport of energy to the outside, where the energy can be distributed into a much larger volume. Thus the growth of bubbles can proceed unhindered.

Thus we must expect that gravitational instability leads to formation of bubbles or voids. This development will stop only, when neighbouring bubbles begin to influence each other, that means, when the pressure between the bubbles increases considerably. In equilibrium the intergalactic plasma will obtain a foamy structure with large voids, separated by filamentary structures of higher density.

To estimate the size of structures, which are formed by this process, we can again look at the balance between pressure and gravitational forces. A bubble can be regarded as stable, if the mean kinetic energy of particles at the boundary of the bubble is no longer sufficient to traverse the bubble against the rising gravitational potential. Assuming as an idealised model a spherical vacuum bubble within a homogeneous environment with properties, as we have

discussed for the intergalactic plasma in the previous chapter, that means, a density of $4 \cdot 10^{-30} \text{g/cm}^3$ and a mean particle energy of 100 keV, the size of the bubbles is of the order 150 Mpc.

To obtain a more accurate estimation one has to regard the fact, of course, that the transition between voids and matter filled space is continuous and not stepwise and that the density in the matter filled filaments is higher than we have estimated from the diffuse x-ray intensity under the assumption of a homogeneous density distribution. A detailed description would require extensive numerical modeling. Here we will confine ourselves to this crude estimation, but discuss, how this estimation is confirmed by observation.

Direct spatially resolved measurements of the x-ray background with sufficient accuracy are not available at present, by which inhomogeneities of the intergalactic plasma could be made visible. But we can expect that with the further development of cosmic x-ray diagnostics this may become possible in the near future.

The distribution of visible matter, especially the distribution of galaxies can be regarded, of course, also as an indication of the inhomogeneous distribution of matter. As we will discuss more in detail in the next section, galaxies will form only in the regions of the highest matter density. Thus the distribution of galaxies can be regarded as kind of an image of the total matter distribution. Indeed this image of the sky shows a cellular structure with large voids. Most galaxies concentrate on a filamentary structure separated by large voids in the order of 100 Mpc, just in the range predicted by our model.

Also the microwave background can give us some hints to the structure of the intergalactic plasma. Though the radiation field due to quantum interference always tends to the equilibrium distribution of a black body spectrum, by interaction with the electrically charged particles of the intergalactic plasma it is continuously disturbed. By inelastic collisions (Compton scattering) kinetic energy is transferred to the photon field. The effective temperature of the microwave radiation thus appears a little bit enhanced after passing through a matter concentration. According to the big bang theory this mechanism, well known as Sunyaev-Zel'dovich effect, is of minor importance. Only the high plasma concentrations in galaxy clusters should exhibit an observable effect. The existence of an intergalactic plasma, which contains the overwhelming fraction of the total mass of the universe, changes this situation, of course.

The probability that a photon is scattered by an electron is not very high. Under the proposition that the size and density of the universe correspond to the values discussed here the probability that a photon is scattered during surrounding the total universe one time is about 20%. But by now the microwave background has been measured with such high accuracy that even inhomogeneities in the order of 10^{-5} can be detected. It is customary to express these inhomogeneities by evaluation of the correlation function with respect to spherical harmonics. The strongest deviation from an homogeneous distribution are found at an angular correlation of $2\pi/220$. Applying a radius of the universe of 4200 Mpc, as it results from the model presented here, this corresponds to a correlation length of 100 Mpc. This is just the same order as the diameter of

the vacuum bubbles, as we have estimated from our simple stability model.

Thus it appears reasonable to assume that the observed intensity fluctuations of the microwave background are an indication of the inhomogeneous distribution of the intergalactic plasma. The explanation, given by the proponents of the big bang model, appear somewhat questionable in comparison. According to this model the fluctuations should be formed at a very early time, when the radiation field was still coupled to the matter field. But the source of the fluctuations remains unclear.

6.2 Thermal instability

From the process of structure formation by gravitational instability, discussed in the last section, we can understand the large scale distribution of matter, but the balance of loss by gravitational radiation and the introduction of energy remains unchanged by these structures, as this balance is nearly independent from the distribution of matter.

But with increasing density an additional loss mechanism occurs: electromagnetic bremsstrahlung. For a plasma with density and temperature corresponding to the mean values of the intergalactic gas this loss amounts only to a small fraction of the loss by gravitational radiation (see appendix), but it has the effect that regions, in which the density was somewhat higher than the mean, cool off faster compared to those of lower density.

By this mechanism the pressure in the denser regions is reduced, so that more gas can flow into these regions of overdensity. The increase of density further enhances the radiation loss, so that the pressure and temperature fluctuations are increased. As in this case the consequence of increasing density is decreasing pressure, contrary to the gravitational instability, now local concentration is preferred compared to density reduction. Thus inside the denser filaments of the gravitationally formed foamy over-structure a clumpy structure is generated by thermal instability.

The size of the overdense regions may be very different in this case. It depends on the fact, if they develop in the narrow wall region between two vacuum bubbles or in the vertex region with more than two adjacent bubbles. The temporal development of the clumps is determined by the fact that not only the thermal energy of the inflowing gas has to be radiated away, but also the temperature enhancement must be compensated, which is generated by transformation of potential to kinetic energy.

Due to the opposite effects of pressure enhancement by concentration and pressure decrease by radiation the formation of this structure is a slow process. It is not an instability in the actual sense. The gas is always close to thermodynamic equilibrium, but tries to adapt to the slowly changing conditions. As long as the density fluctuations are on a length scale, which is small compared to that of gravitational instability, the influence of gravitational forces is weak compared to gas kinetics. Changes are nearly isobaric. That means, the density increases in the same manner as the pressure drops by radiation loss.

To maintain constant pressure there must be a constant inflow of matter, however. This mechanism can work, of course, only if sufficient matter is available and if the inflow does not require a higher velocity than that of free fall.

At the beginning of contraction these conditions are always satisfied. But with increasing density the energy loss is accelerated, so that the time constant of free fall may exceed that of radiative cooling. When the inflow of matter is no longer sufficient to sustain pressure equilibrium, the system becomes unstable. The plasma clouds are disrupted into smaller pieces. This process of fragmentation is again determined by the balance of pressure and gravitational force. But contrary to the large scale formation of voids in this case the formation of matter concentrations is preferred, as also here the pressure decreases with rising density.

The individual matter clumps will further shrink inside the intergalactic plasma, while the cooling goes on. As the individual parts are in motion during fragmentation, they will remain in motion against each other afterwards, while their interior development proceeds autonomously. The different velocities and the mutual gravitational attraction may lead, however, to collisions of individual clouds. They may melt together again, or may gain angular momentum by the action of tidal forces.

6.3 Formation of galaxies

At the end of the last section we have discussed the mechanism, which leads to fragmentation of cooling plasma clouds. As long as the dominant energy loss rate is electromagnetic bremsstrahlung, this process proceeds rather slowly. But as soon as the plasma has cooled so far that recombination into neutral atoms begins, that means, at mean particle energies in the order of 10 eV, the cooling process by recombination radiation and line radiation runs so fast that a massive fragmentation starts.

To get an idea, how large the individual fragments will be, we can again take advantage of the Jeans' stability criterion. Also this fragmentation is controlled by the balance of pressure and gravitational forces. But now enhancement of density does not produce a pressure increase, as this is compensated by increasing radiation loss. Again overdense regions are preferred to the formation of bubbles. Under the assumption that fragmentation starts at about 10 eV mean energy and that up to this point the development has been isobaric, that means, the product of temperature and density has not changed much, the stability limit of fragmentation can be estimated. Based on the density and temperature values, which have been derived from the diffuse x-ray background (energy 100 keV, density $4 \cdot 10^{-30} \text{g/cm}^3$, see appendix) the length scale resulting from the Jeans criterion is 1.7 Mpc, which corresponds to the typical distances between galaxies. The total mass contained inside a volume of this size with $2.5 \cdot 10^{43} \text{g}$ is about that of a small galaxy.

This is only a very crude estimation, of course, as it is based on the assumption of constant density and temperature inside the cloud. Besides an

undisturbed evaluation of the medium has been assumed from the formation of the primary structure up to the beginning of recombination. but the estimation shows that gravitational instability together with radiative cooling leads to the formation of plasma clouds, which can be regarded as the progenitors of galaxies.

From the existence of a stability limit it can be understood, why galaxies cannot have arbitrary size. 'Thermal instability' is independent of the size of the underlying region and there can be matter concentrations of very different size. But the instability, which occurs with the beginning recombination, leads to fragmentation into individual clouds with a rather narrow size distribution.

By this mechanism we can understand that galaxies can develop as individuals, in groups, or even in clusters with thousands of members, but that the size of the galaxies is similar in all cases.

What we have denoted as galaxies are only the progenitors of the strongly radiating objects which we observe in the sky, of course. Even recombination and line radiation is too weak to be observed from very distant objects. Besides the time span of recombination is only a short episode compared to the total development of a galaxy, so that the chance is low to catch one just in this stage. While the time span from injection of high energy particles into the intergalactic medium to the formation of a thermal plasma is a multiple of the Hubble time and the cooling phase, until recombination starts, is in the order of one Hubble time, recombination is terminated within a small fraction of this time.

With the end of recombination also the emission of electromagnetic radiation comes to an end nearly completely. In the beginning there is a rest from vibrational transitions of hydrogen molecules, but also this mechanism can only slowly reduce the temperature. The existence of clouds, formed from neutral hydrogen, is well known from the observation of the absorption of radiation of background sources (mainly quasars) in the spectral range of the hydrogen resonant lines (the Lyman-alpha forest). Proponents of the big bang theory are still searching for an explanation, how such a cloud of neutral hydrogen and helium can develop into a galaxy.

The model described here delivers a plausible explanation, if we take into account that the intergalactic plasma besides of ions and electrons contains considerable amounts of dust grains. During the concentration phase of the hot plasma the motion of these grains follows the contraction of the plasma clouds. Collisions transfer energy to the dust, which is then emitted as infrared radiation from the surface. The temperature of the grains remains so low, however, that even in a very hot environment evaporation does scarcely occur (see appendix).

In the hot plasma radiation from dust grains is negligible in the energy balance. But this changes at the end of the recombination phase, when gravitational radiation is the only competing process. Under these conditions radiation from dust grains becomes the dominant loss mechanism, which is effective even at very low temperatures, as long as this is higher than that of the background radiation field.

This cooling is accompanied by a second effect, for which the dust grains are responsible, too, and which is essential for the further development of galaxies. In

a hot plasma the collision rate between gas atoms and dust grains is so high that the dust grains, apart from a slow Brownian motion, follow the general motion of the plasma. With decreasing temperature collisions with gas atoms become less frequent, so that dust grains can practically move like free particles. They feel their mutual gravitational attraction, forming a nearly pressureless gas, which again obeys the stability criteria of gravitation. Dust grains coagulate into larger clusters and then into clouds of increasing size, a process, as it is assumed in a similar way in big bang theories for the dark matter ('hierarchical clustering'). Only when the dust clouds have reached a sufficient size, they can attract also more and more of the surrounding gas and thus initiate the formation of stars.

Here we will not go into the details of the development of stars, but only emphasise the differences, compared to big bang theory. The primary difference is related to the question, how the first stars in a galaxy can be formed. According to big bang theory the medium, from which the galaxies formed, consisted only of hydrogen and helium. But there is no accepted theory, which can describe, how only from these gases condensations can be formed, which finally develop into stars. Also all observations show that regions, which exhibit strong formation of stars, contain much dust. For stars, which form in later development stages of galaxies, it is not a problem to explain the presence of dust by emission from explosions of earlier generation stars. But for the first generation of stars this explanation cannot be used.

According to the ideas of our model, which are confirmed by the absorption of light from distant supernova explosions, the gas, from which galaxies are made, already contains dust with a mass fraction of about 0.3%, which, as described above, coagulates and is available as condensation seed for the first generation of stars. An essential difference between the formation of the first and later generation of stars is, that in the beginning dust is present in form of larger grains. Later also the gas contains the elements, which have been formed in stellar explosions (the 'metals') as well in form of individual atoms and very small grains.

While in the formation of the first stars first the dust grains coagulate and the actual star of hydrogen and helium forms around this kernel, the gas of later generation stars is enriched with metals, so that the characteristic radiation of these elements is visible in the spectrum.

The observed poor metallicity of the oldest stars in the galaxy is not an indication of little metal content and thus for a formation from a pure hydrogen and helium atmosphere. The metals are only hidden in the inner kernel, which has been surrounded by a metal poor envelop.

There is no reason to assume that the observed low metal content of old stars is an indication of some general development of the universe with time, as it is claimed by big bang theory. On contrary the steadily improving observations of the last years point to the fact that even very far galaxies possess a metal content similar to the nearby ones. This shows that also these galaxies apparently contain young and old stars as well.

6.4 The life of galaxies

While the formation of galaxies appears as a rather straight-line process, their further development is determined by a large number of different processes running on different time scales. In addition the relative motion of galaxies with respect to each other may lead to collisions, by which they may merge or at least exchange part of their matter or angular momentum. The initial conditions for the further development may vary considerably. Varying angular momentum of the protogalaxy apparently is one of the most significant properties, which on a longer time scale determines the different modes of development. On the angular momentum it depends, if the matter distribution remains more or less elliptical, in which the gravitational equilibrium is sustained mainly by the proper motion of stars, or if the contraction perpendicular to the axis of rotation is slowed down by the centrifugal force, so that a flat disk of a spiral galaxy is formed.

Most of the processes, which occur during the life of a galaxy or a cluster of galaxies, for instance the formation of stars, their development and their different stages of life, and finally their final fate by explosion or as a neutron star or white dwarf, do not differ much in the model presented here from what is generally believed. We will not go into the details of these processes, but only highlight a few points, where the interpretation of observed phenomena is different.

In the last chapter one essential difference has already been mentioned: the existence of dust in the intergalactic medium, which enables the formation of the first stars in a young galaxy.

The second important difference is the continuous new production of hot plasma, by which the development of a galaxy or a galaxy cluster cannot be regarded as completely isolated from the rest of the universe. Hot gas can continuously flow in from outside or high energy particles can be captured.

The third difference is that gravitational radiation has to be considered with the consequence that on the scale of the Hubble time there exists practically no stationary motion. All motions relative to the cosmic 'inertial system' must slow down. Kinetic energy is reduced on account of the gravitational potential.

6.5 Galaxy clusters

Let us first look at the hot plasma in clusters of galaxies. The diffuse emission of x-rays from the space between galaxies shows that this space must be filled with hot plasma. The emitted spectra can be explained as bremsstrahlung from a fully ionised plasma with a kinetic temperature in the order of 1 to 10 keV and particle densities up to 10^{-2}cm^{-3} . According to big bang theory this was initially cold gas, which has gained energy while flowing into an existing dip of the gravitational potential. But as also metal lines are observed in the spectrum (the metallicity is of the same order as in the solar spectrum), a large fraction of this gas should already have passed a cycle of star formation and supernova

explosion. But the observation of very far clusters shows that this state must have been reached very shortly after the big bang. With increasing knowledge of the details of distant clusters this interpretation becomes more and more improbable.

If we assume, on contrary, that the universe is all over filled with an intergalactic plasma and that this plasma is regenerated continuously, it can easily be understood that during and after the formation of a galaxy hot matter streams in from outside. By the increasing concentration the energy loss by radiation increases, so that not only the energy gain from the gravitational potential is compensated, but that there is an additional cooling compared to the intergalactic plasma.

Also the observed high metallicity of the cluster plasma can easily be understood. As has been discussed in the last chapter, the metallicity of the intergalactic plasma is high, even if the metals cannot be observed, as they are bound in larger dust grains. But in the cluster gas, where the density is higher by a factor of 1000, the erosion of the grains is strong (see appendix), so that spectral lines of metals can be observed in the gas, too.

From the intensity of the bremsstrahlung we can try to estimate the matter density of the cluster gas. But if we assume that the plasma is an ideal gas in thermodynamic equilibrium, the estimated matter density is far too low to keep the system of galaxies and plasma in balance. The observed velocities of proper motion of the galaxies are far too high to constitute a gravitationally bound system. A five to ten times higher total mass would be required to avoid exceeding the escape velocity.

According to the big bang theory the problem is solved by the assumption, that the overwhelming part of the cluster mass is present in form of so called 'cold dark matter (CDM)', some stuff which interacts with other matter only by gravitation, but otherwise remains completely undetectable. But apart from the fact that up to now there exists no plausible idea, what this 'dark matter' could be, also the overall picture is not consistent in itself.

From model calculations within the CDM model one finds that the density of a matter cloud, which is subject only to gravitational forces, in a spherical cluster should decrease radially as $1/r^3$. But the observed intensity profiles of x-ray emission follow, apart from a narrow central region, a rather exact power law all over the observed range. But for a gas in thermodynamic equilibrium such a profile is compatible only with a matter distribution, which decreases as $1/r^2$, unless the temperature of the gas changes extremely (see appendix).

Also our model does not immediately give an explanation for the 'missing mass', but we can find another interpretation of the observed facts, which makes the search for CDM unnecessary.

Let us start from the assumption that a galaxy cluster consists only of the galaxies and the intracluster gas, which has been concentrated and cooled from the intergalactic plasma. We find immediately that the radial profile of the cluster gas, which results from a thermodynamic equilibrium model, exactly agrees with the profile derived from x-ray emission, if we assume that the plasma temperature shows a slight increase from the centre to the outer region (see

appendix). This increase appears plausible, considering the fact that the cluster is embedded into the much hotter intergalactic medium. An increase of the temperature is observationally found for practically all clusters at least for the inner region.

In the outer regions the various measurements differ considerably. Some authors find constant or decreasing temperatures, others find increasing temperatures. But the error margin of these measurements is very large, as most of them are carried out in the spectral range $< 10\text{keV}$, where the dependence of the intensity distribution on the temperature distribution is weak. In addition the sensitivity of the detectors decreases strongly with increasing energy. For this reason it is scarcely possible with a signal, which consists only of a few photons, to subtract exactly the background resulting from the intergalactic plasma, when this background is higher than the cluster radiation itself. Already an error of 10% of the background correction may change an increasing temperature profile into a decreasing one.

But what is the reason for the discrepancy between the densities determined from the x-ray emission and those, which result from gravitational equilibrium relations? The main reason is the assumption that the cluster plasma can be described using the thermodynamic equilibrium relations of an ideal gas. Reality is quite different from that. Observations with high spatial resolution of clusters in our 'neighbourhood' (e.g. the Coma cluster) show strong density and temperature fluctuation on a length scale of 1 to 10 kpc. This is an indication that the intracluster plasma is not homogeneous, but exhibits a cloudy structure. On the other hand the gas is permeated by turbulent magnetic fields, which change the transport mechanisms.

That the cluster plasma cannot be distributed homogeneously is immediately intelligible, considering the fact that galaxies move through the gas with high velocities and, according to their gravitational attraction, drag with them part of the gas, until by interaction with the next mass concentration, parts of the cloud are separated again. With time this mechanism of continuous drag and separation generates a turbulent system of gas clouds, which move against each other with similar velocities as that of the galaxies. That means that a large fraction of the kinetic energy of the gas is not present as thermal energy of individual particles, but as directed motion of larger clouds.

Directed motion within a cloud does not contribute to collisions and thus to emission of bremsstrahlung. What we observe, is the radiation, which is emitted, when plasma clouds traverse each other. The energy spectrum of the observed radiation in this case is determined by the relative speed of the clouds. If this motion is directly coupled to the motion of galaxies, it becomes clear, why we observe a rather exact proportionality between the velocity dispersion of galaxies and the temperature of the x-ray emission of the cluster gas. But as bremsstrahlung is emitted only, if gas clouds traverse each other, we can understand, why the intensity of x-ray emission is lower than in a homogeneous plasma. At present our knowledge of the details of this mechanism is too poor to make exact statements on the reduction factor.

The second point, as can be supposed, is immediately connected to the first

one. Detailed measurement on radio emission of clusters indicate that they are permeated by magnetic fields. The strength of these fields is in the range of 1 to 10 μG , and they change direction and strength on different length scales from 1kpc to 100 kpc. Only part of these fields is correlated to the motion of galaxies. For the small scale changes it is supposed that they are generated directly inside the plasma. Regarding the fact that the plasma is not at rest but in turbulent motion of individual clouds, the generation of magnetic fields appears as a natural consequence.

Even if the strength of the magnetic fields is only in the range of μG , it has a strong influence on the transport properties in the plasma. The fact that charged particles can move freely only in the direction of the field, while in perpendicular direction they gyrate around the field lines, transport in perpendicular direction is strongly reduced, compared to an ideal gas. Not only thermal conduction, the energy transport by electrons, is reduced, but also momentum transport by ions takes place practically only in direction of the field lines. Pressure gets a tensorial character with different components parallel and perpendicular to the field lines. In a turbulent plasma with statistical distribution of the field direction the mean of pressure gradients is reduced by a factor of three, compared to a plasma without strong fields.

That the effect of magnetic fields is so strong, we can easily get into consciousness, comparing the cluster plasma with the magnetised plasmas, which are used on earth in nuclear fusion experiments. The temperatures of fusion plasmas are similar as those of the cluster gas. The magnetic fields used for confinement are higher by about ten orders of magnitude, but also the particle densities are higher by about twelve orders. Deviation from isotropic behaviour are thus expected to be much higher in the cluster plasma than in fusion plasmas. The radius of gyration is smaller than the mean free path by many orders of magnitude for electrons and ions as well.

For the thermodynamic equilibrium in galaxy clusters that means that the effective pressure gradients are lower by a factor of three, compared to the values one would expect in an ideal gas. Thus the equilibrium between pressure gradients and gravitational forces requires only one third of the mass.

Together with the reduction of x-ray emission due to the partly ordered macroscopic motion of plasma clouds the reduction of the required mass density, compared to an ideal gas in thermodynamic equilibrium, allows an explanation of the observations without the necessity of a dark matter component.

We should emphasise here that the reduced density in the central part of clusters does not imply that the total mass is smaller, too. The definition of a 'total mass' is somewhat questionable, anyway, as clusters are not isolated system in a vacuum environment, but are embedded into the intergalactic plasma. Compared to the model calculations, by which the big bang theory tries to ascribe the matter distribution to some 'dark matter', the density profiles according to our model are much shallower ($1/r^2$ instead of $1/r^3$ for large r). Thus the aberration effects of light from distant background sources (gravitational lensing), which are based on the integrated mass distribution along the line of sight, may give similar results for both theories, even if the density in

the centre of clusters is strongly different.

6.6 Spiral galaxies

Dependent on the size of the initial matter concentrations, which are caused by gravitational instability of the intergalactic plasma, galaxies may be formed as individuals, in groups or in large clusters. But to enable the transformation of a matter concentration into a galaxy with luminous stars, according to our knowledge first individual parts must cool down well below the ionisation limit, so that again on a smaller scale gravitational forces can exceed the pressure forces and the cold matter can contract into individual stars. During this process individual protostars or groups of stars may develop proper motions, which finally lead to a stable state of the galaxy and prevent the matter from collapsing immediately into the centre of the concentration. The stars form kind of a gas, in which the proper motion can be regarded as a pressure, which counteracts gravitational attraction.

But this simple picture is changed, if the protogalaxy possesses a reasonable amount of angular momentum. By the centrifugal force the contraction perpendicular to the axis of rotation is hindered and the matter cloud develops into a more or less flat disc, in which equilibrium is maintained more by the balance of gravitation and centrifugal force than by the proper motion, the 'pressure force' of stars. Only near the centre the pressure force dominates. Thus a spherical kernel, the 'bulge', is formed, surrounded by a flat disc, in which again regions of higher and lower star density develop due to gravitational attraction. The matter cloud develops into the typical image of a spiral galaxy.

But why it is a spiral? If we look at other systems, which are stabilised by the balance of centrifugal force and gravitation, like the motion of planets around the sun or the rotation of moons or dust belts around the planets, we always see that the orbits are approximately circular. This is easy to understand, as circular orbits are the most stable ones against perturbations by neighbouring masses. Why is the picture different with rotating galaxies?

The answer is in the nature of gravity. As has been explained in the first chapters, every motion in the universe feels a retardation by emission of gravitational radiation due to the finite velocity of gravitational interaction, the back reaction of the motion on the metric of space. On the time scale of the Hubble time there exists no true stationary motion. Also with respect to angular momentum there is no 'conservative system' except of the total universe. Only as the orbital time of the planets or moons is very short compared to the Hubble time, we do not recognize the change of angular momentum. The effect would by principle be measurable within the lunar ranging program, which allows to measure the orbit of the moon with centimetre accuracy. But there are other perturbations of the orbit, which are not known with sufficient accuracy to separate them from possible gravitational retardation.

The orbiting time of galaxies is not so short, however, that the loss of angular momentum during on revolution could be neglected. For the sun the loss of

angular momentum in one revolution about the centre of the Milky Way is about 2%. As a consequence circular motion is replaced by spiral motion.

How strong the spiral structure develops, depends on the initial amount of angular momentum, which is transferred to the system at the time of fragmentation. Of course angular momentum can be changed also later by tidal interaction or collision with other galaxies. Thus the range of different galaxy types is very broad, ranging from ellipticals with very low angular momentum to spirals with very extended arms. Formation of new stars takes place mainly in the arms, while old stars gather in the central bulge. But on the time scale of the Hubble time all stars move in the direction of the centre on spiral orbits.

Also within the disc the formation of stars is not homogeneous. Gravitational attraction leads to concentration in individual filaments with higher density of stars with dark spaces between them. Interaction between the stars leads to proper motion of stars within the spiral arms, but the velocity remains low compared to the rotational motion of the disk. Only when the spiral motion approaches the central bulge, the density of stars becomes so high that the mutual interaction changes the rotational motion more into an isotropic distribution, as mentioned at the beginning of this section. The total angular momentum can change, however, only to the extent of loss by gravitational radiation.

6.7 Active galaxies and quasars

What happens, when more and more matter concentrates in the centre of a galaxy, we have already shortly discussed in the second chapter. As the repulsive force of the potential energy with its negative pressure term becomes important, contrary to the ideas of the big bang model, matter does not collapse into a 'black hole' in the sense of a singularity. Under ideal conditions instead an equilibrium configuration would occur, in which the gravitational attraction of matter is in balance with the repulsive action of its own potential energy.

But in reality we cannot expect that the transition into this equilibrium state is completely symmetrical. Normally the matter flowing into the core has a certain amount of angular momentum, so that we must expect that the 'black hole', as we will call it here, too, will rotate very rapidly. With increasing concentration the rotation slows down due to increasing gravitational radiation, as in addition to gravitational bremsstrahlung also quadrupole radiation becomes important. But we must expect that the increase of angular momentum from the inflow of matter exceeds this loss by far, so that the result is a rapidly spinning object.

Also the magnetic fields, which are coupled to the infalling stars and gas clouds, will compensate only by part, so that, if magnetic flux is conserved, the high concentration within a very small volume will lead to extremely high field strength.

With respect to the accretion of matter from the surrounding the scenery in our view does not differ much from conventional theory, only that it contains an additional simple explanation for the loss of angular momentum, which is

necessary to understand, why matter really falls into the centre of mass. The conventional theory assumes that there exists some energy dissipation by internal friction, that means, that there is a kind of viscosity of galactic matter, which slows down the rotation. Of course, this possibility exists, but the loss of angular momentum by gravitational radiation is sufficient to explain the accretion.

An essential difference to conventional models occurs only in those regions, where the gravitational action of potential energy is comparable to that of the matter itself. In the idealised case of a spherical matter distribution this state would be reached at the Schwarzschild radius. The real collapse in the centre of a galaxy is more complex, however. Due to rotation and radial motion of matter towards the centre and in addition also by the energy of the magnetic field the balance between attractive and repulsive forces is far from spherical symmetry.

While in the plane of rotation due to the kinetic pressure of inflowing matter may preponderate, at the same time in the direction of the rotation axis the repulsive action of the negative pressure from potential energy may already dominate. The dynamic equilibrium in this case is characterised by a continuous inflow near the plane of rotation and a simultaneous outflow along the rotational axis. But all the matter goes through an intermediate state, in which the potential energy is of the same order of magnitude as the equivalent energy of its rest mass. During the process of accretion and ejection the energy loss by radiation is negligible. The process can be regarded as adiabatic. Thus the amount of kinetic energy will attain the same order as potential energy.

The intermediate state within the 'black hole' is characterised by a mean energy of the heavy particles or a 'temperature' corresponding to their rest mass energy. This energy is far beyond the binding energy in the atoms, so that the identity of individual atomic species is completely lost. The state of matter is comparable to that, which is called a quark-gluon-plasma by high energy physicists, or, if you want, to the state, which the proponents of the big bang theory assume as the state of matter at the time of elementary synthesis. During the ejection from the 'black hole' all unstable particles will decay. Protons and neutrons will combine into stable light nuclei by part in a similar way as described by conventional models of elementary synthesis.

But dependent on the history of a black hole the conditions may be somewhat different, so that from the ratio of the elements we cannot draw conclusions on the global density of the universe or other parameters like that. Contrary to the picture of big bang theory elementary synthesis takes place still today, so that eventually higher concentrations of some unstable nuclei are expected, for which the time since the big bang was too long to survive.

Characteristic of the collapse in the core of an 'active galaxy' is on one hand the intensive radiation from the compact accretion disk and on the other hand the outflow of matter in high energy plasma beams, the 'jets' in the direction of the rotation axis. The extremely narrow collimation of the jets can be explained by the fact that the emission originates in a region of extremely high magnetic field strength. The motion of the ejected matter, which consists of electrically

charged particles, is thus coupled very strongly to the direction of the field lines. Thus most of the particles are guided back into the accretion disk along the closed field lines.

Only in the direction of the magnetic poles matter can leave the region of very high density and field strength. So it forms those strongly collimated jets, which consist of fully ionised 'primordial matter', that means, of protons, helium nuclei, and electrons, which leave the core with high velocity. Due to the conditions at the origin the kinetic energy of the jet should be of the same magnitude as the energy equivalent of the heavy particles, the velocities should be relativistic.

Direct observation of the jets appears difficult, as emission of radiation from fully ionised matter is weak. The overwhelming fraction of kinetic energy is contained in directional motion of the outward flow, internal interaction inside the jet is comparatively poor. Only if the jet collides with matter at rest in the vicinity, intensive interactions take place. Direct collisions lead to radiation in the gamma and x-ray spectral range, secondary effects produce optical and radio emission. As the matter fields, which are traversed by the jets, normally are not homogeneous, but consist of individual clouds or in some cases of parts of neighbouring galaxies, there occur luminous knots, which are dragged along with the jet by momentum transfer. Far in the outer region, where the energy transferred to the environment is distributed over a larger area, large clouds are observed, from which intense radio emission is observed.

From the linear structure of the jets out to very large distances we must conclude that only a small fraction of the jet energy is lost by interaction with surrounding matter. The main fraction will be distributed into the universe without further interaction. So it is available as the seed of hot matter of primordial composition, from which new structures can be built. Thus the cycle is closed from the source of matter in the black holes and their jets along with the formation of more and more complex structures up to the collapse of galaxies and the generation of new sources.

Chapter 7

The universe - a zero-sum game?

In the last chapters we have demonstrated that it is possible to develop a model of the universe, which is based only on known and experimentally testable physical principles. With this model we can understand the astrophysical observations without applying any unknown or unprovable new effects or constituents.

It is true that also this model brings us no step closer to the answer of the question, why the basic laws of nature are as they are. By principle they might be arbitrarily complex and the universe might contain any number of components, which can by no means be experienced by us. There could be, as some physicists believe, many different universes with different laws and different history. But if these universes do not influence our universe at all, the question of their existence is completely irrelevant for a physicist.

The advantage of the model presented here is, compared to all these considerations, that to understand the cosmic phenomena we rely only on mechanisms, which are, at least by principle, experimentally testable. In addition this model contains no singularities, in contrast to conventional big bang theory. The well proved principles of general relativity are not questioned, but only supplemented. The questions of causality, homogeneity and structure formation are solved quasi automatically without invoking any new 'godhead' like 'dark energy' or 'inflation'.

The singularities of the big bang and of black hole formation, which are mathematically describable, but physically scarcely acceptable, do not occur in the model presented here. Also the attempt to construct a quantum theory of gravitation, becomes superfluous. Matter states, in which gravitational interaction would become so strong that they require a quantum mechanical description, that means, interactions on the scale of the Planck length, do not occur. The repulsive action of potential energy prevents every quantum from penetrating this range.

In addition the attractiveness of the model is enhanced by the fact that, as

already Einstein has emphasised, the size of the universe is finite, if, in contrast to the big bang model, the curvature of the universe is positive. The conception of an infinite space, which continuously expands, and for some unexplained reason even with accelerating velocity, is much more difficult to imagine than a closed, but unlimited continuum, as in this case we know the two-dimensional analog in form of the surface of a sphere.

Also the total mass and the energy content of the universe are finite quantities. Introduction of potential energy, which gives a negative contribution to the total energy content, if we interpret the invariant trace of the energy tensor as the 'total energy', lead to the very simple result that the total energy of the universe is zero. Also quantum fluctuations do not contribute to the energy balance, as the virtual states add the same amount to the positive energy as they add to the negative potential energy. This fact immediately solves the riddle, why the curvature of space is so many orders of magnitude less than quantum mechanics predicts.

But in contrast to Einstein's static universe, in which he tried to establish a balance between the attractive and repulsive forces by introduction of the 'cosmological constant', the equilibrium described here is stable, as the repulsive action increases with increasing curvature and density.

Of course we can put the question: Why is the quantity of mass in the universe just so that there exists equilibrium between potential energy and rest energy of matter? Couldn't we think of a universe, in which there exists only the energy of the quantum vacuum? If we trust the Einstein equations, the answer is: no. These equations lead to contradiction, unless one allows the generation of matter or radiation fields. If on the other side one assumes that matter can be produced, the solution of the time dependent equations always runs into an equilibrium state, in which the gravitational action of quantum fluctuations vanishes.

Considerations, how such a process of matter generation could look like, are highly speculative, of course. According to our present view of high energy and quantum physics no process is known, which leads to the generation of matter. Also if the energy remains in balance due to the simultaneous generation of mass and potential energy, there remains the problem that generation of real particles from vacuum fluctuations always leads to creation of pairs of matter and antimatter particles. But the world, which we know, practically contains only matter, that means leptons and hadrons, but only negligible amounts of the corresponding antiparticles.

But this fact can scarcely be attributed to our understanding of gravitation. From the insight that the total energy of the universe balances to zero it appears more reasonable to suspect that our view of the world of elementary particles is not quite realistic. Maybe that the sum of all elementary particles and their antiparticles is zero indeed, but we have overlooked this symmetry by now.

The proposition that hadrons are composed of three quarks and not of antiquarks and the assumption that quarks and leptons are genuinely different particles makes a symmetrical theory practically impossible. Only if the elementary building stones of electrons and protons are equal, a symmetrical concept

can be realised. Below we will delineate the speculative attempt, to develop such a symmetrical model of elementary particles, which at least describes the basic observed phenomena correctly, but without the claim of representing a confirmed theory. But it may contain a new way of thinking, which could lead to a revision of our the actual standard model.

One basic idea of this speculative model is the assumption that the elementary particles, from which matter is built, are neither bosons nor fermions, but that the spin of particles results from different coupling between these elementary fields. Also in the current theory electrons are described by a system of coupled field functions, which are coupled into a spinor field. The properties related to spin result from this special kind of coupling, which takes place in some abstract 'spin-space', however.

The idea of the model presented here is that the elementary fields - let us call them quarks again for simplicity - are spatial fields, which can interact in different ways: On one hand by superposition in form of a spinor, forming a pointlike system like an electron, on the other hand as spatially extended systems, comparable to the system of proton and electron in a hydrogen atom or the binding between quarks and antiquarks in mesons. Contrary to the 'fermionic' states of spinors such 'bosonic' bound states are possible with different levels of energy or angular momentum.

Four propositions are sufficient to describe the spectrum of known elementary particles in such a symmetrical model:

1. Every particle contains the equal numbers of quarks and antiquarks.
2. Quarks and antiquarks can be bound together 'fermionic' (as a spinor, pointlike, spin quantum number $1/2$) or 'bosonic' (spatially extended, integer spin)
3. Quarks have an electrical charge (the elementary charge) or hypercharge (strangeness) or both, antiquarks the same, but with opposite sign.
4. No two quarks can be in the same quantum state.

Equal to the basic triplet in conventional theory there are three different quarks and the corresponding antiquarks. Thus the spectrum of mesons, the bosonic bond between one quark and an antiquark is identical. The spectrum of leptons results from the corresponding fermionic binding. But in this case there is only one quantum state for every combination of quarks. No excited states or states with different angular momentum are possible (for details see appendix).

With the families of mesons and leptons the possibilities of two-quark states are exhausted. But there is a number of different realisations of states consisting of two quarks and two antiquarks. Hadrons can be interpreted as a system of two fermionically bound quarks, surrounded by two bosonically bound antiquarks. also in this case we find the same spectrum as in conventional theory. As the 'classical' quarks form a complete group, the same quantum states would be obtained, if we replace one quark by the sum of the two other antiquarks. Only the spin properties would be different. This is compensated in our model by the fact that there is one fermionic and two bosonic bonds.

Systems consisting of two fermionically bound quarks, bosonically bound

to a pair of fermionically bound antiquarks, form an additional system of very heavy mesons (J/ψ or Y), W - and Z -particles can be interpreted as fermionically bound states of two quarks and two antiquarks. Details of the spectrum, which result under consideration of the condition of distinguishability (point 4) are discussed in the appendix.

As in conventional theory interactions between elementary particles can be described as exchange reactions. But the processes of strong and weak interaction must now not be regarded as completely different mechanisms. Both have the effect that one quark is replaced by another one. The only difference is that strong interaction relates to a bosonic bond, while in weak interaction a fermionic bond is changed.

It is interesting to remark that in the scheme presented here electron and proton together contain one quark and one antiquark of all three species. The same holds for neutron and electron neutrino. Thus it appears plausible that the universe contains equal amounts of quarks and antiquarks of every type. Thus there exists the possibility that the universe could be created from nothing.

Of course it remains to explain, why the separation into protons and electrons is stable, though the sum of their quarks is equal to the sum of the antiquarks, that means, in total equivalent to zero. That they cannot annihilate each other becomes clear regarding the fact that interactions of electrons and protons take place either with the bosonically bound antiquarks or with the fermionic core built from two quarks. In both cases the total number of the involved quarks is different from the number of antiquarks, so that annihilation is impossible. Only when the density of matter is so high that the bosonic bonds become unstable, that means, e.g. in the interior of black holes, creation or annihilation can be expected.

The concept, which we have outlined here, requires a radical change of thinking, of course. The world is no longer distributed into fermions and bosons and the components of wave functions no longer describe superpositions of possible states, but the superposition of real fields. But maybe that just here we need a radically new ansatz. With quantum theory we have the mathematical tools to describe the phenomena of the microcosmos in an excellent way. But the price is that the underlying physical reality is scarcely intelligible for our imagination.

Maybe that the world on the smallest and largest scales becomes easier to understand, if we are willing to leave the accustomed tracks in spite of the impressive success in the last century. Already for many times the ideas of mankind about the structure of the world, though appearing well confirmed, have been radically changed by new thoughts and new directions of view. This paper should be regarded as an impulse to extend our angle of view.

Appendix A

The definition of density and potential energy

Basis of the theory of general relativity are the Einstein equations ¹, which relate the metric of space-time, expressed by the curvature tensor (or its contracted form, the Ricci tensor R_{ij}) and the energy tensor T_{ij} .

$$R_{ij} - \frac{1}{2}R \times g_{ij} = 8\pi T_{ij} \quad (\text{A.1})$$

(in a unit system, where the speed of light and the gravitational constant have been set to 1). g_{ij} is the fundamental tensor. The first and second derivatives of this tensor determine the curvature tensor. The condition of general covariance, that means, the requirement that the content of the field equations should be independent of the choice of the reference system, leads to certain restrictions for the possible forms of the energy tensor. It must be determined uniquely by scalar or tensorial quantities, which depend only on local invariant properties of matter itself, on the fundamental tensor or on quantities which can be derived from this tensor without differentiation.

In the description of cosmological problems usually the energy tensor is used in the form borrowed from special relativity. It is described as an ideal fluid medium with the total energy expressed by a continuous matter density, with its 4-velocity, and by the pressure tensor.

$$T_{ij} = \rho \frac{dx_i}{ds} \frac{dx_j}{ds} + p \left(g_{ij} + \frac{dx_i}{ds} \frac{dx_j}{ds} \right) \quad (\text{A.2})$$

But in general relativity the definition of density is no longer unique. In Euclidean space the density is defined as the differential of the mass $\rho = dm/dV$. If space is curved, however, the size of the spatial volume element depends on the metric. According to Einstein therefore the energy tensor used in his equations

¹In the mathematical formulations we follow the notation of the standard textbook of R. M. Wald 'General Relativity', The University of Chicago Press, Chicago (1984)

should contain the invariant density, that means, the density, which one would measure in an Euclidean space which is locally tangential to the actual space. The quantity $\rho_0 = dm/dV_0$ (dV_0 is the volume element in tangent space) is a local property of the matter field and thus can be used in a covariant equation.

But the use of the invariant density leads to the problem that the total mass is no longer the result of integration of this density over the actual volume.

$$\int dm \neq \int \rho_0 dV \quad (\text{A.3})$$

By principle it is impossible to define mass density and total mass as invariant properties, if the volume element is not invariant. To obtain invariance of the total mass, we have to introduce a corrective term in the energy tensor. The density in curved space can be expressed by the invariant density as

$$\rho = \rho_0 + \rho_0 \left(\frac{dV_0}{dV} - 1 \right). \quad (\text{A.4})$$

Only the second term in this equation depends on the metric. It is a scalar property, which can be used in a covariant energy tensor, but not together with some vectorial quantity like the 4-velocity. The only reasonable method to change the energy tensor in such a way that the matter term, the quantity T_{00} , is equal to ρ , and that in spite of that the property of covariance is preserved, is to follow the way proposed by Einstein, but to introduce an additional term, which allows for the influence of curvature:

$$T_{ij} = \rho_0 \frac{dx_i}{ds} \frac{dx_j}{ds} + \rho_0 \left(\frac{dV_0}{dV} - 1 \right) \times g_{ij} + p \left(g_{ij} + \frac{dx_i}{ds} \frac{dx_j}{ds} \right) \quad (\text{A.5})$$

The meaning of the additional term can easily be recognised, looking at the Schwarzschild metric, which describes the space-time outside a radially symmetric matter distribution with the total mass M . This metric is characterised by the path element

$$ds^2 = -f(r)dt^2 + h(r)dr^2 + r^2 d\Omega^2 \quad (\text{A.6})$$

with $h(r) = 1/(1 - 2M/r)$ ($d\Omega$ is the element of solid angle). From this we get the volume element

$$dV = \frac{r^2 dr d\Omega}{\sqrt{1 - 2M/r}}, \quad (\text{A.7})$$

from which in the limiting case of weak curvature we find the density

$$\rho = \rho_0 - \frac{\rho_0 M}{r}. \quad (\text{A.8})$$

The difference compared to the density in tangent space is just the density of potential energy in the sense of Newtonian mechanics. To maintain conservation of matter or total energy not only locally but also integrally in curved space, it is necessary to include potential energy in the energy tensor.

The path used here to calculate the potential energy is not applicable generally, of course. Due to the special choice of coordinates the volume element (A.7) is singular at $r = 2M$, so that the expression in (A.4) is no longer meaningful.

Equation (A.5) is valid also inside a matter distribution. But as the size of the volume element depends implicitly on the metric, in the determination of the metric from the Einstein equations (A.1) we have to consider the reactive coupling of the metric to the energy tensor. In the discussion by now we have implicitly assumed that only the mass density determines the metric of space. If there are additional kinetic contributions like pressure in a gas, instead of matter density we must use the density of total energy, expressed by the invariant trace of the energy tensor.

The exact determination of the potential energy may thus become rather intricate. But we can make some general statement, how the solution should look like:

1. Potential energy is proportional to the local density ρ_0 and thus vanishes in empty space.
2. As a positively curved volume element is always larger than the corresponding element in the flat tangent space, the quantity $dV_0/dV - 1$ is always negative.
3. By principle potential energy can attain arbitrarily high negative values, if the size of the curved volume element (like in the limit of the Schwarzschild metric) becomes singular.
4. The local energy density (trace of the energy tensor) can be positive or negative. By the pressure term in the energy tensor potential energy acts like a negative pressure or repulsive force.

Based on these properties of potential energy, formation of singularities in the density distribution, the formation of 'black holes' is impossible. In the conventional view of general relativity kinetic pressure is regarded as an additional positive contribution to the total energy and thus enhances the attractive action of gravitation. Thus stabilisation of a system by pressure gradients becomes impossible beyond a certain mass limit. On contrary potential energy generates a negative pressure and thus a repulsive action, increasing with density, and thus brings every contraction to an halt.

Appendix B

Basic equations for a homogeneous universe

Based on the assumption that the universe does not contain any regions with distinct properties but is homogeneous on large scale, it can be described by a global metric of the form

$$ds^2 = -dt^2 + a(t)^2 \cdot [d\psi^2 + f(\psi) \cdot (d\theta^2 + \sin^2 \theta d\phi^2)]. \quad (\text{B.1})$$

Here one assumes that the metric can vary in time, but is spatially homogeneous. The function $f(\psi)$ takes the values $\sin^2 \psi$, ψ^2 , or $\sinh^2 \psi$ for positive, zero or negative curvature respectively. Here we restrict our discussion to the case of positive curvature.

In the case of a homogeneous matter distribution with density ρ and pressure p the solution of the Einstein field equations leads to the well known Friedmann equations, to determine the temporal evolution of the scale parameter $a(t)$:

$$\frac{3\dot{a}^2}{a^2} + \frac{3}{a^2} = 8\pi\rho \quad (\text{B.2})$$

$$\frac{3\ddot{a}}{a} = -4\pi(\rho + 3p) \quad (\text{B.3})$$

(in a unit system with the speed of light and the constant of gravity set to 1). If there are various fields with different equations of state. that means, different relations between ρ_i and p_i , the right hand sides of (B.2) and (B.3) have to be replaced by the corresponding sum. The equations (B.2) and (B.3) can be satisfied simultaneously, however, only if density and pressure vary in a suitable manner with change of the scale parameter $a(t)$. A pressureless matter field requires $\rho_m \propto 1/a^3$, a radiation field with $p = 1/3 \rho_r$ has to change as $\rho_r \propto 1/a^4$. Global conservation thus exists only for matter. A radiation field changes its total energy on account of a change of the scale variable.

If we postulate global conservation for all kinds of fields, we have to take into account potential energy. To include potential energy in the energy tensor,

any reasonable formula must contain besides of the fundamental tensor only invariant local state variables, the local invariant density of the energy fields and the local curvature of space. Potential energy must vanish, if this density is zero or if the gravitational potential and with this the curvature of space is zero. As the most reasonable ansatz we must consider an expression of the form $\lambda \times g_{ij}$. But in contrast to the 'cosmological constant' introduced in Einstein's static solution, now the quantity λ depends on the density of matter and on the spatial curvature. In analogy to Newtonian mechanics the best choice for λ is an expression of the form

$$\lambda = \frac{K}{a} \sum_k (\rho_k + 3p_k) \quad (\text{B.4})$$

where K is a universal constant, which is related to the gravitational constant. This dependency of λ on density and curvature has far reaching consequences, however:

1. The system (B.2) and (B.3) no longer allows a time dependent solution, but only the trivial solution $a = \text{const.}$. A homogeneous universe necessarily has to be static. The curvature is fixed by the requirement that the attractive force of positive energy is in balance with the repulsive action of potential energy. In a universe, where pressureless matter dominates, we have

$$\frac{3}{a^2} = 8\pi\rho - \frac{K}{a}\rho \quad (\text{B.5})$$

$$0 = -4\pi\rho - \frac{K}{a}\rho, \quad (\text{B.6})$$

leading to

$$K = -4\pi a, \quad a = 1/\sqrt{4\pi\rho} \quad \text{oder} \quad a/\text{cm} = 3.28 \cdot 10^{13} / \sqrt{\rho/\text{cm}^{-3}} \quad (\text{B.7})$$

2. Vacuum energy from quantum fluctuations, as they have been discussed as possible origin of a cosmological constant in conventional theory and which are described by energy fields of the form $C \times g_{ij}$ are without any effect in the field equations, as their contribution to potential energy is just $-C \times g_{ij}$ and thus compensates the field energy exactly.

3. Local changes of fields with zero rest mass like kinetic energy or radiation fields generate changes of curvature and thus changes of the scale parameter a proportional to a^{-4} . At the same time the potential energy of matter is changed with the same proportionality, so that the total energy of the system remains constant. Like in Newtonian theory there is no law of conservation for kinetic or potential energy alone, but only for their sum.

Appendix C

Red shift - energy loss by gravitational radiation

Apart from the geometric interpretation of gravitation the essential difference between Newtonian mechanics and general relativity is the statement that gravitational interaction and thus any change of space-time is limited to the speed of light. As a consequence, similar to electromagnetic interaction, due to the retardation of potentials there exist radiation effects. This gravitational radiation is not restricted to the quadrupole radiation, which reduces the angular momentum of rotating double stars, but also in the 'force free' or geodesic motion in matter filled space there exists a momentum loss by gravitational radiation, analogous to bremsstrahlung in electromagnetism.

An exact derivation of the effect from the field equation is not known by now. A simple estimation can be obtained, however, from a quasi-Newtonian approximation, only considering the retardation of interaction and the curvature of space. To do this we calculate the force on a moving test mass in a homogeneous matter-filled universe under the following assumptions:

1. The velocity of gravitational interaction is limited to the speed of light.
2. Mass and energy create an intrinsic curvature of space. A spatially homogeneous universe can thus be regarded as a three-dimensional surface of constant curvature.
3. Lines of force between masses follow the geodesic lines between the corresponding points.

To determine the motion within a three-dimensional surface it is convenient, to describe it by means of polar coordinates of a four-dimensional sphere of a given radius a .

$$\begin{aligned}x &= r \cos \chi \cos \theta \cos \phi & , & & y &= r \cos \chi \cos \theta \sin \phi, \\z &= r \cos \chi \sin \theta & , & & w &= r \sin \chi,\end{aligned}\tag{C.1}$$

where x, y, z and w is a set of Cartesian coordinates. In this system we determine the force exerted on a mass at the point $P = (a, 0, 0, 0)$ by the mass in some volume element dV of the surface at $r = a$.

The size of the volume element is given by

$$dV = a^3 \cos^2 \chi \cos \theta d\chi d\theta d\phi \quad (\text{C.2})$$

The distance between the element and P , as measured along a straight line through the four-dimensional space is given by

$$r_P = \sqrt{(x-a)^2 + y^2 + z^2 + w^2} = a\sqrt{2 - 2\cos\chi\cos\theta\cos\phi}. \quad (\text{C.3})$$

The distance along the connecting geodesic line is

$$s = 2a \arcsin \frac{r_P}{2a} = 2a \arcsin \sqrt{\frac{1 - \cos\chi\cos\theta\cos\phi}{2}} \quad (\text{C.4})$$

The component of the gravitational force at P in some direction, defined by the unit vector \vec{e} then is

$$dF = \frac{G\rho mdV}{s^2} (\vec{e}_s \vec{e}), \quad (\text{C.5})$$

where ρ is the mass density, G the gravitational constant and \vec{e}_s the unit vector in direction s at point P . As all directions are equivalent, we chose \vec{e} in the direction of the y -coordinate. The projection onto the direction of the force component is

$$(\vec{e}_s \vec{e}) = \frac{\cos\chi\cos\theta\sin\phi}{\sqrt{1 - \cos^2\chi\cos^2\theta\cos^2\phi}}. \quad (\text{C.6})$$

In a universe of constant mass density we find the total force by integration of Eq.(C.5) over all directions and distances:

$$F_y = G\rho ma^3 \int_{-\infty}^{\infty} \int_{-\pi/2}^{\pi/2} \int_{-\pi/2}^{\pi/2} \frac{\cos^3\chi\cos^2\theta\sin\phi}{s^2\sqrt{1 - \cos^2\chi\cos^2\theta\cos^2\phi}} d\chi d\theta d\phi \quad (\text{C.7})$$

The limits of integration over ϕ are set to $\pm\infty$, as the lines of force may extend beyond the reciprocal pole.

Obviously for a mass at rest the integral is zero on symmetry grounds. In Newtonian mechanics this holds also for a moving mass, as the gravitational force acts in every volume element simultaneously. If interaction is limited to the speed of light, however, retarded forces have to be used. In the determination of the force all distances s at time t have to be replaced by $s'(t) = s(t - \tau)$, their value at a point of time, earlier by the running time of the signal $\tau = s'/c$. That means that the integral is no longer symmetrical. Distances in direction of motion appear enlarged, opposite to motion they are reduced. This leads to a resulting force opposite to the direction of motion.

For the numerical evaluation of the force integral it is convenient to change the coordinate system. Instead of changing the distance s and its projection in

Eq.(C.7), it is easier to keep these quantities unchanged and instead to change the integration variable, by introducing a comoving reference system:

$$\phi' = \phi(t') = \phi(t) + \frac{d\phi}{dt}\tau = \phi(t) + \frac{vs'}{ac}, \quad (\text{C.8})$$

where v is the velocity of the moving mass. This is allowed, as ϕ is the only coordinate, which is affected by the motion. Expressing the distance s according to Eq.(C.4) the differential reads:

$$d\phi = d\phi' \left(1 - \frac{v}{c} \frac{\cos \chi \cos \theta \sin \phi'}{\sqrt{1 - \cos^2 \chi \cos^2 \theta \cos^2 \phi'}} \right). \quad (\text{C.9})$$

The limits of integration are not changed by the transformation. The resulting force then is

$$F_y = -2G\rho ma^3 \frac{v}{c} \int_{-\infty}^{\infty} \int_{-\pi/2}^{\pi/2} \int_{-\pi/2}^{\pi/2} \times \frac{\cos^4 \chi \cos^3 \theta \sin^2 \phi' d\chi d\theta d\phi'}{(1 - \cos^2 \chi \cos^2 \theta \cos^2 \phi') \left(\arcsin \sqrt{(1 - \cos \chi \cos \theta \cos \phi')/2} \right)^2} \quad (\text{C.10})$$

From numerical integration of the integral we find the value $Y = 3.0695$. Thus the change of velocity results as

$$F_y = m \frac{dv}{dt} = -mv \frac{2Y\rho Ga}{c}. \quad (\text{C.11})$$

Correspondingly we find for the change of kinetic energy

$$\frac{dE}{dt} = \frac{d}{dt} \frac{mv^2}{2} = -\frac{4Y\rho Ga}{c} E. \quad (\text{C.12})$$

Every motion in mass-filled space is subject to a loss of kinetic energy by emission of gravitational radiation, caused by the finite velocity of gravitational interaction. Here we have calculated the energy loss only for the case of non-relativistic motion, but as the formula for the energy loss does not contain the mass explicitly, it appears reasonable to assume that it is valid also for relativistic motion, in the limiting case also for energy quanta without rest mass. That means that photons experience the corresponding energy loss. Of course in this case the energy loss cannot change the velocity, but instead the wavelength will change.

$$\lambda = \lambda_0 \exp \left(\frac{4Y\rho Ga}{c} t \right). \quad (\text{C.13})$$

As a first approximation this leads to a linear increase of wavelength with distance, exactly the same behaviour as described by big bang theory and as it is confirmed by observations. Differences between the two theories occur on one side, when deviations of the exponential dependence from linear approximation

become important, and on the other side when the different assumptions on the temporal development of expansion are taken into account.

We should emphasise that for the case of Euclidean space the formula leads to similar difficulties as discussed in the well known Olbers paradox. As according to Eq.(C.13) red shift is proportional to the radius of the universe, in this case the red shift would be infinite. This can be regarded as an indication that the size of the universe must be finite.

The quantity $(4Y\rho Ga/c)$ in Eq.(C.13) is identical with the Hubble constant H . On the other hand from the Einstein field equations we have the relation

$$a = \sqrt{\frac{c^2}{4\pi G\rho}} \quad (\text{C.14})$$

(for better clarity, contrary to the last chapter, we have used the traditional writing with the speed of light c and the constant of gravity G) This leads to the relation

$$\rho = \frac{\pi H^2}{4Y^2 G}, \quad a = \frac{cY}{\pi H}, \quad M = \frac{Yc^3}{2GH}. \quad (\text{C.15})$$

M is the total mass of the universe. Assuming a value of 70 km/sec/Mpc for the Hubble constant, the result for the total density is $6.43 \cdot 10^{-30} \text{g/cm}^3$, the radius of the universe is $1.29 \cdot 10^{28} \text{cm}$, and the total mass is $2.73 \cdot 10^{56} \text{g}$.

Appendix D

X-ray emission of the intergalactic plasma

We can prove the existence of the intergalactic plasma and determine its properties from the diffuse x-ray background radiation, if we assume that this plasma is responsible for the emission and not, as assumed in the conventional big bang model, some unresolved individual sources. For the high energy fraction ($E > 3\text{keV}$) this assumption appears justified, as there are no known cosmic processes, which could contribute to the background in quantities comparable to observations.

The emitted spectrum of the x-ray background has been measured with sufficient accuracy to find out the properties of the intergalactic plasma. In a plasma of electron density n_e , the ion density n_i , and kinetic temperature kT the energy emitted per time and frequency interval by bremsstrahlung is given by

$$\frac{d\varepsilon}{d\nu} = \frac{2^5 \pi e^6}{3 m_e c^3} \left(\frac{2\pi}{3 m_e kT} \right)^{1/2} n_e \sum n_i Z_i^2 g e^{-\frac{h\nu}{kT}}. \quad (\text{D.1})$$

Z_i is the charge of the ion species i , g the Gaunt factor, for which we use the approximation $g = 0.74(E/kT)^{-0.4}$ in the following¹. For a fully ionised plasma, consisting of hydrogen and 25% helium for the energy spectrum this leads to the formula

$$\frac{d\varepsilon}{dE} = 7.216 \cdot 10^{-26} \text{cm}^{-3} \text{s}^{-1} \sqrt{\frac{1\text{keV}}{\text{kT}}} \left(\frac{E}{\text{kT}} \right)^{-0.4} \left(\frac{\rho}{10^{-29} \text{gcm}^{-3}} \right)^2 e^{-\frac{E}{\text{kT}}}. \quad (\text{D.2})$$

The intensity of radiation then results by integration along the optical path. Here we must regard the fact that the energy changes by red shift along the path. To determine the intensity at some energy E_0 at a distance s we must use the emissivity at the energy $E = E_0 \times \exp(Hs/c)$. In addition the intensity

¹Itoh, N. et al., *Astrophys.J.Suppl.* 128 (2000)


may be reduced by absorption from dust grains.

$$\frac{di}{dE} = \frac{1}{4\pi} \int_0^\infty \frac{d\varepsilon}{dE} e^{-\kappa s} ds. \quad (\text{D.3})$$

In this equation the absorption coefficient κ has been assumed as independent of frequency. Integration along the path can then be replaced by integration over energies. From $E = E_0 \times \exp(Hs/c)$ we get $ds = c/(HE)dE$. With the abbreviation $x=E/(kT)$ we find

$$\frac{di}{dE} = \frac{1}{4\pi} \frac{c}{H} \int_{x_0}^\infty \frac{d\varepsilon}{dx} \left(\frac{x}{x_0}\right)^{\frac{\kappa c}{H}} \frac{dx}{x}, \quad (\text{D.4})$$

where $d\varepsilon/dx$ according to Eq.(D.2) is given as $kT d\varepsilon/dE$. The numerical solution of this equation can be fitted to the observational data² by suitable choice of the parameters ρ , kT and κ (see figure). The accuracy of the data is not sufficient for an exact determination of the parameters, as the shape of the spectrum varies not much with varying absorption. The effective temperature varies between $\kappa c/H = 0$ and 0.5 from $kT = 110\text{keV}$ to $kT = 90\text{keV}$. The corresponding density values are $\rho = 3.6 \cdot 10^{-30}\text{g/cm}^3$ and $\rho = 4.3 \cdot 10^{-30}\text{g/cm}^3$.



In the range of high energies the calculated intensity distribution decreases considerably steeper than the measured data. This is caused by the contribution of high energy electrons. Due to the decreasing collisional cross section the contribution of the high energy tail is considerably lower than that of the thermal plasma. The dotted line in the figure shows the contribution of the high energy electron under the assumption that the total density of the high energy particles is $1 \cdot 10^{-30}\text{g/cm}^3$ with an energy distribution $dn_e/dE \sim 1/E$ between 1 GeV and 100 keV.

²Gruber et al., *Astrophys.J.* 520, 124 (1999)

Appendix E

Intergalactic dust

Formation of dust needs two conditions to be met: on one hand a relatively high density of heavy elements and secondly a temperature range, low enough to allow the formation of molecules, but high enough that collisions and reactions are sufficiently frequent. To our experience these conditions are found predominantly in the expanding clouds of stellar explosions. Preferably supernova explosions of type Ia (SNIa) must be regarded as sources of intergalactic dust. On one hand these explosions are characterised by very high expansion velocity of the explosion cloud and on the other hand the progenitor star is a 'white dwarf', a highly evolved star, thus containing a large fraction of heavy elements.

All supernovas of type Ia show very similar properties. In the following we consider the formation of dust in a 'standard' supernova of this type.¹ SNIa occur, when a white dwarf exceeds its stability limit (the Chandrasekar limit) by accretion of matter from its surrounding.

E.1 Formation of dust

'White dwarfs' are stars in which the first steps of nuclear fusion, the burning of hydrogen and helium has come to an end. Also the further fusion reactions have practically stopped, so that the star is kept in equilibrium with gravity only by the degeneration pressure of electrons. If the total mass of such a star exceeds a certain limit, it becomes unstable, it collapses. During this collapse in the centre of the star nuclear fusion is resumed up to the final formation of the iron group atoms. This energy production than leads to explosion of the complete star and a matter cloud, consisting preferably of heavy elements expands radially with velocities in the order of 10^4 km/sec. The mass of the expanding cloud is given by the stability limit of the star. It is about 1.4 solar masses ($M_C = 2.8 \cdot 10^{33}$ g). The 'standard' supernova reaches a peak intensity of $M = -19.5$, the decay time per day is $dM/dt = 0.06m$. Only at the beginning

¹Perlmutter, S. et al., ApJ 483(1997)

of the explosion thermal energy is produced in the centre, which is by part emitted in form of high energy neutrinos. Later there is only radiative energy transport in the interior and emission of radiation from the boundary layer of the expanding cloud.

When the emission pattern and the expansion velocity are known, the determination of the temperature in the surface layer is straightforward, if we assume that the plasma is optically thick.

$$L = 4\pi R^2 \sigma T^4 = 10^{\frac{M_{\text{peak}} - M}{2.5}} \times L_{\text{peak}} \quad (\text{E.1})$$

(M is the absolute bolometric magnitude; σ is the Stefan-Boltzmann constant).

With $R = u \cdot t$ and an expansion velocity of $u = 5000$ km/s we find that recombination into neutral atoms (at $T=10000$ K) begins about 20 days after the intensity maximum and is completed after 50 days (4000 K). After this time observations show deviations from the linear decrease of intensity, as the surface layer becomes optically thin. But it appears reasonable to assume that the bolometric magnitude continues its linear decrease and that the observed deviations result only from the shift of emission from visible to infrared.

With this extrapolation the equilibrium composition of the gas can be determined under the assumption that the element abundances are similar as those of the sun, but without hydrogen and helium and a reduced fraction of C, N, and O, as these elements will be consumed by part by the further steps of fusion. The numerical calculations show that the first molecules are formed about 75 days after explosion (SiO and CO at 2500 K).

105 days after maximum intensity the temperature in the surface layer has dropped to 1500 K. Under equilibrium conditions at this time the first condensed phases would occur (MgSiO_3 and SiO_2). (Fe_2SiO_4 would solidify at 1100K).

At this time the radius of the cloud is about $4.5 \cdot 10^{15}$ cm, the particle density is of the order 10^9cm^{-3} . Under these conditions homogeneous nucleation in the gas is scarcely possible. The gas would be strongly supersaturated. But in most cases an explosion blast wave does not expand into vacuum, but into the interstellar medium, which already contains small dust grains, which can serve as a seed for the further growth.

The growth of grains takes place preferably shortly after the gas reaches supersaturation, as later the number of collisions decreases with the third power of the cloud radius. To estimate which final size a dust grain can achieve, we must calculate the number of collisions between a grain and the surrounding molecules. For simplicity we assume that there are only molecules of one species with mass m and that the particle density is constant all over the cloud $n = M_C / (4/3\pi R^3) / m$. Then we get for the temporal change of the size of a spherical grain of density ρ_D :

$$\frac{d}{dt} \left(\frac{4}{3}\pi r^3 \rho_D \right) = \pi r^2 n m v_{\text{th}}. \quad (\text{E.2})$$

$v_{\text{th}} = \sqrt{3kT/m}$ is the mean thermal velocity. The time dependence of the temperature can be approximated by a power law of the form $T/T_0 = (r/r_0)^{-\alpha}$,

where α is close to one for the example considered here. With $R = u \cdot t$ we can integrate Eq.(E.2) from the beginning of supersaturation to infinity with the result

$$r_\infty = r_0 + \frac{3}{40\pi} \frac{M_C v_{th0}}{\rho_D u R_0^2}. \quad (\text{E.3})$$

The index 0 refers to the values at beginning of condensation. With the data of our example we find a final radius of $2.8\mu\text{m}$. Of course this value can be regarded only as a very crude estimation due to the simplifications made above. We have assumed that there is only one species of molecules condensing at 1500 K. For another species condensing only at 800 K there would be only half the final size.

Secondly we have supposed implicitly that the density of molecules is not changed by the condensation itself. If this assumption is justified, depends on the number of seed grains, which are swept up by the expanding cloud. The density of the interstellar medium in the vicinity of the explosion may vary strongly. If we take the conditions in the neighbourhood of the solar system as a guideline, the density of the interstellar medium is of the order $10^{-23}\text{g}/\text{cm}^3$ with a dust fraction of 1%. Assuming a mean grain size of $0.01\mu\text{m}$, the density of seed grains is 10^{-7}cm^{-3} . The main part of dust formation should be finished, before the explosion cloud reaches a volume of 10^{48}cm^3 . That means that the number of seed grains is of the order 10^{41} . If all matter contained in the cloud would condense onto these grains, the radius of one grain would be $15\mu\text{m}$. From this estimation we can conclude that it is the collision rate and not the availability of molecules, which determines the final size of the dust grains. Anyhow we can state that grains, which begin growing with the onset of supersaturation, can easily attain a size of more than $1\mu\text{m}$. Seed grains, which are swept into the cloud at later times, will reach correspondingly smaller size.

E.2 Size selection

At the time of creation the radial velocity of dust grains is equal to the expansion velocity of the cloud. If this expansion would take place into empty space, nearly all grains would leave the galaxy, in which they have produced, as their velocity exceeds the local escape velocity by several times. For instance, for a supernova, which explodes in the spiral arm of a galaxy of say 10^{45}g at a distance from the centre like that of our sun (7.7 kpc), the escape velocity is about 500 km/sec, ten times less than the radial velocity of the grains.

In real galaxies the dust grains must first traverse the interstellar medium, before they reach free space. In every collision with a gas atom they lose part of their initial momentum. Even if the emission is perpendicular to the disk of a galaxy, the column density of the interstellar gas should be in the order of $n_c = 10^{21}\text{cm}^{-2}$.

In every collision with a hydrogen atom the momentum loss on average is $-m_H \cdot u$, as the mean radial velocity component of the gas atoms is zero. The

momentum loss of a spherical dust grain of radius r and the density ρ_D then is

$$\frac{4\pi}{3}\rho_D r^3 \frac{du}{ds} = -n \cdot m_H \cdot \pi r^2 u, \quad (\text{E.4})$$

or by integration along the path of the grain

$$u_\infty = u_0 \cdot \exp[-3m_H n_c / (4\rho_D r)]. \quad (\text{E.5})$$

If, as has been estimated above, the initial velocity is ten times higher than the escape velocity, the exponential factor in the equation above has to be > 0.1 for a grain to leave the galaxy. For a column density of 10^{21}cm^{-2} this means $r > 2.5\mu\text{m}$. Of course, this is only a crude estimation of the order of magnitude. But it clearly demonstrates that only large grains can leave the galaxy, while smaller grains are thermalised within the galaxy and are available there as seed grains for later generations of supernovas.

As the size limit is just in the range, where the absorption properties of dust for optical radiation change from selective to wavelength independent absorption, the dust, which is captured inside the galaxy has highly selective properties and leads to reddening of spectra, while the dust, which escapes into the intergalactic medium, exhibits 'grey' absorption.

E.3 Global distribution and stability

After leaving their parent galaxy dust grains are still exposed to the interaction with the intergalactic plasma. This may lead to erosion or eventually to complete destruction of the grains. To estimate the rate of erosion, we can, similar to dust formation or momentum loss, again start from the collision rate, but we must regard the fact that at the very high velocity of the striking protons ($kT \approx 100\text{keV}$) the sputtering efficiency is low ($\eta < 10^{-2}$). Analogous to Eq.(E.2) we find for a spherical grain with radius r :

$$\frac{d}{dt} \left(\frac{4}{3}\pi r^3 \rho_D \right) = -\eta m_a n \cdot \pi r^2 v_{\text{th}}, \quad (\text{E.6})$$

Here m_a is the mass of the eroded atom. v_{th} is the thermal velocity of the ions in the intergalactic plasma, which is in the order of $5 \cdot 10^8\text{cm/sec}$. By integration we get $\tau = r_0 \rho_D / (\eta m_a n v_{\text{th}})$ as the erosion time constant for grains with radius r_0 .

For a silicate grain of $r_0 = 1\mu\text{m}$ with the data of the intergalactic plasma we obtain a time constant of $> 10^{18}\text{sec}$, more than the Hubble time. This means that intergalactic dust in the form of large grains remains stable and is distributed homogeneously in the universe. As the grains leave their parent galaxy with a considerable fraction of their initial velocity in the order of some 10^3 km/sec , they can traverse even the largest voids of the universe during their lifetime. Thus we can expect that the dust is distributed uniformly all over the universe.

E.4 Detection of intergalactic dust

By principle the existence of dust can be proved by absorption or emission measurements. But due to the low intensity of the signals both measuring methods have large error margins. This is especially true for emission measurements. Though the dust grains are heated by collisions with the surrounding gas particles, which lead to emission in the far infrared, their emission is low compared to the intensity from sources in our galaxy.

An indirect hint on the ejection of dust particles from galaxies into the intergalactic medium can be found in the properties of the gas component in clusters of galaxies. Though also in this case a large amount of dust can escape from the individual galaxies, the velocity is not sufficient, to escape from the complete cluster. On the other hand the density of the intracluster gas is higher by three orders of magnitude, compared to the intergalactic plasma. That means that the erosion time of dust grains is only about 10^{15} sec, only a small fraction of the cluster lifetime. Thus the major part of all metals is in gaseous form, be it produced in the cluster itself or swept in as dust from the intergalactic plasma.

Indeed the observed metallicity of the gas in most clusters is close to that in the solar system, that means, mass fractions in the order of 1%. This high concentration can be understood only, if we assume that inside the galaxies a very efficient mechanism of dust production and ejection exists. But as the corresponding spectral lines are not observed in the intergalactic plasma, the metals must be present there in solid form, bound into dust grains.

The most convincing hint onto the existence of coarse grain intergalactic dust comes from absorption measurements, though even here the interpretation of the observations is controversial due to the lack of spectral selectivity.

Supernovas of type Ia form a very homogeneous group of phenomena and due to their extreme luminosity are observable over large cosmic distances. But what we observe is a stronger decrease of the radiation intensity than would be expected from a pure distance relation ($i \sim 1/s^2$). According to big bang theory one tries to explain this by an accelerated expansion of the universe, caused by the action of some mysterious gravitationally repulsive dark energy.

But absorption by intergalactic dust appears as a much more obvious explanation. The fact that this dust in contrast to interstellar dust inside a galaxy is much coarser, immediately explains, why this absorption does not lead to reddening of the spectra.

In a quantitative determination of the intensity deviation from the normal $1/s^2$ dependence of distant supernovas in addition to absorption the influence of space curvature has to be considered. The connection between the size of surface area and the solid angle element differs from Euclidean geometry. In a homogeneously curved space with a radius of curvature a for a source at distance s the illuminated area is smaller than in Euclidean space by $F/F_0 = \sin^2 \varphi / \varphi^2$, with $\varphi = s/a$. The intensity change of a point source by absorption (absorption


coefficient κ) and curvature thus results as

$$I = I_0 \cdot e^{-\kappa s} \left(\frac{\varphi^2}{\sin^2 \varphi} \right) \quad (\text{E.7})$$

or, expressed by change of magnitude:

$$\Delta m = \frac{2.5}{2.306} (\kappa a \varphi - 2 \ln \varphi + 2 \ln \sin \varphi). \quad (\text{E.8})$$

The relation between red shift z and the angle φ is given by $1+z = \exp(Ha\varphi/c)$. Various observations of SNIa have been carried out by several groups. In the figure below these measurements are plotted together with a fit by Eq.(E.8). In



Solid line: Fit of measurements by Eq.(E.8), dashed line: 'concordance model' $\Omega_M = 0.3, \Omega_\lambda = 0.7$, measurements: crosses: Perlmutter, S. et al., ApJ 517 (1999), spheres: Riess, A.G. et al., ApJ 607 (2004), triangles: Hamuy, M. et al., AJ 112 (1996)

addition the result according to the 'concordance model' of big bang theory is plotted, which ascribes the deviations from the Euclidean distance law to an accelerated expansion of space. From the fit we find a value of 0.98 for κa . With the value $a = 1.29 \cdot 10^{28} \text{cm}$ and $H = 70 \text{km/sec/Mpc}$, as derived from red shift data, assuming a mean grain radius of $1 \mu\text{m}$ and a density of 2g/cm^3 for the dust grains, their total mass fraction in the universe is about 0.3%, just the same order as in the solar system and as it is confirmed by measurements of metallicity of the gas in galaxy clusters.

Appendix F

Jeans-instability

In the year 1902 Jeans has shown that perturbations of the equilibrium between pressure force and gravitational attraction can lead to instabilities. Jeans' considerations were based on the assumption that in homogeneous mass filled space there exists a constant pressure and a constant gravitational potential, a proposition, which at that time, more than a decade before Einstein's theory of general relativity, was in clear contradiction to the Poisson equation of Newtonian theory of gravitation

$$\Delta\Phi = 4\pi G\rho_0, \tag{F.1}$$

which was considered as generally valid and which for $\Phi = \text{const.}$ only allows the trivial solution $\rho_0 = 0$.

The reason is that the Poisson equation does not contain an energy term with repulsive action. All the new theories contain such terms, be it Einstein's cosmological constant, be it the kinetic term represented by the expansion of the universe, or be it, as considered here, the negative pressure of potential energy.

If like Jeans we start from an undisturbed matter distribution of density ρ_0 and discuss the development of a perturbation of the form $\rho = \rho_0 + \rho_1(\mathbf{r})$, The perturbation of the potential $\Phi_1(\mathbf{r})$ is given by

$$\Delta\Phi_1 = 4\pi G\rho_1. \tag{F.2}$$

Together with the equation of continuity

$$\frac{\partial\rho_1}{\partial t} + \rho_0\nabla\mathbf{u}_1 = 0 \tag{F.3}$$

and the equation of motion

$$\rho_0\frac{\partial\mathbf{u}_1}{\partial t} = -c_s^2\nabla\rho_1 - \rho_0\nabla\Phi_1 \tag{F.4}$$

the potential equation forms a complete system to determine the quantities ρ_1 , Φ_1 and the local flow velocity \mathbf{u}_1 . c_s is the velocity of sound. For an isothermal perturbation in an ideal gas we have $c_s^2 = k_B T/m$. Considering in this system

a perturbation with the Fourier component $\exp(i\mathbf{k}\mathbf{x} - i\omega t)$, from Eqs.(F.2-F.4) we get

$$-\omega\rho_1 + \rho_0\mathbf{k}\mathbf{u}_1 = 0, \quad (\text{F.5})$$

$$-\rho_0\omega\mathbf{u}_1 = -c_s^2\mathbf{k}\rho_1 - \rho_0\mathbf{k}\Phi_1, \quad (\text{F.6})$$

$$-\mathbf{k}^2\Phi_1 = 4\pi G\rho_1. \quad (\text{F.7})$$

and from this the dispersion relation

$$\omega^2 = c_s^2(k^2 - k_J^2) \quad \text{with} \quad k_J^2 = \frac{4\pi G\rho_0}{c_s^2}. \quad (\text{F.8})$$

For $k < k_J$ ω is imaginary. This means that the perturbation grows exponentially. For perturbations on a characteristic length scale of more than $l_J = 2\pi/k_J$ pressure forces cannot damp out gravitational action and the system becomes unstable. For a radially symmetric perturbation of the matter distribution with radius $R_J = l_J/4$ and a mean density ρ_0 the minimum involved mass, the so called 'Jeans mass' is

$$M_J = \frac{2}{3}\pi^{5/2} \left(\frac{k_B T}{Gm} \right)^{3/2} \rho_0^{-1/2}. \quad (\text{F.9})$$

From these formulas we find for the stability limit of the intergalactic plasma ($k_B T = 100$ keV, $\rho_0 = 4 \cdot 10^{-30}$ g/cm³) $R_J = 170$ Mpc, which corresponds to the largest observed netlike structures in the universe.

With the assumption that the individual regions cool down isobaric, they reach the recombination limit of about $k_B T = 10$ eV, at which a further fragmentation can be expected, at a density of $4 \cdot 10^{-26}$ g/cm³. This corresponds to a stability limit of $R_J = 1.7$ Mpc and a mass of $2.5 \cdot 10^{43}$ g, the typical mass of a galaxy.

If the gas cools isobarically down to about 10 K, the density would be of the order 10^{-22} g/cm³. The critical mass in this case is $5 \cdot 10^{35}$ g, corresponding to the mass of the largest stars, The distance between these stars would be about $2R_J = 6$ pc, corresponding to the observed regions of star formation. Of course this extrapolation into the range of star formation should be looked at with caution, as here the dynamics of the dust component is of considerable influence.

Appendix G

The mass balance of galaxy clusters

Clusters of galaxies are the largest observable structures in the universe. Besides of galaxies they contain large amounts of hot gas. But if we trust the concept of the 'concordance model' the overwhelming fraction is 'dark matter'. The necessity of a large amount of this dark matter follows from the discrepancy between estimations of the total mass, which results from the sum of luminous matter in galaxies and of the gas mass, radiating in x-rays, and the mass, which appears necessary to explain, why the galaxies form a bound system in spite of their high peculiar velocities.

Basis of most mass estimations are the properties of the cluster gas. The density of the cluster gas can be estimated from the intensity of bremsstrahlung (see Eq.(D.2)). Under the assumption that the matter inside the cluster is in stationary equilibrium from the balance of pressure forces and gravitational attraction, the total mass can be estimated. For a spherically symmetric matter distribution we have

$$\frac{dp}{dr} = -G \frac{\rho_{gas}}{r^2} M(r). \quad (G.1)$$

G is the gravitational constant, $M(r)$ the mass enclosed in a sphere of radius r . The relation between pressure and density for an ideal gas is given by $p = \rho_{gas} k_B T / m$. m is the mean mass per gas particle, for a fully ionised hydrogen gas with 25% mass fraction of helium about $0.6 m_p$ (proton mass).

Already with the application of this equation our view is different from conventional concepts. Especially the experimental determination of the temperature profile is influenced in many cases by the expectations of the model upon which they are based. For the outer region the plasma temperature can scarcely be determined spectroscopically, as the measured signals are small compared to the background level. On the other side measurements always give only a signal, projected through layers with varying properties. Transforming the signals to radial profiles induces additional errors. Besides that, most available measure-

ments are restricted to the energy range $< 10\text{keV}$, in which the dependence of the bremsstrahlung spectrum on temperature is very weak, especially at high temperatures.

Conventional theories assume that the cluster gas has been introduced from outside as cold gas, has gained its thermal energy from the changed gravitational potential and is cooled by radiation only in the dense centre. Thus most derivations assume a decreasing or in some cases constant temperature in the outskirts. If we start with the assumption of a hot intergalactic plasma, we expect a continuously increasing temperature from centre to the outside. Measurements in the central region of most clusters show an increasing temperature, corresponding to $T \propto \rho^{-\gamma}$ with $\gamma = 0.2$ and changing to $\gamma = 0.1$ at larger radii. We suppose that this increase continues also in the region, where accurate measurements are not possible. In contrast, conventional theory assumes values from $\gamma = 0$ (isothermal) to $\gamma = -2/3$ (adiabatic heating) for the outskirts.

Still more important is the assumption that the plasma can be treated as an ideal gas. Measurements of clusters in our 'vicinity' (Coma) show, however, that the plasma is permeated by magnetic fields of the order $B = 1\mu G$. That means that the motion of charged particles is strongly coupled to the direction of the magnetic field lines. The gyration radius $r_B = mvc/eB$ is of the order 10^7cm for electrons, but also for ions with 10^{10}cm many orders of magnitude less than the mean free path, which in a plasma of the density 10^{-3}cm^{-3} and a temperature of 10^8K is more than 10^{23}cm . For all transport phenomena only the component of gradients in direction of the magnetic field contributes to the transport. In a plasma, in which the coherence length of the fields is small compared to the dimensions of the plasma, on average only in one third of the plasma the direction of gradients is identical with the direction of the fields. Transport rates are thus reduced to $1/3$, compared to those in an ideal gas. For energy transport this effect has been discussed in many papers. But it is valid also for mass transport by pressure gradients, as the ions are firmly bound to the field lines, too. Thus, compared to an ideal gas, in a magnetised plasma we have to introduce an additional factor $\xi = 1/3$ in the balance equation (G.1).

To determine the matter distribution of the gas and of the total mass, we start from the measured profiles of x-ray emission. Disregarding the very central part, where the distribution may be strongly disturbed by concentration of gas around a large central galaxy (cD galaxy), the projected emission profiles are well described by a so called ' β profile':

$$I(x) = \frac{I_0}{[1 + (x/r_c)^2]^{\beta_x}} \quad (\text{G.2})$$

The values of β_x normally are between 1.0 and 1.3. In this case inversion of the projected to the radial profile gives such a ' β profile', too, but now with the exponent $\beta_r = \beta_x + 1/2$. From Eq.(D.2) thus, regarding the dependency $T \propto \rho^{-\gamma}$, we find the expression for the density distribution of the gas:

$$\rho(r) = \frac{\rho_0}{[1 + (r/r_c)^2]^\alpha} \quad \text{with} \quad \alpha = \frac{\beta_x + 1/2}{2 - \gamma/2} \quad (\text{G.3})$$

For some historical reason most astrophysicists use the form

$$\rho(r) = \frac{\rho_0}{[1 + (r/r_c)^2]^{\frac{3\beta}{2}}}. \quad (\text{G.4})$$

The values of β for $\gamma = 0.2$ for this definition are between 0.53 and 0.64. For an isothermal gas the values are higher by 5%, in the limiting case of adiabatic heating higher by 40%.

The density ρ_0 follows from integration of Eq.(D.2) over the spectral range of the measurement and resolution of the equation with respect to the density.

$$\left(\frac{\rho_0}{\text{gcm}^{-3}}\right)^2 = 7.3 \cdot 10^{-25} \frac{\varepsilon}{\text{erg/cm}^3\text{s}} / \sqrt{\frac{kT_0}{\text{keV}}} \quad (\text{G.5})$$

The resulting density data are higher by a factor $\sqrt{T/T_0}$ for a temperature profile, increasing to the outside, than for a isothermal profile of temperature T .

The total density follows from Eq.(G.1). Including the reduction of the pressure gradient by the turbulent magnetic field with

$$\frac{dp}{dr} = \xi \frac{d}{dr}(\rho_{gas} kT/m) \quad (\text{G.6})$$

we get

$$M(r) = -\frac{\xi kT_0}{Gm} r^2 (1 + (r/r_c)^2)^\alpha \frac{d}{dr} \frac{1}{(1 + (r/r_c)^2)^{\alpha(1-\gamma)}} \quad (\text{G.7})$$

or

$$M(r) = \frac{2\alpha(1-\gamma)\xi kT_0 r^3}{Gm(1 + (r/r_c)^2)^{1-\alpha\gamma}} \quad (\text{G.8})$$

To fulfil the condition that for large r the gas mass is the dominant component of the total mass, the exponent of Eq.(G.8) and Eq.(G.3) must be the same. That means: $1 - \alpha\gamma = \alpha$. For $\gamma = 0.1$ to 0.2 the resulting values are $\alpha = 0.91$ to 0.83 (or $\beta = 0.61$ to 0.56). This is exactly the range of values, which is confirmed by observations.

Compared to equilibrium of an isothermal ideal gas thus for the central region the total mass is reduced by a factor of $\xi T_0/T$. But now we must put the question, if this mass is sufficient to keep the galaxies with their velocities in the range 1000 km/s bound by the gravitational potential of the cluster plasma. The potential distribution can immediately found by integration from the equilibrium relation Eq.(G.1).

$$\frac{d\Phi}{dr} = -\frac{1}{\rho} \frac{dp}{dr}. \quad (\text{G.9})$$

Introducing the corresponding ' β -profiles' of ρ and p with the abbreviation $x = r/r_c$ we get

$$\frac{d\Phi}{dx} = \frac{\xi kT_0}{m} \frac{2\alpha(1-\gamma)x}{(1+x^2)^{1-\alpha\gamma}}, \quad (\text{G.10})$$

and from this by integration

$$\Phi - \Phi_0 = \frac{\xi k T_0}{m} \frac{\alpha(1-\gamma)}{\alpha\gamma} [(1+x^2)^{\alpha\gamma} - 1]. \quad (\text{G.11})$$

or

$$\Phi - \Phi_0 = \frac{\xi(1-\gamma)}{\gamma} \frac{k(T-T_0)}{m}. \quad (\text{G.12})$$

With $\xi = 1/3$ and $\gamma = 0.1$ we find for a galaxy with an initial radial velocity $\sigma_r = 1000\text{km/s}$ that the potential difference, necessary to slow it down to zero, is

$$k(T-T_0) = \frac{m}{3} \frac{\sigma_r^2}{2}, \quad (\text{G.13})$$

that means a difference of about 1 keV. On the basis of the empirically confirmed relation between velocity dispersion and mean gas temperature, $k\bar{T} = m\sigma_r^2/2$, the condition (G.13) is fulfilled for $\gamma = 0.1$ at $r = 5.3r_c$, for $\gamma = 0.2$ at $r = 3.5r_c$. This is just the region, where the observed density of galaxies in a cluster drops considerably.

Appendix H

Idea of a symmetric model of elementary particles

Under the proposition that spin of elementary particles is not a property of the constituents, but results from the type of coupling between them, the complete system of elementary particles can be described by a symmetric combination of subelementary fields (quarks) and their antifields. To classify all elementary particles the following rules are sufficient:

1. Quarks have a quantum of electric charge or a quantum of strangeness or both.
2. For every quark there exists an antiquark with the opposite quantum numbers.
3. All particles consist of equal numbers of quarks and antiquarks.
4. No two quark can occupy the same quantum state.
5. Quark couplings can be of "fermionic" or 'bosonic' type.

Fermionic coupling means that the constituent fields are superimposed at the same location with the resulting spin quantum number 1/2. Bosonic coupling means that the constituent fields form a spatially extended system, which can attain different integer rotational quantum numbers.

The conditions described above allow three different quarks, which in the following are denoted as a, b, and c. They are defined by the quantum numbers of charge and strangeness:

$$a = \begin{pmatrix} 1 \\ -1 \end{pmatrix} \quad b = \begin{pmatrix} 0 \\ -1 \end{pmatrix} \quad c = \begin{pmatrix} 1 \\ 0 \end{pmatrix} \quad (\text{H.1})$$

As all elementary particles contain equal numbers of quarks and antiquarks, both systems of quantum numbers are not fixed, but may be varied by an

additional constant. Subtraction of 2/3 from the charge quantum numbers and adding 1 to the strangeness quantum numbers leads (apart from spin) to the triplet of antiquarks \bar{u} , \bar{d} , and \bar{s} of conventional theory. The negative sign of the strangeness quantum number has been introduced to remain in agreement with conventional notation.

From combination of two constituents fermionic coupling leads to leptons, bosonic coupling to mesons.

Quark-configuration	Charge	Strangeness	Lepton	Meson(Spin 0)
$a\bar{a}$	0	0	ν_μ	π^0
$a\bar{b}$	1	0	μ^+	π^+
$a\bar{c}$	0	-1	ν_e	$\bar{K}^0, \bar{\kappa}$
$b\bar{a}$	-1	0	μ^-	π^-
$b\bar{b}$	0	0	ν_μ	η
$b\bar{c}$	-1	-1	e^-	K^-
$c\bar{a}$	0	1	$\bar{\nu}_e$	K^0, κ
$c\bar{b}$	1	1	e^+	K^+
$c\bar{c}$	0	0	ν_μ	η'

The meson denoted as κ in the table, a hypothetical meson of very low mass, should exist as a bosonic analog to the electron neutrino. The existence of such a particle would explain the observed violation of parity and strangeness conservation in weak interactions. K^0 mesons can be regarded as an excited state of the κ meson.

The baryon spectrum results from the combination of two quarks and two antiquarks with the two quarks forming a fermionically bound 'nucleus', surrounded by a bosonically bound 'shell' of two antiquarks. For the fermionically bound 'nucleus' the exclusive condition requires that the two quarks have to be different. Thus only the three combinations (ab), (ac), and (bc) are possible. For the antiquarks of the shell the exclusion principle is important only, if both are in the same quantum state. Thus exactly the same combinations of quantum states are possible, which are found in the conventional quark model for the ground state octet and singlet. The excited states form complete decuplets. The ground state multiplet consists of the following particles:

Quark configuration	Charge	Strangeness	Baryon
$(ab)\bar{a}\bar{b}$	0	0	Λ
$(ab)\bar{a}\bar{c}$	-1	-1	Ξ^-
$(ab)\bar{b}\bar{c}$	0	-1	Ξ^0
$(ac)\bar{a}\bar{b}$	1	1	p
$(ac)\bar{a}\bar{c}$	0	0	Σ^0
$(ac)\bar{b}\bar{c}$	1	0	Σ^+
$(bc)\bar{a}\bar{b}$	0	1	n
$(bc)\bar{a}\bar{c}$	-1	0	Σ^-
$(bc)\bar{b}\bar{c}$	0	0	Σ^0

The strangeness numbers in the table differ from conventional notation by 1, they correspond to the quantum number normally denoted as 'hypercharge'.

Excited states are possible with three additional combinations of quantum numbers, which result from the configurations $(ab)\bar{c}\bar{c}'$, $(ac)\bar{b}\bar{b}'$ and $(bc)\bar{a}\bar{a}'$. These configurations have the quantum numbers $(-1,-2)$, $(2,-1)$ und $(-1,-1)$ and correspond to Ω^- , Δ^{++} , and Δ^- .

As an additional combination of four constituents there exists the bosonic bond of a pair of two fermionically bound quarks with a pair of two fermionically bound antiquarks. Such particles should exhibit similar properties as mesons, but with the difference that here the two bosonically bound building stones are real fermions. Due to the double number of constituents the mass of these particles should be much higher than that of normal mesons. To the family of these 'hypermesons' we can assign all the mesons, to which properties as 'charm' and 'beauty' are attributed by conventional theory, as e.g. the mesons J/ψ und Y .

W and Z particles can be interpreted as fermionic bonds of two quarks and two antiquarks.