## Fedosov Quantization of Lagrange–Finsler and Hamilton–Cartan Spaces and Einstein Gravity Lifts on (Co) Tangent Bundles

Mihai Anastasiei\*

Faculty of Mathematics, University "Al. I. Cuza" Iaşi, 11, Carol I Boulevard, Iaşi, Romania, 700506

and

Mathematical Institute "O. Mayer", Romanian Academy Iaşi Branch, 8, Carol I Boulevard, Iaşi, Romania, 700506

#### Sergiu I. Vacaru<sup>†</sup>

The Fields Institute for Research in Mathematical Science 222 College Street, 2d Floor, Toronto M5T 3J1, Canada

and

Faculty of Mathematics, University "Al. I. Cuza" Iaşi, 11, Carol I Boulevard, Iaşi, Romania, 700506

November 9, 2008

#### Abstract

We provide a method of converting Lagrange and Finsler spaces and their Legendre transforms to Hamilton and Cartan spaces into almost Kähler structures on tangent and cotangent bundles. In particular cases, the Hamilton spaces contain nonholonomic lifts of (pseudo) Riemannian / Einstein metrics on effective phase spaces. This allows us to define the corresponding Fedosov operators and develop deformation quantization schemes for nonlinear mechanical and gravity models on Lagrange– and Hamilton–Fedosov manifolds.

**Keywords:** Deformation quantization; quantum gravity; Finsler, Lagrange, Hamilton and Cartan spaces, almost Kähler geometry.

<sup>\*</sup>anastas@uaic.ro

<sup>&</sup>lt;sup>†</sup>Sergiu.Vacaru@gmail.com; http://www.scribd.com/people/view/1455460-sergiu

MSC: 83C99, 53D55, 53B40, 53B35 PACS: 04.20.-q, 02.40.-k, 02.90.+g, 02.40.Yy

## 1 Introduction

To construct a quantum theory for a given classical model is usually understood that it is necessary to elaborate a quantization procedure adapted to certain fundamental field equations and corresponding Lagrangians/ Hamiltonians and theirs symmetries, constraints and locality. In various attempts to develop quantum versions of gravity and nonlinear physical theories, one provided different computation schemes when it is supposed that all constraints can be solved, gauge symmetries can be represented by shift symmetries and there are certain canonical forms for the Poisson brackets. Nevertheless, such quantization usually destroy global symmetries, locality and may result into violation of local Lorentz symmetry.

There were proposed different sophisticate constructions with formal and partial solutions for quantum gravity and field interactions theories. We cite here the BRST quantization methods for non–Abelian and open gauge algebras [1, 2, 3, 4], deformation quantization [5, 6, 7, 8], quantization of general Lagrange structures and, in general, BRST quantization without Lagrangians and Hamiltonians [9, 10], W–geometry and Moyal deformations of gravity via strings and branes [11, 12, 13] and quantum loops and spin networks [14, 15, 16].

In the above–mentioned approaches, it is necessary to quantize curved phase spaces in a manner that is explicitly covariant on phase–space coordinates. Indeed, for instance, the key ingredients of Fedosov and BRST methods, see [9, 10, 17, 18] and references therein, is the embedding of the system into the cotangent bundle over its phase space. There were also elaborated such models following standard geometric constructions over Riemannian manifolds and formal schemes with symplectic groupoids and contravariant connections [17, 19].

A rigorous geometric approach to deformation quantization of gravity, gauge theories and geometric mechanics models with constraints and related generalized Lagrange–Finsler theories, see [20, 21, 22, 23, 24], shows that the quantization schemes have to be developed for nonholonomic manifolds<sup>1</sup> and tangent and cotangent bundles endowed with nonlinear connection (N– connection) structure. The natural step in this direction is to apply the

<sup>&</sup>lt;sup>1</sup>i.e. manifolds endowed with nonholonomic (equivalent, anholonomic, or nonintegrable distributions), see details and references in [25, 26, 27, 28, 29, 30]

methods of the geometry of Hamilton and Cartan spaces and generalizations [31, 32] (such spaces are respectively dual to the Lagrange and Finsler spaces and generalizations [33, 34]).

The aim of this work is to show how Karabegov's approach to Fedosov deformation quantization [19, 35, 36] can be naturally extended for almost Kähler manifolds endowed with canonical geometric structures generated by semi–Riemannian and/or Einstein metrics and Lagrange–Finsler and Hamilton–Cartan fundamental generating functions.

This paper is motivated by the following results: In Refs. [30, 21, 22, 37, 29], we concluded that classical and quantum gravity models on (co) tangent bundles positively result in generalized Finsler like theories with violation of local Lorentz symmetry. The conclusion was supported also by a series of works on definition of spinors and field interactions on (in general, higher order) locally anisotropic spacetimes [38, 39], on low energy limits of (super) string theory [40, 41, 42] and possible Finsler like phenomenological implications and symmetry restriction of quantum gravity [43, 44, 45]. Here, we emphasize that the nonholonomic quantum deformation formalism can be re-defined for nonholonomic (pseudo) Riemannian, or Riemann-Cartan, manifolds with fibred structure. Such nonholonomic spaces, under well defined conditions and for corresponding Lagrange–Finsler variables, possess local Lorentz invariance for classical theories and seem to preserve it for nonholonomic quantum deformations, see results from [23, 24].

The work is organized as follows: In Section 2 we recall the basis of the Lagrange–Finsler and Hamilton–Cartan geometry, nonholonomic lifts of Einstein metrics on cotangent bundles and almost Kähler models of such spaces. We also collect there some geometric constructions that we need for further considerations. Section 3 is devoted to the formalism of canonical nonlinear connections and distinguished connections induced by Lagrange and Hamilton fundamental functions. We introduce the concept of Hamilton–Fedosov spaces and define the corresponding almost symplectic structures. In Section 4 we consider a generalization of the concept of connection to that of connection-pair resulting in Fedosov-Hamilton operatorspairs which is necessary for definition of deformation quantization models being invariant under symplectic morphisms and Legendre transform. We provide Fedosov's theorems for connection-pairs which allows us to develop an approach to geometric quantization of Hamilton and Einstein (generalized on cotangent bundles) spaces in Section 5. We speculate on possible quantum generalizations of gravitational field equations on phase spaces and their deformation quantization. Finally, in Section 6 we conclude the results.

## 2 Lagrange–Finsler and Hamilton–Cartan Geometry and Einstein Spaces

In this section, we outline some results from the geometry of Lagrange– Finsler [33, 34] and Hamilton–Cartan [31, 32] spaces.

# 2.1 Canonical geometric objects on Lagrange and Hamilton spaces

Let us consider a real sufficiently smooth manifold M, dim  $M = n \geq 2$ . We label the local coordinates  $x = \{x^i\}$ , with "base" indices  $i, j, \ldots = 1, 2, \ldots n$ , and write by TM and  $T^*M$ , respectively, the total spaces of tangent and cotangent bundles  $\pi : TM \to M$  and  $\pi^* : T^*M \to M$ , with local coordinates  $u = (x, y) = \{u^{\alpha} = (x^i, y^a)\}$ , for "fiber" indices  $a, b, \ldots = n + 1, \ldots, n + n$  on any local cart  $U \subset TM$ , and  $*u = (x, p) = \{*u^{\alpha} = (x^i, p_a)\}$  on any local cart  $*U \subset T^*M$ , when  $p_a$  are dual to  $y^a$ . In order to apply the Einstein summation rule for contracting base and fiber indices, we shall use identifications of type  $y^i \doteq y^{n+i}$ , i.e. we suppose that indices like  $a, b, \ldots$  can split into respective n + i, n + j.

**Definition 2.1** A generalized Hamilton space is defined by a pair  $GH^n = (M, g^{ij}(x, p))$ , where  $g^{ij}(x, p)$  is a contravariant symmetric tensor field, nondegenerate and of constant signature on  $\widetilde{T^*M} = T^*M/\{0\}$ , for  $\{0\}$  being the null section of  $T^*M$ .

We note that a contravariant tensor of type

$$g^{ij}(x,p) = e^{i}{}_{i'}(x,p)e^{j}{}_{j'}(x,p)g^{i'j'}(x)$$
(1)

includes, for some vielbein fields  $e^i{}_{i'}$ , the dual tensor  $g^{i'j'}(x)$  as the inverse of a (semi) Riemann metric  $g_{i'j'}(x)$  on M. In this paper, we consider that any classical solution of the Einstein equations defines a space-time manifold M and a corresponding deformation quantization procedure on  $T^*M$  defines the values  $e^i{}_{i'}$  and other fundamental quantum geometric objects (like almost symplectic structure and generalized connection) with nontrivial dependence on variables  $p_a$ . In general, we shall provide our constructions for (pseudo) Riemannian spaces with (co) metrics of type  $g^{i'j'}(x)$  and  $g^{ij}(x,p)$ and discuss if there are any important particular properties for Einstein manifolds (spaces) when  $g_{i'j'}(x)$  is subjected to the condition to solve on Mthe gravitational field equations with nonzero cosmological constant  $\lambda$ ,

$$R_{i'j'}(x) = \lambda g_{i'j'}(x), \tag{2}$$

where  $R_{i'j'}(x)$  is the Ricci tensor for the Levi–Civita connection  $\nabla = \{ \Gamma_{j'k'}^{i'}(x) \}$  completely defined by  $g_{i'j'}$ .

For simplicity, we shall work with a more particular class of spaces when  $g^{ij}(x,p)$  is defined by a Hamilton function H(x,p):

**Definition 2.2** A Hamilton space  $H^n = (M, H(x, p))$  is defined by a function  $T^*M \ni (x, p) \to H(x, p) \in \mathbb{R}$ , i.e. by fundamental Hamilton function, which is differentiable on  $\widehat{T^*M}$  and continuous on the null section  $\pi^* : T^*M \to M$  and such that the (Hessian, equivalently, fundamental) tensor field

$$^{*}g^{ab}(x,p) = \frac{\partial^{2}H}{\partial p_{a}\partial p_{b}}$$
(3)

is non-degenerate and of constant signature on  $\widetilde{T^*M}$ .

Let L(x, y) be a regular differentiable Lagrangian on  $U \subset TM$ , with non-degenerate Hessian (equivalently, fundamental tensor field)

$$g_{ab}(x,y) = \frac{\partial^2 L}{\partial y^a \partial y^b}.$$
(4)

**Definition 2.3** A Lagrange space  $L^n = (M, L(x, y))$  is defined by a function  $TM \ni (x, y) \to L(x, p) \in \mathbb{R}$ , i.e. a fundamental Lagrange function, which is differentiable on  $\widetilde{TM}$  and continuous on the null section of  $\pi : TM \to M$  and such that the (Hessian) tensor field  $g_{ab}(x, y)$  (4) is non-degenerate and of constant signature on  $\widetilde{TM}$ .

We can define the Legendre transform  $L \to H$ ,

$$H(x,p) = p_a y^a - L(x,y), \tag{5}$$

where  $y = \{y^a\}$  are solutions of the equations  $p_a = \partial L(x, y) / \partial y^a$ , and (inversely) the Legendre transform  $H \to L$ ,

$$L(x,y) = p_a y^a - H(x,p),$$
(6)

where  $p = \{p_a\}$  is the solution of the equations  $y^a = \partial H(x, p) / \partial p_a$ .<sup>2</sup>

 $<sup>^{2}</sup>$ In some monographs (for instance, see [34, 31]), it is considered the factor 1/2 in the right sides of (3) and (4). We emphasize that in this paper (for simplicity) the Hamilton and Lagrange functions will be supposed to be regular and related mutually by Legendre transforms.

Following terminology from [31], we say that  $N_i^a$  (14) and  $*N_{ij}$  (15) are  $\mathcal{L}$ -dual if L and H are related by Legendre transform (5), or (6). In the following constructions, we shall consider that to Legendre transform there are associated the diffeomorphisms

$$\varphi: TM \supset U \rightarrow {}^{*}U \subset T^{*}M, (x^{i}, y^{a}) \rightarrow \left(x^{i}, p_{a} = \frac{\partial L(x, y)}{\partial y^{a}}\right)$$

and

$$\psi: T^*M \supset \ ^*U \to U \subset TM, (x^i, p_a) \to \left(x^i, y^a = \frac{\partial H(x, p)}{\partial p_a}\right),$$

allowing to define respectively pull-back and push-forward of geometric objects (functions, vectors, differential forms, connections, tensors...) from  ${}^*U$  to U and from U to  ${}^*U$ , i.e. we define  $\mathcal{L}$ -dual geometric objects. For instance, for a differentiable function  ${}_1f$  on U, we define a differentiable function  ${}_1f^* \doteq {}_1f \circ \psi = {}_1f \circ \varphi^{-1}$  on  ${}^*U$  and (inversely) for a differentiable function  ${}^2f$  on  ${}^*U$ , we have a differentiable  $\left({}^2f\right)^0 \doteq {}^2f \circ \varphi = {}^2f \circ \psi^{-1}$  on U. Note that  $H = L \circ \varphi^{-1}$  and  $L = H \circ \psi^{-1}$ . Similarly, for any vector field X on U, we get a vector field  ${}^*X \doteq T\varphi \circ X \circ \varphi^{-1} = T\psi^{-1} \circ X \circ \psi$  on  ${}^*U$  and (inversely) for a vector field  ${}^*X \circ \varphi$  on U, we get a vector field  ${}^*X \circ \varphi = {}^2f \circ \varphi^{-1} = T\psi^{-1} \circ X \circ \psi$  on  ${}^*U$  and (inversely) for a vector field  ${}^*X \circ \varphi = {}^2f \circ \varphi^{-1} = T\psi^{-1} \circ X \circ \psi$  on  ${}^*U$  and (inversely) for a vector field  ${}^*X \circ \varphi = {}^2f \circ \varphi^{-1} = T\psi^{-1} \circ X \circ \psi$  on  ${}^*U$  and (inversely) for a vector field  ${}^*X \circ \varphi = {}^2f \circ \varphi^{-1} = T\psi^{-1} \circ X \circ \psi$  on  ${}^*U$  and (inversely) for a vector field  ${}^*X \circ \varphi = {}^2f \circ \varphi^{-1} = T\psi^{-1} \circ X \circ \psi$  on  ${}^*U$  and (inversely) for any form  ${}^*\omega \circ \varphi = {}^2f \circ \varphi^{-1} = T\psi^{-1} \circ X \circ \psi$  on  ${}^*U$  and (inversely) for any form  ${}^*\omega \circ \varphi = {}^2f \circ \varphi^{-1} = T\psi^{-1} \circ {}^*\omega \circ \psi$  on  ${}^*U$  and (inversely) for any form  ${}^*\omega \circ \varphi = {}^2f \circ \varphi^{-1} = T\psi^{-1} \circ {}^*\omega \circ \psi$  on  ${}^*U$  and (inversely) for any form  ${}^*\omega \circ \varphi = {}^2f \circ \varphi^{-1} = T\psi^{-1} \circ {}^*\omega \circ \psi$  on  ${}^*U$  and (inversely) for any form  ${}^*\omega \circ \varphi = {}^2f \circ \varphi^{-1} = T\psi^{-1} \circ {}^*\omega \circ \psi$  on  ${}^*U$  and (inversely) for any form  ${}^*\omega \circ \varphi = {}^2f \circ \varphi = {}^2f \circ \varphi = {}^2f \circ \varphi = {}^2f \circ \psi = {}$ 

Let vTM and  $vT^*M$  be the vertical distributions on TM and  $T^*M$ , respectively.

#### **Definition 2.4** Any Whitney sums

$$TTM = hTM \oplus vTM \tag{7}$$

and

$$TT^*M = hT^*M \oplus vT^*M \tag{8}$$

define respectively nonlinear connection (N-connection) structures parametrized by the local vector fields

$$e_i = \frac{\partial}{\partial x^i} - N_i^a(x, y) \frac{\partial}{\partial y^a}$$
 on  $TM$ 

and

$$*e_i = \frac{\partial}{\partial x^i} + *N_{ia}(x,p)\frac{\partial}{\partial p_a}$$
 on  $T^*M$ .

One says that a N-connection defines on TM, or  $T^*M$ , a conventional horizontal (h) and vertical (v) splitting (decomposition).

Let consider a regular curve  $c(\tau)$  with real parameter  $\tau$ , when  $c : \tau \in [0,1] \to x^i(\tau) \subset U$ . It can be lifted to  $\pi^{-1}(U) \subset \widetilde{TM}$  as  $\widetilde{c}(\tau) : \tau \in [0,1] \to (x^i(\tau), y^i(\tau) = \frac{dx^i}{d\tau})$  since the vector field  $\frac{dx^i}{d\tau}$  does not vanish on  $\widetilde{TM}$ . Following techniques from variational calculus, one proves:

**Theorem 2.1** The Euler–Lagrange equations,

$$\frac{d}{d\tau}\frac{\partial L}{\partial y^i} - \frac{\partial L}{\partial x^i} = 0,\tag{9}$$

are equivalent to the Hamilton-Jacobi equations,

$$\frac{dx^{i}}{d\tau} = \frac{\partial H}{\partial p_{i}} \text{ and } \frac{dp_{i}}{d\tau} = -\frac{\partial H}{\partial x^{i}}, \tag{10}$$

and to the nonlinear geodesic (semi-spray) equations

$$\frac{d^2x^i}{d\tau^2} + 2G^i(x,y) = 0,$$
(11)

where

$$G^{i} = \frac{1}{2}g^{ij} \left(\frac{\partial^{2}L}{\partial y^{j}\partial x^{k}}y^{k} - \frac{\partial L}{\partial x^{j}}\right),$$

for  $g^{ij}$  being the inverse to  $g_{ij}$  (4).

Let us consider on  $T^*M$  the canonical symplectic structure

$$\theta \doteq dp_i \wedge dx^i. \tag{12}$$

The Hamiltonian H defines an unique vector field on  $T^*M$ :

$$X_H = \frac{\partial H}{\partial p_i} \frac{\partial}{\partial x^i} - \frac{\partial H}{\partial x^i} \frac{\partial}{\partial p_i}$$

by the equation

$$i_{X_H}\theta = -dH,$$

where  $i_{X_H}$  denotes the interior product by  $X_H$ . The same holds for any function on  $T^*M$ .

By Theorem 2.1 one has:

Corollary 2.1 The Hamilton-Jacobi equations (10) are equivalent to

$$\frac{dx^i}{d\tau} = \{H, x^i\} \text{ and } \frac{dp_a}{d\tau} = \{H, p_a\},\$$

where the Poisson structure is defined by brackets

$$\{ {}^{1}f, {}^{2}f \} = \theta(X_{1f}, X_{2f})$$
(13)

for any functions  ${}^{1}f(x,p)$  and  ${}^{2}f(x,p)$  on  $T^{*}M$ .

**Proof.** It can be obtained by a standard calculus in geometric mechanics.  $\Box$ 

The following theorem holds:

**Theorem 2.2** The are canonical N-connections defined respectively by regular Lagrange L(x, y) and/ or Hamilton H(x, p) fundamental functions:

$$N_i^a \doteqdot \frac{\partial G^a}{\partial y^i} \tag{14}$$

and

$$^{*}N_{ij} \doteq \frac{1}{2} \left[ \{ ^{*}g_{ij}, H \} - \frac{\partial^{2}H}{\partial p_{k}\partial x^{(i}} ^{*}g_{j)k} \right],$$
(15)

where,  $*g_{ij}$  is the inverse to  $*g^{ij}(3)$  and, for instance,  $a_{(ij)} = a_{ij} + a_{ji}$  denotes symmetrization of indices.

**Proof.** We can verify respectively that on any open sets  $U \subset TM$  and  $U^* \subset T^*M$  coefficients (14) and (15) satisfy the conditions of Definition 2.4. For details see Ch. 9 in [34] and Ch. 5 in [31].  $\Box$ 

On (co) tangent bundles endowed with N-connection structure, it is convenient to elaborate a covariant calculus adapted to this structure, i.e. preserving the conventional splitting of tensors and other geometric objects (like connections, differential forms etc) into horizontal (h) and vertical (v) components. In brief, such distinguished (by N-connection) components are called respectively d-objects, d-field (for some physical fields of tensor, spinor nature ...), d-tensors, d-vectors, d-forms, d-connections etc, see details in Refs. [31, 32, 33, 34, 38, 29]. **Proposition 2.1** There are canonical frame structures (local N-adapted (co-)bases) defined by canonical N-connections:

$$\mathbf{e}_{\alpha} = (\mathbf{e}_{i} = \frac{\partial}{\partial x^{i}} - N_{i}^{a} \frac{\partial}{\partial y^{a}}, e_{b} = \frac{\partial}{\partial y^{b}}), \text{ on } TM,$$
(16)

$$^{*}\mathbf{e}_{\alpha} = ( ^{*}\mathbf{e}_{i} = \frac{\partial}{\partial x^{i}} + ^{*}N_{ia}\frac{\partial}{\partial p_{a}}, \ ^{*}e^{b} = \frac{\partial}{\partial p_{b}}), \ on \ T^{*}M,$$
(17)

and their dual (coframe) structures

$$\mathbf{e}^{\alpha} = (e^{i} = dx^{i}, \mathbf{e}^{b} = dy^{b} + N^{b}_{i}dx^{i}), \ on \ (TM)^{*},$$
 (18)

$${}^{*}\mathbf{e}^{\alpha} = ({}^{*}e^{i} = dx^{i}, {}^{*}\mathbf{e}_{p} = dp_{b} - {}^{*}N_{ib}dx^{i}), on (T^{*}M)^{*}, (19)$$

when  $\mathbf{e}_{\alpha} \mathbf{e}^{\beta} = \delta_{\alpha}^{\beta}$  and  $\mathbf{e}_{\alpha} \mathbf{e}^{\beta} = \delta_{\alpha}^{\beta}$ , where by  $\mathbf{e}^{\beta}$  we note the interior products and  $\delta_{\alpha}^{\beta}$  being the Kronecker delta symbol.

**Proof.** It follows by construction under the condition that such frames should depend linearly on coefficients of respective N–connections.  $\Box$ 

One says that certain geometric objects are defined on TM (or  $T^*M$ ) in N-adapted form [equivalently, in distinguished form, in brief, d-form] if they are given by coefficients defined with respect to frames  $\mathbf{e}_{\alpha}$  (16) and coframes  $\mathbf{e}^{\alpha}$  (18) and their tensor products (with respect to frames  $*\mathbf{e}_{\alpha}$  (17) and coframes  $*\mathbf{e}^{\alpha}$  (19) and their tensor products). We shall use "boldface" letters in order to emphasize that certain spaces (or geometric objects) are in N-adapted form.

**Definition 2.5** The N-lifts of the fundamental tensor fields  $*g^{ab}$  (3) and  $g_{ab}$  (4) are respectively

$${}^{*}\mathbf{g} = {}^{*}\mathbf{g}_{\alpha\beta} {}^{*}\mathbf{e}^{\alpha} \otimes {}^{*}\mathbf{e}^{\beta} = {}^{*}g_{ij}(x,p)e^{i} \otimes e^{j} + {}^{*}g^{ab}(x,p) {}^{*}\mathbf{e}_{a} \otimes {}^{*}\mathbf{e}_{b}, \quad (20)$$

on  $T^*M$ , where  $*g_{ij}$  is inverse to  $*g^{ab}$ , and

$$\mathbf{g} = \mathbf{g}_{\alpha\beta} \ \mathbf{e}^{\alpha} \otimes \mathbf{e}^{\beta} = g_{ij}(x, y)e^{i} \otimes e^{j} + g_{ab}(x, y)\mathbf{e}^{a} \otimes \mathbf{e}^{b},$$

on TM, where  $g_{ij}$  is stated by  $g_{ab}$  following  $g_{ij} = g_{n+i \ n+j}$ .

The following proposition holds:

**Proposition 2.2** The canonical N-connections N (14) and \*N (15) define respectively the canonical almost complex structures  $\mathbf{J}$ , on TM, and \* $\mathbf{J}$ , on  $T^*M$ . **Proof.** On *TM* one introduces the linear operator **J** acting on  $\mathbf{e}_{\alpha} = (\mathbf{e}_i, e_b)$  (16) as follows:

$$\mathbf{J}(\mathbf{e}_i) = -\mathbf{e}_{n+i}$$
 and  $\mathbf{J}(e_{n+i}) = \mathbf{e}_i$ .

It is clear that **J** defines globally an almost complex structure  $(\mathbf{J} \circ \mathbf{J} = -\mathbf{I})$ for **I** being the unity matrix) on *TM* completely determined for Lagrange spaces by a L(x, y). Now we provide the proof for  $T^*M$ . Let us introduce a linear operator \***J** acting on \* $\mathbf{e}_{\alpha} = (\mathbf{e}_i, \mathbf{e}^b)$  (17) following formulas

$$^{*}\mathbf{J}(^{*}\mathbf{e}_{i}) = -g_{ia}^{*}e^{n+i} \text{ and } ^{*}\mathbf{J}(^{*}e^{n+i}) = ^{*}\mathbf{e}_{i}.$$

Then **\*J** defines globally an almost complex structure ( **\*J**  $\circ$  **\*J** =  $-\mathbf{I}$  for **I** being the unity matrix) on  $T^*M$  completely determined for Hamilton spaces by a H(x, p).  $\Box$ 

**Definition 2.6** The Neijenhuis tensor field for the almost complex structure \*J on  $T^*M$ , or J on TM, defined by a N-connection (equivalently, the curvature of N-connection) is

$${}^{*J}\Omega(\mathbf{X}, \mathbf{Y}) = -[\mathbf{X}, \mathbf{Y}] + [ {}^{*}J\mathbf{X}, {}^{*}J\mathbf{Y}] - {}^{*}J[ {}^{*}J\mathbf{X}, \mathbf{Y}] - {}^{*}J[\mathbf{X}, {}^{*}J\mathbf{Y}],$$

$${}^{J}\Omega(\mathbf{X}, \mathbf{Y}) = -[\mathbf{X}, \mathbf{Y}] + [J\mathbf{X}, J\mathbf{Y}] - J[J\mathbf{X}, \mathbf{Y}] - J[\mathbf{X}, J\mathbf{Y}],$$

$$(21)$$

for any d-vectors  $\mathbf{X}$  and  $\mathbf{Y}$ .

Hereafter, for simplicity and if one shall not result in ambiguities, we shall present the N-adapted component formulas for geometric objects on  $T^*M$  (those for TM being similar), or inversely.

With respect to N-adapted bases, the components of the Neijenhuis tensor  ${}^* {}^{J}\Omega$  involve the coefficients  ${}^*\Omega_{ija}$ :

$$^{*}\Omega_{ija} = \frac{\partial ^{*}N_{ia}}{\partial x^{j}} - \frac{\partial ^{*}N_{ja}}{\partial x^{i}} + ^{*}N_{ib}\frac{\partial ^{*}N_{ja}}{\partial p_{b}} - ^{*}N_{jb}\frac{\partial ^{*}N_{ia}}{\partial p_{b}}.$$
 (22)

They define the coefficients of the N-connection curvature. One gets a complex structure i.e  ${}^* {}^{J}\Omega = 0$  under some quite complicated conditions on  $g^{ab}(x,p)$  and  $N_{ia}$  which will be not written here.

It should be noted here that the N-adapted (co-) bases (16)-(19) are nonholonomic with nontrivial anholonomy coefficients. For instance,

$$[\mathbf{e}_{\alpha}, \mathbf{e}_{\beta}] = \mathbf{e}_{\alpha} \mathbf{e}_{\beta} - \mathbf{e}_{\beta} \mathbf{e}_{\alpha} = W^{\gamma}_{\alpha\beta} \mathbf{e}_{\gamma}$$
(23)

with (antisymmetric) anholonomy coefficients  $W_{ia}^b = \partial_a N_i^b$  and  $W_{ji}^a = \Omega_{ij}^a$ , with :

$$\Omega_{ij}^{a} = \frac{\partial N_{i}^{a}}{\partial x^{j}} - \frac{\partial N_{j}^{a}}{\partial x^{i}} + N_{i}^{b} \frac{\partial N_{j}^{a}}{\partial p_{b}} - N_{j}^{b} \frac{\partial N_{i}^{a}}{\partial p_{b}}$$

#### 2.2 Almost Kähler Lagrange–Hamilton structures

We can adapt to N-connections various geometric structures on TM and  $T^*M$ . For instance, we can consider:

**Definition 2.7** One calls an almost symplectic structure on  $T^*M$  a nondegenerate N-adapted 2-form

$${}^{\mathsf{T}}\theta = \frac{1}{2} {}^{\mathsf{T}}\theta_{\alpha\beta}(u) {}^{*}\mathbf{e}^{\alpha} \wedge {}^{*}\mathbf{e}^{\beta}.$$

The following Proposition holds:

**Proposition 2.3** For any  $\tau\theta$  on  $T^*M$ , there is a unique N-connection  $*\mathbf{N} = \{ *N_{ia} \}$  satisfying the conditions:

$$^{\mathsf{T}}\theta = (h \ ^{*}\mathbf{X}, v \ ^{*}\mathbf{Y}) = 0 \ and \ ^{\mathsf{T}}\theta \doteq h \ ^{\mathsf{T}}\theta + v \ ^{\mathsf{T}}\theta, \tag{24}$$

for any  $*\mathbf{X} = h *\mathbf{X} + v *\mathbf{X}, *\mathbf{Y} = h *\mathbf{Y} + v *\mathbf{Y}$  and  $h \mathsf{T}\theta(*\mathbf{X}, *\mathbf{Y}) \doteq \mathsf{T}\theta(h *\mathbf{X}, h *\mathbf{Y}), v \mathsf{T}\theta(*\mathbf{X}, *\mathbf{Y}) \doteq \mathsf{T}\theta(v *\mathbf{X}, v *\mathbf{Y}).$ 

**Proof.** For  $*\mathbf{X} = *\mathbf{e}_{\alpha} = (*\mathbf{e}_i, *e^a)$  and  $*\mathbf{Y} = *\mathbf{e}_{\beta} = (*\mathbf{e}_l, *e^b)$ , where  $*\mathbf{e}_{\alpha}$  is a N-adapted basis of type (17), we write the first equation in (24) in the form

$${}^{\mathsf{T}}\theta = {}^{\mathsf{T}}\theta( {}^{*}\mathbf{e}_{i}, {}^{*}e^{a}) = {}^{\mathsf{T}}\theta(\frac{\partial}{\partial x^{i}}, \frac{\partial}{\partial p_{a}}) - {}^{*}N_{ib} {}^{\mathsf{T}}\theta(\frac{\partial}{\partial p_{b}}, \frac{\partial}{\partial p_{a}}) = 0.$$

Such conditions uniquely define  $N_{ib}$  because  $T\theta$  is non-degenerate, that is  $rank| T\theta(\frac{\partial}{\partial p_b}, \frac{\partial}{\partial p_a})| = n$ . Setting locally

$${}^{\mathsf{T}}\theta = \frac{1}{2} {}^{\mathsf{T}}\theta_{ij}(u)e^i \wedge e^j + \frac{1}{2} {}^{\mathsf{T}}\theta^{ab}(u) {}^{\mathsf{T}}\mathbf{e}_a \wedge {}^*\mathbf{e}_b,$$
(25)

where the first term is for  $h^{\mathsf{T}}\theta$  and the second term is  $v^{\mathsf{T}}\theta$ , we get the second formula in (24). Finally, we note that in this proposition the constructed N-connection  $*N_{ib}$ , in general, is not a canonical one (15).  $\Box$ 

In a similar form, as in Proposition 2.3, we can construct a unique Nconnection  $\mathbf{N} = \{ N_i^a \}$  for any almost symplectic structure  $\theta$  on TM (from formal point of view, we have to omit in formulas the symbols "\*" and "  $\mathsf{T}$ " and use variables  $y^a$  instead of  $p_a$ ).

A N-connection \*N (8) defines a unique decomposition of a d-vector \* $\mathbf{X} = {}^{*}X^{h} + {}^{*}X^{v}$  on  $T^{*}M$ , for  ${}^{*}X^{h} = h {}^{*}\mathbf{X}$  and  ${}^{*}X^{v} = v {}^{*}\mathbf{X}$ , where the projectors h and v defines respectively the distributions \*N and \*V. They have the properties

$$h + v = \mathbf{I}, h^2 = h, v^2 = v, h \circ v = v \circ h = 0.$$

This allows us to introduce on  $T^*M$  the almost product operator

$$^*\mathbf{P} \doteq I - 2v = 2h - I$$

acting on  $*\mathbf{e}_{\alpha} = (*\mathbf{e}_{i}, *e^{b})$  (17) following formulas

$$^{*}\mathbf{P}(^{*}\mathbf{e}_{i}) = ^{*}\mathbf{e}_{i} \text{ and } ^{*}\mathbf{P}(^{*}e^{b}) = - ^{*}e^{b}.$$

In a similar form, a N–connection  $\mathbf{N}$  (7) induces an almost product structure  $\mathbf{P}$  on TM. One uses also the almost tangent operators

$$\mathbb{J}(\mathbf{e}_i) = e_{n+i} \text{ and } \mathbb{J}(e_a) = 0, \text{ or } \mathbb{J} = \frac{\partial}{\partial y^i} \otimes dx^i;$$

$$^*\mathbb{J}(^*\mathbf{e}_i) = ^*g_{ib} * e^b \text{ and } ^*\mathbb{J}(^*e^b) = 0, \text{ or } \mathbb{J} = ^*g_{ia}\frac{\partial}{\partial p_a} \otimes dx^i$$

The operators  ${}^*\mathbf{P}, {}^*\mathbf{J}$  and  ${}^*\mathbb{J}$  are respectively  $\mathcal{L}$ -dual to  $\mathbf{P}, \mathbf{J}$  and  $\mathbb{J}$  if and only if  ${}^*\mathbf{N}$  and  $\mathbf{N}$  are  $\mathcal{L}$ -dual.

For the above–introduced almost complex and almost product operators, it is straightforward to prove

**Proposition 2.4** Let  $(\mathbf{N}, *\mathbf{N})$  be a pair of  $\mathcal{L}$ -dual N-connections. Then, we can construct canonical d-tensor fields (defined respectively by L(x, y)and H(x, p) related by Legendre transforms (5) and/or (6)):

$$\mathbf{J} = -\delta_i^a e_a \otimes e^i + \delta_a^i \mathbf{e}_i \otimes \mathbf{e}^a, \ ^*\mathbf{J} = - \ ^*g_{ia} \ ^*e^a \otimes \ ^*e^i + \ ^*g^{ia} \ ^*\mathbf{e}_i \otimes \ ^*\mathbf{e}_a$$

corresponding to the  $\mathcal{L}$ -dual pair of almost complex structures  $(\mathbf{J}, *\mathbf{J})$ ,

$$\mathbf{P} = \mathbf{e}_i \otimes e^i - e_a \otimes \mathbf{e}^a, \ ^*\mathbf{P} = \ ^*\mathbf{e}_i \otimes \ ^*e^i - \ ^*e^a \otimes \ ^*\mathbf{e}_a$$

corresponding to the  $\mathcal{L}$ -dual pair of almost product structures  $(\mathbf{P}, *\mathbf{P})$ , and almost symplectic structures

$$\theta = g_{aj}(x, y) \mathbf{e}^a \wedge e^i \quad and \quad {}^*\theta = \delta^a_i \; {}^*\mathbf{e}_a \wedge \; {}^*e^i \tag{26}$$

Let us consider an important example:

A Finsler manifold (space)  $F^n = (M, F(x, y))$  is a particular case of Lagrange space, when the regular Lagrangian  $L = F^2$  is defined by a fundamental Finsler function F(x, y) satisfying the conditions: 1. the positive function F is differentiable function on  $\widetilde{TM}$  continuous on the null section of projection  $\pi : TM \to M$ , 2.  $F(x, \lambda y) = |\lambda| F(x, y)$ , i.e. it is 1-homogeneous on the fibres of TM, and 3. the Hessian (4) defined in this case by  $F^2$  is positively defined on  $\widetilde{TM}$ .

It is used also the notion of Cartan space  $C^n = (M, C(x, p))$  for  $H = C^2(x, p)$  as a particular (1-homogeneous on fiber coordinates) case of Hamilton space when C satisfies the same conditions as F but with respect to coordinates  $p_a$  (in brief, we can say that Cartan spaces are Finsler spaces on  $T^*M$ , see details in [31]). In a similar manner as for Lagrange and Hamilton spaces, we can introduce the concept of  $\mathcal{L}$ -dual geometric objects on Finsler and Cartan spaces. For simplicity, in this work we shall emphasize the bulk constructions for Hamilton spaces considering that by Legendre transform we can generate similar ones for Lagrange spaces and, in particular, for respective Finsler and Cartan geometries.

**Definition 2.8** An almost Hermitian model of a cotangent bundle  $T^*M$ (or tangent bundle TM) equipped with a N-connection structure  $*\mathbf{N}$  (or  $\mathbf{N}$ ) is defined by a triple  $*\mathbf{H}^{2n} = (T^*M, *\theta, *\mathbf{J})$ , where  $*\theta(*\mathbf{X}, *\mathbf{Y}) \doteq *\mathbf{g}(*\mathbf{J}\mathbf{X}, \mathbf{Y})$  (or  $\mathbf{H}^{2n} = (TM, \theta, \mathbf{J})$ , where  $\theta(\mathbf{X}, \mathbf{Y}) \doteq \mathbf{g}(\mathbf{J}\mathbf{X}, \mathbf{Y})$ ). A space  $*\mathbf{H}^{2n}$  is almost Kähler, denoted  $*\mathbf{K}^{2n}$  if  $d * \theta = 0$ .

The following theorem holds:

**Theorem 2.3** The Lagrange and Hamilton spaces can be represented as almost Kähler spaces on, respectively, on TM and  $T^*M$  endowed with canonical N-connection structures  $\mathbf{N}$  (14) and  $*\mathbf{N}$  (15).

**Proof.** It follows from the existence on TM and  $T^*M$  of canonical 1–forms, respectively, defined by a regular Lagrangian L and Hamiltonian H related by a Legendre transform,

$$\omega = \frac{\partial L}{\partial y^i} e^i$$
 and  $*\omega = p_i dx^i$ .

for which

$$\theta = d\omega$$
 and  $^*\theta = d^*\omega$ ,

see (12). As a result, we get that  $d \theta = 0$  and  $d^*\theta = 0$ , which correspond to the Definition 2.8.  $\Box$ 

In this paper, we shall work with almost Kähler models on (co) tangent bundles defined canonically by (pseudo) Riemannian metrics on base manifolds, see (1), and/or (effective, or for regular mechanics) Lagrangians (Hamiltonians). Finally, we emphasize that realistic classical and quantum models are elaborated in explicit form for some classes of linear connections defined to satisfy certain physical principles and constructed geometrically to be adapted, or not, to a N–connection structure. We shall perform such classical and quantum constructions in the following sections.

## 3 Nonlinear Connections and Almost Symplectic Geometry

In this section, we consider the almost symplectic geometry induced by regular Hamiltonians and corresponding canonical N–connections defined naturally, for gravitational and/or geometric mechanics models, on (co) tangent bundles.

#### 3.1 Canonical N–connections and d–connections for Lagrange and Hamilton spaces

Let D be a linear connection on TM when for a  $\mathcal{L}$ -duality between the tangent and corresponding cotangent bundles there are defined pull-back and push-forward maps. We can define a linear connection \*D on  $T^*M$  as follows:

$$^*D_{*X} ^*Y \doteq (D_{\circ X} ^\circ Y)^*,$$

for any vector fields \*X and \*Y on  $T^*M$ . Inversely, for any linear connection \*D on  $T^*M$ , we get a linear connection  $^{\circ}D$  on TM, following the rule

$$^{\circ}D_XY \doteqdot (^*D_{^*X} \ ^*Y)^{\circ},$$

for any vector fields X and Y on TM.

**Definition 3.1** A linear connection  $\mathbf{D}$  or  $(*\mathbf{D})$  on TM (or  $T^*M$ ) is a distinguished connection (*d*-connection) if it is compatible with the almost product structure  $\mathbf{DP} = 0$  (or  $*\mathbf{D} * \mathbf{P} = 0$ ).

For  $\mathcal{L}$ -dual Lagrange and Hamilton spaces, one follows that  $\mathbf{DP} = 0$  induces  $^*\mathbf{D} ^*\mathbf{P} = 0$ , and inversely. The coefficients of d-connections can be defined with respect to N-adapted frames,

$$\mathbf{D}_{\mathbf{e}_{\beta}}\mathbf{e}_{\gamma} \doteqdot \mathbf{\Gamma}^{\alpha}_{\ \beta\gamma}\mathbf{e}_{\alpha} \text{ and } \mathbf{D}_{\mathbf{*e}_{\beta}} \mathbf{*e}_{\gamma} \doteqdot \mathbf{T}^{\alpha}_{\ \beta\gamma} \mathbf{*e}_{\alpha}$$

with corresponding N-adapted splitting,

$$\mathbf{D}_{\mathbf{e}_{k}}\mathbf{e}_{j} \doteq L^{i}_{\ jk}\mathbf{e}_{i}, \mathbf{D}_{\mathbf{e}_{k}}e_{b} \doteq L^{a}_{\ bk}e_{a}, \mathbf{D}_{e_{c}}\mathbf{e}_{j} \doteq C^{i}_{\ jc}\mathbf{e}_{i}, \mathbf{D}_{e_{c}}e_{b} \doteq C^{a}_{\ bc}e_{a}$$

and

$$^{*}\mathbf{D} *_{\mathbf{e}_{k}} ^{*}\mathbf{e}_{j} \stackrel{:}{=} \ ^{*}L^{i}{}_{jk} *^{*}\mathbf{e}_{i}, \ ^{*}\mathbf{D}_{\mathbf{e}_{k}} *e^{b} \stackrel{:}{=} - {}^{*}\dot{L}^{b}{}_{a}{}_{k} *e^{a},$$

$$^{*}\mathbf{D} *_{e^{c}} *\mathbf{e}_{j} \stackrel{:}{=} \ ^{*}\dot{C}^{i}{}_{j}{}^{c} *\mathbf{e}_{i}, \ ^{*}\mathbf{D} *_{e^{c}} *e^{b} \stackrel{:}{=} - {}^{*}C^{bc}{}_{a}{}^{ec} *e^{a},$$

when

$$\Gamma^{\alpha}_{\ \beta\gamma} = \{L^{i}_{\ jk}, \dot{L}^{a}_{\ bk}, \dot{C}^{i}_{\ jc}, C^{a}_{\ bc}\} \text{ and } *\Gamma^{\alpha}_{\ \beta\gamma} = \{*L^{i}_{\ jk}, *\dot{L}^{b}_{a\ k}, *\dot{C}^{i}_{\ j}, *C^{\ bc}_{a}\}$$

define corresponding h- and v-splitting of covariant derivatives

$$\mathbf{D} = \left(\begin{array}{c} {}_{h}\mathbf{D}, {}_{v}\mathbf{D} \right) \text{ and } {}^{*}\mathbf{D} = \left(\begin{array}{c} {}^{*}_{h}\mathbf{D}, {}^{*}_{v}\mathbf{D} \right),$$

where  $_{h}\mathbf{D} = \{L^{i}_{\ jk}, \dot{L}^{a}_{\ bk}\}, \ _{v}\mathbf{D} = \{\dot{C}^{i}_{\ jc}, C^{a}_{\ bc}\} \text{ and } \ _{h}^{*}\mathbf{D} = \{ \ ^{*}L^{i}_{\ jk}, \ ^{*}\dot{L}^{\ b}_{\ a\ k}\}, \ _{v}^{*}\mathbf{D} = \{ \ ^{*}\dot{C}^{i}_{\ j}, \ ^{*}C^{\ bc}_{\ a\ b}\}.$ 

We shall work with a more special class of d-connections:

**Definition 3.2** A linear connection  ${}^{n}\mathbf{D}$  (or  ${}^{*n}\mathbf{D}$ ) on TM (or  $T^{*}M$ ) is Nlinear if it preserves under parallelism the Whitney sum  $\mathbf{N}$  (7) (or  ${}^{*}\mathbf{N}$  (8)) i.e it is a d-connection and is compatible with the almost tangent structure  $\mathbb{J}$  (or  ${}^{*}\mathbb{J}$ ) i.e.  ${}^{n}\mathbf{D}\mathbb{J} = 0$  (or  ${}^{*n}\mathbf{D} {}^{*}\mathbb{J} = 0$ ).

This is a class of N-linear connections characterized by the property:

**Proposition 3.1** The N-adapted coefficients of N-linear connections  ${}^{n}\mathbf{D} \doteq \{ {}^{n}\Gamma^{\alpha}_{\beta\gamma} = (L^{i}_{jk}, \acute{C}^{i}_{jc}) \}$  and  ${}^{*n}\mathbf{D} \doteq \{ {}^{n*}\Gamma^{\alpha}_{\beta\gamma} = ( {}^{*}L^{i}_{jk}, {}^{*}\acute{C}^{i}_{j}{}^{c}) \}$  of respective  $\mathcal{L}$ -dual Lagrange and Hamilton spaces are:

$${}^{n}\Gamma^{\alpha}_{\ \beta\gamma} = \{L^{i}_{\ jk}, \acute{L}^{n+i}_{\ n+jk} = L^{i}_{\ jk}, \acute{C}^{i}_{\ jc}, C^{n+i}_{\ n+jc} = \acute{C}^{i}_{\ jc}\}$$
(27)

and

$${}^{*n}\Gamma^{\alpha}_{\beta\gamma} = \{ {}^{*}L^{i}_{jk}, {}^{*}\dot{L}^{n+i}_{n+j\ k} = {}^{*}L^{i}_{jk}, {}^{*}\dot{C}^{i\ c}_{j}, {}^{*}C^{n+i\ c}_{n+j} = {}^{*}\dot{C}^{i\ c}_{j} \}.$$
(28)

**Proof.** By a straightforward computation for coefficients (27) and (28), when, for instance,  ${}^{*n}\mathbf{D}_{\alpha} = ({}^{*}_{h}D_{i}, {}^{*}_{v}D^{a})$ , for  ${}^{*}_{h}D = \{{}^{*}L^{i}_{jk}\}$  and  ${}^{*}_{v}D = \{{}^{*}C^{i}_{j}{}^{c}\}$ , we can verify that the conditions considered in Definition 3.2 are satisfied.  $\Box$ 

The connection 1–form of N–linear connection  ${}^{*n}\Gamma^{\alpha}_{\ \beta} = {}^{*n}\Gamma^{\alpha}_{\ \beta\gamma} {}^{*}\mathbf{e}^{\gamma} = \{ {}^{*n}\Gamma^{i}_{\ i} \}$  is defined by

$${}^{*n}\Gamma^{i}{}_{j} = {}^{*}L^{i}{}_{jk} {}^{*}e^{k} + {}^{*}\dot{C}^{i}{}_{j}{}^{c} {}^{*}\mathbf{e}_{c},$$
(29)

where the v–components  ${}^{*n}\Gamma^a_b$  are identified with the h–components  ${}^{*n}\Gamma^i_j$  following formulas  ${}^{*n}\Gamma^{n+i}_{n+j} = {}^{*n}\Gamma^a_b$ . The following theorem holds:

**Theorem 3.1** On a Hamilton space, the structure equations for  ${}^{*n}\Gamma^{i}{}_{j}$  are

$$d^{*}e^{k} - {}^{*}e^{j} \wedge {}^{*n}\Gamma^{k}{}_{j} = -{}^{*}\mathcal{T}^{k},$$
  
$$d^{*}\mathbf{e}_{c} + {}^{*}\mathbf{e}_{a} \wedge {}^{*n}\Gamma^{a}{}_{c} = -{}^{*}\mathcal{T}_{c},$$
  
$$d^{*n}\Gamma^{i}{}_{j} - {}^{*n}\Gamma^{i}{}_{k} \wedge {}^{*n}\Gamma^{k}{}_{j} = -{}^{*}\mathcal{R}^{i}{}_{j},$$

where the 2-form of torsion  $\mathcal{T}^{\alpha} = \{\mathcal{T}^k, \mathcal{T}_c\}$  is computed

$${}^{*}\mathcal{T}^{k} = \frac{1}{2} {}^{*}T^{k}_{ij} {}^{*}e^{i} \wedge {}^{*}e^{j} + {}^{*}\dot{C}^{k}_{j}{}^{c} {}^{*}e^{j} \wedge {}^{*}\mathbf{e}_{c},$$

$${}^{*}\mathcal{T}_{a} = \frac{1}{2} {}^{*}\Omega_{ija} {}^{*}e^{i} \wedge {}^{*}e^{j} + \frac{1}{2} {}^{*}P_{aic} {}^{*}e^{i} \wedge {}^{*}\mathbf{e}_{c} + \frac{1}{2} {}^{*}S_{a}{}^{bc} {}^{*}\mathbf{e}_{b} \wedge {}^{*}\mathbf{e}^{c},$$

for the N-connection curvature  $*\Omega_{ija}$  (22) and

$${}^{*}T^{k}_{ij} = {}^{*}L^{k}_{ij} - {}^{*}L^{k}_{ji}, {}^{*}S^{\ bc}_{a} = {}^{*}\dot{C}^{\ b}_{\ a} - {}^{*}\dot{C}^{\ b}_{\ a},$$
  
$${}^{*}P_{aic} = {}^{*}g_{ae} \left( {}^{*}L^{e}_{\ ic} - {}^{*}e^{e} ({}^{*}N_{ic}) \right), \qquad (30)$$

and the 2-form of curvature  ${}^{*}\mathcal{R}^{\alpha}_{\ \beta} = \left( {}^{*}\mathcal{R}^{i}_{\ j}, {}^{*}\mathcal{R}^{a}_{\ b} \right)$ , with  ${}^{*}\mathcal{R}^{i}_{\ j} = {}^{*}\mathcal{R}^{n+i}_{\ n+j}$ , is computed

$${}^{*}\mathcal{R}^{i}{}_{j} = \frac{1}{2} {}^{*}R^{i}{}_{jkm} {}^{*}e^{k} \wedge {}^{*}e^{m} + {}^{*}P^{i}{}_{jk}{}^{c} {}^{*}e^{k} \wedge {}^{*}\mathbf{e}_{c} + \frac{1}{2} {}^{*}S^{i}{}_{j}{}^{bc} {}^{*}\mathbf{e}_{b} \wedge {}^{*}\mathbf{e}_{c},$$

where

$${}^{*}R^{i}{}_{jkm} = {}^{*}e_{m}({}^{*}L^{i}{}_{jk}) - {}^{*}e_{k}({}^{*}L^{i}{}_{jm})$$

$$+ {}^{*}L^{o}{}_{jk} {}^{*}L^{i}{}_{om} - {}^{*}L^{o}{}_{jm} {}^{*}L^{i}{}_{ok} + {}^{*}\acute{C}^{i}{}_{j} {}^{o}{}^{*}\Omega_{oka},$$

$${}^{*}P^{i}{}_{jk} {}^{c} = {}^{*}e^{c}({}^{*}L^{i}{}_{jk}) - {}^{*}_{h}D_{k}({}^{*}\acute{C}^{i}{}_{j} {}^{c}) + {}^{*}\acute{C}^{i}{}_{j} {}^{o}{}^{*}P^{c}{}_{ko},$$

$${}^{*}S^{i}{}_{j} {}^{bc} = {}^{*}e^{c}({}^{*}\acute{C}^{i}{}_{j} {}^{b}) - {}^{*}e^{b}({}^{*}\acute{C}^{i}{}_{j} {}^{c}) + {}^{*}\acute{C}^{k}{}_{j} {}^{b}{}^{*}\acute{C}^{i}{}_{k} {}^{c} - {}^{*}\acute{C}^{k}{}_{j} {}^{c}{}^{*}\acute{C}^{i}{}_{k} {}^{b},$$

$$(31)$$

for "non-boldface"  $*e^c = \partial/\partial p_c$ .

**Proof.** It is a straightforward differential computation for 1–form (29).  $\Box$ 

There is a " $\mathcal{L}$ -dual" Theorem for Lagrange spaces [34], see generalizations of nonholonomic manifolds and deformation quantization of gravity in Ref. [30, 24], similarly to Theorem 3.1. From formal point of view, we have to change H into L and consider the constructions on TM, omitting the labels "\*" and using coordinates  $(x^i, y^a)$  instead of  $(x^i, p_a)$ .

#### 3.2 Hamilton–Fedosov spaces and almost Kähler structures

There are canonical N-linear connections on TM and  $T^*M$  completely defined, respectively, by the fundamental Lagrange (see [34] and, for applications to geometric quantization, [21, 24]) and Hamilton functions.

#### Theorem 3.2 There exists a canonical N-linear connection

 $\label{eq:approx_state_stat$ 

**Proof.** Let us consider the N-adapted coefficients

$$\hat{L}^{i}{}_{jk} = \frac{1}{2} *g^{is} (*e_{j}(*g_{sk}) + *e_{k}(*g_{js}) - *e_{s}(*g_{jk})),$$

$$\hat{C}^{i}{}_{j}^{c} = -\frac{1}{2} *g_{js} *e^{c}(*g^{si})$$

$$(32)$$

defined with respect to N-adapted bases  $*\mathbf{e}_{\alpha}$  (17) and coframes  $*\mathbf{e}^{\alpha}$  (19) defined by  $*N_{ij}$  (15). By a direct computation, we can verify that the conditions of this Theorem are satisfied only for such coefficients and their coordinate/frame transform.  $\Box$ 

The coefficients (32) are just the Christoffel symbols on (co) tangent space  $T^*M$ , defined by H. Nevertheless, for a Hamilton space endowed with canonical N-connection, d-connection and metric structure, there are nontrivial torsion components induced by the nonholonomic distribution defined by H, see  ${}^*P_{aic}$  (30).

From Theorem 3.2, one follows an important property:

**Corollary 3.1** The canonical d-connection \* $\mathbf{D}$  is an almost symplectic d-connection satisfying the conditions

$$^{*}\widehat{\mathbf{D}} ^{*}\theta = 0 \ and \ ^{*}\widehat{\mathbf{D}} ^{*}\mathbf{J} = 0 \tag{33}$$

and being completely defined by a Hamiltonian H(x,p) for  $*\theta(*\mathbf{X}, .) \doteq *\mathbf{g}(*\mathbf{J}\mathbf{X}, .)$ .

**Proof.** A Hamilton space can be equivalently transformed into almost Kähler space  ${}^{*}\mathbf{K}^{2n}$ , see Definition 2.8. Considering the d-tensor fields associated to  ${}^{*}\theta$  and  ${}^{*}\mathbf{J}$ , see Proposition 2.4, and a covariant N-adapted calculus defined by the canonical d-connection coefficients (32) we can verify that the almost symplectic structure compatibility conditions (33) are satisfied.  $\Box$ 

In Ref. [20], it was introduced the concept of Lagrange–Fedosov manifold as nonholonomic manifold with the N–connection and almost symplectic structure defined by a fundamental (in genera, effective) Lagrange function L(x, y). On cotangent bundles, we can consider

**Definition 3.3** A Hamilton–Fedosov space is a cotangent bundle endowed with canonical N–connection and the almost Kähler structure induced by a fundamental Hamilton function H(x, p).

There are Hamilton–Fedosov spaces defined completely by a lift (1) of a (pseudo) Riemannian metric on base M.

**Theorem 3.3** Any Einstein manifold associated to a solution of (2), for a lift (1) on  $T^*M$ , defines canonically a Hamilton–Fedosov space.

**Proof.** We sketch the idea for such constructions. Let us fix any values  $e^{i}_{i'}(x,p)$  in (1) and associate  $g^{ab}(x,p)$  to a  ${}^{*}g^{ab}(x,p)$  (3). This define correspondingly the values  ${}^{*}\mathbf{g}$  (20) and  ${}^{*}\mathbf{N}$  (15). As a result, we construct an effective Hamilton space, which can be modelled as a canonical almost symplectic structure as we described above. For classical configurations, the values  $e^{i}_{i'}$  can be  $\delta^{i}_{i'}$ , but for quantum models they should defined by a scheme of de-quantization, or semi-classical approximation in quantum gravity.

There is a particular case of Cartan–Fedosov spaces with 0-homogeneous on variables "p" components  $e_{i'}^i(x,p)$  resulting in a similar homogeneity for  $g^{ab}(x,p)$  and  $H = g^{ab}p_ap_b$  when \*N is determined from (15) with \*g from (21).

The problem is more sophisticate in the case of general Hamilton–Fedosov spaces. For certain physical important four dimensional spaces (used in general relativity) with nonholonomic splitting of dimensions as 2+2 and, for instance, if dimM = 2, we can fix such local coordinate systems when (3) is integrable for certain solvable partial differential equations for  $e_{i'}^i(x,p)$ , but this may not hold true for other parameterizations and higher dimensions. A general approach should include the case of Eisenhart–Hamilton spaces and their Fedosov quantum deformation analogs, with both symmetric and nonsymmetric components for (3) resulting because of any general quantum nonholonomic Legendre transform. Such constructions should dub on cotangent bundles those for Eisenhart-Lagrange/- Finsler spaces (see Chapter 8 in [33] and Refs. [52,53]). The length of this paper does not allow us to present a detailed proof because it is connected with a sophisticate geometric techniques for nonlinear connections and nonsymmetric metrics arising both in the case of nonholonomic Ricci flows and quantum nonholonomic deformations and/or symplectic transform in gravity and geometric mechanics, see recent results in Refs. [54-56].  $\Box$ 

Finally, we note that similar almost symplectic models can be performed for Cartan spaces when the Hamiltonian is homogeneous on vertical coordinates.

### 4 Fedosov Operator–Pairs for Hamilton Spaces

In this section, we shall apply the method of deformation quantization elaborated in Refs. [5, 6, 35] to define two classes of canonical operators which are necessary to quantize the Hamilton–Fedosov spaces and related subspaces on cotangent bundles defined by lifts of Einstein metrics. We shall address precisely the question how the geometry of cotangent bundles and related deformation quantization change under symplectic transforms and elaborate a formalism which preserves the form of Hamilton–Jacobi equations both on classical and quantum level.

#### 4.1 Canonical Fedosov–Hamilton operators

The formalism of deformation quantization can be developed by using the space  $C^{\infty}({}^{*}\mathbf{K}^{2n})[[v]]$  of formal series in the variable v with coefficients from  $C^{\infty}({}^{*}\mathbf{K}^{2n})$  on a almost Poisson manifold ( ${}^{*}\mathbf{K}^{2n}, \{\cdot, \cdot\}$ ), see the almost symplectic form  ${}^{*}\theta$  and the Poisson brackets (13). Using the associative algebra structure on  $C^{\infty}({}^{*}\mathbf{K}^{2n})[[v]]$  with a v-linear and v-adically continuous star product

$${}^{1}f * {}^{2}f = \sum_{r=0}^{\infty} {}_{r}C({}^{1}f, {}^{2}f) v^{r}, \qquad (34)$$

where  ${}_{r}C, r \geq 0$ , are bilinear operators on  $C^{\infty}({}^{*}\mathbf{K}^{2n})$  with  ${}_{0}C({}^{1}f, {}^{2}f) = {}^{1}f {}^{2}f$  and  ${}_{1}C({}^{1}f, {}^{2}f) - {}_{1}C({}^{2}f, {}^{1}f) = i\{{}^{1}f, {}^{2}f\}$ ; for *i* being the complex unity, we can construct a formal Wick product

$${}^{1}a \circ {}^{2}a(z) \doteq \exp\left(i\frac{v}{2} * \mathbf{\Lambda}^{\alpha\beta} \frac{\partial^{2}}{\partial z^{\alpha} \partial z_{[1]}^{\beta}}\right) \; {}^{1}a(z) \; {}^{2}a(z_{[1]}) \mid_{z=z_{[1]}}, \tag{35}$$

for two elements a and b defined by series of the type

$$a(v,z) = \sum_{r \ge 0, |\{\alpha\}| \ge 0} a_{r,\{\alpha\}}(u) z^{\{\alpha\}} v^r,$$
(36)

where by  $\{\alpha\}$  we label a multi-index and  $*\Lambda^{\alpha\beta} \doteq *\theta^{\alpha\beta} - i * \mathbf{g}^{\alpha\beta}$ , where  $*\theta^{\alpha\beta}$  is the symplectic form (26), with "up" indices, and  $*\mathbf{g}^{\alpha\beta}$  is the inverse to d-tensor (20). This defines a formal Wick algebra  $*\mathbf{W}_u$  associated with the tangent space  $T_u * \mathbf{K}^{2n}$ , for  $u \in *\mathbf{K}^{2n}$ , where the local coordinates on  $*\mathbf{K}^{2n}$  are parameterized in the form  $u = \{u^{\alpha}\}$  and the local coordinates on  $T_u * \mathbf{K}^{2n}$  are labelled  $(u, z) = (u^{\alpha}, z^{\beta})$ , where  $z^{\beta}$  are fiber coordinates.

We trivially extend the fibre product (35) to the space of  $*\mathbf{W}$ -valued N-adapted differential forms  $*\mathcal{W} \otimes \Lambda$ , where by  $\Lambda$  we note the usual exterior product of the scalar forms and  $*\mathcal{W}$  is the sheaf of smooth sections of  $*\mathbf{W}$ . There is a standard grading on  $\Lambda$  noted deg<sub>a</sub>. We also introduce gradings deg<sub>v</sub>, deg<sub>s</sub>, deg<sub>a</sub> on  $*\mathcal{W} \otimes \Lambda$  defined on homogeneous elements  $v, z^{\alpha}, *\mathbf{e}^{\alpha}$ , when deg<sub>v</sub>(v) = 1, deg<sub>s</sub>( $z^{\alpha}$ ) = 1, deg<sub>a</sub>( $*\mathbf{e}^{\alpha}$ ) = 1, and all other gradings of the elements  $v, z^{\alpha}, *\mathbf{e}^{\alpha}$  are set to zero. As a result, the product  $\circ$  from (35) on  $*\mathcal{W} \otimes \Lambda$  is bi-graded, written as w.r.t the grading  $Deg = 2 \deg_v + \deg_s$  and the grading deg<sub>a</sub>.

The canonical d–connection  $\hat{\mathbf{D}} \doteq \{ \hat{\mathbf{\Gamma}}^{\alpha}_{\beta\gamma} = (\hat{L}^{i}_{jk}, \hat{C}^{ic}_{j}) \}$  with coefficients (32) can be extended to an operator

$$^{*}\widehat{\mathbf{D}}\left(a\otimes\lambda\right) \doteq \left( ^{*}\mathbf{e}_{\alpha}(a) - u^{\beta} \ ^{*}\widehat{\mathbf{\Gamma}}_{\alpha\beta}^{\gamma} \ ^{z*}\mathbf{e}_{\alpha}(a) \right) \otimes \left( ^{*}\mathbf{e}^{\alpha}\wedge\lambda\right) + a\otimes d\lambda, \quad (37)$$

on  $*\mathcal{W} \otimes \Lambda$ , where the N-adapted basis  ${}^{z*}\mathbf{e}_{\alpha}$  is  $*\mathbf{e}_{\alpha}$  redefined in z-variables. This canonical almost symplectic d-connection  $*\widehat{\mathbf{D}}$  is a N-adapted deg<sub>a</sub>-graded derivation of the distinguished algebra ( $*\mathcal{W} \otimes \Lambda, \circ$ ), in brief, called d-algebra: this follows from formulas (35) and (37)).

**Definition 4.1** The Fedosov-Hamilton operators  $*\delta$  and  $*\delta^{-1}$  on  $*W \otimes \Lambda$ , are defined

$${}^{*}\delta(a) = {}^{*}\mathbf{e}^{\alpha} \wedge {}^{z*}\mathbf{e}_{\alpha}(a), and {}^{*}\delta^{-1}(a) = \begin{cases} \frac{i}{p+q} z^{\alpha} & {}^{*}\mathbf{e}_{\alpha}(a), & if \ p+q > 0, \\ 0, & if \ p=q = 0, \end{cases}$$
(38)

where any  $a \in {}^*\mathcal{W} \otimes \mathbf{\Lambda}$  is homogeneous w.r.t. the grading deg<sub>s</sub> and deg<sub>a</sub> with deg<sub>s</sub>(a) = p and deg<sub>a</sub>(a) = q.

The d-operators (38) define the formula  $a = (*\delta *\delta^{-1} + *\delta^{-1} *\delta + \sigma)(a)$ , where  $a \mapsto \sigma(a)$  is the projection on the  $(\deg_s, \deg_a)$ -bihomogeneous part of a of degree zero,  $\deg_s(a) = \deg_a(a) = 0$ ;  $*\delta$  is also a  $\deg_a$ -graded derivation of the d-algebra  $(*W \otimes \Lambda, \circ)$ .

#### 4.2 Fedosov–Hamilton operator–pairs

Having defined d-operators (38), we can perform a Fedosov type quantization of Hamilton spaces in a  $\mathcal{L}$ -dual form to Lagrange spaces [21, 24]. Nevertheless, such constructions would not reflect completely the symplectic properties of Hamilton spaces and their quantum deformations. For a cotangent bundle  $\pi^* : T^*M \to M$ , a N-connection **\*N** (15) is a supplementary distribution  $hT^*M$  of the vertical distribution  $vT^*M = \ker \ \ \ \ \pi^*$ , where  $\ \ \ \pi^*$  is the tangent map of  $\pi^*$ . It is often more convenient to consider a N-connection as an almost product structure **\*P**, see Proposition 2.4, such that  $vT^*M = \ker(I + \ \mathbf{P})$ . If  $f \in Diff(T^*M)$ , the push-forward of **\*N** by f generally fails to be a connection. This constrains us to extend the definition of connection, see details in Chapter 8 of [31].

**Definition 4.2** A connection-pair  $*\phi$  is an almost product structure  $*\phi$ on  $T^*M$  such that the horizontal bundle  $hT^*M \doteq \ker(I - *\phi)$  is supplementary to  $vT^*M$  and the oblique bundle  $wT^*M \doteq \ker(I + *\phi)$  define the nonholonomic splitting (Whitney sum)

$$TT^*M = hT^*M \oplus wT^*M.$$
(39)

We can consider preferred  $\phi$ -adapted frame and coframe structures induced by the coefficients of N-connection:

**Proposition 4.1** There are a canonical connection–pair  $*\phi$  and associated frames and coframes:

$${}^{\phi}\mathbf{e}_{\alpha} = \left( {}^{\phi}e_i = \frac{\partial}{\partial x^i} + {}^*N_{ia}\frac{\partial}{\partial p_a}, {}^{w}e^b = \frac{\partial}{\partial p_b} - {}^*g^{bi} {}^{\phi}e_i \right)$$
(40)

$${}^{\phi}\mathbf{e}^{\alpha} = \left( {}^{\phi}e^{i} = dx^{i} + {}^{*}g^{bi} {}^{w}e_{b}, {}^{w}e_{b} = dp_{b} - {}^{*}N_{ib}dx^{i} \right),$$
(41)

where  $*g^{ab}(3)$  and  $*N_{ia}(15)$  are generated by a Hamilton fundamental function H(x, p).

**Proof.** The proof is similar to that for Proposition 2.1 but adapted both the the splitting (39), with respective h- and v-projections,  $2h' = I + *\phi$  and  $2w = I - *\phi$ , and to the splitting (8), with respective h- and v-projections,  $2h = I + *\mathbf{N}$  and  $2v = I - *\mathbf{N}$ .  $\Box$ 

We note that for Hamilton spaces a connection–pair structure  $*\phi$  is symmetric, i.e.  $*N_{ia} = *N_{ai}$  and  $*g^{ab} = *g^{ba}$ . For simplicity, in this work, we shall restrict our considerations only to  $*\phi$ –symmetric configurations.

In order to perform geometric constructions adapted both to the N– connection and almost symplectic structure, it is necessary to work with \* $\phi$ -adapted bases (40) and (41) instead of, respectively, (17) and (19). For instance, every vector field **X** has two components

$$\mathbf{X} = hX + vX = h'X + wX,$$

where  $wX = w(\mathbf{X})$ . This defines the class of  $\phi$ -tensor fields alternatively to that of d-tensor fields considered in the previous sections.

**Definition 4.3** A linear connection  ${}^{\phi}\mathbf{D}$  on  $T^*M$  is a  ${}^*\phi$ -connection if

$${}^{\phi}\mathbf{D} \; {}^{*}\phi = 0 \; and \; {}^{\phi}\mathbf{D} \; w = 0.$$

We can characterize  ${}^{\phi}\mathbf{D} = ({}^{\phi}_{h}D = \{ {}^{\phi}L^{i}_{jk} \}, {}^{\phi}_{w}D = \{ {}^{\phi}C^{j}_{i}{}^{k} \})$  by the coefficients computed with respect to  ${}^{*}\phi$ -adapted frames (40) and (41):

**Theorem 4.1** There is a canonical  ${}^{*}\phi$ -connection  ${}^{\phi}\widehat{\mathbf{D}} = ({}^{\phi}\widehat{L}_{jk}^{i}, {}^{\phi}\widehat{C}_{jk}^{k\,a})$ on  $T^{*}M$  completely defined by a Hamilton fundamental function H(x,p) and satisfying the conditions

$${}^{\phi}_{h}\widehat{D}(\ {}^{*}g^{ab}) = 0 \ and \ {}^{\phi}_{v}\widehat{D}(\ {}^{*}g^{ab}) = 0,$$

$${}^{\phi}\widehat{T}^{k}_{ii} = 0 \ and \ {}^{\phi}\widehat{S}^{\ bc}_{a} = 0.$$

**Proof.** The torsion and curvature of  ${}^{\phi}\widehat{\mathbf{D}}$  are computed as in Theorem 3.1, see formulas (30), but with respect to (40) and (41). By straightforward

computations, we can verify that for

$${}^{\phi}\widehat{L}^{i}_{jk} = \frac{1}{2} {}^{*}g^{km} ({}^{\phi}e_{i} {}^{*}g_{mj} + {}^{\phi}e_{j} {}^{*}g_{im} - {}^{\phi}e_{m} {}^{*}g_{ij}) -\frac{1}{2} {}^{*}g^{km} (A_{jim} + A_{ijm} + A_{mij}), \qquad (42)$$
$${}^{\phi}C^{j}{}^{k}_{i} = -\frac{1}{2} {}^{*}g_{im} ({}^{w}e^{j} {}^{*}g^{mk} + {}^{w}e^{k} {}^{*}g^{im} + {}^{w}e^{m} {}^{*}g^{jk}) -\frac{1}{2} {}^{*}g_{im} (B^{jmk} + B^{kmj} + B^{mkj}),$$

where

$$A_{ijm} = *\Omega_{ijm}$$
 and  $B^{mkj} = *g^{mi}(*g^{ka} w e^{j} *N_{ia} - *g^{ja} w e^{k} *N_{ia}),$ 

the conditions of Theorem are satisfied.  $\Box$ 

The diffeomorphism symmetry of Hamilton mechanical models and possible lifts of Einstein spaces on cotangent bundles is an important characteristic of classical theories. In deformation quantization models, this property can be preserved for regular N–connection structures:

**Definition 4.4** A diffeomorphism  $f \in Diff(T^*M)$  is called  $*\mathbf{N}$ -regular if the restriction of tangent map  $(\pi f)_* : hT^*M \to TM$  to  $hT^*M$  is a diffeomorphism.

For a connection-pair  ${}^*\phi$ , the concept of  ${}^*\mathbf{N}$ -regularity imposes the equivalence of statements: 1) the push-forward map of  ${}^*\phi$  by f, defined as  ${}^*\overline{\phi} = f_*({}^*\phi)f_*^{-1}$ , is a connection-pair and 2) f is  ${}^*\mathbf{N}$ -regular. This follows from the mapping  $\pi_* : \ker(I - {}^*\overline{\phi}) \to TM$  and the equality  $f_*(\ker(I - {}^*\phi)) = \ker(I - {}^*\overline{\phi})$ . The N-connection  ${}^*\overline{\mathbf{N}}$  associated to  ${}^*\overline{\phi}$  is the push-forward of  ${}^*\mathbf{N}$  by f. One says that  ${}^*\phi$  and  ${}^*\overline{\phi}$  are f-related.

Following a calculus with local coordinate transforms, one proves:

**Corollary 4.1** 1) For  ${}^{\phi}\mathbf{e}_{\alpha} = ({}^{\phi}e_i, {}^{w}e^b)$  in (x, p) and  ${}^{\phi}\overline{\mathbf{e}}_{\alpha} = ({}^{\phi}\overline{e}_i, {}^{w}\overline{e}^b)$  in  $(\overline{x}(x, p), \overline{p}(x, p))$  induced respectively by  ${}^{*}\phi$  and  ${}^{*}\overline{\phi}$ , the following formulas hold

$$f_*({}^{\phi}e_i) = \varpi_i^k {}^{\phi}\overline{e}_k and f_*({}^we^b) = \widetilde{\varpi}_c^b {}^we^c,$$

where

$$\varpi_i^{k} * N_{ka} = {}^{\phi} e_i(\overline{p}_a) \text{ and } \widetilde{\varpi}_c^{b} = {}^{w} e^{b}(\overline{p}_c) - {}^{w} e^{b}(\overline{x}^k) * \overline{N}_{kc}.$$

2) The push-forward of a N-connection \*N by a \*N-regular diffeomorphism is a N-connection if f is fiber preserving, i.e. locally  $f(x,p) = (\overline{x}(x), \overline{p}(x,p))$ .

Symplectic morphisms are diffeomorphisms which transform a symplectic form  ${}^*\theta = dp_i \wedge dx^i$  into a symplectic form  ${}^*\overline{\theta} = f^*({}^*\theta) = d\overline{p}_i \wedge d\overline{x}^i$ . By coordinate transforms, one proves:

**Theorem 4.2** For a  $f \in Diff(T^*M)$  being \*N-regular and satisfying the condition  $f^*(\ ^*\theta)(\ ^\phi e_i,\ ^w e^b) = \delta^b_i$ , two from the next statements implies the third: 1) f is a symplectic morphism; 2) \* $\phi$  is symmetric; 3) \* $\overline{\phi}$  is symmetric.

The push-forward of  $*\mathbf{g}$  (20) results in the local form

$${}^{*}\overline{\mathbf{g}} = {}^{*}\overline{\mathbf{g}}_{\alpha\beta} {}^{*}\overline{\mathbf{e}}^{\alpha} \otimes {}^{*}\overline{\mathbf{e}}^{\beta} = {}^{*}\overline{g}_{ij}(x,p)\overline{e}^{i} \otimes \overline{e}^{j} + {}^{*}\overline{g}^{ab}(x,p) {}^{*}\overline{\mathbf{e}}_{a} \otimes {}^{*}\overline{\mathbf{e}}_{b},$$

where  $*\overline{g}^{ij} \circ f = \varpi_k^i \varpi_l^j * g^{kl}$ .

If  $\nabla$  is a linear connection on  $T^*M$ , we define its push–forward by f on  $T^*M$  as

$$\overline{\nabla}_{\overline{X}}\overline{Y} \doteq f_*\left(\nabla_X Y\right)$$

for  $\overline{Y} = f_*(Y)$  and  $\overline{X} = f_*(X)$ . By coordinate parameterizations of diffeomorphisms, we can prove:

**Proposition 4.2** 1) A connection  $\nabla$  is a \* $\phi$ -connection if and only if  $\overline{\nabla}$  is a \* $\overline{\phi}$ -connection; 2)  $\nabla$  is compatible to metric \* $\mathbf{g}$  (almost symplectic \* $\theta$ ) structure if and only if  $\overline{\nabla}$  is compatible to metric \* $\overline{\mathbf{g}}$  (almost symplectic \* $\overline{\theta}$ ) structure.

**Proof.** Locally, we can prove that the coefficients of  $*\phi$ -connection  $\phi \mathbf{D}$  are related to coefficients of  $*\overline{\phi}$ -connection  $\phi \overline{\mathbf{D}}$  by formulas

Using formulas

$$\begin{split} {}^{\phi}\overline{\mathbf{D}}(\ {}^{*}\phi) &= \ 0 \Longleftrightarrow \ {}^{\phi}\mathbf{D}(\ {}^{*}\phi) = 0, \\ {}^{\phi}\overline{\mathbf{D}}\ {}^{*}\overline{\mathbf{g}} &= \ {}^{\phi}\overline{\mathbf{D}}\left[f_{*}(\ {}^{*}\mathbf{g})\right] = f_{*}(\ {}^{\phi}\mathbf{D}\ {}^{*}\mathbf{g}), \\ {}^{\phi}\overline{\mathbf{D}}\ {}^{*}\overline{\theta} &= \ {}^{\phi}\overline{\mathbf{D}}\left[f_{*}(\ {}^{*}\theta)\right] = f_{*}(\ {}^{\phi}\mathbf{D}\ {}^{*}\theta), \end{split}$$

where for  $\phi$ -connections we use the symbol  $\phi \mathbf{D}$  instead of  $\nabla$ , and by local computations, we can verify that there are satisfied the conditions

$${}^{\phi}\overline{\mathbf{D}} {}^{*}\overline{\mathbf{g}} = 0 \Longleftrightarrow {}^{\phi}\mathbf{D} {}^{*}\mathbf{g} = 0,$$
$${}^{\phi}\overline{\mathbf{D}} {}^{*}\overline{\theta} = 0 \Longleftrightarrow {}^{\phi}\mathbf{D} {}^{*}\theta = 0.$$

Summarizing the above–presented results we get the proof of:

**Theorem 4.3** There is a unique canonical  $\phi$ -connection  $\phi \widehat{\mathbf{D}}$  with coefficients (42) generated by a Hamilton fundamental function H(x,p) which is almost symplectic and metric compatible and adapted to symplectic morphism.

We can use  ${}^{\phi}\widehat{\mathbf{D}}$  and frames (40) and (41) to construct instead of (37) an extended on  ${}^{*}\mathcal{W} \otimes \Lambda$  operator

$${}^{\phi}\widehat{\mathbf{D}}\left(a\otimes\lambda\right) \doteq \left( {}^{\phi}\mathbf{e}_{\alpha}(a) - u^{\beta} {}^{\phi}\widehat{\mathbf{\Gamma}}^{\gamma}_{\alpha\beta} {}^{z\phi}\mathbf{e}_{\alpha}(a) \right) \otimes \left( {}^{\phi}\mathbf{e}^{\alpha}\wedge\lambda\right) + a\otimes d\lambda, \quad (43)$$

where the local basis  ${}^{z\phi}\mathbf{e}_{\alpha}$  is  ${}^{\phi}\mathbf{e}_{\alpha}$  redefined in *z*-variables. This allows us to introduce a new class of operators adapted both to the N-connection structure and symplectic morphisms:

**Definition 4.5** The Fedosov-Hamilton operator-pairs  ${}^{\phi}\delta$  and  ${}^{\phi}\delta^{-1}$  on  ${}^*\mathcal{W}\otimes \Lambda$ , are defined

$${}^{\phi}\delta(a) = {}^{\phi}\mathbf{e}^{\alpha} \wedge {}^{z\phi}\mathbf{e}_{\alpha}(a), and {}^{\phi}\delta^{-1}(a) = \begin{cases} \frac{i}{p+q}z^{\alpha} & {}^{\phi}\mathbf{e}_{\alpha}(a), & if \ p+q > 0, \\ 0, & if \ p=q = 0, \end{cases}$$
(44)

where any  $a \in {}^*\mathcal{W} \otimes \mathbf{\Lambda}$  is homogeneous w.r.t. the grading  $\deg_s$  and  $\deg_a$ with  $\deg_s(a) = p$  and  $\deg_a(a) = q$ .

We note that the formulas (44) are different from (38) because they are defined for different nonholonomic distributions and related adapted frame structures.

Using differential calculus of forms on  $^*\mathcal{W} \otimes \Lambda$ , we prove

**Proposition 4.3** The torsion and curvature canonical *d*-operators of  ${}^{\phi}\widehat{\mathbf{D}}(a \otimes \lambda)$  (43) are computed

$${}^{z\phi}\widehat{\mathcal{T}} \stackrel{:}{:=} \frac{z^{\gamma}}{2} {}^{*}\theta_{\gamma\tau} {}^{\phi}\widehat{\mathbf{T}}^{\tau}_{\alpha\beta}(u) {}^{\phi}\mathbf{e}^{\alpha} \wedge {}^{\phi}\mathbf{e}^{\beta}, \tag{45}$$

and

$${}^{z\phi}\widehat{\mathcal{R}} \doteq \frac{z^{\gamma}z^{\varphi}}{4} * \theta_{\gamma\tau} \quad {}^{\phi}\widehat{\mathbf{R}}^{\tau}_{\ \varphi\alpha\beta}(u) \quad {}^{\phi}\mathbf{e}^{\alpha} \wedge {}^{\phi}\mathbf{e}^{\beta}, \tag{46}$$

By straightforward verifications, one gets the proof of

**Theorem 4.4** Any Fedosov–Hamilton operator–pairs (44) and related extended operator–pair (43) are defined by torsion (45) and curvature (46) following formulas:

$$\begin{bmatrix} \ \ \phi \widehat{\mathbf{D}}, \ \ \phi \delta \end{bmatrix} = \frac{i}{v} a d_{Wick} (^{z\phi} \widehat{\mathcal{T}}) \ and \quad \ \ \phi \widehat{\mathbf{D}}^2 = -\frac{i}{v} a d_{Wick} (^{z\phi} \widehat{\mathcal{R}}), \tag{47}$$

where  $[\cdot, \cdot]$  is the deg<sub>a</sub>-graded commutator of endomorphisms of  $*\mathcal{W} \otimes \mathbf{\Lambda}$ and  $ad_{Wick}$  is defined via the deg<sub>a</sub>-graded commutator in  $(*\mathcal{W} \otimes \mathbf{\Lambda}, \circ)$ .

The formulas (47) can be restated for any metric compatible d-connection and  $*\phi$ -connection structures on  $T^*M$ . Finally, we conclude that the operators constructed in this section are invariant under diffeomorphisms and in particular under symplectic morphisms.

## 5 Deformation Quantization of Hamilton and Einstein Spaces

The aim of this section is to provide the main Fedosov's theorems for Hamilton spaces and show how the Einstein manifolds can be encoded into the topological structure of such quantized nonholonomic spaces.

#### 5.1 Fedosov's theorems for connection-pairs

The theorems will be formulated for the canonical  $\phi$ -connection  $\phi \widehat{\mathbf{D}}$  (42).

**Theorem 5.1** A Hamilton fundamental function H(x, p) defines a flat Fedosov \* $\phi$ -connection

$${}^{\phi}\widehat{\mathcal{D}} \doteqdot - {}^{\phi}\delta + {}^{\phi}\widehat{\mathbf{D}} - \frac{i}{v}ad_{Wick}({}^{\phi}r)$$

satisfying the condition  ${}^{\phi}\widehat{\mathcal{D}}^2 = 0$ , where the unique element  ${}^{\phi}r \in {}^*\mathcal{W} \otimes \mathbf{\Lambda}$ ,  $\deg_a({}^{\phi}r) = 1, {}^{\phi}\delta^{-1} {}^{\phi}r = 0$ , is a solution of equation

$${}^{\phi}\delta {}^{\phi}r = {}^{\phi}\widehat{\mathcal{T}} + {}^{\phi}\widehat{\mathcal{R}} + {}^{\phi}\widehat{\mathbf{D}} {}^{\phi}r - \frac{i}{v} {}^{\phi}r \circ {}^{\phi}r.$$

The solution for  $\,^{\phi}r$  can be computed recursively with respect to the total degree Deg in the form

$$\begin{split} {}^{\phi}r^{(0)} &= {}^{\phi}r^{(1)} = 0, \ {}^{\phi}r^{(2)} = {}^{\phi}\delta^{-1} \ {}^{\phi}\widehat{\mathcal{T}}, \\ {}^{\phi}r^{(3)} &= {}^{\phi}\delta^{-1} \left( {}^{\phi}\widehat{\mathcal{R}} + {}^{\phi}\widehat{\mathbf{D}} \ {}^{\phi}r^{(2)} - \frac{i}{v} \ {}^{\phi}r^{(2)} \circ {}^{\phi}r^{(2)} \right), \\ {}^{\phi}r^{(k+3)} &= {}^{\phi}\delta^{-1} \left( {}^{\phi}\widehat{\mathbf{D}} \ {}^{\phi}r^{(k+2)} - \frac{i}{v} \sum_{l=0}^{k} {}^{\phi}r^{(l+2)} \circ {}^{\phi}r^{(l+2)} \right), k \ge 1, \end{split}$$

where  $a^{(k)}$  is the Deg-homogeneous component of degree k of an element  $a \in {}^*\mathcal{W} \otimes \mathbf{\Lambda}$ .

**Proof.** It follows from a local component calculus with  $*\phi$ -adapted coefficients of  $\[\phi \widehat{\mathbf{D}}\]$ , see similar considerations in [5, 6, 35].

For Hamilton spaces, we can define a canonical star–product. By an explicit construction, we prove:

**Theorem 5.2** A \* $\phi$ -adapted star-product for Hamilton spaces is defined on  $C^{\infty}(T^*M)[[v]]$  by formula

$${}^{1}f * {}^{2}f \doteqdot \sigma(\tau({}^{1}f)) \circ \sigma(\tau({}^{2}f)),$$

where the projection  $\sigma: {}^*\mathcal{W} \to C^{\infty}(T^*M)[[v]]$  onto the part of  $\deg_s$ -degree zero is a bijection and the inverse map  $\tau: C^{\infty}(T^*M)[[v]] \to {}^*\mathcal{W}$  can be calculated recursively w.r.t the total degree Deg,

$$\tau(f)^{(0)} = f \text{ and, for } k \ge 0,$$
  
$$\tau(f)^{(k+1)} = \check{\delta}^{-1} \left( \ {}^{\phi} \widehat{\mathbf{D}} \tau(f)^{(k)} - \frac{i}{v} \sum_{l=0}^{k} ad_{Wick} (\ {}^{\phi} r^{(l+2)}) (\tau(f)^{(k-l)}) \right).$$

Let  ${}^{f}X$  the Hamiltonian vector field corresponding to a function  $f \in C^{\infty}(T^*M)$  on space  $(T^*M, {}^*\theta)$  and consider the antisymmetric part

$${}^{-}C({}^{1}f, {}^{2}f) \doteq \frac{1}{2} \left( C({}^{1}f, {}^{2}f) - C({}^{2}f, {}^{1}f) \right)$$

of bilinear operator  $C({}^{1}f, {}^{2}f)$ . A star–product (34) is normalized if  ${}_{1}C({}^{1}f, {}^{2}f) = \frac{i}{2}\{{}^{1}f, {}^{2}f\}$ , where  $\{\cdot, \cdot\}$  is the Poisson bracket, see (13). For the normalized \*, the bilinear operator  ${}_{2}^{-}C$  defines a de Rham–Chevalley 2–cocycle, when there is a unique closed 2–form  ${}^{\phi}\varkappa$  such that

$${}_{2}C({}^{1}f, {}^{2}f) = \frac{1}{2} {}^{\phi}\varkappa({}^{f_{1}}X, {}^{f_{2}}X)$$
(48)

for all  ${}^{1}f, {}^{2}f \in C^{\infty}(T^{*}M)$ . This is used to introduce  $c_{0}(*) \doteq [{}^{\phi}\varkappa]$  as the equivalence class.

Computing  $_2C$  from (48) and using the Theorem 5.2, we get the proof for

**Lemma 5.1** The unique 2-form defined by the unique canonical connection--pair can be computed

$${}^{\phi}\varkappa = -\frac{i}{8} * \mathbf{J}_{\tau}^{\alpha'} \; {}^{\phi}\widehat{\mathcal{R}}_{\alpha'}^{\tau} - \frac{i}{6}d\left( * \mathbf{J}_{\tau}^{\alpha'} \; {}^{\phi}\widehat{\mathbf{T}}_{\alpha'\beta}^{\tau} \; {}^{\phi}\mathbf{e}^{\beta} \right).$$

Let us define the canonical class  ${}^{\phi}\varepsilon$ , for the splitting (39). We can perform a distinguished complexification of such second order tangent bundles in the form  $T_{\mathbb{C}} \left( {}^{\phi}TT^*M \right) = T_{\mathbb{C}} \left( hT^*M \right) \oplus T_{\mathbb{C}} \left( wT^*M \right)$  and introduce  ${}^{\phi}\varepsilon$ as the first Chern class of the distributions  $T'_{\mathbb{C}} \left( {}^{\phi}TT^*M \right) = T'_{\mathbb{C}} \left( hT^*M \right) \oplus$  $T'_{\mathbb{C}} \left( wT^*M \right)$  of couples of vectors of type (1,0) both for the h- and w-parts. We compute  ${}^{\phi}\varepsilon$  using  ${}^{\phi}\widehat{\mathbf{D}}$  and the h- and v-projections  $h\Pi = \frac{1}{2}(Id_h - iJ_h)$ and  $v\Pi = \frac{1}{2}(Id_v - iJ_v)$ , where  $Id_h$  and  $Id_v$  are respective identity operators and  $J_h$  and  $J_v$  are almost complex operators, which are projection operators onto corresponding (1,0)-subspaces. Introducing the matrix  $(h\Pi, v\Pi) \ \widehat{\mathcal{R}} \left( h\Pi, v\Pi \right)^T$ , where  $(...)^T$  means transposition, as the curvature matrix of the N-adapted restriction of  ${}^{\phi}\widehat{\mathbf{D}}$  to  $T'_{\mathbb{C}} \left( {}^{\phi}TT^*M \right)$ , we compute the closed Chern-Weyl form

Using the canonical class  ${}^{\phi}\varepsilon \doteq [{}^{\phi}\gamma]$ , we prove:

**Theorem 5.3** The zero-degree cohomology coefficient  $c_0(*)$  for the almost Kähler model of a Hamilton space  $*\mathbf{K}^{2n}$ , defined by a Hamilton fundamental function H(x, p) is computed  $c_0(*) = -(1/2i)^{\phi} \varepsilon$ .

The coefficient  $c_0(*)$  contains as a particular case the class of zero-degree cohomologies computed for a metric of type  $g_{i'j'}(x)$  on M, defining a solution of the Einstein equations and lifted on cotangent bundle by formula (1). In such cases, this zero-degree coefficient defines certain quantum properties of the gravitational field. A more rich geometric structure should be considered if we define a value similar to  $c_0(*)$  encoding the information about Einstein manifolds lifted to the cotangent bundle.

#### 5.2 Quantum gravitational field equations

Any solution in classical Einstein gravity can be embedded into a Hamilton space model and quantized on cotangent bundle following the Fedosov quantization adapted to  $\phi$  and d-connections. Considering a de-quantization formalism [36], we construct certain quantum deformations of the classical Einstein configurations in the classical limit. Such a model defines a nonholonomic almost Kähler generalization of the Einstein gravity on cotangent bundle. The solutions for the "cotangent" gravity are, in general, with violation of Lorentz symmetry induced by quantum corrections. The nature of such quantum gravity corrections is different from those defined by Finsler-Lagrange models on tangent bundle (see, for instance, [33, 34, 43, 44, 45])), locally anisotropic string gravity [40, 41, 42] with corrections from extradimensions and nonholonomic spinor gravity [37, 38, 39] and noncommutative gravity, see reviews of results in [29, 30]. The aim of this section is to analyze how a generalization of Einstein gravity can be performed on cotangent bundles in terms of canonical  $\phi$ -connections, with geometric structures induced by an effective Hamiltonian fundamental function, when the Fedosov quantization can be naturally performed.

For a canonical \* $\phi$ -connections  ${}^{\phi}\widehat{\mathbf{D}} = \{ {}^{\phi}\mathbf{\Gamma} \}$  (42), we can define the Ricci tensor,

$$Ric(\ ^{\phi}\widehat{\mathbf{D}}) = \{\ ^{\phi}\widehat{\mathbf{R}}_{\ \beta\gamma} \doteq \ ^{\phi}\widehat{\mathbf{R}}_{\ \beta\gamma\alpha}^{\alpha}\},\$$

and the scalar curvature,  ${}^{\phi}R \doteq {}^{*}\mathbf{g}^{\alpha\beta} {}^{\phi}\widehat{\mathbf{R}}_{\alpha\beta}$ . On cotangent bundle  $T^{*}M$ , we postulate the field equations

$${}^{\phi}\widehat{\mathbf{R}}^{\underline{\alpha}}_{\ \beta} - \frac{1}{2} ({}^{\phi}R + \lambda) {}^{\phi}\mathbf{e}^{\underline{\alpha}}_{\ \beta} = 8\pi G \Upsilon^{\underline{\alpha}}_{\ \beta}, \tag{50}$$

where  $\Upsilon_{\beta}^{\underline{\alpha}}$  is the effective energy-momentum tensor,  $\lambda$  is the cosmological constant, G is the Newton constant in the units when the light velocity c = 1, and  ${}^{\phi}\mathbf{e}_{\beta} = {}^{\phi}\mathbf{e}_{\beta}^{\underline{\alpha}}\partial/\partial u^{\underline{\alpha}}$  is the  $*\phi$ -adapted base (40).

We consider the effective source 3–form

$$\overleftarrow{\Upsilon}_{\beta} = \Upsilon^{\underline{\alpha}}_{\ \beta} \ \epsilon_{\underline{\alpha}\underline{\beta}\underline{\gamma}\underline{\delta}} du^{\underline{\beta}} \wedge du^{\underline{\gamma}} \wedge du^{\underline{\delta}},$$

where  $\epsilon_{\alpha\beta\gamma\delta}$  is the absolute antisymmetric tensor, and the action for the "cotangent" gravity and matter fields,

$$S[\mathbf{e}, \mathbf{\Gamma}, \phi] = \ ^{gr}S[\mathbf{e}, \mathbf{\Gamma}] + \ ^{matter}S[\mathbf{e}, \mathbf{\Gamma}, \phi].$$

**Theorem 5.4** The equations (50) can be represented as 3-form equations

$$\epsilon_{\alpha\beta\gamma\tau} \left( \ {}^{\phi}\mathbf{e}^{\alpha} \wedge \ {}^{\phi}\mathcal{R}^{\beta\gamma} + \lambda \ {}^{\phi}\mathbf{e}^{\alpha} \wedge \ {}^{\phi}\mathbf{e}^{\beta} \wedge \ {}^{\phi}\mathbf{e}^{\gamma} \right) = 8\pi G \overleftarrow{\Upsilon} \ _{\tau} \tag{51}$$

following from action by varying the components of  $\mathbf{e}_{\beta}$ , when

$$\overleftarrow{\Upsilon}_{\tau} = \ ^{m}\overleftarrow{\Upsilon}_{\tau} \ and \ \ ^{m}\overleftarrow{\Upsilon}_{\tau} = \ \ ^{m}\underline{\Upsilon}^{\underline{\alpha}}_{\tau}\epsilon_{\underline{\alpha}\underline{\beta}\underline{\gamma}\underline{\delta}}du^{\underline{\beta}} \wedge du^{\underline{\gamma}} \wedge du^{\underline{\delta}}_{\underline{\gamma}}$$

where  ${}^{m}\Upsilon^{\underline{\alpha}}_{\tau} = \delta \;^{matter}S/\delta \mathbf{e}_{\underline{\alpha}}^{\tau}$ .

**Proof.** It is a usual textbook and/or differential form calculus (see, for instance, [46, 14]). In our case, we have to use the \* $\phi$ -adapted bases (40) and (41) for  $\ensuremath{\,}^{\phi}\widehat{\mathbf{D}}$ .  $\Box$ 

The Chern–Weyl 2–form (49) can be used to define the quantum version<sup>3</sup> of Einstein equations (51) in the approaches with deformation quantization:

**Corollary 5.1** The quantum field equations on cotangent bundle generalizing the Einstein gravitations in general relativity are

$${}^{\phi}\mathbf{e}^{\alpha}\wedge {}^{\phi}\gamma = \epsilon^{\alpha\beta\gamma\tau}2\pi G \, {}^{*}\mathbf{J}_{\beta\gamma} \overleftarrow{\Upsilon} \, {}_{\tau} - \frac{\lambda}{4} \, {}^{*}\mathbf{J}_{\beta\gamma} \, {}^{\phi}\mathbf{e}^{\alpha}\wedge {}^{\phi}\mathbf{e}^{\beta}\wedge {}^{\phi}\mathbf{e}^{\gamma}.$$
(52)

**Proof.** Multiplying  ${}^{\phi}\mathbf{e}^{\alpha} \wedge \text{with (49)}$ , taking into consideration the equation (51), and introducing the almost complex operator  ${}^{*}\mathbf{J}_{\beta\gamma}$ , we get the almost symplectic form of the Einstein equations (52).  $\Box$ 

An explicit computation of  $\,^{\phi}\gamma$  for nontrivial matter fields has to be performed for a deformation quantization model with interacting gravitational and matter fields geometrized in terms of an almost Kähler model defined for spinor and fiber bundles on cotangent bundles.

## 6 Conclusions

In this paper we outlined a method of converting any regular Lagrange and Hamilton dynamics into equivalent almost Kähler geometries with canonical nonlinear connection (N-connection) and adapted almost symplectic structures. The formalism was performed to be invariant under symplectic morphisms and adapted to Legendre transforms of Lagrangians into Hamiltonians and inversely. The geometry of cotangent bundles endowed with

<sup>&</sup>lt;sup>3</sup>in the sense of deformation quantization (i.e. when quantum equations are derived following a deformation procedure) but not of perturbative quantum theory

nonholonomic distributions, formulated as certain Hamilton–Cartan spaces being dual to the corresponding Lagrange–Finsler spaces, described in Sections 2 and 3 presents a key prerequisite of this approach to deformation quantization.

Given a regular Lagrange (in particular, Finsler; and related Legendre transforms), or Hamilton (in particular, Cartan), generating function, we completely define the fundamental geometric objects of a Hamilton geometry modelled on cotangent bundle, inducing canonical almost symplectic and compatible symplectic connection. The connection–pair and canonical  $*\phi$ –connection uniquely constructed to preserve the invariance under symplectic morphisms are introduced in Section 4.2. Such geometric objects are crucial for defining Fedosov–Hamilton operator–pairs. This allows us to generalize the Fedosov's theorems for deformation quantization to the case of Hamilton spaces and to show how the Einstein manifolds can be encoded into such a quantization scheme.

Any classical solution of the Einstein equations can be lifted to the cotangent bundle, and embedded into a Hamilton space geometry using frame transforms variables depending both on spacetime and phase (momentum like) coordinates. Such variables can be defined from a de-quantization procedure like that considered in Ref. [36] but in our case re-formulated for nonholonomic cotangent bundles. In general, such constructions result in quasi-classical effects of quantum gravity with violations of local Lorentz symmetry. The surprising result advocated in this paper is that we can model classical and quantum gravitational effects by corresponding effective classical and quantum Hamilton mechanics systems. Nevertheless, certain additional phenomenological and/or experimental data for quantum gravity effects have to be assigned to the scheme in order to define the nonholonomic frame transforms of locally isotropic gravitational fields into quantized and semi-classical ones on cotangent bundle.

On cotangent bundles and curved phase spaces, there were developed different methods of quantization of nonlinear field theories and mechanical systems with nonholonomic constraints. For instance, in Refs. [47, 48], there were constructed in explicit form examples of "cotangent" star-products (in such cases, one derive certain compatible symplectic / Levi Civita type connections). The formalism was revised and developed by introducing auxiliary variables, with further restrictions, and/or higher spin systems, in analogy with gauge theories with generalized symmetries, in the framework of the BRST approach [49, 50, 51, 10, 1]. In general, the methods of quantization of nonlinear physical systems on contangent bundles are very different from that on tangent bundles. The source of this is in the fact that Lagrange and Hamilton (both classical and quantum) schemes may result in very different quantum models for nonlinear classical theories: On phase spaces, we have to consider Legendre transform and additional symmetries related to symplectic morphisms (i.e. morphisms preserving a symplectic structure into a symplectic structure). In such a case, the author of [19] had to work with symplectic groupoids and introduce contravariant connections which modified substantially the Fedosov scheme of quantization.

There are various ideas and approaches to quantize gravity theories, including deformation quantization. In our partner works [23, 24] we elaborated a direction related to effective Lagrange–Finsler geometries by performing nonholonomic deformations to quantum versions of (semi) Riemannian manifolds preserving the local Lorentz invariance at least for semiclassical approximations. Following the same geometric methods developed for Finsler spaces, but extended for nonholonomic manifolds, we proved that there are similar quantization schemes of gravity and Lagrange–Finsler spaces modelled on tangent bundles [21, 22]. Such constructions also result in violations of local Lorentz symmetry but with physical effects which are very different from those for models on cotangent bundles. The general conclusion of this paper is that a deformation quantization scheme for Hamilton spaces and related generalizations of the Einstein gravity on cotangent bundle results in more rich geometric structures and requests more advanced geometric methods with nonlinear connections and connection–pairs. We can elaborate a standard Fedosov formalism also on cotangent bundle (not involving groupoid structures) by introducing canonical nonlinear connections structures and generalizing the concept of linear and distinguished connections to that of connection pairs.

We are planing to compare different approaches to deformation quantization of gravity (preserving or violating the local Lorentz invariance) and other quantization schemes in our further works.

Acknowledgement: M. A. was partially supported by grant CNC-SIS,1158/2007, Romania. S. V. performed his work as a visitor at Fields Institute.

## References

- [1] I. Batalin, E. Fradkin and T. Fradkina, Nucl. Phys. B. 332, (1990).
- [2] B. de Wit and J. W. van Holten, Phys. Lett. B **79**, 389 (1978).

- [3] M. Grigoriev and S. Lyakhovich, Commun. Math. Phys. 218, 437 (2001).
- [4] M. Henneaux, Phys. Rep. **126**, 1 (1985).
- [5] B. Fedosov, J. Diff. Geom. 40, 213 (1994).
- [6] B. Fedosov, Deformation Quantization and Index Theory, Mathematical topics: 9 (Akademie–Verlag, Berlin, 1996).
- [7] M. Kontsevich, Lett. Math. Phys. 66, 157 (2003).
- [8] M. Kontsevich, Lett. Math. Phys. 48, 35 (1999).
- [9] P. Kazinski, S. Lyakhovich and A. Sharapov, JHEP 0507, 076 (2005).
- [10] S. Lyakhovich and A. Sharapov, JHEP **0503**, 011 (2005).
- [11] C. Castro, J. Geom. Phys. **33**, 173 (2000).
- [12] C. Castro, **36**, 2605 (2004).
- [13] C. Castro, Phys. Lett. B 668, 442 (2008).
- [14] C. Rovelli, *Quantum Gravity* (Cambridge University Press, 2004).
- [15] A. Ashtekar and J. Lewandowski, Class. Quant. Grav. 21, R53 (2004).
- [16] T. Thiemann, Modern Canonical Quantum General Relativity (Cambrdigde University Press, Cambridge, 2006).
- [17] I. Gorbunov, S. Lyakhovich and A. Sharapov, J. Geom. Phys. 53, 98 (2005).
- [18] I. Batalin, M. Grigoriev and S. Lyakhovich, J. Math. Phys. 46, 072301 (2005).
- [19] A. Karabegov, Commun. Math. Phys. **258**, 223 (2005).
- [20] F. Etayo, R. Santamaría and S. Vacaru, J. Math. Phys. 46, 032901 (2005).
- [21] S. Vacaru, J. Math. Phys. 48, 123509 (2007).
- [22] S. Vacaru, An. St. Univ. Al. I. Cuza din Iaşi (S.N.), Matematica, vol. LIII, 2007, Supliment, 327-342; e-print arXiv: 0707.1526.

- [23] S. Vacaru, Phys. Lett. A **372**, 2949 (2008).
- [24] S. Vacaru, e-print arXiv: 0709.3609.
- [25] G. Vrănceanu, C. R. Acad. Paris **103**, 852 (1926).
- [26] G. Vrănceanu, Bull. Fac. Șt. Cernăuți 5, 177 (1931).
- [27] G. Vrănceanu, *Leçons de Geometrie Differentielle*, Vol II (Edition de l'Academie de la Republique Populaire de Roumanie, 1957).
- [28] A. Bejancu and H. R. Farran, *Foliations and Geometric Structures* (Springer, 2005).
- [29] S. Vacaru, P. Stavrinos, E. Gaburov and D. Gonţa, Clifford and Riemann-Finsler Structures in Geometric Mechanics and Gravity, Selected Works, Differential Geometry – Dynamical Systems, Monograph 7 (Geometry Balkan Press, 2006); www.mathem.pub.ro/dgds/mono/va-t.pdf and e-print arXiv: gr-qc/0508023.
- [30] S. Vacaru, e-print arXiv: 0707.1524.
- [31] R. Miron, D. Hrimiuc, H. Shimada and V. S. Sabau, *The Geometry of Hamilton and Lagrange Spaces* (Kluwer Academic Publishers, Dordrecht, Boston, London, 2000).
- [32] R. Miron, *The Geometry of Higher–Order Hamilton Spaces* (Kluwer Academic Publishers, Dordrecht, Boston, London, 2003).
- [33] R. Miron and M. Anastasiei, *Vector Bundles and Lagrange Spaces with Applications to Relativity* (Geometry Balkan Press, Bukharest, 1997); translation from Romanian of (Editura Academiei Romane, 1987).
- [34] R. Miron and M. Anastasiei, The Geometry of Lagrange Spaces: Theory and Applications, FTPH no. 59 (Kluwer Academic Publishers, Dordrecht, Boston, London, 1994).
- [35] A. Karabegov and M. Schlichenmaier, Lett. Math. Phys. 57, 135 (2001).
- [36] A. Karabegov, Lett. Math. Phys. 65, 133 (2003).
- [37] S. Vacaru and P. Stavrinos, Spinors and Space–Time Anisotropy (Athens University Press, Greece, 2002); e–print: arXiv: gr-qc/0112028.

- [38] S. Vacaru, J. Math. Phys. **37**, 508 (1996).
- [39] S. Vacaru, JHEP, **09**, 011 (1998).
- [40] S. Vacaru, Ann. Phys. (NY), **256**, 39 (1997).
- [41] S. Vacaru, Nucl. Phys. B, **434**, 590 (1997).
- [42] N. Mavromatos, e-print: arXiv: 0708.2250.
- [43] F. Girelli, S. Liberati and L. Sindoni, Phys. Rev. D 75, 064015 (2007).
- [44] S. Mignemi, Phys. Rev. D 76, 047702 (2007).
- [45] G. Gibbons, J. Gomis and C. N. Pope, Phys. Rev. **D** 76, 081701 (2007).
- [46] C. Misner, K. Thorne and J. Wheeler, *Gravitation* (Freeman, 1973).
- [47] M. Bordemann, N. Neumaier and S. Waldman, Commun. Math. Phys. 198, 363 (1998).
- [48] M. Bordemann, N. Neumaier and S. Waldman, J. Geom. Phys. 29, 199 (1999).
- [49] G. Barnich, M. Grigoriev, A. Semikhatov and I. Tipunin, Commun. Math. Phys. 260, 147 (2005).
- [50] M. A. Vasiliev, Int. J. Geom. Meth. Mod. Phys. 3, 37 (2006).
- [51] M. Grigoriev, e-print arXiv: hep-th/0605089.
- [52] R. Miron and Gh. Atanasiu, Kodai Math. J. 6, 228 (1983).
- [53] Gh. Atanasiu, M. Hashiguchi and R. Miron, Rep. Fac. Sci. Kagoshima Univ. Math. Phys. Chem. No. 18, 19 (1985).
- [54] S. Vacaru, e-print: arXiv: 0806.3810.
- [55] S. Vacaru, SIGMA 4, 071 (2008); arXiv: 0806.3814.
- [56] S. Vacaru, to appear in Int. J. Theor. Phys. (2008); e-print: arXiv: 0806.3812.