# THE SELBERG TRACE FORMULA FOR HECKE OPERATORS ON COCOMPACT KLEINIAN GROUPS

#### JOSHUA S. FRIEDMAN

ABSTRACT. We compute the Selberg trace formula for Hecke operators (also called the trace formula for modular correspondences) in the context of cocompact Kleinian groups with finite-dimentional unitary representations. We give some applications to the distribution of Hecke eigenvalues, and give an analogue of Huber's theorem.

### 1. INTRODUCTION

1.1. **Motivation.** The Selberg trace formula has been well studied for cofinite Kleinian and Fuchsian groups. In the Fuchsian case: [Sel56, Roe66, VKF73, Hej76, Hej83, Ven82, Bus92, Iwa02]; in the Kleinian case: [EGM98, Fri05a, Fri05b]. For the most part (excepting [Hej76, Hej83]) the references above consider the *standard* Selberg trace formula which entails two objects: a discrete group  $\Gamma$ , and possibly a finite-dimensional unitary representation  $\chi$ . A third object to consider is a Hecke operator.

The Selberg trace formula (and the underlying Selberg spectral theory) for Hecke operators (also known as the Selberg trace formula for *modular correspondences*)<sup>1</sup> is far from a trivial extension of the standard Selberg trace formula. By considering the abundance of Hecke operators for the Modular group  $\Gamma = PSL(2, \mathbb{Z})$ , Lindenstrauss and Venkatesh ([LV07], [Gol06, Chap. 4]) gave a new proof<sup>2</sup> for the existence of infinitely many even Maaß forms<sup>3</sup> in  $L^2(\mathbb{H} \setminus \Gamma)$ .

Strömbergsson ([Str01]) made extensive use of the trace formula for Hecke operators to solve a problem in the *well known spectral correspondence for quaternion groups*—the Jacquet-Langlands correspondence ([Hej85]). More specifically, Strömbergsson completely determined the image of the Jacquet-Langlands correspondence.<sup>4</sup> In fact, in [Str98], he extended the Selberg trace formula for modular correspondences from the cocompact Fuchsian case ([Hej76]) to the cofinite Fuchsian case.

The possibility for a trace formula for Hecke operators started with Selberg [Sel89, pp. 444–446, 460–462, 504–505]. In [Hej76] Hejhal gave an explicit trace formula for modular correspondences for cocompact Fuchsian groups with finite-dimensional unitary representations and integer

<sup>&</sup>lt;sup>1</sup>Some authors seem to refer to the subject as *The Selberg trace formula for modular correspondences* and some as *The Selberg trace formula for Hecke operators*. For our purposes these are interchangeable. To be perfectly precise, we are not computing the trace of a Hecke operator, but rather the trace of  $\mathcal{M}h(\Delta)$ , where  $\mathcal{M}$  is a Hecke operator,  $\Delta$  is the Laplacacian, and h is a holomorphic function with a certain decay rate.

<sup>&</sup>lt;sup>2</sup>The old proof is Selberg's original method: The determinant of the scattering matrix is shown to have order that is sufficiently small.

<sup>&</sup>lt;sup>3</sup>Their proof works in much more generality.

<sup>&</sup>lt;sup>4</sup>The bijective correspondence between the nontrivial automorphic forms on the multiplicative group of a division quaternion algebra and certain cusp forms on GL(2).

weight  $k \ge 0$  forms. Akiyama and Tanigawa [AT90] proved the Selberg trace formula for modular correspondences for cofinite Fuchsian groups and gave explicit evaluations of the trace formula for the case of the congruence group  $\Gamma_0(p)$  (p prime). Hoffmann ([Hof94, Hof88]) approached the trace formula for Hecke operator, from the representation-theoretic point of view.

Strömbergsson ([Str98]) gave a very detailed derivation of the Selberg trace formula for modular correspondences in the context of weight-zero forms (functions) and trivial unitary representations on cofinite Fuchsian groups. He also gave explicit formulas for the case of  $\Gamma = \Gamma_0(N)$  (*N* is square-free).

In this work, we derive the Selberg trace formula for Hecke operators for cocompact Kleinian groups with finite-dimensional unitary representations. In order for the formula to apply, the cocompact group  $\Gamma$  needs to have its commensurator **Comm**( $\Gamma$ ) (in PSL(2,  $\mathbb{C}$ )) strictly larger than itself. For each  $\alpha \in$ **Comm**( $\Gamma$ )\ $\Gamma$ , we obtain a Hecke operator  $\mathcal{M}$ . When  $\Gamma$  is *arithmetic*, [**Comm**( $\Gamma$ ) :  $\Gamma$ ] =  $\infty$ ; in fact **Comm**( $\Gamma$ ) is a dense subgroup of PSL(2,  $\mathbb{C}$ ) ([MR03, p. 271, Exer. 8.4.5]).

1.2. **Main results.** In this section we state our main results, and briefly describe the notation and preliminary material needed to state them (for more details see §2).

Let  $\Gamma < PSL(2, \mathbb{C})$  be a cocompact Kleinian group acting on hyperbolic three-space  $\mathbb{H}^3$ . Let V be a finite-dimensional complex inner product space with inner-product  $\langle , \rangle_V$ , and let  $Rep(\Gamma, V)$  denote the set of finite-dimensional unitary representations of  $\Gamma$  in V. Let  $\mathcal{F} \subset \Gamma$  be a (compact) fundamental domain for the action of  $\Gamma$  in  $\mathbb{H}^3$ .

Let  $\chi \in \text{Rep}(\Gamma, V)$ . The Hilbert space of  $\chi$ -*automorphic* functions is the set of measurable functions

$$\begin{split} \mathcal{H}(\Gamma,\chi) &\equiv \{f:\mathbb{H}^3 \to V \mid f(\gamma P) = \chi(\gamma)f(P) \; \forall \gamma \in \Gamma, P \in \mathbb{H}^3, \\ & \text{and} \; \langle f,f \rangle \equiv \int_{\mathcal{F}} \langle f(P),f(P) \rangle_V \; dv(P) < \infty \}. \end{split}$$

Finally, let  $\Delta = \Delta(\Gamma, \chi)$  be the corresponding positive self-adjoint Laplace-Beltrami operator on  $\mathcal{H}(\Gamma, \chi)$ .

The *commensurator* subgroup **Comm**( $\Gamma$ ) is the set of all  $\alpha \in PSL(2, \mathbb{C})$  such that both  $[\Gamma : \Gamma \cap \alpha^{-1}\Gamma\alpha] < \infty$  and  $[\alpha^{-1}\Gamma\alpha : \Gamma \cap \alpha^{-1}\Gamma\alpha] < \infty$ .

Throughout this paper we will consider elements  $\alpha \in \mathbf{Comm}(\Gamma)$  that satisfy:

- (1) the element  $\alpha \in \mathbf{Comm}(\Gamma) \setminus \Gamma$ ;
- (2) the unitary representation  $\chi \in \operatorname{Rep}(\Gamma, V)$  has a single-valued extension from  $\Gamma$  to the set  $\Gamma \alpha \Gamma$ , and satisfies the following properties:
  - (a)  $\chi(g_1 \alpha g_2) = \chi(g_1) \chi(\alpha) \chi(g_2)$  for  $g_1, g_2 \in \Gamma$ ;
  - (b)  $\chi(\alpha)$  is an invertible linear map on the inner product space *V*.

Associated to the element  $\alpha$  is the *Hecke operator*  $\mathcal{M} : \mathcal{H}(\Gamma, \chi) \mapsto \mathcal{H}(\Gamma, \chi)$ . The Hecke operator commutes with  $\Delta$ .

Let  $\mathcal{D}$  be an indexing set for the set of eigenvalues of  $\Delta$ . For each eigenvalue  $\lambda = \lambda_m \ (m \in \mathcal{D})$ , let  $A_{\lambda} \equiv A(\lambda, \Gamma, \chi)$  denote the subspace of  $\mathcal{H}(\Gamma, \chi)$  spanned by  $\{\lambda_m \mid \lambda_m = \lambda\}$ . The subspaces  $A_{\lambda}$ are invariant under  $\mathcal{M}$ . Let  $\lambda_{m_1} \dots \lambda_{m_{k_{\lambda}}}$  generate  $A_{\lambda}$  ( $\lambda$  has multiplicity  $k_{\lambda}$ ). Since  $A_{\lambda}$  is invariant under  $\mathcal{M}$ , the eigenvalues of  $\mathcal{M}|_{A_{\lambda}}$  can be listed as  $\omega_{m_1} \dots \omega_{m_{k_{\lambda}}}$ . Decomposing  $\mathcal{H}(\Gamma, \chi)$  into invariant subspaces  $A_{\lambda}$ , for each  $m \in \mathcal{D}$ , we pair  $\lambda_m$  and  $\omega_m$ . We next state our main result, the Selberg trace formula for Hecke operators. The various notations are described very briefly following the theorem; for more details see §3.

Our main result is the following:

**Theorem.** (Selberg trace formula) Let  $\Gamma$  be a cocompact Kleinian group. Let  $\alpha \in \text{Comm}(\Gamma) \setminus \Gamma$ , and let  $\chi$  be a representation satisfying Assumption 2.1. Let h be a holomorphic function on  $\{s \in \mathbb{C} \mid |\operatorname{Im}(s)| < 2+\delta\}$  for some  $\delta > 0$ , satisfying  $h(1 + z^2) = O(1 + |z|^2)^{3/2-\epsilon}$  as  $|z| \to \infty$ , and let

$$g(x) = \frac{1}{2\pi} \int_{\mathbb{R}} h(1+t^2) e^{-itx} dt.$$

Then

$$\sum_{m \in \mathcal{D}} h(\lambda_m) \omega_m = \sum_{\{R\}_{\text{ell}}} \frac{\operatorname{tr}_V(\chi(R^{-1})^*) g(0) \log N(T_0)}{|\mathcal{E}(R)| |(\operatorname{tr}(R))^2 - 4|} + \sum_{\{T\}_{\text{lox}}} \frac{\operatorname{tr}_V(\chi(T^{-1})^*) g(\log N(T))}{|\mathcal{E}(T)| |a(T) - a(T)^{-1}|^2} \log N(T_0)$$

Here,  $\{\lambda_m\}_{m\in\mathcal{D}}$  are the eigenvalues of  $\Delta$  counted with multiplicity, and  $\omega_m$  are the eigenvalues of  $\mathcal{M}$  (the Hecke operator associated to  $\alpha$ ) with the convention of Equation 2.6. The summation with respect to  $\{R\}_{\text{ell}}$  extends over the finitely many  $\Gamma$ -conjugacy classes of elliptic elements  $R \in \Gamma \alpha^{-1}\Gamma$ , and for such a class,  $N(T_0)$  is the minimal norm of a hyperbolic or loxodromic element of the centralizer  $\mathcal{C}(R) \subset \Gamma$ ;  $\mathcal{E}(R)$  is the maximal finite subgroup contained in  $\mathcal{C}(R)$  (see Lemma 3.5 for more details). The summation with respect to  $\{T\}_{\text{lox}}$  extends over the  $\Gamma$ -conjugacy classes of hyperbolic or loxodromic elements of  $\Gamma \alpha^{-1}\Gamma$ ,  $T_0$  denotes a primitive hyperbolic or loxodromic element of minimal norm in<sup>5</sup>  $\mathcal{C}(T) \subset \Gamma$ ; the element T is conjugate in PSL(2,  $\mathbb{C}$ ) to the transformation described by the diagonal matrix with diagonal entries  $a(T), a(T)^{-1}$  with |a(T)| > 1, and<sup>6</sup>  $N(T) = |a(T)|^2$ ;  $\mathcal{E}(T)$  is the finite cyclic elliptic subgroup of  $\mathcal{C}(T)$  (see Lemma 3.3 for more details). The sum over elliptic elements is finite, and all other sums converge absolutely.

Please keep in mind that the conjugacy classes  $\{R\}_{ell}$  and  $\{T\}_{lox}$ , the operator  $\mathcal{M}$ , and the eigenvalues  $\omega_m$  all depend on  $\alpha \in \mathbf{Comm}(\Gamma) \setminus \Gamma$ .

Our first application of the trace formula is to study the distribution of the Hecke eigenvalues.

Define the *elliptic number of*  $\Gamma$  *with respect to*  $\alpha$ ,  $E_{\alpha}$ , by the finite sum

$$E_{\Gamma}^{\alpha} \equiv \sum_{\{R\}_{\text{ell}}} \frac{\log N(T_0)}{|\mathcal{E}(R)| \left| (\operatorname{tr}(R))^2 - 4 \right|}$$

Here the sum is over *elliptic conjugacy classes* of the set  $\Gamma \alpha^{-1} \Gamma$ , where conjugacy is defined with respect to the group  $\Gamma$ . The notations above are defined in §3.5.

We have

**Theorem.** Let  $\Gamma$  be a cocompact Kleinian group with  $\alpha \in Comm(\Gamma) \setminus \Gamma$ . Then

$$\sum_{m \in \mathcal{D}} \omega_m e^{-\lambda_m t} = \frac{E_{\Gamma}^{\alpha}}{\sqrt{4\pi t}} + O(\sqrt{t}) \quad \text{as } t \to 0^+.$$

<sup>&</sup>lt;sup>5</sup>Note that there is no clear relationship between  $T \in \Gamma \alpha^{-1} \Gamma$  and  $T_0 \in \Gamma$ .

<sup>&</sup>lt;sup>6</sup>Please note that there is a typographical error in the loxodromic and non cuspidal elliptic terms in [EGM98] Theorem 6.5.1; both terms are missing a factor of  $\frac{1}{4\pi}$ .

Here  $E_{\Gamma}^{\alpha}$  is the elliptic number of  $\Gamma$  with respect to  $\alpha$ ,  $\{\lambda_m\}_{m \in D}$  are the eigenvalues of  $\Delta$  counted with multiplicity, and  $\omega_m$  are the eigenvalues of  $\mathcal{M}$  (the Hecke operator associated to  $\alpha$ ) with the convention of Equation 2.6.

# If the set $\Gamma \alpha^{-1} \Gamma$ contains no elliptic elements we have

**Theorem.** Let  $\Gamma$  be a cocompact Kleinian group with  $\alpha \in Comm(\Gamma) \setminus \Gamma$ . Suppose that the set  $\Gamma \alpha^{-1}\Gamma$  contains no elliptic elements. Then

$$\sum_{m \in \mathcal{D}} \omega_m e^{-\lambda_m t} = O\left(t^{-1/2} \exp(-c/t)\right) \quad \text{as } t \to 0^+$$

Hence

$$\lim_{t \to 0^+} \sum_{m \in \mathcal{D}} \omega_m e^{-\lambda_m t} = 0$$

*Here*  $\{\lambda_m\}_{m \in D}$  *are the eigenvalues of*  $\Delta$  *counted with multiplicity, and*  $\omega_m$  *are the eigenvalues of*  $\mathcal{M}$  *(the Hecke operator associated to*  $\alpha$ *) with the convention of Equation 2.6.* 

## Finally, we give an analogue of Huber's theorem.

Following [EGM98], we define the length spectrum of  $\Gamma \alpha^{-1} \Gamma$ . For loxodromic  $T_j \in \Gamma \alpha^{-1} \Gamma$  set  $\mu_j = \log N(T_j)$ . The *length spectrum* of  $\Gamma \alpha^{-1} \Gamma$  is defined to be

$$\mathcal{L}_{\Gamma}^{\alpha} \equiv \left( \mu_j, \sum_{\substack{\{T\}_{lax}\\\log N(T) = \mu_j}} \frac{\log N(T_0)}{|\mathcal{E}(T)| |a(T) - a(T)^{-1}|^2} \right)_{j \ge 1}.$$

In the two-dimensional case the length spectrum simply comprises the lengths of closed geodesics. Here we really need the *complex lengths* a(T) and the order of the elliptic, finite subgroup of C(T),  $|\mathcal{E}(T)|$ .

We define the eigenvalue spectra of  $\Delta$  and  $\mathcal{M}$  ( $\mathcal{M}$  is defined from  $\alpha \in \mathbf{Comm}(\Gamma) \setminus \Gamma$ ) by

$$\mathcal{S}_{\Gamma}^{\alpha} = (\lambda_j, \omega(\lambda_j))_{j \in \mathcal{D}^*}$$

Here the symbol  $D^*$  means that we do not count with multiplicity, and  $\omega(\lambda)$  is the trace of  $\mathcal{M}$  on the invariant subspace generated by all eigenfunctions (of  $\Delta$ ) with eigenvalue  $\lambda$ .

**Theorem.** Let  $\Gamma, \Gamma'$  be cocompact Kleinian groups with  $\alpha \in Comm(\Gamma) \setminus \Gamma$ ,  $\alpha' \in Comm(\Gamma') \setminus \Gamma'$ . Then the following hold:

(1) Suppose that  $S_{\Gamma}^{\alpha}$  and  $S_{\Gamma'}^{\alpha'}$  agree up to at most finitely many terms. Then

$$E_{\Gamma}^{\alpha} = E_{\Gamma'}^{\alpha'}$$
$$S_{\Gamma}^{\alpha} = S_{\Gamma'}^{\alpha'}$$
$$\mathcal{L}_{\Gamma}^{\alpha} = \mathcal{L}_{\Gamma'}^{\alpha'}.$$

(2) Suppose that  $\mathcal{L}^{\alpha}_{\Gamma}$  and  $\mathcal{L}^{\alpha'}_{\Gamma'}$  agree up to at most finitely many terms. Then

$$E_{\Gamma}^{\alpha} = E_{\Gamma'}^{\alpha'}$$
$$\mathcal{S}_{\Gamma}^{\alpha} = \mathcal{S}_{\Gamma'}^{\alpha'}$$
$$\mathcal{L}_{\Gamma}^{\alpha} = \mathcal{L}_{\Gamma'}^{\alpha'}.$$

By applying the above Theorem to the case of a *single* fixed group with two different elements  $\alpha, \alpha' \in \mathbf{Comm}(\Gamma) \setminus \Gamma$ , we obtain:

**Corollary.** Let  $\Gamma$  be a cocompact Kleinian groups with  $\alpha, \alpha' \in Comm(\Gamma) \setminus \Gamma$ . Let  $\omega(\lambda_m) = \omega'(\lambda_m)$  for all but at most finitely many  $m \in \mathcal{D}^*$ . Then  $\omega(\lambda_m) = \omega'(\lambda_m)$  for all  $m \in \mathcal{D}^*$ .

# 2. Preliminaries

In this section we review the basic notions needed to evaluate the Selberg trace formula for modular correspondences. See [Hej76, Chapter 5] for the analogous Fuchsian case<sup>7</sup>.

Let  $\chi \in \text{Rep}(\Gamma, V)$ . The Hilbert space of  $\chi$ -*automorphic* functions is the set of measurable functions

$$\mathcal{H}(\Gamma,\chi) \equiv \{f: \mathbb{H}^3 \to V \mid f(\gamma P) = \chi(\gamma)f(P) \; \forall \gamma \in \Gamma, P \in \mathbb{H}^3, \\ \text{and } \langle f, f \rangle \equiv \int_{\mathcal{F}} \langle f(P), f(P) \rangle_V \; dv(P) < \infty \}.$$

Finally, let  $\Delta = \Delta(\Gamma, \chi)$  be the corresponding positive self-adjoint Laplace-Beltrami operator on  $\mathcal{H}(\Gamma, \chi)$ .

2.2. **Hecke operators.** In this section we define the Hecke operators (or modular correspondences). For more details see [Hej76, Chap. 5] for the cocompact Fuchsian case, [Str98] for the cofinite Fuchsian case.

Let  $\Gamma$  be a cocompact Kleinian group. The *commensurator* subgroup **Comm**( $\Gamma$ ) is the set of all  $\alpha \in PSL(2, \mathbb{C})$  such that both  $[\Gamma : \Gamma \cap \alpha^{-1}\Gamma \alpha] < \infty$  and  $[\alpha^{-1}\Gamma \alpha : \Gamma \cap \alpha^{-1}\Gamma \alpha] < \infty$ .

Let  $\chi \in \operatorname{Rep}(\Gamma, V)$ . By definition,  $\operatorname{Comm}(\Gamma) \subset \Gamma$ . However, in order to define non-trivial Hecke operators, we will need to start with an element  $\alpha \in \operatorname{Comm}(\Gamma)$  that lies outside of  $\Gamma$ . We will also need the unitary representation  $\chi$  to extend from  $\Gamma$  to the set  $\Gamma \alpha \Gamma$ . More precisely:

**Assumption 2.1.** *Throughout this paper we assume:* 

- (1) The element  $\alpha \in Comm(\Gamma) \setminus \Gamma$ ;
- (2) The unitary representation  $\chi \in \text{Rep}(\Gamma, V)$  has a single-valued extension from  $\Gamma$  to the set  $\Gamma \alpha \Gamma$ , and satisfies the following properties:
  - (a)  $\chi(g_1 \alpha g_2) = \chi(g_1)\chi(\alpha)\chi(g_2)$  for  $g_1, g_2 \in \Gamma$ ;
  - (b)  $\chi(\alpha)$  is an invertible linear map on the inner product space V.

See [Hej76, Pages 463-4] for the analogous Fuchsian case.

Let  $d = [\Gamma : \Gamma \cap \alpha^{-1}\Gamma \alpha]$ . It follows, by conjugation, that  $[\alpha^{-1}\Gamma \alpha : \Gamma \cap \alpha^{-1}\Gamma \alpha] = d$ .

<sup>&</sup>lt;sup>7</sup>There is much overlap between the Fuchsian case and our case, the Kleinian case. We will often cite the Fuchsian case, as a reference when proofs in the two cases are identical.

Let the symbol [] denote disjoint union. Then

$$\Gamma = \bigsqcup_{i=1}^{d} \left( \Gamma \cap \alpha^{-1} \Gamma \alpha \right) \epsilon_i \quad \text{iff} \quad \Gamma \alpha \Gamma = \bigsqcup_{i=1}^{d} \Gamma \alpha \epsilon_i.$$

Setting  $\alpha_i = \alpha \epsilon_i$  (*i* = 1...*d*), we define the operator  $\mathcal{M} : \mathcal{H}(\Gamma, \chi) \mapsto \mathcal{H}(\Gamma, \chi)$  by

$$(\mathcal{M}f)(P) = \sum_{i=1}^{d} \chi(\alpha_i)^* f(\alpha_i P).$$

The operator  $\mathcal{M}$  does not depend on the above choice of coset representatives, and it is a bounded linear operator on  $\mathcal{H}(\Gamma, \chi)$  ([Hej76, p. 467] or [Str98]).

The adjoint of  $\mathcal{M}$  can be constructed as follows: Since  $\alpha \in \mathbf{Comm}(\Gamma)$ , and  $\mathbf{Comm}(\Gamma)$  is a group;  $\alpha^{-1}$  also satisfies Assumption 2.1. As before, we can write

$$\Gamma \alpha^{-1} \Gamma = \bigsqcup_{i=1}^{\circ} \Gamma \beta_i$$

It follows that e = d ([Hej76, p. 469]). Define  $\mathcal{M}^* : \mathcal{H}(\Gamma, \chi) \mapsto \mathcal{H}(\Gamma, \chi)$  by

$$(\mathcal{M}^*f)(z) = \sum_{i=1}^d \chi(\beta_k^{-1}) f(\beta_k P).$$

For every  $f, g \in \mathcal{H}(\Gamma, \chi)$  we have ([Hej76, p. 470])

$$\langle \mathcal{M}f,g\rangle = \langle f,\mathcal{M}^*g\rangle.$$

2.3. Interplay between  $\Delta$  and  $\mathcal{M}$ . For P = z + rj,  $P' = z' + r'j \in \mathbb{H}^3$  set

(2.1) 
$$\delta(P,P') \equiv \frac{|z-z'|^2 + r^2 + r'^2}{2rr'}.$$

It follows that  $\delta(P, P') = \cosh(d(P, P'))$ , where *d* denotes the hyperbolic distance in  $\mathbb{H}^3$ . Next, for  $k \in \mathcal{S}([1, \infty))$  a Schwartz-class function, define the *point-pair invariant*  $K(P, Q) \equiv k(\delta(P, Q))$ . Note that for all  $\gamma \in PSL(2, \mathbb{C})$  we have  $K(\gamma P, \gamma Q) = K(P, Q)$ . Define the Poincaré series

$$K_{\Gamma}(P,Q) \equiv \sum_{T \in \Gamma} \chi(T) K(P,TQ).$$

The series above converges absolutely and uniformly on compact subsets of  $\mathbb{H}^3 \times \mathbb{H}^3$ , and is the kernel of a bounded operator  $\mathcal{K} : \mathcal{H}(\Gamma, \chi) \mapsto \mathcal{H}(\Gamma, \chi)$ . The Selberg theory allows us to *diagonalize*  $\mathcal{K}$  with respect to a basis of eigenfunctions of  $\Delta$ .

We first need the following spectral expansion: Each  $f \in \mathcal{H}(\Gamma, \chi)$  has an expansion of the form

(2.2) 
$$f(P) = \sum_{m \in \mathcal{D}} \langle f, e_m \rangle e_m(P)$$

Here the sum converges in the Hilbert space  $\mathcal{H}(\Gamma, \chi)$ , and  $\mathcal{D}$  is an indexing set of the eigenfunctions  $e_m$  of  $\Delta$  with corresponding eigenvalues  $\lambda_m$ .

Now, let *h* be the Selberg–Harish-Chandra transform of *k*; explicitly, for  $\lambda = 1 - s^2 \in \mathbb{C}$ ,

(2.3) 
$$h(\lambda) = h(1-s^2) \equiv \frac{\pi}{s} \int_1^\infty k\left(\frac{1}{2}\left(t+\frac{1}{t}\right)\right) (t^s - t^{-s})\left(t-\frac{1}{t}\right) \frac{dt}{t};$$

#### MODULAR CORRESPONDENCES

and let g be the Fourier transform of h:

(2.4) 
$$g(x) = \frac{1}{2\pi} \int_{\mathbb{R}} h(1+t^2) e^{-itx} dt.$$

For  $v, w \in V$  let  $v \otimes \overline{w}$  be a linear operator in V defined by  $v \otimes \overline{w}(x) = \langle x, w \rangle v$ . An immediate application of the spectral expansion (2.2) and the Selberg–Harish-Chandra transform (2.3) gives us (see [Fri05a, Fri05b] or [EGM98, Equation 6.4.10, page 278]):

**Lemma 2.2.** Let  $k \in S$  and  $h : \mathbb{C} \to \mathbb{C}$  be the Selberg–Harish-Chandra Transform of k. Then with  $K_{\Gamma}$  defined above, we have

(2.5) 
$$K_{\Gamma}(P,Q) = \sum_{m \in \mathcal{D}} h(\lambda_m) e_m(P) \otimes \overline{e_m(Q)}$$

*The sum converges absolutely and uniformly on compact subsets of*  $\mathbb{H}^3 \times \mathbb{H}^3$ *.* 

Since the Laplace operator  $\Delta$  commutes with the action of  $PSL(2, \mathbb{C})$  on  $\mathbb{H}^3$ , it follows that  $\Delta$  commutes with  $\mathcal{M}$ . From Selberg's original ideas ([Sel56]), the operator  $\mathcal{K} = h(\Delta)$ , and thus  $\mathcal{M}$  must also commute with  $\mathcal{K}$ . A direct proof of this fact is given in [Hej76, p. 473]:

**Lemma 2.3.** Let  $\mathcal{K}$  be the linear operator defined from the kernel function  $K_{\Gamma}(P,Q)$ . Then, on  $\mathcal{H}(\Gamma,\chi)$ 

$$\mathcal{KM} = \mathcal{MK},$$

and on  $\mathcal{H}(\Gamma, \chi) \cap C^2(\mathbb{H}^3)$ 

$$\mathcal{M} \Delta = \Delta \mathcal{M}.$$

For each eigenvalue  $\lambda = \lambda_m$   $(m \in D)$ , let  $A_{\lambda} \equiv A(\lambda, \Gamma, \chi)$  denote the subspace of  $\mathcal{H}(\Gamma, \chi)$  spanned by  $\{\lambda_m \mid \lambda_m = \lambda\}$ . Since  $\Delta$  commutes with  $\mathcal{M}$ , it follows that the subspaces  $A_{\lambda}$  are invariant under  $\mathcal{M}$ . Let  $m_1 \dots m_{k_{\lambda}} \in D$  generate  $A_{\lambda}$  ( $\lambda$  has multiplicity  $k_{\lambda}$ ). Since  $A_{\lambda}$  is invariant under  $\mathcal{M}$ , the eigenvalues of  $\mathcal{M}|_{A_{\lambda}}$  can be listed as  $\omega_{m_1} \dots \omega_{m_{k_{\lambda}}}$ . Next, we define

(2.6) 
$$\omega(\lambda) \equiv \operatorname{tr} \mathcal{M}|_{A_{\lambda}} = \sum_{j=1}^{k_{\lambda}} \omega_{m_j}.$$

Decomposing  $\mathcal{H}(\Gamma, \chi)$  into invariant subspaces  $A_{\lambda}$ , for each  $m \in \mathcal{D}$  we pair  $\lambda_m$  and  $\omega_m$ . Since  $\mathcal{M}$  is a bounded operator, we conclude that the  $\omega_m$  ( $m \in \mathcal{D}$ ) are uniformly bounded.

### 3. THE SELBERG TRACE FORMULA FOR HECKE OPERATORS

The standard Selberg trace formula is an explicit evaluation of both the integral and spectral of the operator  $\mathcal{K}$ . In our case of interest we evaluate the trace of  $\mathcal{KM}$ .

3.1. The kernel of operator  $\mathcal{KM}$ . The first step is to construct a Poincaré series for the kernel of  $\mathcal{KM}$ .

For all  $f \in \mathcal{H}(\Gamma, \chi)$ , by Lemma 2.3, and by definition,

$$\mathcal{KM}f(P) = \mathcal{MK}f(P) = \int_{\mathcal{F}} \mathcal{M}_P K_{\Gamma}(P,Q) f(Q) \, dv(Q).$$

Here  $\mathcal{M}_P K_{\Gamma}(P, Q)$  is the action of  $\mathcal{M}$ , treating  $K_{\Gamma}(P, Q)$  as a function of P. Set

(3.1) 
$$K_{\mathcal{M}}(P,Q) \equiv \sum_{T \in \Gamma \alpha^{-1} \Gamma} \chi(T^{-1})^* K(P,TQ),$$

it<sup>8</sup> then follows ([Hej76, p. 476]) that  $K_{\mathcal{M}}(P,Q) = \mathcal{M}_P K_{\Gamma}(P,Q)$ . Hence:

**Lemma 3.1.** For all  $f \in \mathcal{H}(\Gamma, \chi)$ ,

$$\mathcal{KM}f(P) = \mathcal{MK}f(P) = \int_{\mathcal{F}} K_{\mathcal{M}}(P,Q)f(Q) \, dv(Q).$$

Upon applying  $\mathcal{M}$  to (2.5), we obtain

(3.2) 
$$K_{\mathcal{M}}(P,Q) = \sum_{m \in \mathcal{D}} h(\lambda_m) \left( \mathcal{M}e_m(P) \right) \otimes \overline{e_m(Q)}.$$

The Selberg trace formula will result from a careful evaluation of

$$\operatorname{tr}(\mathcal{K}\mathcal{M}) \equiv \int_{\mathcal{F}} \operatorname{tr}_{V} K_{\mathcal{M}}(P, P) \, dv(P)$$

3.2. Spectral trace. Since  $K_{\mathcal{M}}(P,Q)$  is a continuous function on the compact set  $\mathcal{F} \times \mathcal{F}$ , the integral

$$\int_{\mathcal{F}} \operatorname{tr}_V K_{\mathcal{M}}(P, P) \, dv(P)$$

converges absolutely. Since M is bounded, the derivation of the standard Selberg trace formula for cofinite Kleinian groups [EGM98, §6.5] implies that

(3.3) 
$$\int_{\mathcal{F}} \operatorname{tr}_{V} K_{\mathcal{M}}(P, P) \, dv(P) = \sum_{m \in \mathcal{D}} h(\lambda_{m}) \omega_{m}.$$

3.3. Integral trace. Selberg's clever method of evaluating the integral (3.3) entails decomposing the Poincaré series  $K_{\mathcal{M}}(P,Q)$  into conjugacy classes of various types of isometries (the types being: elliptic, parabolic, hyperbolic, ...). A similar method works here, though conjugacy classes must be defined in the context of the set  $\Gamma \alpha^{-1} \Gamma$ . One also must deal with centralizer subgroups of various isometries, which lead to complications not found in the standard Selberg trace formula derivations.

For 
$$T \in \Gamma \alpha^{-1} \Gamma$$
 set

$$\{T\} \equiv \{T\}_{\Gamma} \equiv \{\sigma^{-1}T\sigma \mid \sigma \in \Gamma\},\$$

and set

$$\mathcal{C}(T) \equiv \mathcal{C}_{\Gamma}(T) \equiv \{ \sigma \in \Gamma \mid \sigma T = T\sigma \}.$$

<sup>&</sup>lt;sup>8</sup>Note that  $\chi$  is not necessarily unitary when extended to  $\Gamma \alpha \Gamma$ , so  $\chi(T^{-1})^*$  can not be simplified.

It now follows ([Hej76, pp. 480-482]) that

(3.4) 
$$\operatorname{tr}(\mathcal{KM}) = \sum_{\{T\}} \operatorname{tr}_V(\chi(T^{-1})^*) \int_{\mathcal{F}(\mathcal{C}(T))} K(P, TP) \, dv(P)$$

Here the sum is over all conjugacy classes of  $\Gamma \alpha^{-1} \Gamma$  with respect to  $\Gamma$ ;  $\mathcal{F}(\mathcal{C}(T))$  is a fundamental domain for the subgroup  $\mathcal{C}(T)$ .

Since the integral in (3.4) is invariant when T is conjugated within  $PSL(2, \mathbb{C})$ , it suffices to choose a conjugate of T which has the simplest form to allow one to explicitly evaluate the following integral: For  $T \in \Gamma \alpha^{-1} \Gamma$  set

(3.5) 
$$I(T) \equiv I_{\Gamma}(T) = \int_{\mathcal{F}(\mathcal{C}(T))} K(P, TP) \, dv(P).$$

There are four cases to consider:

- (1) T = id (this case does not occur by Assumption 2.1)
- (2) T is loxodromic
- (3) T is elliptic
- (4) T is parabolic.

In fact, we must first compute the structure of each centralizer subgroup C(T).

See [EGM98, chap. 1] for details on the action of  $PSL(2, \mathbb{C})$  on  $\mathbb{H}^3$ , and for the definitions of loxodromic, parabolic, ...

3.4. Loxodromic case. Suppose that  $T \in \Gamma \alpha^{-1} \Gamma$  is loxodromic. We can then assume (replacing *T* by its unique conjugate within PSL(2,  $\mathbb{C}$ )) that *T* has the form

$$T = \left(\begin{array}{cc} a(T) & 0\\ 0 & a(T)^{-1} \end{array}\right)$$

such that  $a(T) \in \mathbb{C}$  has |a(T)| > 1. Let N(T) denote the *norm* of *T*, defined by

$$N(T) \equiv |a(T)|^2.$$

For  $z + rj \in \mathbb{H}^3$ (3.6)  $T(z + rj) = a(T)^2 z + N(T)rj.$ 

The element *T* fixes two points: 0 and  $j\infty \in \widehat{\mathbb{C}}$ . It follows from elementary calculations that any  $\sigma \in \mathcal{C}(T) \subset \Gamma$  must be of the form

$$\left(\begin{array}{cc} u & 0\\ 0 & u^{-1} \end{array}\right).$$

Since  $\Gamma$  is discrete and cocompact, it follows ([EGM98, chap. 5]) that C(T) has at least one element of minimal norm. Choose one, say  $T_0$ . The other minimal norm elements are obtained by multiplying by the elliptic elements of the finite cyclic elliptic subgroup  $\mathcal{E}(T)$  of order m(T), generated by an element  $E_{T_0}$ . We now have

$$\mathcal{C}(T) = \langle T_0 \rangle \times \mathcal{E}(T) \,.$$

Here  $\langle T_0 \rangle = \{ T_0^n \mid n \in \mathbb{Z} \}$ . The elliptic element  $E_{T_0}$  is conjugate in  $PSL(2, \mathbb{C})$  to an element of the form

$$\left(\begin{array}{cc}\zeta(T_0) & 0\\ 0 & \zeta(T_0)^{-1}\end{array}\right),\,$$

where here  $\zeta(T_0)$  is a primitive 2m(T)-th root of unity.

Note that (by definition) if  $T \notin \Gamma$ ,  $T \notin C(T)$ . It seems conceivable then that C(T) could be a finite group. However, in the next lemma we will show that  $N(T_0) > 1$  which implies that C(T) contains an infinite cyclic loxodromic subgroup<sup>9</sup>.

**Lemma 3.2.** Let  $T \in \Gamma \alpha^{-1} \Gamma$ . Then C(T) contains a loxodromic element  $T_0$  of minimal norm  $N(T_0) > 1$ .

*Proof.* Let  $k \in S[1, \infty)$  be a smooth, positive bump function with  $k \equiv 1$  on [1, 2], (zero everywhere else) and let  $\chi$  be the trivial representation. The right-hand side of (3.3) converges absolutely, and by positivity the left side does, too. Hence (3.4) and (3.5) imply that

(3.7) 
$$\int_{\mathcal{F}(\mathcal{C}(T))} K(P, TP) \, dv(P) < \infty.$$

For  $\varphi \in \mathbb{R}$  set

$$R(\varphi) = \left(\begin{array}{cc} e^{\frac{i\varphi}{2}} & 0\\ 0 & e^{-\frac{i\varphi}{2}} \end{array}\right).$$

 $R(\varphi)$  is a rotation on  $\mathbb{H}^3$  that fixes  $0, \infty j \in \widehat{\mathbb{C}}$ . For  $z + rj \in \mathbb{H}^3$ ,

$$R(\varphi)(z+rj) = e^{i\varphi}z + rj.$$

Next, suppose there are not loxodromic elements in C(T), that is, that  $C(T) = \langle R(\varphi) \rangle$ , where  $\varphi \equiv \frac{2\pi}{m}$ , and  $m = |R(\varphi)| = |C(T)|$ . A fundamental domain for C(T) is given by (see [EGM98, p. 193])

$$\mathcal{F}_1 = \left\{ \rho e^{i\varphi} + rj \mid r > 0, \ \rho \ge 0, \ 0 \le \varphi \le \frac{2\pi}{m} \right\}$$

We claim that

(3.8) 
$$\int_{\mathcal{F}_1} K(P, TP) \, dv(P) = \infty$$

To see this, first note that using (3.6) and (2.1), we can simplify K(P, TP). We obtain

$$\int_{\mathcal{F}_1} K(P, TP) \, dv(P) = \int_{\mathcal{F}_1} k\left(\frac{|a(t)^2 - 1|^2 |z|^2 + (N(T)^2 + 1)^2 r^2}{2N(T)r^2}\right) \, \frac{dx \, dy \, dr}{r^3}$$

Applying the elementary substitution  $x \mapsto rx$ ,  $y \mapsto ry$ ; the integral becomes

$$\begin{split} \int_{\mathcal{F}_1} k \left( \frac{|a(t)^2 - 1|^2 |z|^2 + (N(T)^2 + 1)^2}{2N(T)} \right) \frac{dx \, dy \, dr}{r} \\ &= \int_0^\infty \frac{dr}{r} \int_{z(\mathcal{F}_1)} k \left( \frac{|a(t)^2 - 1|^2 |z|^2 + (N(T)^2 + 1)^2}{2N(T)} \right) \, dx \, dy = \infty. \end{split}$$

Here  $z(\mathcal{F}_1)$  is the standard projection of  $\mathcal{F}_1 \subset \mathbb{H}^3$  onto  $\mathbb{C}$ . Hence  $\mathcal{C}(T)$  must contain at least one loxodromic element, and since  $\Gamma$  is discrete,  $\mathcal{C}(T)$  must contain a minimally normed element.  $\Box$ 

<sup>&</sup>lt;sup>9</sup>This is closely analogous to the Fuchsian group case, where the centralizer of a hyperbolic element is infinite [Hej76, p. 483]. Another analogous result: "... In dimension 3, the centralizer of an elliptic element of a group with compact quotient never is a finite cyclic group. " [EGM98, p. 194].Meanwhile in dimension two, it is a finite cyclic group.

An element R in  $\Gamma$  is called *primitive* if there *does not* exist  $S \in \Gamma$  with  $S^m = R$  for some positive integer m. Since the  $T_0$  from Lemma 3.2 has minimal norm in C(T), it is necessarily primitive.

Now that we see that the structure of C(T) is identical to the case where  $T \in \Gamma$  ([EGM98, p. 193]), hence we can evaluate the integral (3.5), obtaining:

**Lemma 3.3.** Let  $T \in \mathbf{Comm}(\Gamma)$  be loxodromic (hyperbolic elements are considered loxodromic). Let

 $\mathcal{C}(T) = \langle T_0 \rangle \times \mathcal{E}(T)$ 

with  $T_0$  loxodromic<sup>10</sup> and primitive, and  $\mathcal{E}(T)$  a finite cyclic subgroup of elliptic elements. Then

$$\int_{\mathcal{F}(\mathcal{C}(T))} K(P, TP) \, dv(P) = \frac{g(\log N(T)) \log N(T_0)}{|\mathcal{E}(T)| \, |a(T) - a(T)^{-1}|^2}$$

Here, g comes from (2.4) and (2.3).

3.5. Elliptic case. Suppose that  $R \in \Gamma \alpha^{-1} \Gamma$  is elliptic. We can then assume (replacing *R* by its unique conjugate within  $PSL(2, \mathbb{C})$ ) that *R* has the form

$$R = R(\varphi) = \begin{pmatrix} e^{\frac{i\varphi}{2}} & 0\\ 0 & e^{-\frac{i\varphi}{2}} \end{pmatrix} \quad (\varphi \in \mathbb{R}).$$

 $R(\varphi)$  is a rotation on  $\mathbb{H}^3$  that fixes  $0, \infty j \in \widehat{\mathbb{C}}$ . For  $z + rj \in \mathbb{H}^3$ ,

$$R(\varphi)(z+rj) = e^{i\varphi}z + rj.$$

An elementary calculation shows that

$$\left(\begin{array}{cc}a&b\\c&d\end{array}\right)\in \mathrm{PSL}(2,\mathbb{C})$$

commutes with  $R(\varphi)$  iff one of the two conditions are satisfied:

(1) 
$$b = c = 0$$
  
(2)  $\exp \frac{i\varphi}{2} = \pm i$  and  $a = d = 0$ 

In both cases, a similar argument to Lemma 3.2 (see [EGM98, pp. 193–194]) shows that C(R) contains loxodromic elements. In particular C(R) contains a loxodromic element of minimal norm  $T_0$ . Following [EGM98, pp. 193–197] and generalizing to  $R \in \text{Comm}(\Gamma)$  we have ([EGM98, Thm. 5.2.1])<sup>11</sup>

**Lemma 3.4.** Let  $R \in \text{Comm}(\Gamma)$  be elliptic. Then there exists a primitive elliptic element  $R_0 \in C(R) \subset \Gamma$  (it is possible that  $R_0 = \text{id}$ ), an element  $T_0 \in C(R)$  loxodromic, of minimal norm in C(R) so that one of the following cases holds:

(1) Either all the elliptic elements of C(R) are contained in  $\mathcal{E}(R) \equiv \langle R_0 \rangle$ , and

$$\mathcal{C}(R) = \langle T_0 \rangle \times \mathcal{E}(R).$$

<sup>&</sup>lt;sup>10</sup>Note that there is *no* clear relationship between *T* and *T*<sub>0</sub> other than the fact that they have the same fixed points and commute. It seems reasonable to conjecture that  $T = T_0^{k/n}$  for  $k, n \in \mathbb{N}$ .

<sup>&</sup>lt;sup>11</sup>We modified their hypothesis to allow R to be in **Comm**( $\Gamma$ ) instead of in  $\Gamma$ . We are justified by the *proof* of Lemma 3.2.

(2) Or R is elliptic of order 2, and there exists an elliptic element  $S \in C(R)$  so that

$$\mathcal{E}(R) = \langle R_0 \rangle \cup \langle R_0 \rangle S$$

and

$$\mathcal{C}(R) = \langle T_0 \rangle \times \mathcal{E}(R).$$

See [EGM98, pp. 191-198] for more details.

Now that we see that the structure of C(T) is identical to the case where  $T \in \Gamma$  ([EGM98, p. 193]), we can evaluate the integral (3.5), obtaining:

**Lemma 3.5.** Let  $R \in \mathbf{Comm}(\Gamma)$  be elliptic. Let

$$\mathcal{C}(R) = \langle T_0 \rangle \times \mathcal{E}(R)$$

as in Lemma 3.4. Then

$$\int_{\mathcal{F}(\mathcal{C}(R))} K(P, RP) \, dv(P) = \frac{g(0) \log N(T_0)}{|\mathcal{E}(R)| \, |(\operatorname{tr}(R))^2 - 4|}$$

Here, g comes from (2.4) and (2.3).

We will also need the following:

**Lemma 3.6.** Let  $R \in \text{Comm}(\Gamma)$  be elliptic. Then there are only finitely many elliptic conjugacy classes of  $\Gamma \alpha^{-1} \Gamma$ , where conjugacy is defined with respect to  $\Gamma$ .

*Proof.* Let  $k \in S[1, \infty)$  be non-negative and let  $\chi$  be the trivial representation. By (3.4) and (3.3),

$$\sum_{\{R\}_{\rm ell}} \int_{\mathcal{F}(\mathcal{C}(R))} K(P,RP) \, dv(P) < \infty.$$

Hence by Lemma 3.5,

$$g(0) \sum_{\{R\}_{\text{ell}}} \frac{\log N(T_0)}{|\mathcal{E}(R)| \, |(\operatorname{tr}(R))^2 - 4|} < \infty.$$

Now, since  $N(T_0) \ge \delta > 1$  (independent of  $T_0$ ) and since  $-2 < \operatorname{tr}(R) < 2$ , in order for there to be infinitely many conjugacy classes, it is necessary for  $|\mathcal{E}(R)|$  to be unbounded as R goes through all conjugacy classes. But by Lemma 3.4 that would imply the order of each  $R_0$  is unbounded; a contradiction since cofinite Kleinian groups have only finitely many conjugacy classes of elliptic elements (and two elements in the same conjugacy class have the same order; see[EGM98]).

3.6. **Parabolic case.** Suppose that  $T \in \text{Comm}(\Gamma)$  is parabolic. We will soon see that this case can not occur when  $\Gamma$  is assumed to be cocompact. That is:  $\text{Comm}(\Gamma)$  contains no parabolic elements when  $\Gamma$  is cocompact. To see this, suppose that T is already conjugated into the simple form

$$T = \left(\begin{array}{cc} 1 & \tau \\ 0 & 1 \end{array}\right).$$

Then, an elementary calculation shows that

$$S \equiv \left(\begin{array}{cc} a & b \\ c & d \end{array}\right) \in \mathrm{PSL}(2,\mathbb{C})$$

commutes with *T* iff c = 0 and  $a = d = \pm 1$ . By definition, *S* must be parabolic (or the identity). But, since  $\Gamma$  is cocompact, it contains no parabolic elements. So C(T) = id. An argument similar to Lemma 3.2 shows that this can not happen. Hence there are no parabolic elements in **Comm**( $\Gamma$ )\ $\Gamma$ .

3.7. The Selberg trace formula. By putting together (3.4), Lemma 3.3, Lemma 3.5, Assumption 2.1, and the results of §3.6; and by applying standard approximation techniques in order to allow more general growth conditions on the function k (and hence on h and g) (see [Hej76, pp. 32–34] for the details), we obtain:

**Theorem 3.7.** (Selberg trace formula) Let  $\Gamma$  be a cocompact Kleinian group. Let  $\alpha \in \text{Comm}(\Gamma) \setminus \Gamma$ , and let  $\chi$  be a representation satisfying Assumption 2.1. Let h be a holomorphic function on  $\{s \in \mathbb{C} \mid |\operatorname{Im}(s)| < 2 + \delta\}$  for some  $\delta > 0$ , satisfying  $h(1 + z^2) = O(1 + |z|^2)^{3/2 - \epsilon}$  as  $|z| \to \infty$ , and let

$$g(x) = \frac{1}{2\pi} \int_{\mathbb{R}} h(1+t^2) e^{-itx} dt.$$

Then

$$\sum_{m \in \mathcal{D}} h(\lambda_m) \omega_m = \sum_{\{R\}_{\text{ell}}} \frac{\operatorname{tr}_V \left( \chi(R^{-1})^* \right) g(0) \log N(T_0)}{|\mathcal{E}(R)| \, |(\operatorname{tr}(R))^2 - 4|} + \sum_{\{T\}_{\text{lox}}} \frac{\operatorname{tr}_V \left( \chi(T^{-1})^* \right) g(\log N(T))}{|\mathcal{E}(T)| \, |a(T) - a(T)^{-1}|^2} \log N(T_0)$$

Here,  $\{\lambda_m\}_{m\in\mathcal{D}}$  are the eigenvalues of  $\Delta$  counted with multiplicity, and  $\omega_m$  are the eigenvalues of  $\mathcal{M}$  (the Hecke operator associated to  $\alpha$ ) with the convention of Equation 2.6. The summation with respect to  $\{R\}_{\text{ell}}$  extends over the finitely many  $\Gamma$ -conjugacy classes of elliptic elements  $R \in \Gamma \alpha^{-1} \Gamma$ , and for such a class,  $N(T_0)$  is the minimal norm of a hyperbolic or loxodromic element of the centralizer  $\mathcal{C}(R) \subset \Gamma$ ;  $\mathcal{E}(R)$  is the maximal finite subgroup contained in  $\mathcal{C}(R)$  (see Lemma 3.5 for more details). The summation with respect to  $\{T\}_{\text{lox}}$  extends over the  $\Gamma$ -conjugacy classes of hyperbolic or loxodromic elements of  $\Gamma \alpha^{-1} \Gamma$ ,  $T_0$  denotes a primitive hyperbolic or loxodromic element of minimal norm  $\ln^{12} \mathcal{C}(T) \subset \Gamma$ ; the element T is conjugate in PSL(2,  $\mathbb{C}$ ) to the transformation described by the diagonal matrix with diagonal entries  $a(T), a(T)^{-1}$  with |a(T)| > 1, and<sup>13</sup>  $N(T) = |a(T)|^2$ ;  $\mathcal{E}(T)$  is the finite cyclic elliptic subgroup of  $\mathcal{C}(T)$  (see Lemma 3.3 for more details). The sum over elliptic elements is finite, and all other sums converge absolutely.

Please keep in mind that the conjugacy classes  $\{R\}_{ell}$  and  $\{T\}_{lox}$ , the operator  $\mathcal{M}$ , and the eigenvalues  $\omega_m$  all depend on  $\alpha \in \mathbf{Comm}(\Gamma) \setminus \Gamma$ .

3.8. Self-Adjoint Hecke Operators. Throughout this work we were under Assumption 2.1. By imposing stronger conditions on  $\alpha$  and  $\chi$  we can ensure that  $\mathcal{M}$  is self-adjoint.

**Assumption 3.8.** In addition to Assumption 2.1

(1) 
$$\Gamma \alpha \Gamma = \Gamma \alpha^{-1} \Gamma$$
  
(2)  $\chi(\alpha^{-1}) = \chi(\alpha)^*$ .

Assumption 3.8 implies that  $\mathcal{M}$  is self-adjoint. Since  $\mathcal{M}$  commutes with  $\Delta$  we can choose a simultaneous diagonalization  $\{e_m\}_{m \in \mathcal{D}}$  so that

$$\Delta e_m = \lambda_m e_m$$
 and  $\mathcal{M} e_m = \omega_m e_m$ .

<sup>&</sup>lt;sup>12</sup>Note that there is no clear relationship between  $T \in \Gamma \alpha^{-1} \Gamma$  and  $T_0 \in \Gamma$ .

<sup>&</sup>lt;sup>13</sup>Please note that there is a typographical error in the loxodromic and non cuspidal elliptic terms in [EGM98] Theorem 6.5.1; both terms are missing a factor of  $\frac{1}{4\pi}$ .

## 4. Application of the Trace Formula to the distribution $\omega_m$ .

In the Selberg trace formula (Theorem 3.7), the eigenvalues  $\lambda_m$  of  $\Delta$  do not depend on  $\mathcal{M}$ . The distribution of these eigenvalues has been well studied [Hub59, Sel56, Hej76, EGM98, Ven82]. More specifically, in the case of cocompact Kleinian groups we have

**Theorem 4.1.** [EGM98, p. 308] Let  $\Gamma$  be a cocompact Kleinian group. Then

$$\sum_{m \in \mathcal{D}} e^{-\lambda_m t} \sim \frac{\operatorname{vol}(\Gamma)}{8\pi^{3/2}} t^{-3/2} \quad \text{as } t \to 0^+.$$

*Here*  $vol(\Gamma)$  *is the volume of a fundamental domain for the action of*  $\Gamma$  *in*  $\mathbb{H}^3$ *.* 

For  $\Gamma$  a cocompact Kleinian group with  $\alpha \in \mathbf{Comm}(\Gamma) \setminus \Gamma$ , define  $E_{\alpha}$  the *elliptic number of*  $\Gamma$  *with respect to*  $\alpha$  by

(4.1) 
$$E_{\Gamma}^{\alpha} \equiv \sum_{\{R\}_{\text{ell}}} \frac{\log N(T_0)}{|\mathcal{E}(R)| \, |(\operatorname{tr}(R))^2 - 4|}$$

(see §3.5 for the definitions of the above notation).

Our result is as follows:

**Theorem 4.2.** Let  $\Gamma$  be a cocompact Kleinian group with  $\alpha \in Comm(\Gamma) \setminus \Gamma$ . Then

$$\sum_{m \in \mathcal{D}} \omega_m e^{-\lambda_m t} = \frac{E_{\Gamma}^{\alpha}}{\sqrt{4\pi t}} + O(\sqrt{t}) \quad \text{as } t \to 0^+$$

Here  $E_{\Gamma}^{\alpha}$  is the elliptic number of  $\Gamma$  with respect to  $\alpha$ ,  $\{\lambda_m\}_{m\in\mathcal{D}}$  are the eigenvalues of  $\Delta$  counted with multiplicity, and  $\omega_m$  are the eigenvalues of  $\mathcal{M}$  (the Hecke operator associated to  $\alpha$ ) with the convention of Equation 2.6.

*Proof.* It follows from Lemma 4.3 (below) that the loxodromic sum in (4.2) is bounded by  $O(t^{-1/2} \exp(-c/t))$  as  $t \to 0^+$ . Hence

$$\sum_{m \in \mathcal{D}} \omega_m e^{-t\lambda_m} = \frac{\exp(-t)}{\sqrt{4\pi t}} \sum_{\{R\}_{\text{ell}}} \frac{\log N(T_0)}{|\mathcal{E}(R)| \left| (\operatorname{tr}(R))^2 - 4 \right|} + O\left(t^{-1/2} \exp(-c/t)\right) = \frac{E_{\Gamma}^{\alpha}}{\sqrt{4\pi t}} + O(\sqrt{t}) \quad \text{as } t \to 0^+.$$

**Lemma 4.3.** Let  $\Gamma$  be a cocompact Kleinian group with  $\alpha \in Comm(\Gamma) \setminus \Gamma$ . Then there exists a constant  $c_0 > 1$  so that  $N(T) > c_0$  for all conjugacy class  $\{T\} \in \{T\}_{lox}$ .

*Proof.* Recall that  $\{T\}_{lox}$  is the set of conjugacy classes of loxodromic elements of  $\Gamma \alpha^{-1} \Gamma$ , where conjugacy is taken with respect to the group  $\Gamma$ . An application of the Selberg trace formula, for the pair of functions

$$h(z) = \exp(-zt), \quad g(r) = \frac{\exp(-t)}{\sqrt{4\pi t}} \exp\left(-\frac{r^2}{4t}\right),$$

yields

(4.2) 
$$\sum_{m \in \mathcal{D}} \omega_m e^{-t\lambda_m} = \frac{\exp(-t)}{\sqrt{4\pi t}} \sum_{\{R\}_{\text{ell}}} \frac{\log N(T_0)}{|\mathcal{E}(R)| \, |(\operatorname{tr}(R))^2 - 4|} + \frac{\exp(-t)}{\sqrt{4\pi t}} \sum_{\{T\}_{\text{lox}}} \frac{\exp\left(-\frac{(\log N(T))^2}{4t}\right)}{|\mathcal{E}(T)| \, |a(T) - a(T)^{-1}|^2} \log N(T_0).$$

For t > 0 the left hand side of (4.2) converges absolutely. Fix t = 1. Now, the sum over loxodromic terms is comprised of positive terms. Hence each term of the form

$$\frac{\exp\left(-\frac{(\log N(T))^2}{4}\right)}{|\mathcal{E}(T)| |a(T) - a(T)^{-1}|^2} \log N(T_0)$$

must be bounded by a constant *independent of* T and  $T_0$ . Since  $\Gamma$  is a discrete group, there exists a constant  $d_0 > 1$  so that  $N(T_0) > d_0$  for all loxodromic  $T_0 \in \Gamma$ . Hence  $\log N(T_0)$  is bounded above zero, uniformly (since  $T_0 \in \Gamma$ ). It thus follows that  $|a(T) - a(T)^{-1}|$  must be uniformly bounded above zero. In other words, there is a constant  $c_0 > 1$ , independent of T, with  $|a(T)| > c_0$ . Noting that  $N(T) = |a(T)|^2$ , the Lemma follows.

Theorem 4.2 immediately implies:

**Theorem 4.4.** Let  $\Gamma$  be a cocompact Kleinian group with  $\alpha \in Comm(\Gamma) \setminus \Gamma$ . Suppose that the set  $\Gamma \alpha^{-1}\Gamma$  contains no elliptic elements. Then

$$\sum_{m \in \mathcal{D}} \omega_m e^{-\lambda_m t} = O\left(t^{-1/2} \exp(-c/t)\right) \quad \text{as } t \to 0^+.$$

Hence

$$\lim_{t \to 0^+} \sum_{m \in \mathcal{D}} \omega_m e^{-\lambda_m t} = 0.$$

*Here*  $\{\lambda_m\}_{m \in D}$  *are the eigenvalues of*  $\Delta$  *counted with multiplicity, and*  $\omega_m$  *are the eigenvalues of*  $\mathcal{M}$  *(the Hecke operator associated to*  $\alpha$ *) with the convention of Equation 2.6.* 

## 5. ANALOGUES OF HUBER'S THEOREM FOR HECKE OPERATORS

Following [EGM98, p. 202], we define the length spectrum of  $\Gamma \alpha^{-1} \Gamma$ . For loxodromic  $T_j \in \Gamma \alpha^{-1} \Gamma$  set  $\mu_j = \log N(T_j)$ . The *length spectrum* of  $\Gamma \alpha^{-1} \Gamma$  is defined to be (see §3 for the notation)

\

$$\mathcal{L}_{\Gamma}^{\alpha} \equiv \left( \mu_j, \sum_{\substack{\{T\}_{lax}\\\log N(T) = \mu_j}} \frac{\log N(T_0)}{|\mathcal{E}(T)| |a(T) - a(T)^{-1}|^2} \right)_{j \ge 1}$$

In the two-dimensional case the length spectrum is simply comprises the lengths of closed geodesics. Here we really need the *complex lengths* a(T) and the order of the elliptic, finite subgroup of C(T),  $|\mathcal{E}(T)|$ .

We define the eigenvalue spectra of  $\Delta$  and  $\mathcal{M}$  ( $\mathcal{M}$  is defined from  $\alpha \in \mathbf{Comm}(\Gamma) \setminus \Gamma$ ) by

1

$$\mathcal{S}_{\Gamma}^{\alpha} = \left(\lambda_j, \omega(\lambda_j)\right)_{j \in \mathcal{D}^*};$$

here the symbol  $D^*$  means that we do not count with multiplicity: recall that  $\omega(\lambda)$  is the trace of  $\mathcal{M}$  on the invariant subspace generated by all eigenfunctions (of  $\Delta$ ) with eigenvalue  $\lambda$ , so multiplicity is already encoded into  $\omega(\lambda)$ .

**Theorem 5.1.** Let  $\Gamma, \Gamma'$  be cocompact Kleinian groups with  $\alpha \in Comm(\Gamma) \setminus \Gamma$ ,  $\alpha' \in Comm(\Gamma') \setminus \Gamma'$ . Then the following hold:

(1) Suppose that  $S_{\Gamma}^{\alpha}$  and  $S_{\Gamma'}^{\alpha'}$  agree up to at most finitely many terms. Then

$$E_{\Gamma}^{lpha} = E_{\Gamma'}^{lpha'}$$
  
 $\mathcal{S}_{\Gamma}^{lpha} = \mathcal{S}_{\Gamma'}^{lpha'}$   
 $\mathcal{L}_{\Gamma}^{lpha} = \mathcal{L}_{\Gamma'}^{lpha'}$ 

(2) Suppose that  $\mathcal{L}^{\alpha}_{\Gamma}$  and  $\mathcal{L}^{\alpha'}_{\Gamma'}$  agree up to at most finitely many terms. Then

$$\begin{split} E_{\Gamma}^{\alpha} &= E_{\Gamma'}^{\alpha'} \\ \mathcal{S}_{\Gamma}^{\alpha} &= \mathcal{S}_{\Gamma'}^{\alpha'} \\ \mathcal{L}_{\Gamma}^{\alpha} &= \mathcal{L}_{\Gamma'}^{\alpha'}. \end{split}$$

*Proof.* Upon applying the Selberg trace formula (with trivial representations) to the pair of functions,

$$h(w) = \frac{1}{s^2 + w - 1} - \frac{1}{B^2 + w - 1}$$
$$g(x) = \frac{1}{2s}e^{-s|x|} - \frac{1}{2B}e^{-B|x|},$$

where  $1 < \operatorname{Re}(s) < \operatorname{Re}(B)$ , we obtain

(5.1) 
$$\frac{1}{2s} \sum_{\{T\}_{\text{lox}}} \frac{\log N(T_0)}{m(T)|a(T) - a(T)^{-1}|^2} N(T)^{-s} - \frac{1}{2B} \sum_{\{T\}_{\text{lox}}} \frac{\log N(T_0)}{m(T)|a(T) - a(T)^{-1}|^2} N(T)^{-B} = \sum_{n \in D^*} \omega(\lambda_n) \left(\frac{1}{s^2 - s_n^2} - \frac{1}{B^2 - s_n^2}\right) - \left(\frac{1}{2s} - \frac{1}{2B}\right) \sum_{\{R\}_{\text{ell}}} \frac{\log N(T_0)}{|\mathcal{E}(R)| |(\operatorname{tr}(R))^2 - 4|}$$

Here  $s_n^2 = 1 - \lambda_n$  and  $s_n$  is chosen to lie in  $\{z \in \mathbb{C} \mid \text{Im } z \ge 0, \text{Re } z \ge 0\}$ . The Theorem follows by comparing asymptotics. Simply apply the proof of the [EGM98, Theorem 3.3, p. 203] replacing [EGM98, Equation 5.2.35, p. 198] with our (5.1). Note that our symbol  $w(\lambda_n)$  is the analogue of the multiplicity of  $\lambda_n$  as considered in [EGM98].

By applying the above Theorem to the case of a *single* fixed group with two different elements  $\alpha, \alpha' \in \mathbf{Comm}(\Gamma) \setminus \Gamma$ , we obtain:

**Corollary 5.2.** Let  $\Gamma$  be a cocompact Kleinian groups with  $\alpha, \alpha' \in Comm(\Gamma) \setminus \Gamma$ . Let  $\omega(\lambda_m) = \omega'(\lambda_m)$  for all but at most finitely many  $m \in \mathcal{D}^*$ . Then  $\omega(\lambda_m) = \omega'(\lambda_m)$  for all  $m \in \mathcal{D}^*$ .

#### MODULAR CORRESPONDENCES

### REFERENCES

- [AT90] Shigeki Akiyama and Yoshio Tanigawa, The Selberg trace formula for modular correspondences, Nagoya Math. J. 117 (1990), 93–123. MR MR1044938 (91c:11028)
- [Bus92] Peter Buser, *Geometry and spectra of compact Riemann surfaces*, Progress in Mathematics, vol. 106, Birkhäuser Boston Inc., Boston, MA, 1992.
- [EGM98] J. Elstrodt, F. Grunewald, and J. Mennicke, *Groups acting on hyperbolic space*, Springer Monographs in Mathematics, Springer-Verlag, Berlin, 1998, Harmonic analysis and number theory.
- [Fri05a] Joshua S. Friedman, The Selberg trace formula and Selberg zeta-function for cofinite Kleinian groups with finite-dimensional unitary representations, Ph.D. thesis, Stony Brook University, 2005, http://arxiv.org/abs/math.NT/0612807.
- [Fri05b] \_\_\_\_\_, The Selberg trace formula and Selberg zeta-function for cofinite Kleinian groups with finite-dimensional unitary representations, Math. Z. 250 (2005), no. 4, 939–965.
- [Gol06] Dorian Goldfeld, Automorphic forms and L-functions for the group  $GL(n, \mathbf{R})$ , Cambridge Studies in Advanced Mathematics, vol. 99, Cambridge University Press, Cambridge, 2006, With an appendix by Kevin A. Broughan. MR MR2254662
- [Hej76] Dennis A. Hejhal, The Selberg trace formula for PSL(2, R). Vol. I, Springer-Verlag, Berlin, 1976, Lecture Notes in Mathematics, Vol. 548.
- [Hej83] \_\_\_\_\_, The Selberg trace formula for PSL(2, R). Vol. 2, Lecture Notes in Mathematics, vol. 1001, Springer-Verlag, Berlin, 1983.
- [Hej85] \_\_\_\_\_, A classical approach to a well-known spectral correspondence on quaternion groups, Number theory (New York, 1983–84), Lecture Notes in Math., vol. 1135, Springer, Berlin, 1985, pp. 127–196. MR MR803353 (87h:11045)
- [Hof88] Werner Hoffmann, On a trace formula for Hecke operators, Seminar Analysis of the Karl-Weierstrass-Institute of Mathematics, 1986/87 (Berlin, 1986/87), Teubner-Texte Math., vol. 106, Teubner, Leipzig, 1988, pp. 237–246. MR MR1066744 (91g:22017)
- [Hof94] \_\_\_\_\_, An invariant trace formula for the universal covering group of SL(2, R), Ann. Global Anal. Geom. 12 (1994), no. 1, 19–63. MR MR1261526 (95e:11060)
- [Hub59] H. Huber, Zur analytischen theorie hyperbolischer raumformen und bewegungsgruppen, Math. Ann. 138 (1959), 1–26.
- [Iwa02] Henryk Iwaniec, Spectral methods of automorphic forms, second ed., Graduate Studies in Mathematics, vol. 53, American Mathematical Society, Providence, RI, 2002.
- [LV07] Elon Lindenstrauss and Akshay Venkatesh, Existence and Weyl's law for spherical cusp forms, Geom. Funct. Anal. 17 (2007), no. 1, 220–251. MR MR2306657
- [MR03] Colin Maclachlan and Alan W. Reid, *The arithmetic of hyperbolic 3-manifolds*, Graduate Texts in Mathematics, vol. 219, Springer-Verlag, New York, 2003. MR MR1937957 (2004i:57021)
- [Roe66] Walter Roelcke, Das Eigenwertproblem der automorphen Formen in der hyperbolischen Ebene. I, II, Math. Ann. 167 (1966), 292–337; ibid. 168 (1966), 261–324.
- [Sel56] A. Selberg, Harmonic analysis and discontinuous groups in weakly symmetric Riemannian spaces with applications to Dirichlet series, J. Indian Math. Soc. (N.S.) 20 (1956), 47–87.
- [Sel89] Atle Selberg, Collected papers. Vol. I, Springer-Verlag, Berlin, 1989, With a foreword by K. Chandrasekharan.
- [Str98] Andreas Strömbergsson, *The Selberg trace formula for modular correspondences*, Master's thesis, Uppsala University, Licentiat thesis, 1998, http://www.math.uu.se/~andreas/.
- [Str01] Andreas Strömbergsson, Some remarks on a spectral correspondence for Maass waveforms, Internat. Math. Res. Notices (2001), no. 10, 505–517. MR MR1832538 (2002h:11045)
- [Ven82] A. B. Venkov, Spectral theory of automorphic functions, Proc. Steklov Inst. Math. (1982), no. 4(153), ix+163 pp. (1983), A translation of Trudy Mat. Inst. Steklov. 153 (1981).
- [VKF73] A. B. Venkov, V. L. Kalinin, and L. D. Faddeev, A nonarithmetic derivation of the Selberg trace formula, Zap. Naučn. Sem. Leningrad. Otdel. Mat. Inst. Steklov. (LOMI) 37 (1973), 5–42, Differential geometry, Lie groups and mechanics. MR MR0506043 (58 #21949)

DEPARTMENT OF MATHEMATICS AND SCIENCES, UNITED STATES MERCHANT MARINE ACADEMY, 300 STEAMBOAT ROAD, KINGS POINT, NY 11024

E-mail address: CrownEagle@gmail.com

E-mail address: friedmanj@usmma.edu

E-mail address: joshua@math.sunysb.edu