

Solution of the periodic discrete Toda equation associated with BBSs with M kinds of balls

Shinsuke Iwao

Graduate School of Mathematical Sciences,
The University of Tokyo,
3-8-1 Komaba Meguro-ku, Tokyo 153-8914, Japan

Abstract

A box-ball system with more than one kind of balls is obtained by the generalized periodic discrete Toda equation (pd Toda eq.). We study the pd Toda equation in view of algebraic geometry. The time evolution of pd Toda eq. is linearized on an algebraic variety, and theta function solutions are obtained.

1 Preface

A cellular automaton (CA) is a discrete dynamical system in which the dependent variables take on a finite set of discrete values. Although CAs evolve in time by simple evolution rules, they often show very complicated behavior.

The box-ball system (BBS) is an important CA in which finitely many balls move in an array of boxes under a certain evolution rule. This discrete dynamical system is obtained from a discrete analogue of the Toda equation through the limiting procedure “ultradiscretization”, and displays the behavioral characteristics of nonlinear integrable equations. In fact, the BBS has soliton-like solutions and a large number of conserved quantities. Moreover, the solution of the initial value problem of the periodic box-ball system (pBBS) with one kind of ball has been obtained by ultradiscretizing the solution of the periodic discrete Toda equation (pd Toda).

In 1999, Tokihiro, Nagai and Satsuma pointed out that the pBBS with M ($M \geq 1$) kind of balls (and with capacity one) is obtained by ultradiscretizing the generalized periodic discrete Toda equation [1]. Tokihiro and the author performed the ultradiscretization of theta function solutions of the pd Toda [2]. It is to be expected that the solution of the initial value problem of the pBBS with M kinds of balls can be obtained from the solution of the generalized Toda equation, as was the case for $M = 1$.

In this paper, we study the generalized pd Toda ($M \geq 1$):

$$I_n^{t+M} = I_n^t + V_n^t - V_{n-1}^{t+1}, \quad (1.1)$$

$$V_n^{t+1} = \frac{I_{n+1}^t V_n^t}{I_n^{t+M}}, \quad (1.2)$$

with periodicity condition $I_{n+N}^t = I_n^t, V_{n+N}^t = V_n^t$, where $N, n, t \in \mathbb{N}$.

The time evolution of this system is linearized on an algebraic variety. For example, the time evolution of pd Toda with $M = 1$ is linearized on Jacobi varieties of hyperelliptic curves [3]. For general M , the algebraic curves which appear in the linearization are more complicated than hyperelliptic curves.

These algebraic curves were analyzed by P. van Moerbeke and D. Mumford in 1978 [4]. In their work, they clarify the algebro-geometric properties of the curves which are defined by so-called *regular periodic operator* matrices. Although our matrices are not always regular operators, their results are essentially also applicable to our case.

The special feature of the discrete system (1.1-1.2) is that we can explicitly determine the action of the unit time evolution $t \mapsto t+1$ on the algebraic variety (proposition 2.6). Using this result, we extended the theta function expression of the solution of pd Toda to the $M \geq 1$ case.

The paper is organized as follows. In section 2, we introduce a *spectral curve* associated to the discrete system (1.1-1.2) and discuss its algebro-geometric properties. Our aim in this section is to determine the actions of the index shift $n \mapsto n+1$ and the time shift $t \mapsto t+1$ on a Picard group $\text{Pic}^d(C)$ of this spectral curve C . In section 3, we give the theta function expression (theorem 3.2) which is the extension of the formula obtained by Kimijima and Tokihiro [3].

2 Spectral curve associated with pd Toda

2.1 The nature of the spectral curve

The pd Toda equation (1.1), (1.2) is equivalent to the following matrix form:

$$L_{t+1}(y)R_{t+M}(y) = R_t(y)L_t(y), \quad (2.1)$$

where $L_t(y)$ and $R_t(y)$ are given by

$$L_t(y) = \begin{pmatrix} 1 & & & V_N^t \cdot 1/y \\ V_1^t & 1 & & \\ & \ddots & \ddots & \vdots \\ & & V_{N-1}^t & 1 \end{pmatrix}, \quad R_t(y) = \begin{pmatrix} I_1^t & 1 & & \\ & I_2^t & \ddots & \\ & & \ddots & 1 \\ y & & & I_N^t \end{pmatrix},$$

and y is a complex variable. Let us introduce a new matrix $X_t(y)$ defined by

$$X_t(y) := L_t(y)R_{t+M-1}(y) \cdots R_{t+1}(y)R_t(y). \quad (2.2)$$

From (2.1) and (2.2), we obtain

$$X_{t+1}(y)R_t(y) = R_t(y)X_t(y), \quad (2.3)$$

which implies that the eigenvalues of $X_t(y)$ are conserved quantities under the time evolution. Let $\tilde{\Phi}(x, y) := \det(X_t(y) - xE)$ be the characteristic polynomial of $X_t(y)$ (E is the unit matrix). By definition, $\Phi(x, y) := y \times \tilde{\Phi}(x, y)$ is a polynomial in x and y .

Remark 2.1 *The matrix $X_t(y)$ is a $N \times N$ matrix and is of the form*

$$X(y) = \begin{pmatrix} \alpha_1^{(1)} & \alpha_2^{(2)} & \cdots & \alpha_M^{(M)} & 1 & 0 & & & \\ \beta_1 & \alpha_2^{(1)} & \alpha_3^{(2)} & \cdots & \alpha_{M+1}^{(M)} & 1 & 0 & & \\ 0 & \beta_2 & \alpha_3^{(1)} & \alpha_4^{(2)} & \cdots & \alpha_{M+2}^{(M)} & 1 & 0 & \\ & 0 & \ddots & \ddots & \ddots & & \ddots & \ddots & \\ & & & & & & & & 1 \\ & & & & & & & & \alpha_N^{(M)} \\ & & & & & & & \ddots & \vdots \\ & & & & & & 0 & \beta_{N-1} & \alpha_N^{(1)} \end{pmatrix} + \begin{pmatrix} 0 \\ \vdots \\ 1 \\ \alpha_1^{(M)} & \ddots \\ \vdots & \ddots \\ \alpha_1^{(3)} & \ddots & \ddots \\ \alpha_1^{(2)} & \alpha_2^{(3)} & \cdots & \alpha_{M-1}^{(M)} & 1 & \cdots & 0 \end{pmatrix} \times y + \begin{pmatrix} & \beta_N \end{pmatrix} \times 1/y, \quad (2.4)$$

where $\alpha_j^{(i)}, \beta_j, (1 \leq i \leq M, 1 \leq j \leq N)$ are polynomials in $\{V_n^t, I_n^t, I_n^{t+1}, \dots, I_n^{t+M-1}\}_{n=1}^N$.

Remark 2.2 *When the greatest common divisor $(N, M) \neq 1$, this particular matrix (2.4) is not a regular periodic difference operator which is analyzed in [4].*

Let us define the algebraic curve $\tilde{C}$ by the algebraic equation

$$\tilde{\Phi}(x, y) = 0. \quad (2.5)$$

A *spectral curve* C of the pd Toda equation is the projective curve defined as the completion of $\tilde{C}$. C is a $(M+1)$ -fold ramified covering over $\mathbb{P}^1$:

$$p_x : C \rightarrow \mathbb{P}^1, \quad (2.6)$$

and is also a N -fold ramified covering $p_y : C \rightarrow \mathbb{P}^1$. It goes without saying that C is conserved under the time evolution and is completely determined by the initial values $\{V_n^0, I_n^0, I_n^1, \dots, I_n^{M-1}\}_{n=1}^N$. Note that C may fail to be smooth in certain situations. We restrict ourselves to the case where C is smooth.

We now analyze the points contained in $C \setminus \tilde{C}$. We calculate the polynomial expression of the function $\tilde{\Phi}(x, y)$ in x , y and y^{-1} :

$$\tilde{\Phi}(x, y) = A_0(x)y^{M+1} + A_1(x)y^M + \dots + A_M(x) + A_{M+1}(x)y^{-1} = 0. \quad (2.7)$$

For the analysis of the behavior of C , we need to analyze $A_j(x)$, ($j = 0, 1, \dots, M+1$). The following lemma is established by van Moerbeke and Mumford [4]. They analyzed the determinant $\det(X(y) - xE)$ by direct calculation.

Lemma 2.1 *Let $m := (N, M)$, $N = mN_1$ and $M = mM_1$. The polynomial $A_j(x)$ is a polynomial of degree k_j satisfying*

$$k_j \leq \frac{jN}{M}, \quad (0 \leq j \leq M), \quad k_{M+1} = 0. \quad (2.8)$$

The equality in (2.8) holds if and only if the right hand side is an integer. Moreover, $A_0(x), A_{M_1}(x), A_{2M_1}(x), \dots, A_{mM_1}(x)$ are expressed as:

$$A_{rM_1}(x) = (-1)^{M(N-M)+r} \binom{m}{r} x^{rN_1} + \dots, \quad (r = 0, 1, \dots, m).$$

We start from the polynomial (2.7). Let $\gamma := x^{N_1}y^{-M_1}$. By lemma 2.1, we obtain the expression

$$\begin{aligned} y^{-mM_1} \tilde{\Phi}(x, y) &= (-1)^{M(N-M)} \left(\gamma^m - \binom{m}{1} \gamma^{m-1} + \binom{m}{2} \gamma^{m-2} + \dots + (-1)^m \right) \\ &\quad + \text{lower order terms when } |x|, |y| \rightarrow \infty \\ &\sim (-1)^{M(N-M)} (\gamma - 1)^m. \end{aligned}$$

This implies $x^{N_1}y^{-M_1} = \gamma \sim 1$ near $(x, y) = (\infty, \infty)$. By $(N_1, M_1) = 1$, there exists the local coordinate t equipped with the completion $C \supset \tilde{C}$ such that

$$x \sim t^{-mM_1} = t^{-M}, \quad y \sim t^{-mN_1} = t^{-N}, \quad (x, y) \sim (\infty, \infty).$$

In particular, there exists *only one* point $P \in C$ which is expressed as $P = (\infty, \infty)$.

In a similar manner, there exists a point $Q \in C$ which is expressed as $Q = (\infty, 0)$. The local coordinate t near Q satisfies

$$x \sim t^{-1}, \quad y \sim t^N.$$

Using these fact, the divisors $(x), (y) \in \text{Div}(C)$ are

$$(x) = -MP - Q + (\text{a positive divisor on } \tilde{C}), \quad (2.9)$$

$$(y) = -NP + NQ. \quad (2.10)$$

Remark 2.3 *The existence of the unique point $P(\infty, \infty)$ is a special property of $X(y)$. In fact, there exist m points $P_j(x, y) = (\infty, \infty)$ ($j = 1, \dots, m$) on the algebraic curve associated to a regular operator matrix [4].*

Although the concrete calculations in [4] should be applied only to the case that $X(y)$ is a regular operator, these results are also applicable to our case on condition that C is smooth. Precisely, these calculations become true for our case by substituting

$$P_1 = P_2 = \dots = P_m (= P). \quad (2.11)$$

2.2 The eigenvector mapping

We now define the *isolevel set* $\mathcal{T}_C$ as the set of matrices $X(y)$ associated with the spectral curve C . The following proposition is fundamental to the algebro-geometric method for integrable systems.

Proposition 2.2 *Let C be smooth and $X(y) \in \mathcal{T}_C$. There is a unique line bundle $V \subset C \times \mathbb{C}^N$ s.t.*

$\pi^{-1}(x, y) = \{\text{the eigenspace of } X(y) \text{ corresponding to the eigenvalue } x\} \subset \mathbb{C}^N$,
for all $(x, y) \in C \setminus \{x = 0, \infty\}$ where

$$C \times \mathbb{C}^N \supset V \xrightarrow{\pi} C$$

is called the *canonical projection*.

By virtue of this proposition, we obtain the map

$$\begin{array}{ccc} \varphi_C : & \mathcal{T}_C & \rightarrow \{U \rightarrow \mathbb{P}^{N-1}\}_{U \subset C} \\ & X(y) & \mapsto V^\vee \end{array},$$

where $V^\vee$ is the dual bundle of V . The section of $V^\vee$ is a component of the eigenvector of $X(y)$. By the Grothendieck-Riemann-Roch theorem,

$$\text{Im } \varphi_C \subset \text{Pic}^d(C),$$

where

$$d = g + N - 1 \quad (2.12)$$

and g is the genus of C . (See [5]) Hence,

Definition 2.1 *For smooth C , the eigenvector mapping associated to the equation (2.3) is the mapping*

$$\varphi_C : \mathcal{T}_C \rightarrow \text{Pic}^d(C)$$

defined as above. We shall call $V^\vee$ the eigenvector bundle.

The eigenvector mapping is an important tool to analyze the various integrable systems [6]. The following proposition is essential to our arguments in the present paper.

Proposition 2.3 *The eigenvector mapping*

$$\varphi_C : \mathcal{T}_C \rightarrow \text{Pic}^d(C)$$

is an isomorphism to $\text{Im}\varphi_C$.

This proposition is a straightforward result of the following theorem provided by van Moerbeke and Mumford.

Theorem 2.4 *There is a one-to-one correspondence between the two sets of data:*

- (a) *a multi diagonal matrix $X(y)$ of the form (2.4) such that $\det(X(y) - xE) = 0$ gives a smooth curve.*
- (b) *a smooth curve C , two points P, Q , two functions x, y on C and a divisor $\mathcal{D}$ which satisfies*

$$\varphi_C(X(y)) = \mathcal{D} + (N - 1)Q. \quad (2.13)$$

C has genus:

$$g = \frac{(N - 1)(M + 1) - m + 1}{2},$$

and $\deg \mathcal{D} = g$.

Remark 2.4 *Although the equation (2.13) does not appear in the original paper [4], we easily derive this equation from the relation (p.107)*

$$(g_k) + \mathcal{D} \geq \sum_{i=k+1}^N P_i - \sum_{i=k}^{N-1} Q_i, \quad k = 1, 2, \dots, N - 1, \quad (2.14)$$

where $(g_1, \dots, g_{N-1}, 1)^T$ is a section of the eigenvector bundle $V^\vee$. In fact, (2.14) yields

$$(g_k)_\infty \leq \mathcal{D} + (N - k)Q, \quad (2.15)$$

which implies $d = \deg(\text{Im}\varphi_C) \leq \deg \mathcal{D} + N - 1 = g + N - 1$. Because of the equality (2.12), $(g_1)_\infty$ must satisfy

$$(g_1)_\infty = \mathcal{D} + (N - 1)Q. \quad (2.16)$$

(2.15) and (2.16) imply (2.13).

Definition 2.2 *A finite component $\varphi_{\text{fn}}(X(y))$ of the eigenvector mapping is a positive divisor $\mathcal{D}$ of degree g which appears in (2.13).*

The divisor $\varphi_{\text{fn}}(X(y))$ satisfies

$$\varphi(X(y)) = \varphi_{\text{fn}}(X(y)) + (N - 1)Q$$

by definition.

2.3 The action of the evolutions on the eigenvector bundle

In this section, we represent the two actions — index evolution ($n \mapsto n+1$) and time evolution ($t \mapsto t+1$) — on the eigenvector bundle.

Proposition 2.5 *Let D_n be the divisor*

$$D_n = P - Q.$$

Then the following diagram is commutative.

$$\begin{array}{ccc} \mathcal{T}_C & \rightarrow & \text{Pic}^d(C) \\ n \mapsto n+1 & \downarrow & \downarrow +D_n \\ \mathcal{T}_C & \rightarrow & \text{Pic}^d(C) \end{array}$$

Proof. Let us denote $\sigma : n \mapsto n+1$. A straightforward calculation leads to

$$X(y) \begin{pmatrix} v_1 \\ \vdots \\ v_{N-1} \\ v_N \end{pmatrix} = x \begin{pmatrix} v_1 \\ \vdots \\ v_{N-1} \\ v_N \end{pmatrix} \Leftrightarrow \sigma^{-1}(X(y)) \begin{pmatrix} y^{-1}v_N \\ v_1 \\ \vdots \\ v_{N-1} \end{pmatrix} = x \begin{pmatrix} y^{-1}v_N \\ v_1 \\ \vdots \\ v_{N-1} \end{pmatrix}. \quad (2.17)$$

It is enough to prove

$$(y^{-1}g_{N-1}^{-1})_\infty - (g_1)_\infty \sim -D_n. \quad (2.18)$$

By (2.14), we have

$$(y g_{N-1}) + \mathcal{D} \geq (N-1)Q - (N-1)P. \quad (2.19)$$

An argument similar to remark 2.4 allows to conclude

$$(y^{-1}g_{N-1}^{-1})_\infty \sim (y g_{N-1})_\infty = \mathcal{D} + (N-1)P,$$

which completes the proof. ■

In order to determine the action of the time evolution on the eigenvector bundle, we need to prove the following proposition.

Proposition 2.6 *Let D_t be the divisor*

$$D_t = A - Q,$$

where $A = (0, y_0)$ and y_0 is the complex number which satisfies $\det R_t(y_0) = 0$. Then, the following diagram is commutative.

$$\begin{array}{ccc} \mathcal{T}_C & \rightarrow & \text{Pic}^d(C) \\ t \mapsto t+1 & \downarrow & \downarrow +D_t \\ \mathcal{T}_C & \rightarrow & \text{Pic}^d(C) \end{array}$$

To prove this proposition, we introduce the concepts of *Bloch solution* and *transposed operator*.

We identify the eigenvectors of $X(y)$ with the *Bloch solutions* of the periodic infinite matrix

$$\tilde{X} = \begin{pmatrix} \ddots & \ddots & & \ddots & \ddots & & & & & \\ \ddots & \alpha_N^{(1)} & \alpha_1^{(2)} & \cdots & \alpha_{M-1}^{(M)} & 1 & 0 & & & \\ & \beta_N & \alpha_1^{(1)} & \alpha_2^{(2)} & \cdots & \alpha_M^{(M)} & 1 & 0 & & \\ & 0 & \beta_1 & \alpha_2^{(1)} & \alpha_3^{(2)} & \cdots & \alpha_{M+1}^{(M)} & 1 & 0 & \\ & & 0 & \ddots & \ddots & & & \ddots & \ddots & \end{pmatrix},$$

which are the infinite vectors $\tilde{\mathbf{v}} = (\cdots, v_{n-1}, v_n, v_{n+1}, \cdots)^T$ such that

$$\tilde{X}\tilde{\mathbf{v}} = x\tilde{\mathbf{v}} \quad (2.20)$$

and

$$v_{n+N} = y v_n. \quad (2.21)$$

Equation (2.20) is equivalent to

$$\beta_{n-1}v_{n-1} + \alpha_n^{(1)}v_n + \alpha_{n+1}^{(2)}v_{n+1} + \cdots + \alpha_{n+M-1}^{(M)}v_{n+M-1} + v_{n+M} = xv_n \quad (2.22)$$

for $n \in \mathbb{Z}$ ($\alpha_{n+N}^{(j)} = \alpha_n^{(j)}$, $\beta_{n+N} = \beta_n$). Because the l.h.s. of equation (2.22) is a linear combination of

$$v_{n-1}, v_n, \cdots, v_{n+M-1},$$

the Bloch solution associated with $(x, y) \in C$ are also be written as a linear combination of

$$\tilde{\mathbf{v}}^{(1)}, \tilde{\mathbf{v}}^{(2)}, \dots, \tilde{\mathbf{v}}^{(M+1)},$$

where

$$\begin{aligned} \tilde{\mathbf{v}}^{(1)} &= (\dots, 1, 0, \dots, 0, v_{M+2}^{(1)}, v_{M+3}^{(1)}, \dots)^T \\ \tilde{\mathbf{v}}^{(2)} &= (\dots, 0, 1, \dots, 0, v_{M+2}^{(2)}, v_{M+3}^{(2)}, \dots)^T \\ &\vdots \\ \tilde{\mathbf{v}}^{(M+1)} &= (\dots, 0, 0, \dots, 1, v_{M+2}^{(M+1)}, v_{M+3}^{(M+1)}, \dots)^T. \end{aligned}$$

More precisely, let $\psi(x, y)$ be the Bloch solution:

$$\psi(x, y) = a_1\tilde{\mathbf{v}}^{(1)} + a_2\tilde{\mathbf{v}}^{(2)} + \cdots + a_{M+1}\tilde{\mathbf{v}}^{(M+1)}, \quad (2.23)$$

where

$$a_i = a_i(x, y) \quad \text{and} \quad \tilde{\mathbf{v}}^{(i)} = \tilde{\mathbf{v}}^{(i)}(x).$$

Recalling the equation (2.3) and proposition 2.2 the eigenvector $\mathbf{v}_t(x, y)$ at time t satisfies

$$\mathbf{v}_{t+1}(x, y) = R_t(y) \cdot \mathbf{v}_t(x, y). \quad (2.24)$$

Equivalently, the Bloch solution $\psi_t(x, y)$ satisfies

$$\psi^{t+1}(x, y) = \tilde{R}_t \cdot \psi^t(x, y), \quad (2.25)$$

where

$$\tilde{R}_t = \begin{pmatrix} \ddots & \ddots & & & & \\ & I_N^t & 1 & & & \\ & & I_1^t & 1 & & \\ & & & I_2^t & \ddots & \\ & & & & \ddots & 1 \\ & & & & & I_N^t & \ddots \\ & & & & & & \ddots \end{pmatrix}.$$

(2.23) and (2.25) yield

$$a_j^{t+1} = a_1^t(\tilde{R}_t \tilde{\mathbf{v}}^{(1)})_j + \cdots + a_{M+1}^t(\tilde{R}_t \tilde{\mathbf{v}}^{(M+1)})_j, \quad j = 1, 2, \dots, M+1, \quad (2.26)$$

where $(\mathbf{v})_j$ is a j th-component of the vector $\mathbf{v}$. Equation (2.26) is equivalent to

$$\begin{pmatrix} a_1^{t+1} \\ a_2^{t+1} \\ \vdots \\ a_M^{t+1} \\ a_{M+1}^{t+1} \end{pmatrix} = H_t \cdot \begin{pmatrix} a_1^t \\ a_2^t \\ \vdots \\ a_M^t \\ a_{M+1}^t \end{pmatrix}, \quad (2.27)$$

where

$$H_t = \begin{pmatrix} I_1^t & 1 & & & \\ & I_2^t & \ddots & & \\ & & \ddots & 1 & \\ & & & I_M^t & 1 \\ v_{M+2}^{(1)} & v_{M+2}^{(2)} & \cdots & v_{M+2}^{(M)} & I_{M+1}^t + v_{M+2}^{(M+1)} \end{pmatrix}. \quad (2.28)$$

Using this equation, we easily obtain

$$\begin{pmatrix} v_1^{t+1} \\ v_2^{t+1} \\ \vdots \\ v_M^{t+1} \\ v_{M+1}^{t+1} \end{pmatrix} = H_t \cdot \begin{pmatrix} v_1^t \\ v_2^t \\ \vdots \\ v_M^t \\ v_{M+1}^t \end{pmatrix}, \quad (2.29)$$

where

$$\psi^t(x, y) = (\dots, v_1^t, v_2^t, \dots, v_{M+1}^t, \dots)^T.$$

Lemma 2.7 *For fixed generic x , the $M + 1$ Bloch solutions associated with x are linearly independent.*

Proof. For generic x , associated y_j ($j = 1, 2, \dots, M + 1$) are all distinct. ■

Lemma 2.8

$$\det H_t = (-1)^{M+1} I_1^t x.$$

This lemma is proved by an elementary calculation, which we shall give in Appendix.

Let $\mathcal{F}$ be the invertible sheaf associated with the eigenvector bundle $V^\vee$. Let us consider the direct image

$$(p_x)_* \mathcal{F},$$

where p_x is a projection of C defined by (2.6). The sheaf $(p_x)_* \mathcal{F}$ is a locally free sheaf of rank $M + 1$. By lemma 2.7, the fiber of this direct sheaf is $\mathbb{C}^{M+1}$.

The equation (2.29) is regarded as the time action to the space $\mathbb{C}^{M+1}$. Recall that the components of the Bloch solutions of $\tilde{X}$ are the section of the sheaf $\mathcal{F}$. For $x \in \mathbb{P}^1$, we denote the k -th component of the infinite vector $\psi^t(x, y_j)$ by $v_{jk}^t(x)$, and $\psi^t(x, y_j)$ by $\psi_j^t(x)$. By (2.23), the finite vectors

$$\hat{\psi}_j^t := (v_{j,1}^t, v_{j,2}^t, \dots, v_{j,M+1}^t)^T$$

have the property

$$\{\psi_j^t\}_{j=1}^{M+1} \text{ are linearly independent} \iff \{\hat{\psi}_j^t\}_{j=1}^{M+1} \text{ are linearly independent.} \quad (2.30)$$

Equation (2.29) implies that

$$(\hat{\psi}_1^{t+1}, \hat{\psi}_2^{t+1}, \dots, \hat{\psi}_{M+1}^{t+1}) = H_t \cdot (\hat{\psi}_1^t, \hat{\psi}_2^t, \dots, \hat{\psi}_{M+1}^t). \quad (2.31)$$

On the other hand, lemma 2.7, 2.8 and (2.30) imply that there exists at least one vector $\hat{\psi}_j^t$ which satisfies

$$\text{mult}_x \hat{\psi}_j^{t+1} > \text{mult}_x \hat{\psi}_j^t, \quad (2.32)$$

where

$$\text{mult}_x(z_1, z_2, \dots, z_N)^T = \min [\text{mult}_x z_1, \text{mult}_x z_2, \dots, \text{mult}_x z_N].$$

The same discussion can be repeated for the projection

$$p_y : C \rightarrow \mathbb{P}^1,$$

which is an N -fold ramified covering over $\mathbb{P}^1$.

The following two facts are then obvious to prove.

Lemma 2.9 *For fixed generic y , the N eigenvectors of $X_t(y)$ are linearly independent.*

Lemma 2.10 *$\det R_t(y)$ is a polynomial of degree one in y .*

Let us consider the direct image

$$(p_y)_* \mathcal{F}.$$

From lemma 2.9 we then find that the fiber of this direct image is $\mathbb{C}^N$.

The equation (2.24) can be regarded as the time action to the space $\mathbb{C}^N$. For fixed y , we denote $p_y^{-1}(x) = \{(x_1, y), \dots, (x_N, y)\}$. Then we obtain that

$$\exists j \quad \text{s.t.} \quad \text{mult}_{y_0} \mathbf{v}_{t+1}(x_j, y) > \text{mult}_{y_0} \mathbf{v}_t(x_j, y), \quad (2.33)$$

where $\det R_t(y_0) = 0$.

2.4 The transposed operator

In this subsection, we introduce the *transposed operator*, and give the proof of proposition 2.6.

The section of the eigenvector bundle $V^\vee$ can be described as a rational function of x and y . We denote the set of these sections by $\Gamma(V^\vee)$. Let $\Delta_{i,j} := (-1)^{i+j} \times (i, j)$ -th minor of $X(y) - xE$. We have

$$g_k = \frac{\Delta_{N,k}}{\Delta_{N,N}}, \quad k = 1, 2, \dots, N-1,$$

where $(g_1, \dots, g_{N-1}, 1)^T \in \Gamma(V^\vee)$. We are interested in the divisor $(g_1)_\infty$ and hence, it is important to explore the common zeros of $\Delta_{N,1}$ and $\Delta_{N,N}$.

Equation (2.3) is equivalent to

$$X_t(y)^T R_t(y)^T = R_t(y)^T X_{t+1}(y)^T. \quad (2.34)$$

Let us define $s := -t$, and $A^* := JA^T J$ for a regular matrix A where

$$J = \begin{pmatrix} 0 & & & 1 \\ & & \ddots & \\ & 1 & & \\ 1 & & & 0 \end{pmatrix}.$$

Then (2.34) becomes

$$X_{s+1}^* R_{s+1}^* = R_{s+1}^* X_s^*. \quad (2.35)$$

Note that the matrix $X^*(y)$ is also of the form (2.4). We call this new matrix X^* the *transposed operator* of X . A careful analysis of the eigenvector bundle of the transposed operator gives more information on the original matrix X .

Lemma 2.11 (van Moerbeke, Mumford) (1) *There exist positive regular divisors $\mathcal{D}_1$, $\mathcal{D}_2$, and $\mathcal{D}_3$ such that*

$$\left(\frac{\Delta_{N,1}}{\Delta_{N,N}} \right) = \mathcal{D}_1 + (N-1)P - \varphi_{\text{fn}}(X) - (N-1)Q, \quad (2.36)$$

and

$$\left(\frac{\Delta_{1,N}}{\Delta_{N,N}} \right) = \mathcal{D}_2 + (N-1)Q - \mathcal{D}_3 - (N-1)P. \quad (2.37)$$

Moreover, $\deg \mathcal{D}_i = g$ ($i = 1, 2, 3$).

(2) *The divisor $(\Delta_{N,N})$ satisfies*

$$(\Delta_{N,N}) = \varphi_{\text{fn}}(X) + \mathcal{D}_3 - (NM - M - n + 1)P - NQ. \quad (2.38)$$

Remark 2.5 *Lemma 2.11 (1) and Remark 2.16 imply that $\deg \frac{\Delta_{1,N}}{\Delta_{N,N}} = g + N - 1$, and that $\mathcal{D}_2$ and $\mathcal{D}_3$ do not have common points.*

Lemma 2.12

$$\varphi_{\text{fn}}(X^*) = \varphi_{\text{fn}}(X^T) = \mathcal{D}_2.$$

Proof. Note that X^* and X^T give the same spectral curve C . By definition, it follows that:

$$X^T(J\mathbf{v}) = x(J\mathbf{v}) \Leftrightarrow X^*\mathbf{v} = x\mathbf{v},$$

which implies the first equality of the lemma.

The second equality is easily obtained from (2.13), (2.37) and the trivial fact that the $(N, 1)$ -th minor of X is the $(1, N)$ -th minor of X^T . ■

Lemma 2.12 and equation (2.37) yield

$$\begin{aligned} \mathcal{D}_3 &\sim \varphi_{\text{fn}}(X^*) + (N-1)Q - (N-1)P \\ &\sim \varphi_{\text{fn}}(X^*) - Q + P. \end{aligned}$$

Using proposition 2.5, we obtain $\mathcal{D}_3 \sim \varphi_{\text{fn}}(\sigma(X^*)) = \varphi_{\text{fn}}((\sigma^{-1}X)^*)$. In fact, by virtue of the Riemann-Roch theorem, we obtain the following stronger result:

$$\mathcal{D}_3 = \varphi_{\text{fn}}((\sigma^{-1}X)^*)$$

because $\mathcal{D}_3$ is a regular divisor of degree g .

Proposition 2.13 $\Delta_{N,N}$ *has $2g$ zeros in $C \setminus \{x = \infty\}$. And the corresponding divisor is*

$$\varphi_{\text{fn}}(X) + \varphi_{\text{fn}}((\sigma^{-1}X)^*).$$

As a corollary of this proposition, the zeros of $\Delta_{1,1}$, $\Delta_{1,N}$, $\Delta_{N,1}$ are also determined:

Corollary 2.14 $\Delta_{1,1}, \Delta_{1,N}, \Delta_{N,1}$ have $2g$ zeros in $C \setminus \{x = \infty\}$ respectively. The corresponding divisors are:

$$\begin{aligned} \Delta_{1,1} &: \varphi_{\text{fn}}(\sigma^{-1}X) + \varphi_{\text{fn}}(X^*) \\ \Delta_{1,N} &: \varphi_{\text{fn}}(X) + \varphi_{\text{fn}}(X^*) \\ \Delta_{N,1} &: \varphi_{\text{fn}}(\sigma^{-1}X) + \varphi_{\text{fn}}((\sigma^{-1}X)^*). \end{aligned} \quad (2.39)$$

Proof. The divisor of $\Delta_{1,1}$ is obtained from $\Delta_{N,N}$ by the change of the index: $n \rightarrow n-1$. The remaining divisors are determined by the identity $\Delta_{1,1}\Delta_{N,N} = \Delta_{1,N}\Delta_{N,1}$ and (2.16). ■

Let us define the set $\mathcal{A} := \{\alpha_1^{(1)}, \dots, \alpha_N^{(1)}; \dots; \alpha_1^{(M)}, \dots, \alpha_N^{(M)}; \beta_1, \dots, \beta_N\}$ and the map $\tau : \mathcal{A} \rightarrow \mathcal{A}$ which is associated with the correspondence $X \rightarrow (\sigma^{-1}X)^*$, as in the example below. Note that $\tau^2 = \text{id}$.

Example ($N = 4, M = 2$)

$$\begin{aligned} X &= \begin{pmatrix} \alpha_1^{(1)} & \alpha_2^{(2)} & 1 & \beta_4 1/y \\ \beta_1 & \alpha_2^{(1)} & \alpha_3^{(2)} & 1 \\ y & \beta_2 & \alpha_3^{(1)} & \alpha_4^{(2)} \\ \alpha_1^{(2)} y & y & \beta_3 & \alpha_4^{(1)} \end{pmatrix}, X^* = \begin{pmatrix} \alpha_4^{(1)} & \alpha_4^{(2)} & 1 & \beta_4 1/y \\ \beta_3 & \alpha_3^{(1)} & \alpha_3^{(2)} & 1 \\ y & \beta_2 & \alpha_2^{(1)} & \alpha_2^{(2)} \\ \alpha_1^{(2)} y & y & \beta_1 & \alpha_1^{(1)} \end{pmatrix}, \\ \sigma^{-1}X &= \begin{pmatrix} \alpha_4^{(1)} & \alpha_1^{(2)} & 1 & \beta_3 1/y \\ \beta_4 & \alpha_1^{(1)} & \alpha_2^{(2)} & 1 \\ y & \beta_1 & \alpha_2^{(1)} & \alpha_3^{(2)} \\ \alpha_4^{(2)} y & y & \beta_2 & \alpha_3^{(1)} \end{pmatrix}, (\sigma^{-1}X)^* = \begin{pmatrix} \alpha_3^{(1)} & \alpha_3^{(2)} & 1 & \beta_3 1/y \\ \beta_2 & \alpha_2^{(1)} & \alpha_2^{(2)} & 1 \\ y & \beta_1 & \alpha_1^{(1)} & \alpha_1^{(2)} \\ \alpha_4^{(2)} y & y & \beta_4 & \alpha_4^{(1)} \end{pmatrix}. \end{aligned}$$

The correspondence $X \leftrightarrow (\sigma^{-1}X)^*$ is equivalent to the correspondence:

$$\alpha_i^{(1)} \leftrightarrow \alpha_{4-i}^{(1)}, \quad \alpha_i^{(2)} \leftrightarrow \alpha_{1-i}^{(2)}, \quad \beta_i \leftrightarrow \beta_{3-i}.$$

Our aim is to analyze the distribution of common zeros of $\Delta_{N,1}$ and $\Delta_{N,N}$. This set of zeros has the following characteristic property.

Lemma 2.15

$$\begin{aligned} &\{ \text{The common zeros of } \Delta_{N,1} \text{ and } \Delta_{N,N} \} \\ &= \{ \text{The common zeros of } \Delta_{N,k} \text{ (} k = 1, 2, \dots, N \text{)} \}. \end{aligned}$$

Proof. Consider the following open subset on C : $\tilde{C} = C \setminus \{x = \infty\}$. By (2.15) and (2.16), we obtain, if $k \geq 2$,

$$(g_k)_\infty|_{\tilde{C}} < (g_1)_\infty|_{\tilde{C}}. \quad (2.40)$$

This is equivalent to

$$\left(\frac{\Delta_{N,k}}{\Delta_{N,N}} \right)_\infty < \left(\frac{\Delta_{N,1}}{\Delta_{N,N}} \right)_\infty, \quad (2.41)$$

which proves the lemma. ■

We can now proceed with the proof of proposition 2.6. Proposition 2.13 and corollary 2.14 imply

$$\{\text{Common zeros of } \Delta_{N,1} \text{ and } \Delta_{N,N}\} = \varphi_{\text{fn}}((\sigma^{-1}X)^*).$$

Using this fact, it becomes clear that the equations (2.32) and (2.33) give the time evolution action of the transposed operator X^* . In fact, by virtue of lemma 2.15, equation (2.32) requires the existence of a point $(x, y) = (0, y')$ which becomes the common zero of $\Delta_{N,1}$ and $\Delta_{N,N}$ when $t \mapsto t + 1$. Then equation (2.33) also requires the existence of a point $(x, y) = (x', y_0)$ which becomes the common zero of $\Delta_{N,1}$ and $\Delta_{N,N}$. Using the same argument about poles, it follows that the divisor

$$\mathcal{D}'_t = Q - A$$

obeys the following relation:

$$\psi_{\text{fn}}(X_{t+1}^*) = \psi_{\text{fn}}(X_t^*) + \mathcal{D}'_t. \quad (2.42)$$

Recalling the equation (2.35), the two time evolutions defined by X_t and X_s^* are opposite: $s = -t$. Hence the divisor $\mathcal{D}_t = -\mathcal{D}'_t = A - Q$ gives the time evolution of original system. ■

3 Theta function solutions

3.1 The distribution of the points $\varphi_{\text{fn}}(X)$

Let $\varphi_{\text{fn}}(X) = P_1 + \dots + P_g$, $P_j = (x_j, y_j) \in C$. We are interested in the numbers $x_j \in \mathbb{C}$, ($j = 1, 2, \dots, g$). Now, consider the resultant of the polynomials $\Phi(x, y) = y \det(X(y) - xE)$ and $\Delta_{N,N}$ as polynomials in y . Let us denote this resultant by $\text{Rest}_y(\Phi, \Delta_{N,N}) =: R(x)$, which is a polynomial in x . More precisely, $R(x)$ is an element of $\mathbb{C}[\alpha_1^{(1)}, \dots, \alpha_N^{(1)}; \dots; \alpha_1^{(M)}, \dots, \alpha_N^{(M)}; \beta_1, \dots, \beta_N][x]$. From proposition 2.13, it follows that $\deg_x R(x) = 2g$.

We also consider the resultant $S(x) := \text{Rest}_y(\Phi, \Delta_{1,N})$. Due to proposition 2.13 and corollary 2.14 the common divisor of $R(x)$ and $S(x)$ is an element of $\mathbb{C}(\alpha_1^{(1)}, \dots, \alpha_N^{(1)}; \dots; \alpha_1^{(M)}, \dots, \alpha_N^{(M)}; \beta_1, \dots, \beta_N)[x]$, the degree of which (as a polynomial in x) is equal to g . Multiplying divisors (if needed), we obtain the monic polynomial

$$\Upsilon(x) \in \mathbb{C}(\alpha_1^{(1)}, \dots, \alpha_N^{(1)}; \dots; \alpha_1^{(M)}, \dots, \alpha_N^{(M)}; \beta_1, \dots, \beta_N)[x],$$

the roots of which are the common roots of $R(x)$ and $S(x)$, i.e., the common roots of $\Delta_{1,N}$ and $\Delta_{N,N}$. Recalling that the set of the common zeros of $\Delta_{N,N}$ and $\Delta_{1,N}$ contained in $C \setminus \{x = \infty\}$ is $\{P_1, \dots, P_g\}$, we conclude:

$$\deg_x \Upsilon(x) = g, \quad \Upsilon(x_i) = 0, \quad i = 1, 2, \dots, g. \quad (3.1)$$

3.2 Theta function solutions

For a complex curve (or Riemann surface) C of genus g , one usually considers a *canonical basis* of $H_1(C, \mathbb{Z})$. We denote the canonical basis by $a_1, \dots, a_g; b_1, \dots, b_g \in H_1(C, \mathbb{Z})$. Let $\omega_1, \dots, \omega_g$ be the holomorphic differential of C which satisfies $\int_{a_j} \omega_i = \delta_{j,i}$. A *period matrix* of the Riemann surface C is a $g \times g$ matrix $B = (\int_{b_j} \omega_i)$. Let $\theta(z, B)$ be the *theta function*; $\mathbb{C}^g \rightarrow \mathbb{C}$, and

$$\mathbf{A} : \text{Pic}^g \xrightarrow{\sim} J(C) (:= \mathbb{C}^g / (\mathbb{Z}^g + B\mathbb{Z}^g))$$

the *Abelian mapping*. The following theorem is a classical and fundamental result.

Theorem 3.1 (Riemann) *Let C be a Riemann surface of genus g , and let $\mathcal{D} = P_1 + \dots + P_g$ be a regular positive divisor. Then the function*

$$F(p) = \theta(\mathbf{A}(p) - \mathbf{A}(\mathcal{D}) - \mathbf{K}, B), \quad p \in C$$

has exactly g zeros $p = P_1, \dots, P_g$ on C , where $\mathbf{K}$ is the Riemann constant of C .

To obtain the solution to the pd Toda equation, we consider the following integral:

$$I = \frac{1}{2\pi i} \int_{\partial C^\circ} x(p) \frac{dF(p)}{F(p)} \left(=: \frac{1}{2\pi i} \int_{\partial C^\circ} d\mathcal{Z}(p) \right). \quad (3.2)$$

Here the integral path ∂C° goes along the edge of the simply connected domain C° obtained from the Riemann surface by cutting it along $a_1, \dots, a_g; b_1, \dots, b_g$.

The integral I can be rewritten as

$$\begin{aligned} I &= \frac{1}{2\pi i} \sum_{k=1}^g \left(\int_{a_k} + \int_{a_k^{-1}} + \int_{b_k} + \int_{b_k^{-1}} \right) d\mathcal{Z}(p) \\ &= \frac{1}{2\pi i} \sum_{k=1}^g \left(\int_{a_k} \{d\mathcal{Z}(p) - d\mathcal{Z}(p + b_k)\} + \int_{b_k} \{d\mathcal{Z}(p) - d\mathcal{Z}(p + a_k)\} \right). \end{aligned}$$

Recalling the classical fact

$$F(p + a_k) = F(p), \quad (3.3)$$

$$F(p + b_k) = \exp(-2\pi i(\mathbf{A}(p) - \mathbf{A}(\mathcal{D}) - \mathbf{K})_k) F(p), \quad (3.4)$$

and

$$d(\mathbf{A}(p))_k = \omega_k(p), \quad (3.5)$$

the integral I is transformed to

$$I = \sum_{k=1}^g \int_{a_k} x(p) \omega_k(p). \quad (3.6)$$

On the other hand, by the residue theorem, the integral I also has the expression:

$$I = \sum_{i=1}^g x(P_i) + \text{Res}_P(d\mathcal{Z}) + \text{Res}_Q(d\mathcal{Z}). \quad (3.7)$$

Let t_P and t_Q be local coordinates around P and Q respectively. These satisfy

$$x \sim 1/(t_P)^M, \text{ (neighbor of } P), \quad x \sim 1/t_Q, \text{ (neighbor of } Q).$$

In a neighborhood of P , one has:

$$d\mathcal{Z} \sim \frac{1}{(t_P)^M} \frac{d \log F}{dt_P} dt_P \quad (3.8)$$

$$= \frac{1}{(t_P)^M} \sum_{l=1}^g (\partial_l \log F) \left(\frac{d(\mathbf{A})_l}{dt_P} \right) dt_P \quad (3.9)$$

To calculate the residue of the differential (3.9), we explore the behavior of

$$\frac{d(\mathbf{A})_l}{dt_P} dt_P = \omega_l$$

around points P and Q . Let $c_l := \text{Res}_P(\omega_l/(t_P)^M)$. Then we obtain the expression:

$$\text{Res}_P(d\mathcal{Z}) = \sum_{l=1}^g c_l (\partial_l \log(F(P))). \quad (3.10)$$

In the similar manner, we also conclude

$$\text{Res}_Q(d\mathcal{Z}) = \sum_{l=1}^g c'_l (\partial_l \log(F(Q))), \quad (3.11)$$

where $c'_l := \text{Res}_Q(\omega_l/t_Q)$.

By (3.6) and (3.7), we obtain

$$\sum_{l=1}^g x(P_l) = \sum_{l=1}^g \int_{a_l} x(p) \omega_l(p) - \sum_{l=1}^g c_l (\partial_l \log(F(P))) - \sum_{l=1}^g c'_l (\partial_l \log(F(Q))). \quad (3.12)$$

Using this equation, we obtain the following theorem which is a generalization of the well-known result concerning the theta function solution to the pd Toda ($M = 1$) equation and which is the main theorem in the present paper:

Theorem 3.2 *Let $\Upsilon(x)$ be the monic polynomial of degree g obtained by (3.1):*

$$\Upsilon(x) = x^g - a_1 x^{g-1} + \cdots + (-1)^g a_g,$$

with $a_1, \dots, a_g \in \mathbb{C}(\alpha_1^{(1)}, \dots, \alpha_N^{(1)}; \dots; \alpha_1^{(M)}, \dots, \alpha_N^{(M)}; \beta_1, \dots, \beta_N)$. Then we find

$$a_1 = \sum_{l=1}^g \int_{a_l} x(p) \omega_l(p) - \sum_{l=1}^g c_l \{ \partial_l \log \theta(n\mathbf{k} + t\boldsymbol{\nu} + \mathbf{c}_0, B) \} \\ - \sum_{l=1}^g c'_l \{ \partial_l \log \theta((n+1)\mathbf{k} + t\boldsymbol{\nu} + \mathbf{c}_0, B) \}, \quad (3.13)$$

where $\mathbf{k} = \mathbf{A}(P - Q)$, $\boldsymbol{\nu} = \mathbf{A}(A - Q)$, $\mathbf{c}_0 = \mathbf{A}(Q - \mathcal{D}_0 + \Theta)$, and

$$c_l = \text{Res}_P(\omega_l / (t_P)^M), \quad c'_l = \text{Res}_Q(\omega_l / t_Q).$$

Here Θ is the theta divisor: $\mathbf{A}(\Theta) = -\mathbf{K}$. The divisor $\mathcal{D}_0$ is the initial value

$$\varphi_{\text{fn}}(X_{t=0}(y)) = \mathcal{D}_0.$$

Acknowledgments

The author is very grateful to Professor Teshuji Tokihiro and Professor Ralph Willox for helpful comments on this paper.

A Proof of lemma 2.8

In this section, we give the proof of lemma 2.8. Let

$$U_k := R_{t+k-1}(y) R_{t+k-2}(y) \dots R_t(y), \quad 1 \leq k \leq M.$$

Define the row vector $\mathbf{u}^{(k)} = (u_{1,1}^{(k)}, u_{1,2}^{(k)}, \dots, u_{1,N}^{(k)})$, where $U_k = (u_{i,j}^{(k)})$, and the following homomorphism of rings:

$$\sigma : \mathbb{Z}[\{V_n^t, I_n^t\}_{n \in \mathbb{Z}}] \rightarrow \mathbb{Z}[\{V_n^t, I_n^t\}_{n \in \mathbb{Z}}] \quad ; \quad V_n^t \mapsto V_{n+1}^t, \quad I_n^t \mapsto I_{n+1}^t.$$

We rewrite the j -th component of $\mathbf{u}^{(k)}$ as $u_j^{(k)}$ for short. By definition of $R_t(y)$, we obtain

$$u_j^{(k+1)} = \sigma(u_{j-1}^{(k)}) + I_1^{t+k} u_j^{(k)}, \quad (u_{-1}^{(k)} = 0). \quad (\text{A.1})$$

On the other hand, the 2nd row of the matrix $X_t(y)$, which is of the form

$$(\beta_1, \alpha_2^{(1)}, \alpha_3^{(2)}, \dots, \alpha_{M+1}^M, 1, 0, \dots, 0),$$

satisfies

$$\beta_1 = V_1^t u_1^{(M)}, \quad \alpha_{j+1}^{(j)} = \sigma(u_j^{(M)}) + V_1^t u_{j+1}^{(M)}. \quad (\text{A.2})$$

Figure 1 displays the algorithm which we shall use to obtain the row vector $\mathbf{u}^{(k)}$ expressed by (A.1).

Let us introduce the signs $\swarrow$ and $\searrow$ to describe the terms which appear in the figure 1. Let us define the set of arrows $Ar := \{\swarrow, \searrow\}$ and the set of




Figure 1: The $j + 1$ -th row of this diagram displays the first $j + 1$ non-zero components of the row vector $\mathbf{u}^{(j)}$.

sequences $Ar^r := \{(a_1, a_2, \dots, a_r) \mid a_j \in Ar\}$, $r \in \mathbb{N}$. We define the map of sets $\{\cdot\} : \bigcup_r Ar^r \rightarrow \mathbb{Z}[\{I_n^{t+l}\}_{1 \leq n \leq N, 1 \leq l \leq M}]$ as follows:
For $r = 0$, define $\{\emptyset\} := 1$. If $\{a_1, a_2, \dots, a_r\} \in \mathbb{Z}[\{I_n^{t+l}\}_{n,l}]$ is given, we define $\{a_1, a_2, \dots, a_r, a_{r+1}\}$ by

$$\{a_1, \dots, a_r, \swarrow\} := I_1^{t+k} \{a_1, a_2, \dots, a_r\}, \quad \{a_1, \dots, a_r, \searrow\} := \sigma(\{a_1, \dots, a_r\}),$$

inductively. For example,

$$\{\swarrow\} = I_1^t, \quad \{\swarrow\swarrow\} = I_1^t I_1^{t+1}, \quad \{\swarrow\searrow\} = I_2^t, \quad \{\searrow\swarrow\} = I_1^{t+1}.$$

By definition, the j -th component of $\mathbf{u}^{(k)}$ satisfies

$$u_j^{(k)} = \sum_{\# \searrow = j-1, \# \swarrow = k-j+1} \{a_1, \dots, a_k\}. \quad (\text{A.3})$$

Lemma A.1 $\{\swarrow, a_2, \dots, a_k\} = I_{l+1}^t \{\searrow, a_2, \dots, a_k\}$,
where $l = \# \{\text{the arrow } \searrow \text{ included in } \{a_2, \dots, a_k\}\}$.

Proof. It is obvious by proof of induction of k . ■

Lemma A.2 $u_1^{(M)} + \sum_{j=1}^M (-1)^j u_{j+1}^{(M)} I_1^t \cdots I_j^t = 0$.

Proof.

$$\begin{aligned}
& u_1^{(M)} + \sum_{j=1}^M (-1)^j u_{j+1}^{(M)} I_1 \cdots I_j \\
&= \{\swarrow \swarrow \cdots \swarrow\} + \sum_{j=1}^M (-1)^j \left[\sum_{\# \searrow = j, \# \swarrow = M-j} \{\swarrow, *, \dots, *\} + \{\searrow, *, \dots, *\} \right] I_1 \cdots I_j \\
&= \{\swarrow \swarrow \cdots \swarrow\} + \sum_{j=1}^{M-1} (-1)^j \sum_{\# \searrow = j+1, \# \swarrow = M-j-1} \{\searrow, *\} I_1 \cdots I_j I_{j+1} \\
&\quad + \sum_{j=1}^M (-1)^j \sum_{\# \searrow = j, \# \swarrow = M-j} \{\searrow, *\} I_1 \cdots I_j \quad (\because \text{Lemma A.1}) \\
&= \{\swarrow \swarrow \cdots \swarrow\} - I_1 \{\searrow \swarrow \cdots \swarrow\} \\
&= 0 \quad \blacksquare
\end{aligned}$$

Lemma A.3 $\sum_{j=1}^M (-1)^j \sigma(u_j^{(M)}) I_1^t \cdots I_j^t = (-1)^M I_1^t \cdots I_{M+1}^T.$

Proof. We start from lemma A.2.

$$\begin{aligned}
0 &= \sigma \left(u_1^{(M)} + \sum_{j=1}^M (-1)^j u_{j+1}^{(M)} I_1 \cdots I_j \right) \\
&= \sigma(u_1^{(M)}) + \sum_{j=1}^M (-1)^j \sigma(u_{j+1}^{(M)}) I_2 \cdots I_{j+1}
\end{aligned}$$

Multiplying I_1 , we obtain

$$\begin{aligned}
0 &= \sum_{j=1}^{M+1} (-1)^j \sigma(u_j^{(M)}) I_1 \cdots I_j \\
&= \sum_{j=1}^M (-1)^j \sigma(u_j^{(M)}) I_1 \cdots I_j + (-1)^{M+1} I_1 \cdots I_{M+1},
\end{aligned}$$

which complete the proof. $\blacksquare$

proof of lemma 2.8

Calculate $\det H_t$ by the definition of H_t (2.28). The components $v_{M+2}^{(j)}$, ($j = 1, 2, \dots, M+1$) are rewritten as

$$v_{M+2}^{(1)} = -\beta_1, \quad v_{M+2}^{(2)} = x - \alpha_2^{(1)}, \quad v_{M+2}^{(j)} = -\alpha_j^{(j-1)}, \quad (j = 3, 4, \dots, M+1).$$

by virtue of (2.22). The expansion of the determinant with respect to the $(M + 1)$ -st row yields

$$\begin{aligned}
\det H_t &= (-1)^M \{ -\beta_1 + (\alpha_2^{(1)} - x)I_1^t - \alpha_3^{(2)}I_1^t I_2^t + \cdots \\
&\quad + (-1)^M (I_{M+1}^t - \alpha_{M+1}^{(M)})I_1^t \cdots I_M^t \} \\
&= (-1)^M \left\{ -\beta_1 - xI_1^t - \sum_{j=1}^M (-1)^j \alpha_{j+1}^{(j)} I_1^t \cdots I_j^t + (-1)^M I_1^t \cdots I_M^t I_{M+1}^t \right\} \\
&= (-1)^M \left\{ -V_1^t u_1^{(M)} - xI_1^t - \sum_{j=1}^M (-1)^j \{ \sigma(u_j^{(M)}) + V_1^t u_{j+1}^{(M)} \} I_1^t \cdots I_j^t \right. \\
&\quad \left. + (-1)^M I_1^t \cdots I_M^t I_{M+1}^t \right\} \\
&= (-1)^{M+1} I_1^t x. \quad (\because \text{Lemma A.2 and A.3}) \quad \blacksquare
\end{aligned}$$

References

- [1] T Tokihiro, A Nagai and J Satsuma 1999 *Inverse Problems* **15** 1639–62
- [2] Iwao S and Tokihiro T 2007 *J.Phys A. Math. Theor.* **40** 12987–13021
- [3] Kimijima T and Tokihiro T 2002 *Inverse Problems* **18** 1705–32
- [4] van Moerbeke P and Mumford D 1979 *Acta math.* **143** 94–154
- [5] Audin M 1996 *Spinning tops (Cambridge Studies in Advanced Mathematics vol 51)* ed Bolladais B and Fulton W et al (Cambridge: Cambridge university press)
- [6] Griffith P A 1985 *Amer. J. math.* **107** 1445–84
- [7] Prasolov V V 2001 *Polynomials (Algorithms and Computation in Mathematics vol 11)* ed Bronstein M and Cohen A M et al (New York: Springer)