

Homogeneous quasimorphisms on the symplectic linear group

Gabi Ben Simon Dietmar A. Salamon
ETH-Zürich

11 December 2007

Let G be a group. A **quasimorphism** on G is a map $\rho : G \rightarrow \mathbb{R}$ satisfying

$$|\rho(gh) - \rho(g) - \rho(h)| \leq C$$

for all $g, h \in G$ and a suitable constant C . It is called **homogeneous** if $\rho(g^k) = k\rho(g)$ for every $g \in G$ and every integer $k \geq 0$. Let

$$\mathrm{Sp}(2n) := \{ \Psi \in \mathbb{R}^{2n \times 2n} \mid \Psi J_0 \Psi^T = J_0 \}, \quad J_0 := \begin{pmatrix} 0 & -\mathbb{1} \\ \mathbb{1} & 0 \end{pmatrix},$$

denote the group of symplectic matrices and $\widetilde{\mathrm{Sp}}(2n)$ its universal cover. Think of an element of $\widetilde{\mathrm{Sp}}(2n)$ as a homotopy class $[\Psi]$ (with fixed endpoints) of a smooth path $\Psi : [0, 1] \rightarrow \mathrm{Sp}(2n)$ satisfying $\Psi(0) = \mathbb{1}$.

Theorem 1. *There is a unique homogeneous quasimorphism μ on $\widetilde{\mathrm{Sp}}(2n)$ that descends to the determinant homomorphism on $\mathrm{U}(n)$ in the sense that*

$$\det(X + iY) = \exp(2\pi i \mu([\Psi])), \quad \begin{pmatrix} X & -Y \\ Y & X \end{pmatrix} := \Psi(1),$$

for every $[\Psi] \in \widetilde{\mathrm{Sp}}(2n)$ with $\Psi(1) \in \mathrm{Sp}(2n) \cap \mathrm{O}(2n) \cong \mathrm{U}(n)$.

The quasimorphism of Theorem 1 plays a central role in [2] and this motivated the present note. Two explicit constructions of the quasimorphism can be found in [1] and [4]. The construction in [1] uses the unitary part in a polar decomposition and homogenization. The construction in [4] uses the

eigenvalue decomposition of a symplectic matrix (but does not mention the term *quasimorphism*).

Lemma 1 *If $\rho : G \rightarrow \mathbb{R}$ is a homogeneous quasimorphism then ρ is invariant under conjugation and $\rho(g^{-1}) = -\rho(g)$ for every $g \in G$.*

Proof of Lemma 1. Let C be the constant in the definition of quasimorphism. By homogeneity, we have $\rho(1) = 0$. Hence $|\rho(g^k) + \rho(g^{-k})| \leq C$ for every $g \in G$ and every integer $k \geq 0$. By homogeneity, we obtain $|\rho(g) + \rho(g^{-1})| \leq C/k$ for every k and so $\rho(g^{-1}) = -\rho(g)$. Hence

$$|\rho(ghg^{-1}) - \rho(h)| = |\rho(ghg^{-1}) - \rho(g) - \rho(h) - \rho(g^{-1})| \leq 2C.$$

Using homogeneity again we obtain $\rho(ghg^{-1}) = \rho(h)$ for all $g, h \in G$. \square

Proof of Theorem 1. Let $\mathcal{P} \subset \text{Sp}(2n)$ denote the set of symmetric positive definite symplectic matrices. This space is contractible and hence there is a natural injection $\iota : \mathcal{P} \rightarrow \widetilde{\text{Sp}}(2n)$. Explicitly, the map ι assigns to a matrix $P \in \mathcal{P}$ the unique homotopy class of paths $\Phi : [0, 1] \rightarrow \mathcal{P}$ with endpoints $\Phi(0) = \mathbb{1}$ and $\Phi(1) = P$.

Let $\mu : \widetilde{\text{Sp}}(2n) \rightarrow \mathbb{R}$ be a homogeneous quasimorphism that descends to the determinant homomorphism on $U(n)$. It suffices to prove that the restriction of μ to $\iota(\mathcal{P})$ is bounded. (If μ' is another quasimorphism satisfying the requirements of Theorem 1 and μ, μ' are bounded on $\iota(\mathcal{P})$ then, by polar decomposition and the determinant assumption, their difference is bounded and so, by homogeneity, they are equal.) We prove that μ vanishes on $\iota(\mathcal{P})$. For every unitary matrix $Q \in U(n) \subset \text{Sp}(2n)$ and every $P \in \mathcal{P}$ we have

$$(1) \quad \mu(\iota(QPQ^T)) = \mu(\iota(P)).$$

To see this, choose two paths $\Phi : [0, 1] \rightarrow \mathcal{P}$ and $\Psi : [0, 1] \rightarrow U(n)$ such that $\Phi(0) = \Psi(0) = \mathbb{1}$ and $\Phi(1) = P$, $\Psi(1) = Q$. Then $\mu([\Phi]) = \mu([\Psi\Phi\Psi^{-1}])$, by Lemma 1, and so (1) follows from the fact that $\Psi^{-1} = \Psi^T$. Now let $P \in \mathcal{P}$. Since P is a symmetric symplectic matrix we have $PJ_0P = J_0$ and hence

$$\mu(\iota(P)) = \mu(\iota(J_0P^{-1}J_0^{-1})) = \mu(\iota(P^{-1})) = \mu(\iota(P)^{-1}) = -\mu(\iota(P)).$$

Here the second equation follows from (1) and the last from Lemma 1. This shows that $\mu(\iota(P)) = 0$ for every $P \in \mathcal{P}$. \square

Remark 1. Lemma 1 is well known to the experts. We included a proof to give a self-contained exposition, and because we didn't find an explicit reference.

Remark 2. Related results, obtained with different methods, are contained in [1] and [3]. Our main theorem can in fact be deduced from these results.

Remark 3. The determinant homomorphism $\det : U(n) \rightarrow S^1$ is uniquely determined by the condition that it induces an isomorphism on fundamental groups. Hence it follows from Theorem 1 that the homogeneous quasimorphism $\mu : \widetilde{\mathrm{Sp}}(2n) \rightarrow \mathbb{R}$ is uniquely determined by the condition that it restricts to an isomorphism of the fundamental group of $\mathrm{Sp}(2n)$ to the integers.

References

- [1] J. Barge and E. Ghys, Cocycle D'Euler et de Maslov, *Math. Ann.* **294** (1992), 235-265.
- [2] G. Ben Simon, The geometry of partial order on contact transformations of prequantization manifolds, in *Arithmetic and Geometry Around Quantization*, edited by O. Ceyhan, Y.I. Manin and M. Marcolli, Progress in Mathematics, Birkhäuser.
- [3] M. Burger, A. Iozzi, A. Wienhard, Surface group representations with maximal Toledo invariant, Preprint, ETH-Zürich, 2006.
- [4] D.A. Salamon and E. Zehnder, Floer homology, the Maslov index, and periodic solution of Hamiltonian equations *Analysis et cetera*, edited by P.H. Rabinowitz and E. Zehnder, Academic Press, 1990, pp. 573-600.