

# One metric result about analytic continuation of some Dirichlet series.

I.S. Rezvyakova <sup>1</sup>

We consider

$$f(s, \theta) = \prod_p \left( 1 - \frac{e^{2\pi i \theta p}}{p^s} \right)^{-1}, \quad \operatorname{Re} s > 1, \quad 0 \leq \theta \leq 1,$$

where the product is taken over all prime numbers, i.e.

$$f(s, \theta) = \sum_{n=1}^{+\infty} a_n(\theta) n^{-s},$$

and for  $n = p_1^{\alpha_1} \dots p_k^{\alpha_k}$

$$a_n(\theta) = e^{2\pi i(\alpha_1 p_1 + \dots + \alpha_k p_k)\theta}.$$

**Theorem 1.** *For almost all  $\theta$ , the function  $f(s, \theta)$  has analytic continuation to the half plane  $\operatorname{Re} s > \frac{1}{2}$ , where it doesn't vanish.*

This theorem was proved by different authors. Here we give a proof that is based on the estimates of special trigonometric sums.

Let  $\operatorname{Re} s > 1$ . Then

$$\begin{aligned} \ln f(s, \theta) &= - \sum_p \ln \left( 1 - \frac{e^{2\pi i \theta p}}{p^s} \right), \\ - \frac{f'(s, \theta)}{f(s, \theta)} &= \sum_p \sum_{m=1}^{+\infty} \frac{e^{2\pi i m \theta p} \ln p}{p^{ms}} = \\ &= \sum_{n=2}^{+\infty} \frac{e^{2\pi i \theta n} \Lambda(n)}{n^s} + \sum_p \sum_{m \geq 2} \frac{e^{2\pi i m \theta p} \ln p}{p^{ms}} - \sum_p \sum_{m \geq 2} \frac{e^{2\pi i \theta p^m} \ln p}{p^{ms}}, \end{aligned}$$

---

<sup>1</sup>This work was supported by grant RFBR 06-01-00518a and grant of the President of Russian Federation MK-3370.2007.1

where  $\Lambda(n)$  — is von Mangoldt function. Note, that the last two series define analytic functions for  $\operatorname{Re} s > \frac{1}{2}$ . We show, that for almost all  $\theta$  the series

$$F(s, \theta) = \sum_{n=2}^{+\infty} \frac{e^{2\pi i \theta n} \Lambda(n)}{n^s}$$

defines analytic function on the half plane  $\operatorname{Re} s > \frac{1}{2}$ . It implies that  $(\ln f(s, \theta))'$  is analytic for  $\operatorname{Re} s > \frac{1}{2}$ , which yields the desired result.

To show that  $F(s, \theta)$  is analytic function for  $\operatorname{Re} s > \frac{1}{2}$ , it is sufficient to consider the sum

$$S_N(\theta) = \sum_{n=2}^N \Lambda(n) e^{2\pi i \theta n}$$

and estimate it by  $|S_N(\theta)| \ll_{\varepsilon} N^{\frac{1}{2} + \varepsilon}$  for every  $\varepsilon > 0$ , where the constant implied by the Vinogradov symbol  $\ll_{\varepsilon}$  may depend on  $\varepsilon$ . Indeed, let  $s = \sigma + it$ ,  $\sigma > \frac{1}{2} + 2\varepsilon$ . Then for  $N > 1$ , Abel's partial summation formula yields

$$\begin{aligned} \sum_{n=2}^N \Lambda(n) e^{2\pi i \theta n} n^{-s} &= - \int_1^N S_u(\theta) d \frac{1}{u^s} + S_N(\theta) N^{-s} \\ &\ll |s| \int_1^N u^{-1 - \sigma + \frac{1}{2} + \varepsilon} du + N^{-\sigma + \frac{1}{2} + \varepsilon} < +\infty, \end{aligned}$$

which implies, that the series for  $F(s, \theta)$  defines analytic function in the region  $\operatorname{Re} s > \frac{1}{2}$ . To complete the proof of the theorem, we prove the following lemma.

**Lemma 1.** *Let*

$$S_N(\theta) = \sum_{n=2}^N \Lambda(n) e^{2\pi i \theta n}.$$

Then for almost all  $\theta \in [0; 1]$  and every  $\varepsilon > 0$  we have

$$|S_N(\theta)| \ll_{\varepsilon} N^{\frac{1}{2}}(\ln N)^{\frac{5}{2}+\varepsilon}.$$

The proof of Lemma 1 is based on

**Lemma 2.** (Erdős - Gál - Koksma, see [1], p.154) Let  $F(M, N, \theta)$  be non-negative functions in  $L^p[0; 1]$  ( $p \geq 1$ ) for  $M, N = 0, 1, \dots$ ,  $0 \leq \theta \leq 1$  such that  $F(M, 0, \theta) = 0$  and

$$F(M, N, \theta) \leq F(M, N', \theta) + F(M + N', N - N', \theta)$$

for all  $M = 0, 1, \dots$  and all  $0 \leq N' \leq N$ . Suppose further

$$\int_0^1 F(M, N, \theta)^p d\theta = O(g(N)),$$

uniformly in  $M = 0, 1, \dots$ , where  $g(N)/N$  is a non-decreasing function.

Then for arbitrary  $\varepsilon > 0$

$$F(0, N, \theta)^p = O(g(N)(\ln N)^{p+1+\varepsilon})$$

for almost all  $\theta \in [0; 1]$ .

**Proof of lemma 1.** Set

$$F(M, N, \theta) = \left| \sum_{M < n \leq M+N} \frac{\Lambda(n)}{\ln n} e^{2\pi i \theta n} \right| \quad \text{if } N \geq 1$$

and

$$F(M, 0, \theta) = 0.$$

The non-negative functions  $F(M, N, \theta) \in L^2[0; 1]$  satisfy conditions of lemma 2. From the definition of  $F(M, N, \theta)$  we find

$$\int_0^1 F(M, N, \theta)^2 d\theta = \sum_{M < n \leq M+N} \frac{\Lambda^2(n)}{\ln^2 n} \leq \sum_{M < n \leq M+N} 1 = N.$$

According to lemma 2, for every  $\varepsilon > 0$  and almost all  $\theta \in [0; 1]$  we have

$$\left| \sum_{2 \leq n \leq N} \frac{\Lambda(n)}{\ln n} e^{2\pi i \theta n} \right| \ll_{\varepsilon} N^{\frac{1}{2}} (\ln N)^{\frac{3}{2} + \varepsilon}.$$

By Abel's partial summation formula we find

$$S_N(\theta) = \sum_{2 \leq n \leq N} \frac{\Lambda(n)}{\ln n} e^{2\pi i \theta n} \ln n = - \int_1^N \mathbb{C}(u) d \ln u + \mathbb{C}(N) \ln N,$$

where

$$\mathbb{C}(u) = \sum_{1 < n \leq N} \frac{\Lambda(n)}{\ln n} e^{2\pi i \theta n}.$$

Thus, for almost all  $\theta \in [0; 1]$  we have

$$|S_N(\theta)| \ll \max_{1 \leq u \leq N} |\mathbb{C}(u)| \ln N \ll N^{\frac{1}{2}} (\ln N)^{\frac{5}{2} + \varepsilon}.$$

Obviously, once the estimate of Lemma 1 is proved for every  $\varepsilon > 0$  and almost all  $\theta$ , it is also valid for almost all  $\theta$  and every  $\varepsilon > 0$ . This completes the proof of lemma 1.  $\square$

## References

- [1] *M.Drmota, R.F.Tichy. Sequences, Discrepancies and Applications. Lecture Notes in Mathematics 1651. Springer, Berlin, 1997.*