

# ON THE BERSTEIN-SVARC THEOREM IN DIMENSION 2

ALEXANDER N. DRANISHNIKOV<sup>1</sup> AND YULI B. RUDYAK<sup>2</sup>

ABSTRACT. We prove that for any group  $\pi$  with  $\text{cd } \pi = n$  the  $n$ th power of the Berstein class is nontrivial,  $\mathfrak{b}^n \neq 0$ . This allows us to prove the following Berstein-Svarc theorem for all  $n$ :

**Theorem.** *For a connected complex  $X$  with  $\dim X = \text{cat } X = n$ ,  $\mathfrak{b}_X^n \neq 0$  where  $\mathfrak{b}_X$  is the Berstein class of  $X$ .*

Previously it was known for  $n \geq 3$  [Sva62, Ber76].

We also prove that, for every map  $f : M \rightarrow N$  of degree  $\pm 1$  of closed orientable manifolds, the fundamental group of  $N$  is free provided that the fundamental group of  $M$  is.

## CONTENTS

1.	Introduction	1
2.	The Berstein class	2
3.	The powers of the Berstein class	4
4.	Applications to the Lusternik–Schnirelmann category	6
5.	Applications to maps of degree 1	7
	References	8

## 1. INTRODUCTION

We follow the normalization of the Lusternik-Schnirelmann category (LS category) used in the recent monograph [CLOT03], i.e. the category of the point is equal to zero. The category of a space  $X$  is denoted by  $\text{cat } X$ .

Given a group  $\pi$  we denote by  $\mathbb{Z}[\pi]$  the group ring of  $\pi$  and by  $I(\pi)$  the augmentation ideal of  $\pi$ , i.e. the kernel of the augmentation map  $\varepsilon : \mathbb{Z}[\pi] \rightarrow \mathbb{Z}$ . The cohomological dimension of  $\pi$  is denoted by  $\text{cd } \pi$ .

---

*Date:* November 22, 2018.

<sup>1</sup>Supported by NSF, grant DMS-0604494.

<sup>2</sup>Supported by NSF, grant DMS-0406311.

Consider a pointed connected  $CW$ -space  $(X, x_0)$ . Let  $p : \tilde{X} \rightarrow X$  be the universal covering and put  $\tilde{X}_0 = p^{-1}(x_0)$ . Put  $\pi = \pi_1(X, x_0)$  and let  $I(\pi)$  denote the augmentation ideal of  $\mathbb{Z}[\pi]$ . The exactness of the sequence

$$0 \rightarrow H_1(\tilde{X}, \tilde{X}_0) \rightarrow H_0(\tilde{X}_0) \rightarrow H_0(\tilde{X})$$

yields an isomorphism  $H_1(\tilde{X}, \tilde{X}_0) \cong I(\pi)$ . So, we get isomorphisms

$$H^1(X, x_0; I(\pi)) \cong \text{Hom}_{\mathbb{Z}[\pi]}(H_1(\tilde{X}, \tilde{X}_0), I(\pi)) \cong \text{Hom}_{\mathbb{Z}[\pi]}(I(\pi), I(\pi)).$$

We denote by  $\bar{\mathbf{b}} \in H^1(X, x_0; I(\pi))$  the element that corresponds to the identity  $1_{I(\pi)}$  in the right hand side. Finally, we set

$$(1.1) \quad \mathbf{b} = \mathbf{b}_X = j^* \bar{\mathbf{b}} \in H^1(X; I(\pi))$$

where  $j : X = (X, \emptyset) \rightarrow (X, x_0)$  is the inclusion of pairs. Consider the cup power  $\mathbf{b}^n \in H^n(X; I(\pi) \otimes \cdots \otimes I(\pi))$  of  $\mathbf{b}$ . The following theorem was proved by I. Berstein [Ber76, Theorem A] and A. Svarc [Sva62], see also [CLOT03, Proposition 2.1]

**1.1. Theorem.** *If  $\dim X = \text{cat } X = n \geq 3$ , then  $\mathbf{b}^n \neq 0$ . (For the case  $n = \infty$  this means that  $\mathbf{b}^k \neq 0$  for all  $k$ .)*

The proof in [Ber76, Sva62] is topological. It is based on the obstruction theory and the Whitehead approach to the Lusternik-Schnirelmann category. Thus, it cannot be extended to  $n = 2$ . In this paper we give an algebraic prove of the theorem for all  $n$  when  $X = K(\pi, 1)$ . In this case the condition  $\dim X = \text{cat } X = n$  can be replaced by a formally weaker condition  $\text{cd } \pi = n$ . Though we note that the equality  $\text{cat } K(\pi, 1) = \text{cd } \pi$  holds for all  $n$ . Then using the Stallings-Swan theorem we apply our result to extend the Berstein-Svarc theorem to the case  $n = 2$ .

For  $n = 1$  the Theorem holds for trivial reasons.

## 2. THE BERSTEIN CLASS

Here we present alternative constructions of the Berstein class and prove its universal properties. We regard  $X$  as a  $CW$ -space with one vertex and consider the chain complex

$$\dots \longrightarrow C_1(\tilde{X}) \xrightarrow{\partial_1} C_0(\tilde{X}) \longrightarrow \mathbb{Z} \longrightarrow 0$$

of the universal covering space  $\tilde{X}$ . This complex is a free resolution of the  $\mathbb{Z}[\pi]$ -module  $Z$  with  $C_0(X) \cong \mathbb{Z}[\pi]$ . Then  $\text{Im } \partial_1 = I(\pi)$ .

**2.1. Proposition.** *The class  $\mathbf{b}_X$  is the cohomology class of the cocycle  $f : C_1 \rightarrow I(\pi)$ ,  $f(c) = \partial_1(c)$ .*

*Proof.* Consider the diagram

$$\begin{array}{ccccccc} C_2(\tilde{X}) & \xrightarrow{\partial_2} & C_1(\tilde{X}) & \xrightarrow{\partial_1} & C_0(\tilde{X}) & \longrightarrow & 0 \\ \parallel & & \parallel & & & & \\ C_2(\tilde{X}, \tilde{X}_0) & \xrightarrow{\bar{\partial}_2} & C_1(\tilde{X}; \tilde{X}_0) & \xrightarrow{\bar{\partial}_1} & C_0(\tilde{X}, \tilde{X}_0) & = & 0 \end{array}$$

We can compute the groups  $H^*(X; I(\pi))$  and  $H^*(X, x_0; I(\pi))$  by applying the functor  $\text{Hom}_{\mathbb{Z}[\pi]}(-, I(\pi))$  to the first and second sequences, respectively, and then taking the cohomology of the cochain complexes obtained. Now,

$$\begin{aligned} H_1(\tilde{X}; \tilde{X}_0) &= \text{Ker } \bar{\partial}_1 / \text{Im } \bar{\partial}_2 = C_1(\tilde{X}, \tilde{X}_0) / \text{Im } \bar{\partial}_2 \\ &= C_1(\tilde{X}) / \text{Im } \partial_2 \cong I(\pi). \end{aligned}$$

The class  $\bar{\mathbf{b}}$  is given by the quotient homomorphism

$$C_1(\tilde{X}, \tilde{X}_0) \rightarrow C_1(\tilde{X}, \tilde{X}_0) / \text{Im } \partial_2 \rightarrow I(\pi).$$

Thus, the class  $\mathbf{b}$  is given by the same homomorphism  $C_1(\tilde{X}) = C_1(\tilde{X}, \tilde{X}_0) \rightarrow I(\pi)$ , but it is considered modulo

$$\text{Im}\{\delta_1 : \text{Hom}_{\mathbb{Z}[\pi]}(C_0(\tilde{X}), I(\pi)) \rightarrow \text{Hom}_{\mathbb{Z}[\pi]}(C_1(\tilde{X}), I(\pi))\}.$$

Clearly, this homomorphism coincides with  $f$ .  $\square$

The exact sequence  $0 \rightarrow I(\pi) \rightarrow \mathbb{Z}[\pi] \rightarrow \mathbb{Z} \rightarrow 0$  of  $\mathbb{Z}[\pi]$ -modules yields the cohomology exact sequence

$$\dots \rightarrow H^0(X; \mathbb{Z}) \xrightarrow{\delta} H^1(X; I(\pi)) \longrightarrow H^1(X; \mathbb{Z}[\pi]) \rightarrow \dots$$

**2.2. Proposition.** *We have the equality  $\mathbf{b}_X = \delta(1)$  where 1 is the generator of  $H^0(X; \mathbb{Z}) = \mathbb{Z}$ .*

*Proof.* The class  $\delta(1)$  is given by any cocycle  $h : C_1(\tilde{X}) \rightarrow I(\pi)$  such that diagram below commutes.

$$\begin{array}{ccccc} C_1(\tilde{X}) & \xrightarrow{\partial_1} & C_0(\tilde{X}) & \xlongequal{\quad} & C_0(\tilde{X}) \\ h \downarrow & & \parallel & & 1 \downarrow \\ I(\pi) & \longrightarrow & \mathbb{Z}[\pi] & \longrightarrow & \mathbb{Z} \end{array}$$

Clearly, the map  $f$  from Proposition 2.1 satisfies this property.  $\square$

**2.3. Theorem.** *Let  $X$  be a space with  $\pi_1(X) = \pi$  and put  $\mathbf{b} = \mathbf{b}_X$ . Let  $A$  be a local coefficient system on  $X$ .*

- (i) *For any  $a \in H_1(X; A)$ ,  $a \neq 0$ , we have  $\langle a, \mathbf{b}_X \rangle \neq 0$  in  $A \otimes_{\mathbb{Z}[\pi]} I(\pi)$ .*
- (ii) *For every  $u \in H^1(X; A)$  there exists a  $\mathbb{Z}[\pi]$ -homomorphism  $h : I(\pi) \rightarrow A$  such that  $u = h_* \mathbf{b}$ .*

*Proof.* We use the interpretation of  $\mathfrak{b}$  from Proposition 2.1. Consider the part

$$C_2(\tilde{X}) \xrightarrow{\partial_2} C_1(\tilde{X}) \xrightarrow{\partial_1} C_0(X) \longrightarrow \mathbb{Z} \longrightarrow 0$$

of the  $\mathbb{Z}[\pi]$ -resolution of  $\mathbb{Z}$ .

(i) Since the tensor product is right exact, we obtain the diagram

$$\begin{array}{ccccccc} A \otimes_{\mathbb{Z}[\pi]} C_2(\tilde{X}) & \xrightarrow{1 \otimes \partial_1} & A \otimes_{\mathbb{Z}[\pi]} C_1(\tilde{X}) & \xrightarrow{1 \otimes f} & A \otimes_{\mathbb{Z}[\pi]} I(\pi) & \longrightarrow & 0 \\ & & & & g \downarrow & & \\ & & & & A \otimes_{\mathbb{Z}[\pi]} C_0(\tilde{X}) & & \end{array}$$

where the row is exact. The composition

$$A \otimes_{\mathbb{Z}[\pi]} C_1(\tilde{X}) \xrightarrow{1 \otimes f} A \otimes_{\mathbb{Z}[\pi]} I(\pi) \xrightarrow{g} A \otimes_{\mathbb{Z}[\pi]} C_0(\tilde{X})$$

coincides with  $1 \otimes \partial_1$ . We represent the class  $a$  by a cycle

$$z \in A \otimes_{\mathbb{Z}[\pi]} C_1(\tilde{X}).$$

Since  $z \notin \text{Im}(1 \otimes \partial_2)$ , we conclude that

$$(1 \otimes f)(z) \neq 0 \text{ in } A \otimes_{\mathbb{Z}[\pi]} I(\pi) = H_0(X; \mathcal{A} \otimes I(\pi)).$$

Thus,  $\langle a, \mathfrak{b} \rangle \neq 0$ .

(ii) The class  $u$  is given by a homomorphism  $\varphi : C_1(\tilde{X}) \rightarrow A$  with  $\varphi \partial_2 = 0$ . So, there exists  $h : I(\pi) = \text{Im } \partial_1 \rightarrow M$  with  $h \partial_1 = \varphi$ .  $\square$

**2.4. Corollary.** *Suppose that there exist  $u_i \in H^1(X; A_i), i = 1, \dots, n$  such that  $u_1 \cup \dots \cup u_n \neq 0$  in  $H^n(X; A_1 \otimes \dots \otimes A_n)$ . Then  $\mathfrak{b}^n \neq 0$ .*

### 3. THE POWERS OF THE BERSTEIN CLASS

Throughout this section we fix a group  $\pi$  and denote  $\mathfrak{b}_\pi$  by  $\mathfrak{b}$ . We note that the exact sequence of  $\mathbb{Z}[\pi]$ -modules

$$(3.1) \quad 0 \rightarrow I(\pi) \rightarrow \mathbb{Z}\pi \xrightarrow{\varepsilon} \mathbb{Z} \rightarrow 0$$

stays an exact sequence of  $\mathbb{Z}[\pi]$ -modules after tensoring over  $\mathbb{Z}$  with any  $\mathbb{Z}[\pi]$ -module  $M$ :

$$0 \rightarrow I(\pi) \otimes M \rightarrow \mathbb{Z}\pi \otimes M \xrightarrow{\varepsilon \otimes 1} M \rightarrow 0.$$

Here the  $\mathbb{Z}$ -tensor product  $M \otimes N$  is taken with the diagonal  $\pi$ -action.

**3.1. Proposition.** *For every short exact sequence of left  $\mathbb{Z}[\pi]$ -modules  $0 \rightarrow N' \rightarrow N \rightarrow N'' \rightarrow 0$  and every right  $\mathbb{Z}[\pi]$ -module  $M$  such that the sequence*

$$(3.2) \quad 0 \rightarrow N' \otimes M \rightarrow N \otimes M \rightarrow N'' \otimes M \rightarrow 0$$

is exact, there is the formula  $\delta(u \cup v) = \delta u \cup v$  where  $u \in H^p(\pi; N'')$ ,  $v \in H^q(\pi; M)$ ,  $u \cup v \in H^{p+q}(\pi; N' \otimes M)$  and

$$\delta : H^{p+q}(\pi; N'' \otimes M) \rightarrow H^{p+q+1}(\pi; N' \otimes M)$$

is the connecting homomorphism in the long exact sequence obtained from the coefficient sequence (3.2).

*Proof.* see [Bro94, V,(3.3)].  $\square$

**3.2. Corollary.** *The connecting homomorphism*

$$\delta : H^i(\pi; M) \rightarrow H^{i+1}(\pi; I(\pi) \otimes M)$$

*induced by the coefficient exact sequence*

$$(3.3) \quad 0 \rightarrow I(\pi) \otimes M \rightarrow \mathbb{Z}\pi \otimes M \xrightarrow{\alpha} \mathbb{Z} \otimes M \rightarrow 0$$

*has the form  $\delta(u) = \mathfrak{b} \cup u$ .*

*Proof.* Note that every  $u \in H^1(\pi; \mathbb{Z} \otimes M)$  can be presented as  $1 \cup u$  where  $1 \in H^0(\pi; \mathbb{Z})$ . Since by Proposition 2.2  $\delta(1) = \mathfrak{b}$ , the result follows from Proposition 3.1.  $\square$

Let  $I = I(\pi)$ . Consider the exact sequence

$$0 \rightarrow I^{k+1} \xrightarrow{\beta_k} \mathbb{Z}\pi \otimes I^k \xrightarrow{\alpha_k} I^k \rightarrow 0$$

obtained from the sequence (3.1) via tensoring by  $I^k$ .

**3.3. Lemma.** (i) *The  $\mathbb{Z}[\pi]$ -module  $\mathbb{Z}[\pi]^{\otimes k} = \mathbb{Z}[\pi] \otimes \cdots \otimes \mathbb{Z}[\pi]$  is free for all  $k$ ;*

(ii) *the  $\mathbb{Z}[\pi]$ -module  $\mathbb{Z}[\pi] \otimes I^k$  is projective for all  $k$ .*

*Proof.* (i) Clearly,  $\mathbb{Z}[\pi] \otimes \cdots \otimes \mathbb{Z}[\pi] = \mathbb{Z}[\pi \times \cdots \times \pi]$  is a free  $\mathbb{Z}[\pi \times \cdots \times \pi]$ -module. Regarding  $\pi$  as the diagonal subgroup of  $\pi \times \cdots \times \pi$ , we conclude that  $\mathbb{Z}[\pi]^{\otimes k}$  is a free  $\mathbb{Z}[\pi]$ -module. (If  $H$  is a subgroup of  $G$  then  $\mathbb{Z}[G]$  is a free  $\mathbb{Z}[H]$ -module with the base  $G/H$ .)

(ii) We tensor the sequence (3.1) by  $I^{k-1} \otimes \mathbb{Z}[\pi]$  and get the exact sequence

$$0 \rightarrow I^k \otimes \mathbb{Z}[\pi] \rightarrow I^{k-1} \otimes \mathbb{Z}[\pi] \otimes \mathbb{Z}[\pi] \rightarrow I^{k-1} \otimes \mathbb{Z}[\pi] \rightarrow 0.$$

Now induct on  $k$ , using (i) as the basis of induction.  $\square$

**3.4. Theorem.** *If  $\text{cd } \pi = n$ , then  $\mathfrak{b}^n \neq 0$ .*

*Proof.* For  $n = 1$  this is trivial. Let  $n > 1$ . In view of Lemma 3.3 there is a projective resolution

$$(3.4) \quad C_n \xrightarrow{\partial_n} C_{n-1} \rightarrow \cdots \rightarrow C_1 \rightarrow C_0 \rightarrow \mathbb{Z} \rightarrow 0$$

such that  $C_k = \mathbb{Z}\pi \otimes I^k$  for all  $k$  and with

$$\partial_k = \beta_{k-1} \circ \alpha_k : \mathbb{Z}\pi \otimes I^k \xrightarrow{\alpha_k} \mathbb{Z} \otimes I^k \xrightarrow{\beta_{k-1}} \mathbb{Z}\pi \otimes I^{k-1}.$$

Claim:  $\mathfrak{b}^k = [\alpha_k]$ . We apply the induction on  $k$ .

For  $k = 1$ , note that in Proposition 2.1 we can construct  $\mathfrak{b}$  using any  $\mathbb{Z}[\pi]$ -resolution of  $\mathbb{Z}$  provided  $C_0 = \mathbb{Z}[\pi]$ . Hence,  $\mathfrak{b} = [\alpha_1]$ . Furthermore, it is easy to see that  $\delta([\alpha_k]) = [\alpha_{k+1}]$ , and hence  $\mathfrak{b}^k = [\alpha_k]$  by Corollary 3.2

Now we prove that  $\mathfrak{b}^n \neq 0$ . Suppose the contrary. Then  $\alpha_n = \varphi \circ \partial_n = \varphi \circ \beta_{n-1} \circ \alpha_n$  for some  $\varphi : C_{n-1} \rightarrow I^n$ . Since  $\alpha_n$  is an epimorphism, we conclude that  $\varphi$  yields a  $\mathbb{Z}[\pi]$ -splitting of the sequence

$$0 \rightarrow I^n \xrightarrow{\beta_{n-1}} \mathbb{Z}\pi \otimes I^{n-1} \xrightarrow{\alpha_{n-1}} I^{n-1} \rightarrow 0.$$

This implies that  $I^{n-1}$  is projective. Thus the above resolution can be shortened by one. Therefore  $\text{cd } \pi \leq n - 1$ . Contradiction.  $\square$

The following universal property was stated as [Sva62, Proposition 3.4] without proof.

**3.5. Corollary.** *Given a  $\mathbb{Z}[\pi]$ -module  $A$  and a class  $a \in H^k(\pi; A)$ , there exists a  $\mathbb{Z}[\pi]$ -homomorphism  $\mu : I^k \rightarrow A$  such that  $\mu_*(\mathfrak{b}) = a$ .*

*Proof.* Consider the resolution (3.4). The class  $a$  is given by a homomorphism  $h : \mathbb{Z}[\pi] \otimes I^k \rightarrow A$ . Since  $h$  is a cocycle,  $h \circ \partial_{k+1} = 0$ , we conclude that  $h = \mu \circ \alpha_k$  for some  $\mu : I^k \rightarrow A$ .  $\square$

**3.6. Remarks.** 1. Of course, Corollary 3.5 implies Theorem 3.4. However, we decided to prove Theorem 3.4 here since Svarc [Sva62] stated it without proof.

2. For  $n \neq 2$  Theorem 3.4 follows from Theorem 1.1 in view of the equality  $\text{cat } K(\pi, 1) = n$  and existence of an  $n$ -dimensional  $CW$ -space  $K(\pi, 1)$ , [EG57]. So, the new point is the case  $n = 2$ .

#### 4. APPLICATIONS TO THE LUSTERNIK–SCHNIRELMANN CATEGORY

Now we can prove the Berstein-Svarc Theorem for  $n = 2$ .

**4.1. Proposition.** *If  $X$  is a connected 2-dimensional  $CW$ -complex with  $\pi_1(X)$  free, then  $\text{cat } X \leq 1$ .*

*Proof.* Since  $\pi_1(X)$  is free, we can construct a space

$$Y = (\vee_{\alpha} S_{\alpha}^1) \bigvee (\vee_{\beta} S_{\beta}^2)$$

and a map  $f : Y \rightarrow X$  such that  $f_* : \pi_1(Y) \rightarrow \pi_1(X)$  is an isomorphism and  $f_* : \pi_2(Y) \rightarrow \pi_2(X)$  is an epimorphism. Since the homotopy fiber

of  $f$  is simply connected, there exists a map (homotopy section)  $s : X \rightarrow Y$  such that  $fs \cong 1_X$ . Hence,  $\text{cat } X \leq \text{cat } Y = 1$ .  $\square$

**4.2. Theorem.** *Given a connected 2-dimensional CW-complex  $X$ , the equality  $\text{cat } X = 2$  holds if and only if  $\mathfrak{b}_X^2 \neq 0$ .*

*Proof.* Let  $\pi$  denote  $\pi_1(X)$ . First, note that  $\text{cat } X \leq \dim X = 2$ . Clearly, if  $\mathfrak{b}_X^2 \neq 0$  then  $\text{cat } X \geq 2$ , and so  $\text{cat } X = 2$ . Furthermore, if  $\text{cat } X = 2$  then  $\pi$  is not free by Proposition 4.1. Hence, according to Stallings [Sta68] and Swan [Swan69],  $\text{cd } \pi > 1$ . So, by Theorem 3.4,  $\mathfrak{b}_\pi^2 \neq 0$ . Let  $f : X \rightarrow K(\pi, 1)$  be a map that induces an isomorphism on the fundamental groups. Then  $f^* : H^2(\pi; M) \rightarrow H^2(X; M)$  is injective for any  $\mathbb{Z}[\pi]$ -module  $M$ , and the theorem follows since  $f^*(\mathfrak{b}_\pi) = \mathfrak{b}_X$ .  $\square$

## 5. APPLICATIONS TO MAPS OF DEGREE 1

Let  $X$  be a closed oriented  $n$ -dimensional manifold with the fundamental class  $[X]$ . Then for a local coefficient system  $G$  on  $X$  we have the Poincaré duality isomorphism  $\Delta : H^i(X; G) \rightarrow H_{n-i}(X; G)$ ,  $\Delta(x) = x \cap [X]$ , see e.g. [Bre97]. Given a map  $f : M^n \rightarrow N^n$  of two closed manifolds and a local coefficient system  $G$  on  $N$ , we define the homomorphism  $f^! : H^*(M; f^*(G)) \rightarrow^*(N; G)$  by setting

$$f^!(x) = \Delta_N^{-1} f_* \Delta_M(x)$$

For future needs we state the following known proposition.

**5.1. Proposition.** *The index  $[\pi_1(N) : f_*(\pi_1(M))]$  does not exceed  $|\deg f|$  provided that  $\deg f \neq 0$ , and  $f^! f^*(x) = (\deg f)x$  for any  $x \in H^*(M, f^*(G))$ .*

*Proof.* The first claim follows since  $f$  passes through the covering map  $\tilde{N} \rightarrow N$  corresponding to the subgroup  $f_*(\pi_1(M))$  of  $\pi_1(N)$ . For the second claim, we have

$$\begin{aligned} f^!(f^*x) &= \Delta_N^{-1} f_* \Delta_M(f^*x) = \Delta_N^{-1} f_*((f^*x \cap M) = \Delta_N^{-1}(x \cap f_*M) \\ &= \deg f (\Delta_N^{-1}(x) \cap N) = (\deg f)x \end{aligned}$$

$\square$

**5.2. Theorem.** *If  $f : M \rightarrow N$  is a map of degree 1 and  $\pi_1(M)$  is free then  $\pi_1(N)$  is free.*

*Proof.* Let  $\pi = \pi_1(M)$ ,  $\tau = \pi_1(N)$  and assume that  $\tau$  is not free. Then  $\text{cd } \tau > 1$  according to Stallings [Sta68] and Swan [Swan69]. So,  $\mathfrak{b}_\tau^2 \neq 0$  by Theorem 3.4. Hence  $\mathfrak{b}_N^2 \neq 0$ , and so  $f^* \mathfrak{b}_N^2 \neq 0$  by Proposition 5.1.

So,  $\mathfrak{b}_M^2 \neq 0$  by Corollary 2.4. Hence,  $\mathfrak{b}_\pi^2 \neq 0$ , and so  $\text{cd } \pi > 1$ . Thus,  $\pi$  is not free.  $\square$

**5.3. Remarks.** 1. Note that  $f_*; \pi_1(M) \rightarrow \pi_1(N)$  is an epimorphism for any map of degree 1 because of Proposition 5.1

2. Every epimorphism  $h : F_r \rightarrow F_s$  of free finitely generated groups can be realized as  $h = f_* : \pi_1(M) \rightarrow \pi_1(N)$  for a map  $f : M \rightarrow N$  of degree 1. Indeed, take  $M$  and  $N$  to be the connected sums of  $r$  and  $s$  copies of  $S^1 \times S^n$ , respectively, and put  $f$  to be the map that shrinks  $r - s$  copies of  $S^1 \times S^n$ .

## REFERENCES

- [Ber76] Berstein, I: On the Lusternik-Schnirelmann category of Grassmannians. *Math. Proc. Camb. Philos. Soc.* **79** 404(1976) 129–134.
- [Bro94] Brown, K.: Cohomology of groups. *Graduate Texts in Mathematics*, **87** Springer, New York Heidelberg Berlin, 1994.
- [Bre97] Bredon, G.: Sheaf Theory. *Graduate Text in Mathematics*, **170**, Springer, New York Heidelberg Berlin, 1997.
- [CLOT03] Cornea, O.; Lupton, G.; Oprea, J.; Tanré, D.: Lusternik-Schnirelmann category. *Mathematical Surveys and Monographs*, **103**. American Mathematical Society, Providence, RI, 2003.
- [EG57] Eilenberg, S; Ganea, T.: On the Lusternik-Schnirelmann category of abstract groups. *Ann. of Math.* (2) **65** (1957), 517–518.
- [Sta68] Stallings, J.: Groups of dimension 1 are locally free. *Bull. Amer. Math. Soc.* **74** (1968), 361–364.
- [Sva62] Švarc, A.: The genus of a fibered space. *Trudy Moskov. Mat. Obšč* **10**, **11** (1961 and 1962), 217–272, 99–126, (in *Amer. Math. Soc. Transl. Series 2*, vol **55** (1966)).
- [Swan69] Swan, R.: Groups of cohomological dimension one. *J. Algebra* **12** (1969), 585–610.

ALEXANDER N. DRANISHNIKOV, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF FLORIDA, LITTLE HALL, GAINESVILLE, FL 1-10, USA

*E-mail address:* dranish@math.ufl.edu

YULI B. RUDYAK, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF FLORIDA, LITTLE HALL, GAINESVILLE, FL 1-10, USA

*E-mail address:* rudyak@math.ufl.edu