

THE DIV-CURL LEMMA REVISITED

DAN POLIŠEVSKI †

Abstract. The Div-Curl Lemma, which is the basic result of the compensated compactness theory in Sobolev spaces(see [2]-[6]), was introduced by [1] with distinct proofs for the $L^2(\Omega)$ and $L^p(\Omega), p \neq 2$, cases. In this note we present a slightly different proof, relying only on a Green-Gauss integral formula and on the usual Rellich-Kondrachov compactness properties.

Keywords: Compensated compactness; Weak convergence; Sobolev spaces.

2000 Mathematics Subject Classifications: 49J45; 46E40; 47B07.

1 An integral formula

From now on, for any vector distribution w in \mathbb{R}^N , $N \geq 2$, we denote:

$$D(w) = \sum_{i=1}^n \frac{\partial w_i}{\partial x_i} \quad \text{and} \quad C_{ij}(w) = \frac{\partial w_i}{\partial x_j} - \frac{\partial w_j}{\partial x_i}, \quad 1 \leq i, j \leq N.$$

Theorem 1.1. *Let Ω be an open bounded set of \mathbb{R}^N , $N \geq 2$. Let $p, q > 1$ such that $1/p + 1/q = 1$. If $u \in [W_{loc}^{2,p}(\Omega)]^N$ and $v \in [W_{loc}^{2,q}(\Omega)]^N$, then for any $\varphi \in \mathcal{D}(\Omega)$ we have:*

$$\begin{aligned}
& \int_{\Omega} \varphi \Delta u \Delta v = -\langle D(\Delta u), \varphi D(v) \rangle_p - \int_{\Omega} D(u) (\nabla \varphi) \Delta v + \\
(1.1) \quad & + \int_{\Omega} D(u) (\nabla \varphi) \nabla D(v) - \int_{\Omega} D(v) (\nabla \varphi) \nabla D(u) - \\
& - \sum_{1 \leq i, j \leq N} \int_{\Omega} \Delta v_i C_{ij}(u) \frac{\partial \varphi}{\partial x_j} - \frac{1}{2} \sum_{1 \leq i, j \leq N} \langle C_{ij}(\Delta v), \varphi C_{ij}(u) \rangle_q,
\end{aligned}$$

where $\langle \cdot, \cdot \rangle_p$ denotes the pairing of $W^{-1,p}(\Omega)$ with $W_0^{1,q}(\Omega)$ and $\langle \cdot, \cdot \rangle_q$ denotes the pairing of $W^{-1,q}(\Omega)$ with $W_0^{1,p}(\Omega)$.

Proof. For any $u \in [W_{loc}^{2,p}(\Omega)]^N$, $v \in [W_{loc}^{2,q}(\Omega)]^N$ and $\varphi \in \mathcal{D}(\Omega)$ we have

$$\begin{aligned}
& \langle D(\Delta u), \varphi D(v) \rangle_p + \frac{1}{2} \sum_{1 \leq i, j \leq N} \langle C_{ij}(\Delta v), \varphi C_{ij}(u) \rangle_q = \\
(1.2) \quad & = - \int_{\Omega} \Delta u \nabla (\varphi D(v)) + \int_{\Omega} \Delta v (\nabla D(u) - \Delta u) \varphi - \\
& - \sum_{1 \leq i, j \leq N} \int_{\Omega} \Delta v_i C_{ij}(u) \frac{\partial \varphi}{\partial x_j},
\end{aligned}$$

which implies:

$$\begin{aligned}
& \int_{\Omega} \varphi \Delta u \Delta v = -\langle D(\Delta u), \varphi D(v) \rangle_p - \int_{\Omega} D(u) (\nabla \varphi) \Delta v - \\
(1.3) \quad & - \int_{\Omega} \Delta u \nabla (\varphi D(v)) + \int_{\Omega} \Delta v \nabla (\varphi D(u)) - \\
& - \sum_{1 \leq i, j \leq N} \int_{\Omega} \Delta v_i C_{ij}(u) \frac{\partial \varphi}{\partial x_j} - \frac{1}{2} \sum_{1 \leq i, j \leq N} \langle C_{ij}(\Delta v), \varphi C_{ij}(u) \rangle_q.
\end{aligned}$$

As $(\nabla D(u) - \Delta u) \in [L_{loc}^p(\Omega)]^N$ is divergence-free and as $\varphi D(v) \in W_0^{1,q}(\Omega)$, it follows:

$$\begin{aligned}
(1.4) \quad & \int_{\Omega} \Delta u \nabla (\varphi D(v)) = - \int_{\Omega} (\nabla D(u) - \Delta u) \nabla (\varphi D(v)) + \\
& + \int_{\Omega} \nabla D(u) \nabla (\varphi D(v)) = \int_{\Omega} \nabla D(u) \nabla (\varphi D(v)).
\end{aligned}$$

By treating similarly the next term of (1.3), that is $\int_{\Omega} \Delta v \nabla (\varphi D(u))$, we get straightly (1.1).

2 The Div-Curl Lemma

Theorem 2.1. *Let Ω be an open bounded set of \mathbb{R}^N , $N \geq 2$. Let $p, q > 1$ such that $1/p + 1/q = 1$. For any $\varepsilon > 0$, let $a^\varepsilon \in [L^p(\Omega)]^N$ and $b^\varepsilon \in [L^q(\Omega)]^N$ with the properties:*

- (i) $a^\varepsilon \rightharpoonup a$ weakly in $[L^p(\Omega)]^N$, as $\varepsilon \rightarrow 0$,
- (ii) $b^\varepsilon \rightharpoonup b$ weakly in $[L^q(\Omega)]^N$, as $\varepsilon \rightarrow 0$,
- (iii) $\{D(a^\varepsilon)\}_\varepsilon$ lies in a compact subset of $W^{-1,p}(\Omega)$,
- (iv) $\{C(b^\varepsilon)\}_\varepsilon$ lies in a compact subset of $[W^{-1,q}(\Omega)]^{N \times N}$,

Then $a^\varepsilon b^\varepsilon \rightarrow ab$ in the sense of distributions.

Proof. For any $\varepsilon > 0$, let u^ε and v^ε be the weak solutions of the following problems:

$$(2.1) \quad -\Delta u^\varepsilon = a^\varepsilon \quad \text{in } \Omega, \quad u^\varepsilon = 0 \quad \text{on } \partial\Omega,$$

$$(2.2) \quad -\Delta v^\varepsilon = b^\varepsilon \quad \text{in } \Omega, \quad v^\varepsilon = 0 \quad \text{on } \partial\Omega.$$

From hypotheses (i)-(ii) it follows that

$$(2.3) \quad u^\varepsilon \rightharpoonup u \quad \text{weakly in } [W^{1,p}(\Omega)]^N,$$

$$(2.4) \quad v^\varepsilon \rightharpoonup v \quad \text{weakly in } [W^{1,q}(\Omega)]^N,$$

where u and v are uniquely defined by the problems:

$$(2.5) \quad -\Delta u = a \quad \text{in } \Omega, \quad u = 0 \quad \text{on } \partial\Omega,$$

$$(2.6) \quad -\Delta v = b \text{ in } \Omega, \quad v = 0 \text{ on } \partial\Omega.$$

For $\varphi \in \mathcal{D}(\Omega)$, we denote by Ω_φ a smooth open set with the property:

$$(2.7) \quad \text{supp } \varphi \subseteq \Omega_\varphi \subseteq \overline{\Omega_\varphi} \subseteq \Omega.$$

Then, the integral formula (1.1) yields:

$$(2.8) \quad \begin{aligned} \int_{\Omega} a^\varepsilon b^\varepsilon \varphi &= \langle D(a^\varepsilon), \varphi D(v^\varepsilon) \rangle_p + \int_{\Omega_\varphi} b^\varepsilon D(u^\varepsilon) \nabla \varphi + \\ &+ \int_{\Omega_\varphi} D(u^\varepsilon) (\nabla \varphi) \nabla D(v^\varepsilon) - \int_{\Omega_\varphi} D(v^\varepsilon) (\nabla \varphi) \nabla D(u^\varepsilon) + \\ &+ \sum_{1 \leq i, j \leq N} \int_{\Omega_\varphi} b_i^\varepsilon C_{ij}(u^\varepsilon) \frac{\partial \varphi}{\partial x_j} + \frac{1}{2} \sum_{1 \leq i, j \leq N} \langle C_{ij}(b^\varepsilon), \varphi C_{ij}(u^\varepsilon) \rangle_q. \end{aligned}$$

From hypotheses (iii)-(iv) we find that there exists a subsequence (still denoted with ε) for which the following properties hold:

$$(2.9) \quad D(a^\varepsilon) \rightarrow D(a) \text{ strongly in } W^{-1,q}(\Omega),$$

$$(2.10) \quad C(b^\varepsilon) \rightarrow C(b) \text{ strongly in } [W^{-1,p}(\Omega)]^{N \times N},$$

The theorem about the interior regularity of the solutions of the problems (1)-(2)(see [7] §6.3) implies:

$$(2.11) \quad \{u_\varepsilon\}_\varepsilon \text{ is bounded in } [W^{2,p}(\Omega_\varphi)]^N,$$

$$(2.12) \quad \{v_\varepsilon\}_\varepsilon \text{ is bounded in } [W^{2,q}(\Omega_\varphi)]^N.$$

In the light of (2.9)-(2.10) and using the appropriate Rellich-Kondrachov compactness results (see [7] §5.7) that follow from (2.11)-(2.12), we see that every term of the right-hand side of (2.8) represents a product of a strongly convergent (sub)sequence by a weakly convergent one. Hence, the right-hand

side of (2.8) is converging to the corresponding formula which represents $\int_{\Omega} ab\varphi$. By the uniqueness of the limit, the convergence holds on the entire sequence.

References.

- [1] Murat F., *Compacité par compensation*, Ann. Scuola Norm. Sup. Pisa Cl. Sci., **5(4)**, 1978.
- [2] Murat F., *Compacité par compensation II, Recent Methods in Nonlinear Analysis* (De Giorgi E., Magenes E., Mosco U. eds.), Pitagora Editrice, Bologna, 1979.
- [3] Tartar L., *Compensated compactness and applications to partial differential equations*, Heriot-Watt Sympos., vol. IV, Pitman, New York, 1979.
- [4] Dacorogna B., *Weak Continuity and Weak Lower Semicontinuity of Nonlinear Functionals*, Springer-Verlag, Berlin, 1982.
- [5] Tartar L., *The compensated compactness method applied to systems of conservation laws*, *Systems of Nontinear PDE* (Ball J. ed.), D. Reidel Pub. Co., Dordrecht, 1983.
- [6] Evans L.C., *Weak Convergence Methods for Nonlinear Partial Differential Equations* (CBMS Regional Conference Series in Mathematics, vol.74), American Mathematical Society, Providence, 1991.
- [7] Evans L.C., *Partial Differential Equations* (Graduate Studies in Mathematics, vol.19), American Mathematical Society, Providence, 1998.

† I.M.A.R., P.O. BOX 1-764, BUCHAREST, ROMANIA