

One- and two-dimensional Coulomb Green's function matrices in parabolic Sturmians basis

S. A. Zaytsev*

Pacific National University, Khabarovsk, 680035, Russia

Abstract

One- and two-dimensional operators which originate from the asymptotic form of the three-body Coulomb wave equation in parabolic coordinates are treated within the context of square integrable basis set. The matrix representations of Green's functions corresponding to these operators are obtained.

PACS numbers: 02.30Gp, 03.65Ca, 03.65.Nk

*E-mail: zaytsev@fizika.khstu.ru

I. INTRODUCTION

The three Coulomb (3C) wave functions [1, 2] have been introduced to approximate the three-body continuum Coulomb wave function. The three-body Coulomb wave equation is asymptotically separable in terms of parabolic coordinates introduced by Klar [3]

$$\xi_j = r_{ls} + \hat{\mathbf{k}}_{ls} \cdot \mathbf{r}_{ls}, \quad \eta_j = r_{ls} - \hat{\mathbf{k}}_{ls} \cdot \mathbf{r}_{ls}, \quad (1)$$

where \mathbf{r}_{ls} and \mathbf{k}_{ls} , $l, s = 1, 2, 3$ and $l \neq s$, are the relative coordinate and momentum vectors between the particles l and s . The 3C wave functions are proportional to solutions Ψ_{3C} of the asymptotic equation [3] (atomic units are used):

$$\left\{ \sum_{j=1}^3 \frac{1}{\mu_{ls}(\xi_j + \eta_j)} \left[\hat{h}_{\xi_j} + \hat{h}_{\eta_j} + 2k_{ls}t_{ls} \right] \right\} \Psi_{3C} = 0, \quad (2)$$

where $t_{ls} = \frac{Z_l Z_s \mu_{ls}}{k_{ls}}$, $\mu_{ls} = \frac{m_l m_s}{m_l + m_s}$ are the reduced masses. Here the operators \hat{h}_{ξ_j} and \hat{h}_{η_j} are defined by

$$\hat{h}_{\xi_j} = -2 \left(\frac{\partial}{\partial \xi_j} \xi_j \frac{\partial}{\partial \xi_j} + ik_{ls} \xi_j \frac{\partial}{\partial \xi_j} \right), \quad (3)$$

$$\hat{h}_{\eta_j} = -2 \left(\frac{\partial}{\partial \eta_j} \eta_j \frac{\partial}{\partial \eta_j} - ik_{ls} \eta_j \frac{\partial}{\partial \eta_j} \right). \quad (4)$$

A solution Ψ_{3C} with pure outgoing behavior can be written as a product of three two-body Coulomb wave functions:

$$\Psi_{3C} = \prod_{j=1}^3 u_j(\xi_j) \quad (5)$$

with

$$u_j(\xi_j) = {}_1F_1(it_{ls}, 1; -ik_{ls}\xi_j). \quad (6)$$

The solutions (5) are correct only when all particles are far from each other. In the last years many works have been published with different proposal to improve the simply 3C (see [4, 5, 6] and references therein). In this paper we seek to consider the possibilities for computing the three-body continuum Coulomb wave function which are afforded by expansion in a set of square-integrable functions. In principle, the short-range part of the Hamiltonian, which contains mixed second derivative terms in the expression for the kinetic energy [3], can be approximated by a finite order matrix. The long-range part (2) of the Hamiltonian consists of two-dimensional operators

$$\hat{\mathbf{h}}_j = \hat{h}_{\xi_j} + \hat{h}_{\eta_j} + 2k_{ls}t_{ls} + C_j(\xi_j + \eta_j), \quad (7)$$

which are related by the constraint [3]

$$C_1 + C_2 + C_3 = 0. \quad (8)$$

To treat the one-dimensional operators

$$\hat{h}_\xi + 2kt + C\xi, \quad \hat{h}_\eta + 2kt + C\eta \quad (9)$$

within the context of square integrable basis set, the J -matrix method [7, 8] or the tools of the “Tridiagonal Physics” program (see [9] and reference therein) can be employed. In the framework of these methods a special basis set is used which supports an infinite tridiagonal matrix representation of the operator. Thus the resulting three-term recursion is analytically solved.

In particular, in this paper we obtain matrices of Green’s functions (resolvent) corresponding to the one-dimensional operators (9) and the two-dimensional operator

$$\hat{\mathbf{h}} = \hat{h}_\xi + \hat{h}_\eta + 2kt_0 + C(\xi + \eta) \quad (10)$$

in the set of parabolic Sturmian functions,

$$\phi_{n_1 n_2}(\xi, \eta) = \varphi_{n_1}(\xi) \varphi_{n_2}(\eta), \quad (11)$$

$$\varphi_n(x) = \sqrt{2b}e^{-bx} L_n(2bx), \quad (12)$$

where b is the scale parameter. This basis set have been used in the analysis of the Coulomb potential within the parabolic formulation of the J -matrix method [10]. The basis functions (12) are orthonormal:

$$\int_0^\infty dx \varphi_n(x) \varphi_m(x) = \delta_{nm}. \quad (13)$$

In Sec. II using the tridiagonal matrix representations of one-dimensional operators (9) in the bases (12), we construct the corresponding Green’s function matrices. In this case we do not seek to determine the Green’s matrices uniquely. In Sec. III the weight function is obtained for the orthogonal polynomials satisfying the three-term recurrence relation. The two-dimensional Green’s function matrix elements are expressed as convolution of one-dimensional Green’s matrix elements in Sec. III. An orthogonality relation employed in the two-dimensional Green’s matrix construction is derived in the Appendix.

II. ONE-DIMENSIONAL COULOMB GREEN'S FUNCTION MATRICES

a) $C = 0$

The matrix representation $\mathbf{h}_\xi + 2kt\mathbf{I}_\xi$ [\mathbf{I}_ξ is the unit matrix] of the operator $\hat{h}_\xi + 2kt$ in the basis set $\{\varphi_n(\xi)\}_{n=0}^\infty$ (12) is tridiagonal

$$\mathbf{h}_\xi + 2kt\mathbf{I}_\xi = \begin{pmatrix} b_0 & d_1 & & & \\ a_1 & b_1 & d_2 & & 0 \\ & a_2 & b_2 & d_3 & \\ & & a_3 & \times & \times \\ 0 & & & \times & \times & \times \end{pmatrix}, \quad (14)$$

where

$$b_n = (b + ik) + 2bn + 2kt, \quad (15)$$

$$a_n = (b - ik)n, \quad d_n = (b + ik)n.$$

To construct the Green's function matrix \mathbf{g}_ξ , which is matrix inverse of the infinite tridiagonal matrix (14), consider the three-term recurrence relation

$$a_n w_{n-1} + b_n w_n + d_{n+1} w_{n+1} = 0, \quad n \geq 1. \quad (16)$$

It can be easily verified that

$$p_n(t; \zeta) = \frac{(-1)^n \Gamma(n+1-it)}{n! \Gamma(1-it)} {}_2F_1(-n, it; -n+it; \zeta), \quad (17)$$

where $\zeta = \frac{b-ik}{b+ik}$, is the “regular” solution of (16) which satisfies the initial conditions:

$$p_0(t; \zeta) = 1, \quad b_0 p_0(t; \zeta) + d_1 p_1(t; \zeta) = 0. \quad (18)$$

Suffice it to say that apart from the factor $\sqrt{\frac{2}{b}} \left(\frac{\zeta+1}{2}\right)^{it}$, p_n is the coefficient of the n th basis function $\varphi_n(\xi)$ (12) in the expansion of $u(\xi)$ (6). Note that p_n are polynomials of degree n in t .

The second solution q_n of the recursion (16) can be obtained from the condition that q_n satisfies the same differential equation as the p_n [8]. In other words, if $p_n \sim {}_2F_1(a, b; c; z)$, then $q_n \sim z^{1-c} {}_2F_1(a-c+1, b-c+1; 2-c; z)$. It is readily verified that an appropriate $q_n(t; \zeta)$ is

$$\begin{aligned} q_n(t; \zeta) &= -\frac{n!\Gamma(1-it)}{\Gamma(n+2-it)} (-\zeta)^{n+1} {}_2F_1(1-it, n+1; n+2-it; \zeta) \\ &= -\frac{n!\Gamma(1-it)}{\Gamma(n+2-it)} \left(\frac{\zeta}{\zeta-1}\right)^{n+1} {}_2F_1(n+1, n+1; n+2-it; \frac{\zeta}{\zeta-1}) \end{aligned} \quad (19)$$

This satisfies the initial condition

$$b_0 q_0(t; \zeta) + d_1 q_1(t; \zeta) = b - ik. \quad (20)$$

Multiplying $[\mathbf{h}_\xi + 2kt\mathbf{I}_\xi]$ by the diagonal matrix $\mathbf{Z} = [1, \zeta^{-1}, \dots, \zeta^{-n}, \dots]$, we obtain the symmetric tridiagonal matrix \mathbf{T} :

$$\mathbf{T} = \mathbf{Z} [\mathbf{h}_\xi + 2kt\mathbf{I}_\xi] = \begin{pmatrix} \beta_0 & \alpha_1 & & & \\ \alpha_1 & \beta_1 & \alpha_2 & & 0 \\ & \alpha_2 & \beta_2 & \alpha_3 & \\ & & \alpha_3 & \times & \times \\ 0 & & & \times & \times & \times \end{pmatrix} \quad (21)$$

with nonzero elements

$$\beta_n = b_n/\zeta^n, \quad \alpha_n = d_n/\zeta^{n-1}. \quad (22)$$

Notice that p_n and q_n satisfy the three term recurrence relation

$$\alpha_n w_{n-1} + \beta_n w_n + \alpha_{n+1} w_{n+1} = 0, \quad n \geq 1. \quad (23)$$

Thus, to invert the symmetric tridiagonal matrix \mathbf{T} , one can draw on the standard method [11, 12]. Namely, the elements of a Green's matrix \mathbf{g}_T , which is the matrix inverse to \mathbf{T} , can be determined by

$$g_{nm}^T(t) = \frac{p_\nu(t; \zeta) q_\mu(t; \zeta)}{W(q, p)}, \quad \nu = \min(n, m), \quad \mu = \max(n, m), \quad (24)$$

where the Wronskian W is defined as

$$\begin{aligned} W(q, p) &= \alpha_n [q_n(t; \zeta) p_{n-1}(t; \zeta) - q_{n-1}(t; \zeta) p_n(t; \zeta)] \\ &= b - ik = -2ik \left(\frac{\zeta}{\zeta-1} \right). \end{aligned} \quad (25)$$

The Green's matrix $\mathbf{g}_\xi = [\mathbf{h}_\xi + 2kt\mathbf{I}_\xi]^{-1}$ is related to \mathbf{g}_T : $\mathbf{g}_\xi = \mathbf{g}_T \mathbf{Z}$. Therefore, we can express the matrix \mathbf{g}_ξ elements in the form

$$g_{nm}^\xi(t) = \frac{i}{2k} \left(\frac{\zeta-1}{\zeta} \right) \frac{1}{\zeta^m} p_\nu(t; \zeta) q_\mu(t; \zeta). \quad (26)$$

From (3) and (4) it follows that the Green's matrix $\mathbf{g}_\eta = [\mathbf{h}_\eta + 2kt\mathbf{I}_\eta]^{-1}$ and \mathbf{g}_ξ are complex conjugates (for real k and t):

$$g_{nm}^\eta(t) = (g_{nm}^\xi(t))^*. \quad (27)$$

Note that there is an ambiguity in determining the matrix \mathbf{g}_ξ , since the solution $q_n(t; \zeta)$ is not unique:

$$\tilde{q}_n(t; \zeta) = q_n(t; \zeta) + y(t)p_n(t; \zeta), \quad (28)$$

where $y(t)$ is an arbitrary function of t , also satisfies (16).

b) $C \neq 0$

It is not difficult to convince oneself that the differential equation

$$\left[\hat{h}_\xi + 2kt + C\xi \right] u(\xi) = 0 \quad (29)$$

is satisfied by the function

$$u(\xi) = e^{\lambda\xi} {}_1F_1(i\tau, 1; -i\gamma\xi), \quad (30)$$

where

$$\lambda = ik \frac{\sqrt{1 - \frac{2C}{k^2}} - 1}{2}, \quad \tau = \frac{2t + i \left(1 - \sqrt{1 - \frac{2C}{k^2}} \right)}{2\sqrt{1 - \frac{2C}{k^2}}}, \quad \gamma = k\sqrt{1 - \frac{2C}{k^2}}. \quad (31)$$

Clearly, in the limit $C = 0$ the function (30) reduces to (6).

Because the operator ξ , evaluated in the basis (12), has the symmetric tridiagonal form

$$Q_{n,n'} = \begin{cases} -\frac{1}{2b} n, & n' = n - 1, \\ \frac{1}{2b} (2n + 1), & n' = n, \\ -\frac{1}{2b} (n + 1), & n' = n + 1, \end{cases} \quad (32)$$

the matrix representation of the operator $\hat{h}_\xi + 2kt + C\xi$ is also tridiagonal. Thus,

$$\mathbf{h}_\xi + 2kt\mathbf{I}_\xi + C\mathbf{Q}_\xi = \begin{pmatrix} b_0 & d_1 & & & \\ a_1 & b_1 & d_2 & & 0 \\ & a_2 & b_2 & d_3 & \\ & & a_3 & \times & \times \\ 0 & & & \times & \times & \times \end{pmatrix}, \quad (33)$$

where

$$\begin{aligned} b_n &= (b + \frac{C}{2b} + ik) + 2(b + \frac{C}{2b})n + 2kt, \\ a_n &= (b - \frac{C}{2b} - ik)n, \quad d_n = (b - \frac{C}{2b} + ik)n. \end{aligned} \quad (34)$$

Then, if we consider the coefficients of the regular solution (30) expansion in the basis set (12), we obtain that

$$s_n(t; C) = \theta^n p_n(\tau; \zeta), \quad (35)$$

where

$$\theta = \frac{b + \lambda}{b - \lambda}, \quad \zeta = \frac{b + \frac{C}{2b} - ik\sqrt{1 - \frac{2C}{k^2}}}{b + \frac{C}{2b} + ik\sqrt{1 - \frac{2C}{k^2}}}, \quad (36)$$

is the solution of the three term recurrence relation

$$a_n w_{n-1} + b_n w_n + d_{n+1} w_{n+1} = 0, \quad n \geq 1. \quad (37)$$

It is easy to verify that a second solution of Eq. (37) can be expressed in the form

$$c_n(t; C) = \theta^{n+1} q_n(\tau; \zeta). \quad (38)$$

Notice that the matrix $\mathbf{h}_\xi + 2kt\mathbf{I}_\xi + C\mathbf{Q}_\xi$ inversion procedure is simplified if we introduce the symmetric tridiagonal matrix \mathbf{T} :

$$\mathbf{T} = \mathbf{Z} [\mathbf{h}_\xi + 2kt\mathbf{I}_\xi + C\mathbf{Q}_\xi] = \begin{pmatrix} \beta_0 & \alpha_1 & & & \\ \alpha_1 & \beta_1 & \alpha_2 & & 0 \\ & \alpha_2 & \beta_2 & \alpha_3 & \\ & & \alpha_3 & \times & \times \\ 0 & & & \times & \times & \times \end{pmatrix}. \quad (39)$$

Here \mathbf{Z} is the diagonal matrix: $\mathbf{Z} = [1, \chi^{-1}, \dots, \chi^{-n}, \dots]$, $\chi = \frac{b - \frac{C}{2b} - ik}{b - \frac{C}{2b} + ik}$. The elements β_n and α_n are given by

$$\beta_n = b_n / \chi^n, \quad \alpha_n = d_n / \chi^{n-1}. \quad (40)$$

Then, it is easy to check that s_n and c_n also satisfy the recursion equation

$$\alpha_n w_{n-1} + \beta_n w_n + \alpha_{n+1} w_{n+1} = 0, \quad n \geq 1, \quad (41)$$

and the Wronskian

$$W(c, s) = \alpha_n [c_n(t; C)s_{n-1}(t; C) - c_{n-1}(t; C)s_n(t; C)] \quad (42)$$

is independent of n . Namely,

$$W(c, s) = b - \frac{C}{2b} - ik = -2ik \left(\frac{\chi}{\chi - 1} \right). \quad (43)$$

Finally, given the two linearly independent solutions s_n and c_n of Eq. (37), we can express the Green's matrix $\mathbf{g}_\xi = [\mathbf{h}_\xi + 2kt\mathbf{I}_\xi + C\mathbf{Q}_\xi]^{-1}$ elements in the form

$$g_{nm}^\xi(t; C) = \frac{s_\nu(t; C) c_\mu(t; C)}{W(c, s)} \chi^{-m} = \frac{i}{2k} \left(\frac{\chi - 1}{\chi} \right) \frac{\theta^{n+m+1}}{\chi^m} p_\nu(\tau; \zeta) q_\mu(\tau; \zeta). \quad (44)$$

Clearly, the Green's matrix $\mathbf{g}_\eta = [\mathbf{h}_\eta + 2kt\mathbf{I}_\eta + C\mathbf{Q}_\eta]^{-1}$ is obtained from \mathbf{g}_ξ by replacing $t \rightarrow -t$, $k \rightarrow -k$ [this leaves kt unchanged].

III. ORTHOGONAL POLYNOMIALS p_n

In this section we obtain the weight function with respect to which the polynomials p_n (17) are orthonormal.

The appropriate Kummer's relation [(15.3.7) in Ref. [13]] expresses the solution q_n as

$$q_n(t; \zeta) = -\frac{n!\Gamma(-n-it)}{\Gamma(1-it)} {}_2F_1(it, n+1; n+1+it; \zeta^{-1}) - \Gamma(1-it)\Gamma(it)(-\zeta)^{it} p_n(t; \zeta), \quad (45)$$

$$|\arg(-\zeta)| < \pi. \quad (46)$$

Then introduce the function

$$\rho(t; \zeta) = \frac{\Gamma(1-it)\Gamma(it)(-\zeta)^{it}}{2\pi i} \quad (47)$$

and evaluate the integral

$$\mathcal{I}(\zeta) = \int_{\mathcal{C}} dt \rho(t; \zeta), \quad (48)$$

where \mathcal{C} runs along real axis, except for an infinitesimal indentation on the underside of the pole $t = 0$; see Fig. 1.

To evaluate $\mathcal{I}(\zeta)$ for $|\zeta| > 1$, change the variable of integration to s : $t = \frac{-is}{s+i}$, transforming the path of integration into the circle \mathcal{C}_1 in the complex s -plane; see Fig. 2. Thus the interior of \mathcal{C}_1 corresponds to the upper half of the t -plane and hence the integral

$$\mathcal{I}(\zeta) = \int_{\mathcal{C}_1} \frac{ds}{(s+i)^2} \rho_1(s; \zeta), \quad (49)$$

where $\rho(t; \zeta) = \rho(\frac{-is}{s+i}; \zeta) = \rho_1(s; \zeta)$, is reduced to the sum of the residues at the poles [corresponding to $t_n = in$, $n = 0, 1, \dots$] inside the circle contour \mathcal{C}_1 :

$$\mathcal{I}(\zeta) = i \sum_{n=0}^{\infty} \frac{1}{\zeta^n}, \quad (50)$$

i. e.

$$\mathcal{I}(\zeta) = \frac{i\zeta}{1-\zeta}. \quad (51)$$

In the case $|\zeta| < 1$ it is convenient to change variables from t to s : $t = \frac{is}{s+i}$. The integration contour \mathcal{C} is transformed into the circle \mathcal{C}_2 [see Fig. 3] and the lower half of the t -plane is mapped onto the interior of \mathcal{C}_2 . Then introducing $\rho_2(s; \zeta) = \rho(\frac{is}{s+i}; \zeta) = \rho(t; \zeta)$, we obtain

$$\mathcal{I}(\zeta) = \int_{\mathcal{C}_2} \frac{-ds}{(s+i)^2} \rho_2(s; \zeta) = i \sum_{n=1}^{\infty} \zeta^n = \frac{i\zeta}{1-\zeta}. \quad (52)$$

Note that for $Im(\zeta) \neq 0$ (46) implies that

$$\rho(t; \zeta) = \begin{cases} \frac{\zeta^{it}}{1 - e^{2\pi t}}, & -\pi < \arg(\zeta) < 0, \\ \frac{\zeta^{it}}{e^{-2\pi t} - 1}, & 0 < \arg(\zeta) < \pi. \end{cases} \quad (53)$$

Using the Sokhotsky formula we obtain

$$\mathcal{I}(\zeta) = \mathcal{P} \int_{-\infty}^{\infty} dt \rho(t; \zeta) - \frac{i}{2}. \quad (54)$$

In this case (53) provides the convergence of the principal-value integral on the right hand side of (54).

Clearly, integrals

$$\int_{\mathcal{C}} dt \rho(t; \zeta) p_n(t; \zeta) p_m(t; \zeta) \quad (55)$$

can be expanded in terms of derivatives of (48). It is now easy to obtain that

$$\frac{i}{\zeta^n} \left(\frac{\zeta - 1}{\zeta} \right) \int_{\mathcal{C}} dt \rho(t; \zeta) p_n(t; \zeta) p_m(t; \zeta) = \delta_{nm}, \quad (56)$$

i. e. ρ (47) is the weight function for the orthogonal polynomials p_n (17).

For $Im(\zeta) \neq 0$ we can rewrite the orthonormality relation (56) as

$$\frac{i}{\zeta^n} \left(\frac{\zeta - 1}{\zeta} \right) \left(\mathcal{P} \int_{-\infty}^{\infty} dt \rho(t; \zeta) p_n(t; \zeta) p_m(t; \zeta) - \frac{i}{2} (-1)^{n+m} \right) = \delta_{nm}, \quad (57)$$

where the weight function ρ in the form (53) is used. Note that in the case $C = 0$ we have $\zeta = e^{i\varphi}$, $\varphi < 0$, and the weight function reduces to

$$\rho_0(t; \zeta) = \frac{e^{-\varphi t}}{1 - e^{2\pi t}}. \quad (58)$$

It may be remarked that the polynomials p_n are discrete analogues of the charge parabolic Coulomb Sturmians introduced in [14].

IV. TWO-DIMENSIONAL GREEN'S FUNCTION MATRICES

In this section we obtain a matrix representation $\mathbf{G}(t_0; C)$ of the Green's function (resolvent) of the operator (10). Formally $\mathbf{G}(t_0; C)$ is the matrix inverse of the infinite matrix

$$\mathbf{h} = \mathbf{h}_\xi \otimes \mathbf{I}_\eta + \mathbf{I}_\xi \otimes \mathbf{h}_\eta + 2kt_0 \mathbf{I}_\xi \otimes \mathbf{I}_\eta + C (\mathbf{Q}_\xi \otimes \mathbf{I}_\eta + \mathbf{I}_\xi \otimes \mathbf{Q}_\eta), \quad (59)$$

i. e.

$$\mathbf{h} \mathbf{G}(t_0; C) = \mathbf{I}_\xi \otimes \mathbf{I}_\eta. \quad (60)$$

a) $C = 0$

The elements of $\mathbf{G}(t_0) \equiv \mathbf{G}(t_0; 0)$ can be expressed as a convolution integral (see, e. g. [15])

$$G_{n_1 n_2, m_1 m_2}(t_0) = \int_{\mathcal{C}_0} dt \tilde{g}_{n_1 m_1}^\xi(t) \tilde{g}_{n_2 m_2}^\eta(t_0 - t), \quad (61)$$

where the integrand contains functions $\tilde{g}_{n m}^\xi$ and $\tilde{g}_{n m}^\eta$ that are proportional to the one-dimensional Green's function matrix elements [the non-uniqueness (28) of the solution q_n is taken into account]:

$$\tilde{g}_{n m}^\xi(t) = \frac{i}{2k} \left(\frac{\zeta - 1}{\zeta} \right) \frac{1}{\zeta^m} p_\nu(t; \zeta) [A_\xi q_\mu(t; \zeta) + x_\mu(t) p_\mu(t; \zeta)] B_m \quad (62)$$

and

$$\tilde{g}_{n m}^\eta(t) = \frac{i}{2k} (\zeta - 1) \zeta^m p_\nu(-t; \zeta^{-1}) [A_\eta q_\mu(-t; \zeta^{-1}) + y_\mu(t) p_\mu(-t; \zeta^{-1})] D_m. \quad (63)$$

Inserting (61)-(63) into (60) then gives the restriction on the factors A_ξ and A_η , functions $\{x_n(t)\}_{n=0}^\infty$ and $\{y_n(t)\}_{n=0}^\infty$, diagonal matrices $\mathbf{B} = [B_0, B_1, \dots]$ and $\mathbf{D} = [D_0, D_1, \dots]$, and the path of integration \mathcal{C}_0 , namely,

$$A_\eta D_{m_2} \int_{\mathcal{C}_0} dt \tilde{g}_{n_1 m_1}^\xi(t) = U_{n_1 m_2} \delta_{n_1 m_1}, \quad (64)$$

$$A_\xi B_{m_1} \int_{\mathcal{C}_0} dt \tilde{g}_{n_2 m_2}^\eta(t_0 - t) = V_{m_1 n_2} \delta_{n_2 m_2}, \quad (65)$$

$$U_{n_1 m_2} + V_{m_1 n_2} = 1. \quad (66)$$

If we choose the contour \mathcal{C} (see Fig. 1) as the path of integration in (61) [and (64), (65)], we can draw on the orthonormality relation (56) to determine the rest of parameters.

In this case we readily check that, in particular, the set: $A_\xi = 0$, $x_n(t) = 2\pi i \rho_0(t; \zeta)$, $A_\eta = \frac{k}{i\pi}$, $y_n(t) \equiv 0$, $B_n = D_n = 1$, satisfies the conditions (64)-(66), and hence the elements $G_{n_1 n_2, m_1 m_2}$ can be expressed in the form

$$G_{n_1 n_2, m_1 m_2}(t_0) = \frac{i}{\zeta^{m_1}} \left(\frac{\zeta-1}{\zeta} \right) \left(\mathcal{P} \int_{-\infty}^{\infty} dt \rho_0(t; \zeta) p_{n_1}(t; \zeta) p_{m_1}(t; \zeta) g_{n_2 m_2}^\eta(t_0 - t) - \frac{i}{2} (-1)^{n_1+m_1} g_{n_2 m_2}^\eta(t_0) \right). \quad (67)$$

Further, combining (56) with (A1) and (27), we can obtain another allowable sets of the parameters. For instance, $A_\xi = 1$, $x_n(t) = 2\pi i \rho_0(t; \zeta)$, $A_\eta = \frac{k}{i\pi}$, $y_n(t) \equiv 0$, $B_n = D_n = 1$.

b) $C \neq 0$

The elements of the matrix $\mathbf{G}(t_0; 0)$ may be written as the convolution integral

$$G_{n_1 n_2, m_1 m_2}(t_0; C) = \int_{\mathcal{C}_3} d\tau \tilde{g}_{n_1 m_1}^\xi(t; C) g_{n_2 m_2}^\eta(t_0 - t; C). \quad (68)$$

Here, the non-uniqueness of only \mathbf{g}_ξ is taken into consideration for simplicity, i. e. \tilde{g}_{nm}^ξ and g_{nm}^η are taken to be

$$\tilde{g}_{nm}^\xi(t; C) = \frac{i}{2k} \left(\frac{\chi-1}{\chi} \right) \frac{\theta^{n+m+1}}{\chi^m} p_\nu(\tau; \zeta) [A_\xi q_\mu(\tau; \zeta) + x_\mu(\tau) p_\mu(\tau; \zeta)] B_m, \quad (69)$$

$$g_{nm}^\eta(t_0 - t; C) = \frac{i}{2k} \frac{(\chi-1)\chi^m}{\theta^{n+m+1}} p_\nu \left(\tau - \frac{t_0}{\sqrt{1 - \frac{2C}{k^2}}}; \zeta^{-1} \right) q_\mu \left(\tau - \frac{t_0}{\sqrt{1 - \frac{2C}{k^2}}}; \zeta^{-1} \right). \quad (70)$$

The path of integration \mathcal{C}_3 in the complex τ -plane (see Fig. 4) is identical to the contour \mathcal{C} in the complex t -plane. The integration over τ (36) is performed on the assumption that τ and C are independent of one another.

Allowable parameters A_ξ , $x_n(\tau)$ and B_m can be determined by substitution (68)-(70) into (60). For instance, putting $A_\xi = 0$ and $x_n(\tau) = 2k\rho(\tau; \zeta)$, we obtain $B_m = \frac{1}{\theta^{2m+1}} \frac{(\zeta-1)}{(\chi-1)} \left(\frac{\chi}{\zeta} \right)^{m+1}$. Thus the matrix elements of the two-dimensional Green's function can be written in the form

$$G_{n_1 n_2, m_1 m_2}(t_0; C) = \frac{i}{\zeta^{m_1}} \left(\frac{\zeta-1}{\zeta} \right) \theta^{n_1-m_1} \int_{\mathcal{C}_3} d\tau \rho(\tau; \zeta) p_{n_1}(\tau; \zeta) p_{m_1}(\tau; \zeta) g_{n_2 m_2}^\eta(t_0 - t; C). \quad (71)$$

Appendix A: ONE USEFUL ORTHOGONALITY RELATION

In this appendix we derive the orthogonality relation

$$\frac{2ik}{\pi} \int_{-\infty}^{\infty} dt g_{nm}^{\xi}(t) = \delta_{nm}. \quad (\text{A1})$$

Using the integral representation [Eq. (15.3.1) in Ref. [13]] of the hypergeometric function in (19), we can rewrite the integral on the left hand side of (A1) in the form

$$\frac{1}{\pi} \left(\frac{\zeta}{\zeta-1} \right)^{\mu} \frac{1}{\zeta^m} \int_0^1 dx \frac{x^{\mu}}{\left(1 - x \frac{\zeta}{\zeta-1}\right)^{\mu+1}} \int_{-\infty}^{\infty} dt p_{\nu}(t; \zeta) e^{-it \ln(1-x)}. \quad (\text{A2})$$

Notice that the integral over t in (A2) consists of integrals

$$\int_{-\infty}^{\infty} dt (-it)^{\ell} e^{-it \ln(1-x)} = 2\pi \left[(x-1) \frac{d}{dx} \right]^{\ell} \delta(x), \quad \ell \leq \nu \leq \mu, \quad (\text{A3})$$

i. e. is expressed in terms of derivatives of the delta function. Further, introducing the function

$$y_j(x) = \frac{x^{\mu}(x-1)^j}{\left(1 - x \frac{\zeta}{\zeta-1}\right)^{\mu+1}}, \quad (\text{A4})$$

we obtain

$$\mathcal{I}_{j\ell} \equiv \int_0^1 dx y_j(x) \delta^{(\ell)}(x) = \begin{cases} \frac{1}{2} (-1)^{\ell} y_j^{(\ell)}(0) = \frac{\mu!}{2} (-1)^{\ell+j}, & \ell = \mu, \\ 0, & \ell < \mu. \end{cases} \quad (\text{A5})$$

From (A5) it follows that the integral in (A2) is nonzero if $n = m$ ($= \nu = \mu$). In this case the only contribution to the integral comes from the leading term of the polynomial $p_n(t; \zeta)$ [which is equal to $\frac{1}{n!} (\zeta-1)^n (-it)^n$]. Thus, we obtain for (A2):

$$\frac{1}{\pi} \left(\frac{\zeta}{\zeta-1} \right)^n \frac{1}{\zeta^n} \frac{1}{n!} (\zeta-1)^n 2\pi \mathcal{I}_{nn} = 1. \quad (\text{A6})$$

[1] Dz. Belkic, J. Phys. B **11**, 3529 (1978).

[2] C. R. Garibotti, J. E. Miraglia, Phys. Rev. A **21**, 572 (1980).

[3] H. Klar, Z. Phys. D **16**, 231 (1990).

- [4] G. Gasaneo, F. D. Colavecchia, S. Otranto and C. R. Garibotti, J. Phys. A **34**, 8531 (2001).
- [5] G. Gasaneo, F. D. Colavecchia, Nuclear Instruments and Methods in Physics Research B **192**, 150 (2002).
- [6] A. M. Mukhamedzhanov, A. S. Kadyrov, F. Pirlepesov, Phys. Rev. A, **73** 012713 (2006).
- [7] E. J. Heller and H. A. Yamani, Phys. Rev. A **9**, 1201 (1974).
- [8] H. A. Yamani and L. Fishman, J. Math. Phys. **16**, 4102 (1975).
- [9] A. D. Alhaidari, J. Phys. A **40**, 14843 (2007).
- [10] P. C. Ojha, J. Math. Phys. **28**, 392 (1987).
- [11] K. M. Case, M. Kac, J. Math. Phys. **14**, 594 (1973).
- [12] B. Simon, Journal of Functional Analysis **178**, 396 (2000).
- [13] M. Abramowitz and I. A. Stegun, Handbook of mathematical functions (New York: Dover), 1970.
- [14] G. Gasaneo and F. D. Colavecchia, J. Phys. A **36** 8443 (2003).
- [15] R. Shakeshaft, Phys. Rev. A, **70**, 042704 (2004).




Figure 1: The path of integration C in (48).




Figure 2: The integration contour \mathcal{C}_1 in (49).




Figure 3: The integration contour C_2 in (52).




Figure 4: The path of integration \mathcal{C}_3 of the convolution integral in (68.)