

FULL TITLE
*ASP Conference Series, Vol. **VOLUME**, **YEAR OF PUBLICATION***
NAMES OF EDITORS

Variations of the Radio Synchrotron Spectral Index in M33

F. S. Tabatabaei¹, R. Beck¹, E. Krügel¹, M. Krause¹,
 E. M. Berkhuijsen¹, K. D. Gordon², K. M. Menten¹

Abstract. We determine the variation in the nonthermal radio spectral index in the nearby spiral galaxy M33. We separate the thermal and nonthermal components of the radio continuum emission without the assumption of a constant nonthermal spectral index. Using the Spitzer FIR data at 70 and 160 μm and a standard dust model, we de-redden the $\text{H}\alpha$ emission. The extinction-corrected $\text{H}\alpha$ emission serves as a template for the thermal free-free radio emission. Subtracting from the observed 3.6 and 20cm emission (Effelsberg and the VLA) this free-free emission, we obtain maps of the nonthermal intensity and spectral index.

1. Introduction

The problem of separating the two components of the radio continuum emission, free-free (thermal) and synchrotron (nonthermal) emission, dates back to the beginnings of radio astronomy. The usually applied technique is based on the assumption of a constant nonthermal spectral index. Although this assumption may be reasonable for global studies, it does not lead to a correct thermal/nonthermal distribution in detailed studies, and it is not possible to investigate the origin and energy-loss processes of the electron component of cosmic rays (CRs).

2. Separation of Thermal and Nonthermal Emission from M33

The dust color temperature obtained between MIPS 70 μm and 160 μm (Tabatabaei et al. 2007a) shows variations between 19 K and 28 K with a most probable value at 21.5 K. Using the 160 μm column density and the dust temperature maps, we mapped the dust optical depth at 160 μm and then at $\text{H}\alpha$ wavelength ($\tau_{\text{H}\alpha}$). In the extended central region and in the two main arms IN and IS, $\tau_{\text{H}\alpha} > 0.5$ and between 0.2 and 0.4 in other arms. We found a radial gradient for the mean extinction in rings given by: $\tau_{\text{H}\alpha}(\text{R}) = (-0.009 \pm 0.002) \text{R} + (0.24 \pm 0.03)$. Assuming a homogeneous distribution of the emitting and attenuating medium in the line of sight (for ~ 0.4 kpc spatial resolution), we de-reddened the observed $\text{H}\alpha$ emission (Tabatabaei et al. 2007a). We found that less than 25% of the total $\text{H}\alpha$ emission is obscured by dust in M33. The intrinsic $\text{H}\alpha$ emission mea-

¹Max-Planck Institut für Radioastronomie, Auf dem Hügel 69, 53121 Bonn, Germany

²Space Telescope Science Institute, Baltimore, MD 21218




Figure 1. Mean nonthermal spectral index in rings of 0.5 kpc width in the galactic plane versus galactocentric radius. The total (observed) spectral index is also shown for comparison. The minima at $3 < R < 4$ kpc is due to the HII complex NGC604.

sure converted to the thermal radio emission (assuming recombination case-B, e.g. Dickinson et al. 2003). Subtracting the thermal emission from the observed 3.6 cm and 20 cm emission (Tabatabaei et al., 2007b), the nonthermal maps were derived. We obtained a total thermal fraction of $\simeq 17\%$ at 20 cm and $\simeq 50\%$ at 3.6 cm for galactocentric radii $R < 7.5$ kpc. The nonthermal emission is more smoothly distributed than the thermal emission. Furthermore, the structure of the nonthermal emission is smoother at 20 cm than at 3.6 cm, indicating a stronger diffuse component of the synchrotron emission (in the form of a diffuse disk or halo) at 20 cm (see also Tabatabaei et al. 2007c).

3. Distribution of the Nonthermal Spectral Index

Using the nonthermal maps at 20 and 3.6 cm, the nonthermal spectral index map was obtained. In the star forming regions, the nonthermal spectrum is relatively flat with an average value of 0.6 ± 0.1 , the typical spectral index of supernova remnants, but it increases to 1.2 ± 0.2 in the interarm regions and outer parts of the galaxy. This indicates energy losses of the relativistic electrons while they diffuse away from their origin in starforming regions towards the interarm regions and the outer parts of the galaxy. For the first time, a nonthermal spectral index map can be used to achieve more realistic models for the propagation of CR electrons. Figure. 1 shows that the total radio emission is mostly nonthermal at $R > 4.5$ kpc in M33, where the spectral index is dominated by synchrotron and inverse Compton loss.

References

- Dickinson, C., Davis, R. D., & Davis, R. J. 2003, MNRAS, 341, 369
 Tabatabaei, F. S., Beck, R., Krause, M., et al. 2007a, A&A, 466, 509
 Tabatabaei, F. S., Krause, M., & Beck, R. 2007b, A&A, 472, 785
 Tabatabaei, F. S., Beck, R., Krügler, E., et al. 2007c, A&A, 475, 133