

# Q-SYSTEMS AS CLUSTER ALGEBRAS II: CARTAN MATRIX OF FINITE TYPE AND THE POLYNOMIAL PROPERTY

P. DI FRANCESCO AND R. KEDEM

ABSTRACT. We define the cluster algebra associated with the  $Q$ -system for the Kirillov-Reshetikhin characters of the quantum affine algebra  $U_q(\widehat{\mathfrak{g}})$  for any simple Lie algebra  $\mathfrak{g}$ , generalizing the simply-laced case treated in [Kedem 2007]. We describe some special properties of this cluster algebra, and explain its relation to the deformed  $Q$ -systems which appeared on our proof of the combinatorial-KR conjecture. We prove that the polynomiality of the cluster variables in terms of the “initial cluster seeds”, including solutions of the  $Q$ -system, is a consequence of the Laurent phenomenon and the boundary conditions.

## 1. INTRODUCTION

The  $Q$ -system is a recursion relation for the characters of certain finite-dimensional representations of the quantum affine algebra  $U_q(\widehat{\mathfrak{g}})$  or the Yangian  $Y(\mathfrak{g})$ , where  $\mathfrak{g}$  is a simple Lie algebra.

$Q$ -systems were introduced in [12] for the classical algebras. They were later generalized by [7] for the exceptional algebras and later to more complicated situations, such as twisted quantum affine algebras [8] and double affine algebras [9].

The special modules related to  $Q$ -systems are called Kirillov-Reshetikhin modules. The fact that their characters satisfy the  $Q$ -system was proved by Kirillov and Reshetikhin [12] for  $\mathfrak{g} = A_r$ , by Nakajima [14] for the simply-laced algebras and by Hernandez [10, 9] in further generality.

Cluster algebras were introduced by Fomin and Zelevinsky [3] in 2000, and are a very general algebraic tool which has since been applied in various algebraic, combinatorial and geometric contexts. In particular, they have been used to study  $Y$ -systems [5], which are related to  $Q$ -systems in the sense that both can be derived from  $T$ -systems [13]. The  $T$ -systems are a consequence of the fusion relations for Yangian or quantum algebra modules.

The form of the  $Q$ -system suggests that it should be possible to recast it as part of a cluster algebra. The first step in this reformulation, for the case where  $\mathfrak{g}$  is simply-laced, was derived in [11]. In the current article, we give a generalization of this case to non-simply laced  $\mathfrak{g}$ . In this formulation, the  $Q$ -system appears in a very simple and easily generalizable form. We note that it does not appear to be directly related to the cluster algebra coming from the  $Y$ -system studied by Fomin and Zelevinsky [5].

In [2], we proved a combinatorial identity (“the  $M = N$ ” conjecture of [7]) which implies the proof of the combinatorial Kirillov-Reshetikhin conjecture. In our proof, we introduced what we called the deformed  $Q$ -system, depending on an increasing number of formal variables. A specialization of this system can be expressed as the  $Q$ -system with general boundary conditions (not corresponding to Kirillov-Reshetikhin characters), or as a cluster algebra. It is essential for the formulation as a cluster algebra to have this more general system.

Our proof of the  $M = N$  identity depends crucially on the fact that the KR-characters are polynomials in the fundamental KR-characters. This can be rephrased in terms of the cluster algebra, by considering a specialization of the initial cluster variables to the special point which gives the KR-characters at the other nodes of the cluster graph. We call this specialization of the initial parameters the KR point.

On the other hand, it is known that cluster variables obey the Laurent phenomenon [4]. We show that, under the specialization of the cluster variables to the KR point, this becomes what we called in [11] the strong Laurent phenomenon. That is, the cluster variables on the entire cluster graph, not just the subgraph corresponding to the  $Q$ -system, are polynomials in the initial cluster variables.

The paper is organized as follows. We recall the definition of  $Q$ -systems for any simple Lie algebra in Section 2, as well as the definition of normalized cluster algebras without coefficients. Section 3 deals with the formulation of each  $Q$ -system as a subgraph in a cluster algebra. In Section 3.1, we review the results of [11] about the formulation of  $Q$ -systems as cluster algebras for simply-laced Lie algebras. In Sections 3.2 and 3.3, we formulate the cluster algebras corresponding to the non-simply laced simple Lie algebras. In Section 4, we prove that the special boundary conditions which give solutions of the  $Q$ -system as characters of Kirillov-Reshetikhin modules imply the polynomiality of the cluster variables as functions of the seed variables at the boundary node. Section 5 is a discussion of the results.

**Acknowledgements.** We thank Bernhard Keller, Hugh Thomas and especially Sergei Fomin for their valuable input. RK thanks CEA-Saclay IPhT for their hospitality. We thank the organizers of the MSRI program on “Combinatorial Representation Theory” for their hospitality. RK is supported by NSF grant DMS-05-00759. PDF acknowledges the support of European Marie Curie Research Training Networks ENIGMA MRT-CT-2004-5652, ENRAGE MRTN-CT-2004-005616, ESF program MISGAM, and of ANR program GIMP ANR-05-BLAN-0029-01.

## 2. DEFINITIONS

### 2.1. The $Q$ -system.

2.1.1. *KR-modules.* Let  $\mathfrak{g}$  be a simple Lie algebra of rank  $r$  and let  $I_r = \{1, \dots, r\}$  be the parameterizing set for the simple roots of  $\mathfrak{g}$ . Let  $C$  be the Cartan matrix of  $\mathfrak{g}$ .

For any such algebra, there is a corresponding quantum affine algebra  $U_q(\widehat{\mathfrak{g}})$  and a Yangian algebra  $Y(\mathfrak{g})$ . In the study of the finite-dimensional modules of these algebras, there are certain special modules, called Kirillov-Reshetikhin (KR) modules [12]. These can be defined in terms of their Drinfeld polynomials [1], for example, which have a particularly simple form for these modules.

The modules are parameterized by  $\mathfrak{g}$ -highest weights of the form  $k\omega_\alpha$ , that is, a multiple of one of the fundamental weights, as well as a complex parameter  $\zeta$ . We refer to such a module by  $W_{\alpha,k}(\zeta)$  where  $\alpha \in I_r$ ,  $k \in \mathbb{Z}_+$  and  $\zeta \in \mathbb{C}^*$ . These are finite-dimensional modules, which are analogs of evaluation modules of affine algebras, which are obtained from inducing from the irreducible representation of the finite-dimensional subalgebra  $\mathfrak{g}$ , which does not depend on  $\zeta$ . Unlike such evaluation modules, however, they are not necessarily irreducible when restricted to the subalgebra  $U_q(\mathfrak{g}) \subset U_q(\widehat{\mathfrak{g}})$  or  $\mathfrak{g} \in Y(\mathfrak{g})$ , except in special cases, such as when  $\mathfrak{g} \neq A_r$ . The idea is that  $W_{\alpha,k}(\zeta)$  is the “smallest” finite-dimensional  $\mathfrak{g}[t]$ -module, with largest component  $V(k\omega_\alpha)$  (the irreducible  $\mathfrak{g}$ -module), which deforms to a Yangian module.

KR-modules have been extensively studied in recent years, and much is known about their properties. In particular, their decomposition, and the decomposition of their tensor products, into irreducible  $U_q(\mathfrak{g})$ -modules is known in terms of explicit multiplicity formulas. These formulas can be derived [12, 7, 10, 2] from the  $Q$ -system, a recursion relation for the characters

$$Q_{\alpha,k} = \text{char Res}_{U_q(\mathfrak{g})}^{U_q(\widehat{\mathfrak{g}})} W_{\alpha,k}(\zeta),$$

in addition to a certain asymptotic property (see (C) of Theorem 8.1 of [7]). We leave out the parameter  $\zeta$  in what follows, because it only affects the action of the affine part of the algebra and not  $U_q(\mathfrak{g})$  or the characters  $Q_{\alpha,k}$ .

The formulas for the decomposition coefficients and the statement that the characters of KR-modules satisfy the  $Q$ -system were both known (until they were proven) as the Kirillov-Reshetikhin conjecture.

2.1.2. *Q-systems.* We introduce the  $Q$ -system in two steps. First, let us write down a completely general relation for a family of variables

$$\{Q_{\alpha,k} \mid \alpha \in I_r, k \in \mathbb{Z}\}.$$

Let  $\mathfrak{g}$  be a simple Lie algebra with Cartan matrix  $C$ . We denote the simple roots  $\alpha$  by the corresponding integers in  $I_r = \{1, \dots, r\}$ . The  $Q$ -system associated with  $\mathfrak{g}$  is a recursion relation of the form

$$(2.1) \quad Q_{\alpha,k+1}Q_{\alpha,k-1} = Q_{\alpha,k}^2 - \prod_{\beta \sim \alpha} \mathcal{J}_k^{(\alpha,\beta)}, \quad \alpha \in I_r, k \in \mathbb{Z},$$

where  $\alpha \sim \beta$  means that  $\alpha$  is connected to  $\beta$  in the Dynkin diagram, and

$$(2.2) \quad \mathcal{J}_k^{(\alpha,\beta)} = \prod_{i=0}^{|C_{\alpha,\beta}|-1} Q_{\beta, \lfloor \frac{t_\beta k+i}{t_\alpha} \rfloor},$$

where  $\lfloor a \rfloor$  is the integer part of  $a$ . Here,  $t_\alpha$  are the integers which symmetrize the Cartan matrix. Namely,  $t_r = 2$  for  $B_r$ ,  $t_\alpha = 2$  ( $\alpha < r$ ) for  $C_r$ ,  $t_3 = t_4 = 2$  for  $F_4$  and  $t_2 = 3$  for  $G_2$ , and  $t_\alpha = 1$  otherwise.

This is a recursion relation which can be used to express any  $Q_{\alpha,k}$  in terms of  $2r$  initial data points. Usually, one chooses the initial conditions  $\{Q_{\alpha,0}, Q_{\alpha,1} \mid \alpha \in I_r\}$ . Then

$$(2.3) \quad Q_{\alpha,k+1} = \frac{Q_{\alpha,k}^2 - \prod_{\beta \sim \alpha} \mathcal{J}_k^{\alpha,\beta}}{Q_{\alpha,k-1}}, \quad k \geq 1,$$

and

$$(2.4) \quad Q_{\alpha,k-1} = \frac{Q_{\alpha,k}^2 - \prod_{\beta \sim \alpha} \mathcal{J}_k^{\alpha,\beta}}{Q_{\alpha,k+1}}, \quad k \leq 0.$$

Consider Equation (2.3) in the particular case of the initial conditions that for all  $\alpha$ ,  $Q_{\alpha,0} = 1$  and  $Q_{\alpha,1} = \text{Char}W_{\alpha,1}$ , the character of the fundamental Kirillov-Reshetikhin module, with highest  $\mathfrak{g}$ -weight  $\omega_\alpha$ . In this case, the solutions  $Q_{\alpha,k}$  of (2.3) are known to be the characters of KR-modules  $W_{\alpha,k}$  with highest  $\mathfrak{g}$ -weight  $k\omega_\alpha$  [14, 10]. The recursion relation (2.3) with these initial conditions is what is usually called the  $Q$ -system in the literature.

We do not impose these initial conditions in the definition of the cluster algebra. As we will see, this is a special (singular) point in the parameter space for this algebra. One way to see that

it is singular is that equation (2.4) cannot be used, in this case, to define all the variables  $Q_{\alpha,k}$  with  $k < 0$  because, for example,  $Q_{\alpha,-1} = 0$  in this case.

More general initial conditions, of the form

$$(2.5) \quad Q_{\alpha,0} = b_\alpha, \quad Q_{\alpha,1} = a_\alpha$$

for some formal variables  $a_\alpha, b_\alpha$  give a solution of (a special case of) the “deformed Q-system” introduced in [2]. We will explain this point in Section 5.

**2.1.3. Normalized Q-systems.** In this paper we deal with cluster algebras without coefficients. To see that it is possible to reformulate the Q-system in this way, a version of the following Lemma was given in [11]:

**Lemma 2.1.** *There exist complex numbers  $\epsilon_1, \dots, \epsilon_r$  (all of them roots of 1) such that the variables  $R_{\alpha,i} = \epsilon_\alpha Q_{\alpha,i}$  satisfy the normalized Q-system*

$$(2.6) \quad R_{\alpha,k+1}R_{\alpha,k-1} = R_{\alpha,k}^2 + \prod_{\beta \sim \alpha} \tilde{\mathcal{J}}_k^{(\alpha,\beta)},$$

where

$$\tilde{\mathcal{J}}_k^{(\alpha,\beta)} = \prod_{i=0}^{|C_{\alpha,\beta}|-1} R_{\beta, \lfloor \frac{t_\beta(k+i)}{t_\alpha} \rfloor}.$$

*Proof.* From Equation (2.2) we note that the total degree of  $Q_{\beta,j'}$  (for various  $j'$ ) in  $\mathcal{J}_k^{(\alpha,\beta)}$  is always  $|C_{\alpha,\beta}|$ , where  $C$  is the Cartan matrix of  $\mathfrak{g}$ . Therefore  $R_{\alpha,k} = \epsilon_\alpha Q_{\alpha,k}$  satisfy the following system:

$$R_{\alpha,k+1}R_{\alpha,k-1} = R_{\alpha,k}^2 - \left( \prod_{\beta} \epsilon_\beta^{C_{\alpha,\beta}} \right) \prod_{\beta \sim \alpha} \tilde{\mathcal{J}}_k^{(\alpha,\beta)},$$

where  $\tilde{\mathcal{J}}$  is just  $\mathcal{J}$  with the variables  $Q$  replaced by  $R$ .

Thus, if the variables  $\epsilon_\alpha$  satisfy

$$\prod_{\beta=1}^r \epsilon_\beta^{C_{\alpha,\beta}} = -1$$

then (2.6) is satisfied. But since  $C$  is a Cartan matrix of finite type, it is invertible, and this system has a solution. Namely if we set  $\epsilon_\beta = e^{i\pi\mu_\beta}$ , then  $\mu_\alpha = \sum_{\beta} C_{\beta,\alpha}^{-1}$ .  $\square$

In fact, we find  $\epsilon_\beta^4 = 1$  in general. We note that this choice of normalization is done in order to make contact with cluster algebras without coefficients below. We do not believe that it is essential for the final result.

We will refer to (2.6) as the normalized Q-system or simply the Q-system when it is clear that we are considering the evolution of the variables  $R$ .

**2.2. Cluster algebras.** We use here the definition of [6] to define a cluster algebra without coefficients.

**Definition 2.2.** *A cluster algebra. Fix  $n$  and define a labeled  $n$ -ary tree  $\mathbb{T}_n$ , with nodes labeled by some parameter  $t$  and  $n$  edges labeled  $1, \dots, n$  emanating from each node.*

*To each node  $t$  we associate a seed  $(\mathbf{x}, B)_t$ , where  $\mathbf{x} = (x_1, \dots, x_n)$  (the cluster variables) and  $B$  (the mutation matrix) is a skew-symmetric  $n \times n$  matrix with integer coefficients.*

If the node  $t$  is connected to the node  $t'$  by an edge labeled by  $k$ , then  $(\mathbf{x}[t'], B[t']) = \mu_k(\mathbf{x}[t], B[t])$  where we have the following relation (mutations or evolution equations along the edge  $k$ ) between the cluster at  $t$  and  $t'$ :

$$(2.7) \quad \mu_k(x_i) = \begin{cases} x_i, & i \neq k; \\ x_i^{-1}(\prod_j x_j^{[B_{jk}]_+} + \prod_j x_j^{[-B_{jk}]_+}), & i = k. \end{cases}$$

$$(2.8) \quad \mu_k(B_{ij}) = \begin{cases} -B_{ij}, & \text{if } i = k \text{ or } j = k; \\ B_{ij} + \text{sgn}(B_{ik})[B_{ik}B_{kj}]_+, & \text{otherwise.} \end{cases}$$

where  $[n]_+$  denotes the positive part of  $n$  and

$$\text{sgn}(n) = \begin{cases} 0, & n = 0; \\ 1, & n > 0; \\ -1, & n < 0. \end{cases}$$

One can check the following properties of cluster mutations:

- The mutations (2.8) preserve the skew-symmetry of  $B$ .
- If  $B_{ij} = 0$  then the mutations  $\mu_i$  and  $\mu_j$  commute.
- If  $B_{ij} = 0$  and  $k, l$  are two integers distinct from both  $i$  and  $j$ , then

$$(2.9) \quad \mu_i \circ \mu_j(B_{kl}) = B_{kl} + \text{sgn}(B_{ki})[B_{ki}B_{il}]_+ + \text{sgn}(B_{kj})[B_{kj}B_{jl}]_+.$$

This statement can be extended to any finite sequence of commuting mutations.

- Mutations are defined in such a way that  $\mu_i^2 = 1$ . Thus, cluster graphs are not oriented.

**Definition 2.3.** We define  $\mathcal{A}$  to be the family of variables  $\{R_{\alpha,i}\}_{i \in \mathbb{Z}, \alpha \in I_r}$  which satisfy the normalized  $Q$ -system (2.6).

In what follows we are interested in subgraphs of  $\mathbb{T}_n$  which encode the  $Q$ -system, that is, the cluster variables at the nodes are elements of  $\mathcal{A}$  and the mutations between nodes are  $Q$ -system evolutions.

More precisely, we will define the quotient graphs  $\mathcal{G}_{\mathfrak{g}}$ , by identifying nodes corresponding to the same cluster variables and mutation matrices. These are infinite graphs, but they have certain periodicity properties and are therefore easy to describe explicitly.

Our starting goal is to find a graph  $\mathcal{G}_{\mathfrak{g}}$  for each algebra, and the corresponding cluster variables, which encode the full  $Q$ -system, in the following sense: The union over all nodes of  $\mathcal{G}_{\mathfrak{g}}$  of the cluster variables is the full set  $\mathcal{A}$ .

Of course once the seed  $(\mathbf{x}, B)$  is given at any node  $t$ , then the cluster algebra is defined, that is, the cluster at any other node is defined via the evolution equations. Therefore, our definition gives a cluster algebra on the full tree  $\mathbb{T}_n$ . We will have something to say about special properties of the cluster variables on  $\mathbb{T}_n$  in Section 4, but thus far they have no representation-theoretical interpretation to our knowledge.

This definition of  $\mathcal{G}_{\mathfrak{g}}$  allows us to encode the  $Q$ -system inside a cluster algebra. The graphs  $\mathcal{G}_{\mathfrak{g}}$  which we describe are usually subgraphs of a larger component of  $\mathbb{T}_n$  with the same property, and therefore our definition is not unique. There may be other nodes of  $\mathbb{T}_n$  with cluster variables in  $\mathcal{A}$  which are not in  $\mathcal{G}_{\mathfrak{g}}$ . This happens if the rank of  $\mathfrak{g}$  is greater than two. We will give the example of  $A_3$  to make this point clearer in the discussion.

### 3. THE $Q$ -SYSTEMS AS CLUSTER ALGEBRAS FOR SIMPLE $\mathfrak{g}$

Let  $C$  be a Cartan matrix of a simple Lie algebra  $\mathfrak{g}$  of rank  $r$ , and let  $B$  be the  $2r \times 2r$  matrix defined in the block form:

$$(3.1) \quad B = \begin{pmatrix} A & -C^t \\ C & 0 \end{pmatrix},$$

where  $A = C^t - C$ . This is a skew-symmetric matrix of integers.

For each  $k \in \mathbb{Z}$ , define  $\mathbf{x}[k] = (x_1, \dots, x_{2r})$ , where

$$(3.2) \quad x_\alpha = \begin{cases} R_{\alpha, 2t_\alpha k}, & \alpha \leq r \\ R_{\alpha-r, 2t_\alpha k+1}, & r < \alpha \leq 2r \end{cases}$$

Here,  $t_\alpha = 2$  for the short roots of  $B_r$ ,  $C_r$  and  $F_4$ ,  $t_\alpha = 3$  for the short root of  $G_2$ , and  $t_\alpha = 1$  otherwise.

In our cluster algebra graph, nodes labeled by  $k \in \mathbb{Z}$  have associated with them cluster  $(\mathbf{x}[k], B)$ . We will prove the following theorem:

**Theorem 3.1.** *There exists a cluster graph  $\mathcal{G}_{\mathfrak{g}}$ , which includes all nodes labeled by  $k \in \mathbb{Z}_+$ , with corresponding cluster variables  $\mathbf{x}[k]$  as in (3.2) and  $B$  as in (3.1), such that all mutations in the graph are  $Q$ -system evolutions. Moreover, the set  $\mathcal{A}$  is equal to the union of the cluster variables over all nodes in  $\mathcal{G}_{\mathfrak{g}}$ .*

We will describe such a graph for each algebra  $\mathfrak{g}$  below.

In all cases considered below, if the mutations  $\{\mu_{i_1}, \dots, \mu_{i_n}\}$  commute starting from node  $t$  (that is, if  $B_{i_k, i_j} = 0$  for all  $k, j$  at the node  $t$ ), then we may act with these mutations in any order, as long as they are distinct, and nodes corresponding to acting with a certain set of mutations, regardless of order, can be identified with each other in the quotient graph  $\mathcal{G}_{\mathfrak{g}}$  (this is similar in spirit to the bipartite graph of [6]). These mutations give rise to an  $n$ -simplex (e.g. a cube in the case  $n = 3$ ) in  $\mathcal{G}_{\mathfrak{g}}$ .

We will in general describe only the cluster variables reached via certain compound mutations  $\mu^{(i)} = \mu_{i_1} \circ \dots \circ \mu_{i_n}$ , where the corresponding mutations commute. The cluster graph  $\mathcal{G}_{\mathfrak{g}}$  is recovered by replacing a compound mutation with the corresponding  $n$ -simplex.

**3.1. The simply laced case.** This case was considered in [11]. We review the results here in order to introduce the notations.

If  $\mathfrak{g}$  is simply-laced, the Cartan matrix is symmetric,  $C^t = C$ , and therefore the matrix  $A$  vanishes and  $B$  is block-diagonal:

$$(3.3) \quad B = \begin{pmatrix} 0 & -C \\ C & 0 \end{pmatrix}.$$

The graph  $\mathcal{G}_{\mathfrak{g}}$  has a particularly simple form in this case. It includes the special nodes  $k$  ( $k \in \mathbb{Z}_+$ ) with cluster variables

$$(3.4) \quad \mathbf{x}[k] = (R_{1,2k}, \dots, R_{r,2k}, R_{1,2k+1}, \dots, R_{r,2k+1}).$$

In addition, it includes the special nodes  $k'$ , which are reached from  $k$  by the compound mutation  $\mu^{(1)} = \mu_1 \circ \dots \circ \mu_r$ . These commute due to the block-diagonal structure of  $B$ . The following was proved in [11]:

**Theorem 3.2.** *For  $k \in \mathbb{Z}_+$ , Let  $\mathbf{x}[k]$  be as in (3.4) and  $B$  be the matrix in (3.3). Then*

- (1) At the node  $k$ , the mutations  $\mu_1, \dots, \mu_r$  commute among themselves, and evolution according to these mutations corresponds to the  $Q$ -system evolutions  $R_{\alpha, 2k} \mapsto R_{\alpha, 2k+2}$ ,  $\alpha \in I_r$ .
- (2) At the same node, the mutations  $\mu_{r+1}, \dots, \mu_{2r}$  also commute among themselves, and applying them to the variables  $\mathbf{x}[k]$  corresponds to the  $Q$ -system evolution  $R_{\alpha, 2k+1} \mapsto R_{\alpha, 2k-1}$ ,  $\alpha \in I_r$ .
- (3) Define the compound mutation

$$\mu^{(I_r)} = \mu_1 \circ \dots \circ \mu_r$$

and denote the node reached by this compound mutation acting on the cluster variables at  $k$  by  $k'$ . Then the mutation matrix  $B' = \mu^{(I_r)}(B)$  is

$$B' = -B = \begin{pmatrix} 0 & C \\ -C & 0 \end{pmatrix}.$$

and

$$\mu^{(I_r)}(\mathbf{x}[k]) = (R_{1, 2k+2}, \dots, R_{r, 2k+2}, R_{1, 2k+1}, \dots, R_{r, 2k+1}) = \mathbf{x}[k'].$$

- (4) At the point  $k'$ , the mutations  $\mu_{r+1}, \dots, \mu_{2r}$  commute among themselves, the evolution according to these mutations corresponds to the  $Q$ -system evolution  $R_{\alpha, 2k+1} \mapsto R_{\alpha, 2k+3}$ .
- (5) The compound mutation


$$\mu^{(I_r')} = \mu_{r+1} \circ \dots \circ \mu_{2r}$$

maps the seed at  $k'$  to the seed at  $k+1$  with cluster variables  $\mathbf{x}[k+1]$  and mutation matrix  $B$ .

This theorem is proved by direct calculation.


Let us illustrate some examples of graphs which correspond to  $\mathcal{G}_{\mathfrak{g}}$ .

**Example 3.3.** The graph  $\mathcal{G}_{\mathfrak{sl}_2}$  is linear:



The cluster variables at the primed and unprimed nodes are described by  $\mathbf{x}[k] = (R_{1, 2k}, R_{1, 2k+1})$  and  $\mathbf{x}[k'] = (R_{1, 2k+2}, R_{1, 2k+1})$  of Theorem 3.2. The numbers above each edge correspond to the mutations  $\mu_1$  and  $\mu_2$ . This is the full tree  $\mathbb{T}_2$ .


**Example 3.4.** The graph  $\mathcal{G}_{\mathfrak{sl}_3}$  is slightly more interesting:



The cluster algebra has four variables, and four possible mutations, which are the labels along the edges. The special nodes  $k$  and their primed versions appear on the center line of the graph. The other nodes have intermediate cluster variables.

The full cluster algebra graph of  $\mathfrak{sl}_3$  would include, for example, an edge labeled 4 emanating from the node  $\mu_1(k)$ . However the cluster mutation in this direction is not a  $Q$ -system evolution.

**Example 3.5.** The graph  $\mathcal{G}_{\mathfrak{sl}_4}$  described in Theorem 3.2 is as follows:




The graphs  $\mathcal{G}_{\mathfrak{g}}$  which we describe are not, in general, the complete subgraph of  $\mathbb{T}_n$  corresponding to  $Q$ -system evolutions, except when  $\text{rank}(\mathfrak{g}) \leq 2$ . We chose the graphs  $\mathcal{G}_{\mathfrak{g}}$  so that the union of all cluster variables at their nodes corresponds to the complete set of normalized characters of the  $KR$ -modules. These graphs are not unique, and their extensions are of interest also (see section 5.2).

Clearly, the cluster graphs  $\mathcal{G}_{\mathfrak{g}}$  are difficult to draw as the rank becomes high. Therefore, in treating the higher-rank algebras, it is useful to adopt a pictorial notation which is similar in spirit to the bipartite graph notation of [6].


First, we define the notion of a compound mutation to be a product of  $n$  distinct commuting mutations. Let  $I = \{i_1, \dots, i_n\}$  be the index set of these mutations, and define  $\mu^{(I)} = \mu_{i_1} \circ \dots \circ \mu_{i_n}$ . The order of mutations is irrelevant by definition. We may also use the notation  $\mu^{(I)} = \prod_{i \in I} \mu_i$ . Obviously, since  $\mu_i^2 = 1$ , we preserve the property that  $(\mu^{(I)})^2 = 1$  for the product of commuting notations.

The edge:



corresponds to the compound mutation  $t' = \mu^{(I)}(t)$ . In the graph  $\mathcal{G}_{\mathfrak{g}}$ , this corresponds to an  $n$ -simplex with node  $t$  at one end and node  $t'$  at the other. It is a diamond in the case for  $\mathcal{G}_{\mathfrak{sl}_3}$  or a cube in the case of  $\mathcal{G}_{\mathfrak{sl}_4}$ .

Thus the graph  $\mathcal{G}_{\mathfrak{g}}$  for  $\mathfrak{g}$  simply-laced with the sequence of nodes and compound mutations as follows:






where  $\Pi = I_r$  and  $\Pi' = \{r + 1, \dots, 2r\}$ . The cluster variables and mutation matrices at the nodes  $k, k'$  are given by Theorem 3.2.

**3.2. The  $Q$ -system for the algebras  $B_r, C_r, F_4$ .** In the case of non-simply laced algebras, we describe the graph  $\mathcal{G}_{\mathfrak{g}}$  for algebras of types  $B_r, C_r, F_4$  separately from  $G_2$ .

Let  $\mathfrak{g}$  be a simple algebra of type  $B_r, C_4$  or  $F_4$ . In the Dynkin diagram of  $\mathfrak{g}$ , let  $\ell$  be the long root connected to the short root  $s$ :



In each case, in the normalized  $Q$ -system (2.6) we have

$$(3.5) \quad \tilde{\mathcal{J}}_{2k+1}^{(s,\ell)} = R_{\ell,k} R_{\ell,k+1} = R_{\ell,k}^{[-C_{s\ell}/2]_+} R_{\ell,k+1}^{[-C_{s\ell}/2]_+};$$

$$(3.6) \quad \tilde{\mathcal{J}}_{2k}^{(s,\ell)} = R_{\ell,k}^2 = R_{\ell,k}^{[-C_{s\ell}]_+};$$

$$(3.7) \quad \tilde{\mathcal{J}}_k^{(\ell,s)} = R_{s,2k} = R_{s,2k}^{[-C_{\ell,s}]_+}.$$


In case  $(\alpha, \beta)$  are not the pair  $(\ell, s)$  or  $(s, \ell)$  then

$$(3.8) \quad \tilde{\mathcal{J}}_k^{(\alpha,\beta)} = R_{\beta,k}^{[-C_{\beta,\alpha}]_+}.$$

Define the sets

- $\Pi' = I'_r = \{r + 1, \dots, 2r\}$ ;
- $\Pi_{<} :$  The subset in  $I_r$  corresponding to the short roots of  $\mathfrak{g}$ ;
- $\Pi_{>} :$  The subset in  $I_r$  corresponding to the long roots of  $\mathfrak{g}$ ;
- $\Pi'_{<} = \{\alpha + r \mid \alpha \in \Pi_{<}\}$ ;
- $\Pi'_{>} = \{\alpha + r \mid \alpha \in \Pi_{>}\}$ ;

**Theorem 3.6.** *The graph  $\mathcal{G}_{\mathfrak{g}}$  is obtained from the sequence*



Here, the cluster at the node labeled  $k$  is  $\mathbf{x}[k]$  as in (3.2) with mutation matrix  $B$  as in (3.1). All the compound mutations are products of mutually commuting mutations, and the cluster variables at each node are in  $\mathcal{A}$ .

To prove this Theorem, we need only list the cluster variables  $(\mathbf{x}[k^{(i)}], B^{(i)})$  at  $k^{(i)}$  – at intermediate nodes, the cluster variables are a mixture of cluster variables from  $k^{(i-1)}$  and  $k^{(i)}$ . Note that in this subsection, we read the superscript of  $k$  modulo 5, so  $k^{(0)} := k$  etc.

We then observe that the relevant columns of  $B^{(i)}$  describe  $Q$ -system evolutions when applied to the relevant entries of  $\mathbf{x}[k^{(i)}]$ . We use the fact that when  $\mu_i$  and  $\mu_j$  commute, the  $j$ th column of  $\mu_i(B)$  is equal to the  $j$ th column of  $B$ .

We recall that at the node  $k$ , we have  $\mathbf{x}[k] = (x_1, \dots, x_{2r})$ , where

$$x_\alpha = \begin{cases} R_{\alpha, 2t_\alpha k}, & \alpha \in I_r; \\ R_{\alpha, 2t_\alpha k+1}, & \text{otherwise.} \end{cases}$$

Moreover, according to (3.1), at the node  $k$  we have

$$B = \begin{pmatrix} A & -C^t \\ C & 0 \end{pmatrix}.$$

Here, the submatrix  $A$  is no longer the zero matrix. Instead, it has zeros everywhere except for two entries,  $A_{\ell, s} = -1$  and  $A_{s, \ell} = 1$ . Thus, the mutations  $\{\mu_\alpha \mid \alpha \in \Pi_{<}\}$  commute among themselves, as do the all mutations  $\{\mu_i \mid i \in \Pi'\}$ .

It is clear from the form of the  $Q$ -system (see Equations (3.5) and (3.8)) that the mutations  $\{\mu_\alpha \mid \alpha \in \Pi_{<}\}$  describe the evolutions  $\mu_\alpha : R_{\alpha, 2k} \mapsto R_{\alpha, 2k+2}$  ( $\alpha \in \Pi_{<}$ ). This is the same evolution as in the simply-laced case if  $\alpha \neq s$ , and for  $\alpha = s$ , it follows from (3.5).

Thus, edges labeled by  $\alpha \in \Pi_{<}$  connected to node  $k$  lead to nodes in which all the cluster variables are in  $\mathcal{A}$ . The compound mutation  $\mu^{(\Pi_{<})}$  leads to the node  $k^{(1)}$ .

(We also note that the mutations columns  $r+1, \dots, 2r$  describe the  $Q$ -system evolutions  $R_{\alpha, 2k+1} \mapsto R_{\alpha, 2k-1}$ , from Equations (3.6), (3.7) and (3.8); However due to the periodic structure of the graph, we do not need to consider this evolution separately).

**Lemma 3.7.** (1) *The cluster variables at the node  $k^{(1)}$  are  $\mathbf{x}[k^{(1)}] = (x_1, \dots, x_{2r}) = \mu^{(\Pi_{<})}(\mathbf{x}[k])$  with*

$$x_\alpha = \begin{cases} R_{\alpha, 4k+2} & \text{if } \alpha \text{ is a short root;} \\ R_{\alpha, 2k} & \text{if } \alpha \text{ is a long root;} \\ R_{\alpha, 2t_\alpha k+1} & \text{if } \alpha > r; \end{cases}$$

(2) *The mutation matrix  $B^{(1)}$  at  $k^{(1)}$  has the following form:*

$$B^{(1)} = \left( \begin{array}{c|c} -A & -D^t \\ \hline D & 2A \end{array} \right),$$

where

$$D_{ij} = \begin{cases} C_{ij}, & \text{if } i \text{ and } j \text{ are both long roots;} \\ -C_{ij}, & \text{if } j \text{ is short;} \\ 0 & \text{otherwise.} \end{cases}$$

*Proof.* The first statement follows from the discussion above.

The second statement involves several cases.

- Because the mutations  $\mu_\alpha$  with  $\alpha < r$  and  $\alpha \in \Pi_{<}$  mutually commute, their effect on the columns and rows of  $B$  indexed by  $\{\alpha : \alpha \in \Pi_{<}\}$  is only to change their sign. Thus,

$$B_{i, \alpha}^{(1)} = -B_{\alpha, i}^{(1)} = -B_{i, \alpha}, \quad \alpha \in \Pi_{<}.$$

This accounts for the appearance of  $-A$  in the upper left hand corner of  $B^{(1)}$ , as well as the columns of the matrix  $D$  which correspond to short roots.

- Let  $\beta \in \Pi_{>}$  be a long root and consider the change column  $\beta$  of  $B$ . If  $i \notin \Pi_{<}$ , then

$$B_{i,\beta}^{(1)} = B_{i,\beta} + \sum_{\alpha \in \Pi_{<}} \operatorname{sgn}(B_{i,\alpha}) [B_{i,\alpha} B_{\alpha,\beta}]_+.$$

But  $B_{\alpha,\beta} = 0$  unless  $(\alpha, \beta) = (s, \ell)$ . In that case,  $B_{s,\ell} = 1$ . We have that  $B_{i,s} > 0$  if and only if  $i = s + r$ , in which case it is equal to 2. Thus, we have

$$B_{s+r,\ell}^{(1)} = 0,$$

and all other elements in the long-root columns of the lower left hand corner of  $B^{(1)}$  are unchanged. This gives the matrix  $D$  in this quadrant, and  $-D^t$  in the upper right block of  $B^{(1)}$  since mutations preserve skew-symmetry.

- Finally, elements in the lower right quadrant mutate as follows:

$$B_{i+r,j+r}^{(1)} = \sum_{\alpha \in \Pi_{<}} \operatorname{sgn}(B_{i+r,\alpha}) [B_{i+r,\alpha} B_{\alpha,j+r}]_+ = 2A.$$

□

By examination of the matrix  $B^{(1)}$  we conclude that at the node  $k^{(1)}$ , the mutations  $\{\mu_\alpha : \alpha \in \Pi_{>}\}$  all commute among themselves.

**Lemma 3.8.** (1) *The cluster variables  $\mathbf{x}[k^{(2)}] = (x_1, \dots, x_{2r}) = \mu^{(\Pi_{>})}(\mathbf{x}[k^{(1)}])$  are*

$$x_\alpha = \begin{cases} R_{2t_\alpha k+2} & \alpha \leq r; \\ R_{2t_\alpha k+1} & \alpha > r. \end{cases}$$

(2) *The mutation matrix  $B^{(2)}$  has the form*

$$\begin{pmatrix} A & C - A \\ -C^t - A & 2A \end{pmatrix}.$$

*Proof.* Again, by inspection, the columns corresponding to long roots in the matrix  $B^{(1)}$  correspond to the evolution of the  $Q$ -system according to Equations (3.7) and (3.8). This proves the first statement in the Lemma.

We look at the evolution of the matrix  $B^{(1)}$  under  $\mu^{(\Pi_{>})}$ .

- If  $\alpha \in \Pi_{>}$  then  $B_{i,\alpha}^{(2)} = -B_{i,\alpha}^{(1)}$ . This accounts for the appearance of  $A$  in the upper left-hand block of  $B^{(2)}$  and for the columns corresponding to the long roots in the lower left-hand quadrant of  $B^{(2)}$ .
- If  $\alpha \in \Pi_{<}$  and  $i > r$  then

$$B_{i,\alpha}^{(2)} = B_{i,\alpha}^{(1)} + \sum_{\beta \in \Pi_{>}} \operatorname{sgn}(B_{i,\beta}^{(1)}) [B_{i,\beta}^{(1)} B_{\beta,\alpha}^{(1)}]_+$$

The summation has a contribution only when  $(\beta, \alpha) = (\ell, s)$  in which case  $B_{\ell,s}^{(1)} = 1$ . Then the only non-vanishing contribution occurs if  $i = \beta + r$ , in which case  $B_{\beta+r,\beta}^{(1)} = 2$  and  $B_{\ell+r,s} = 3$ . This gives the matrix in the lower left quadrant of  $B^{(2)}$ .

- Finally we have

$$B_{i+r,j+r}^{(2)} = 2A_{i,j} + \sum_{\alpha \in \Pi_{>}} \operatorname{sgn}(B_{i+r,\alpha}^{(1)}) [B_{i+r,\alpha}^{(1)} B_{\alpha,j+r}^{(1)}]_+ = 2A_{ij}.$$

□

At this point, it is evident that the mutations  $\{\mu_{\alpha+r} : \alpha \in \Pi_{<}\}$  commute among themselves, due to the form of the matrix  $2A$  appearing in the lower right quadrant of  $B^{(2)}$ . Moreover, the corresponding columns of  $B^{(2)}$  give the  $Q$ -system evolution for  $R_{\alpha,2k+1} \mapsto R_{\alpha,2k+3}$  if  $\alpha$  is short. As a result we have

**Lemma 3.9.** (1) *The cluster variables at  $k^{(3)}$  are  $\mu^{(\Pi_{<})}(\mathbf{x}[k^{(2)}]) = \mathbf{x}[k^{(3)}] = (x_1, \dots, x_i)$  where*

$$x_{\alpha+r} = R_{\alpha,4k+3}, \quad \alpha \in \Pi_{<},$$

*and the other cluster variables are unchanged from  $\mathbf{x}[k^{(2)}]$ .*

(2) *The mutation matrix  $B^{(3)} = -B^{(1)}$*

The proof is by a calculation analogous to the previous two lemmas.

The columns  $\{\alpha : \alpha \in \Pi_{<}\}$  of  $-B^{(1)}$  describe the mutations corresponding to the evolutions  $R_{\alpha,4k+2} \rightarrow R_{\alpha,4k+4}$  as they are identical to the corresponding columns in the matrix  $B$ . The mutations corresponding to the short roots again commute. We have the compound evolution  $\mu^{(\Pi_{<})}$  which brings us to the node  $k^{(4)}$ , with

**Lemma 3.10.** (1) *The cluster variables at the point  $k^{(4)}$  are  $\mu^{(\Pi_{<})}(\mathbf{x}[k^{(3)}]) = \mathbf{x}[k^{(4)}] = (x_1, \dots, x_{2r})$ , with*

$$x_{\alpha} = \begin{cases} R_{\alpha,2t_{\alpha}(k+2)}, & \alpha \leq r \\ R_{\alpha,2k+1}, & \alpha - r \in \Pi_{>} \\ R_{\alpha,2k+3}, & \alpha - r \in \Pi_{<}. \end{cases}$$

(2) *The mutation matrix at the node  $k^{(4)}$  is  $B^{(4)} = -B$ .*

Again, the proof is analogous to that of the previous lemmas.

Finally we note that at the node  $k^{(4)}$ , not only do the mutations  $\{\mu_{\alpha+r} : \alpha \in \Pi\}$  all commute among themselves, the corresponding columns of  $B^{(4)}$  describe the evolution of the  $Q$ -system for any  $R_{\alpha,i}$  with  $i$  odd. Therefore the compound evolution  $\mu^{(\Pi')}$  makes sense at this node. The node  $k^{(5)}$  is reached from node  $k^{(4)}$  by acting with  $\mu^{(\Pi')}$ . In fact, we find that the node  $k^{(5)}$  is the node  $k+1$  in  $\mathcal{G}_{\mathfrak{g}}$ :

**Lemma 3.11.** (1) *The cluster variables  $\mathbf{x}[k^{(5)}] = \mu^{(\Pi')}(\mathbf{x}[k^{(4)}]) = \mathbf{x}[k+1]$ , where  $\mathbf{x}[k]$  for any  $k \in \mathbb{Z}$  was defined in equation (3.2).*

(2) *The mutation matrix at the node  $k^{(5)}$  is  $B$ .*


The proof is by direct calculation. Thus, there is a periodicity to the graph  $\mathcal{G}_{\mathfrak{g}}$ , where the matrix  $B^{(i)}$  is repeated along the “band” corresponding to the subgraph every fifth step.

**3.3. The  $Q$ -system for  $G_2$  as a cluster algebra.** In this case the normalized  $Q$ -system (2.6) has terms  $\tilde{\mathcal{T}}_{\alpha,k}$  as follows:


$$\begin{aligned} \tilde{\mathcal{T}}_k^{(1,2)} &= R_{2,3k} \\ \tilde{\mathcal{T}}_{3k}^{(2,1)} &= R_{1,k}^3 \\ \tilde{\mathcal{T}}_{3k+1}^{(2,1)} &= R_{1,k}^2 R_{1,k+1} \\ \tilde{\mathcal{T}}_{3k+2}^{(2,1)} &= R_{1,k} R_{1,k+1}^2. \end{aligned}$$

Again we will describe a graph  $\mathcal{G}_{G_2}$ , a quotient graph of a subgraph of  $\mathbb{T}_4$ , which contains at its nodes cluster variables with all the normalized characters  $\{R_{\alpha,k} : \alpha \in I_2, k \in \mathbb{Z}_+\}$ .

The graph  $\mathcal{G}_{G_2}$  contains all nodes labeled by  $k \in \mathbb{Z}_+$ , and also the nodes  $k^{(i)}$  with  $i = 1, \dots, 6$ . At these nodes we have cluster variables  $\mathbf{x}[k^{(k)}]$  and mutation matrices  $B^{(i)}$ . We will describe the clusters at those nodes explicitly. As in the other cases, we choose to evolve along nodes of the graph when the corresponding mutations describe one of the  $Q$ -system equations. The graph  $\mathcal{G}_{G_2}$  is the following graph:



With the notation of the previous subsection, we can also depict this graph as follows:



The following can be easily verified by explicit calculation:

**Theorem 3.12.** *The mutations along the graph  $\mathcal{G}_{G_2}$  describe the evolution of the  $Q$ -system of  $G_2$ , the cluster variables at each point  $k^{(i)}$  are defined as follows:*

$i$	$\mathbf{x}[k^{(i)}]$	$B^{(i)}$
0	$(R_{1,2k}, R_{2,6k}, R_{1,2k+1}, R_{2,6k+1})$	$\begin{pmatrix} 0 & -2 & -2 & 3 \\ 2 & 0 & 1 & -2 \\ 2 & -1 & 0 & 0 \\ -3 & 2 & 0 & 0 \end{pmatrix}$
1	$(R_{1,2k}, R_{2,6k+2}, R_{1,2k+1}, R_{2,6k+1})$	$\begin{pmatrix} 0 & 2 & -2 & -1 \\ -2 & 0 & -1 & 2 \\ 2 & 1 & 0 & -2 \\ 1 & -2 & 2 & 0 \end{pmatrix}$
2	$(R_{1,2k}, R_{2,6k+2}, R_{1,2k+1}, R_{2,6k+3})$	$\begin{pmatrix} 0 & -C \\ C^t & -A \end{pmatrix}$
3	$(R_{1,2k+2}, R_{2,6k+4}, R_{1,2k+1}, R_{2,6k+3})$	$-B^{(2)}$
4	$(R_{1,2k+2}, R_{2,6k+4}, R_{1,2k+1}, R_{2,6k+5})$	$-B^{(1)}$
5	$(R_{1,2k+2}, R_{2,6k+6}, R_{1,2k+1}, R_{2,6k+5})$	$-B$
6	$\mathbf{x}[k+1] = (R_{1,2k+2}, R_{2,6k+6}, R_{1,2k+3}, R_{2,6k+7})$	$B$

Note that  $k^{(0)} := k$  and  $k^{(6)} = k+1$  as the cluster variables at that node correspond to the point  $k+1$ .

To prove this theorem, one simply compares the  $Q$ -system evolution along in the direction of the label of the edge emanating from the node  $k^{(i)}$ , and the corresponding columns of the mutation matrices at the points  $k^{(i)}$ . That is, if the an edge labeled  $n$  emanates from  $k^{(i)}$  in the graph  $\mathcal{G}_{G_2}$ , then the  $n$ th variable of  $x[k^{(i)}]$  satisfies the  $Q$ -system described by column  $n$  of  $B^{(i)}$ .

The “diamonds” in  $\mathcal{G}_{G_2}$  occur when two mutations commute. Since this is a finite system it can be done by a direct calculation.

#### 4. POLYNOMIALITY PHENOMENON

We now consider the specialization of the cluster algebra to the point  $\{Q_{\alpha,0} = 1\}_\alpha$  (so that  $R_{\alpha,0} = \epsilon_\alpha$ ), which we call the Kirillov-Reshetikhin point, because solutions to the  $Q$ -system with this boundary condition, for  $m > 0$ , are characters of Kirillov-Reshetikhin modules.

Note that at the KR point,  $Q_{\alpha,-1} = R_{\alpha,-1} = 0$  for all  $\alpha$ . However due to the Laurent phenomenon, which states, among other things, that all cluster variables are Laurent polynomials in  $R_{\alpha,0}$  and  $R_{\alpha,1}$ , all cluster variables are well-defined at this point. They have no singularity since at the KR point  $R_{\alpha,0} \neq 0$ .

In our proof of the combinatorial Kirillov-Reshetikhin conjecture [2], a crucial ingredient was the following Lemma:

**Lemma 4.1.** *For any algebra  $\mathfrak{g}$ , any solution to the KR  $Q$ -system  $Q_{\alpha,m}$  is a polynomial in the  $r$  variables  $\{Q_{\alpha,1}\}$ .*

This follows from the fact that KR-modules are all in the Grothendieck group of the trivial and fundamental KR-modules.

One of the reasons for recasting the  $Q$ -system as a cluster algebra is the availability of the Laurent phenomenon theorem. It implies a purely algebraic proof for the polynomiality lemma. (The idea for this proof is due to Sergei Fomin.) We will explain the general situation which gives rise to this property in the  $Q$ -system.

Consider a cluster algebra with a special node, which we call 0 or the origin, with cluster variables

$$\mathbf{x} = (a_1, \dots, a_n; b_1, \dots, b_m) = (\mathbf{a}, \mathbf{b})$$

and mutation matrix  $B$  such that

$$B_{ij} = 0 \quad \text{if } i, j < n.$$

We note that, as a result of this assumption about the form of the  $B$ -matrix, we have

$$\mu_i(a_i) = \frac{N_i(\mathbf{b})}{a_i}, \quad 1 \leq i \leq n,$$

where

$$N_i(\mathbf{b}) = \prod_j b_j^{[B_{j+n,i}]_+} + \prod_j b_j^{[-B_{j+n,i}]_+}$$

is a function of  $\{b_1, \dots, b_m\}$  only, because  $B_{j,i} = 0$  if  $i, j \leq n$ .

We assume that there is a special point in the parameter space  $\mathbf{b}$ , which we call the KR point in analogy with the  $Q$ -system situation, such that for all  $i \leq n$ ,  $\mu_i(a_i) = 0$ . Evaluation of  $\mathbf{b}$  at the KR point means the evaluation at the point  $N_i(\mathbf{b}) = 0$  ( $i \leq n$ ). This point depends only on  $\mathbf{b}$ , not on  $\mathbf{a}$ , which are left as formal variables.

We will show that for a cluster variable  $y(\mathbf{a}, \mathbf{b})$  at any node of the cluster tree  $\mathbb{T}_{m+n}$ , considered as a function of  $(\mathbf{a}, \mathbf{b})$  (a Laurent polynomial, due to the Laurent property), terms with singularities at any  $a_i = 0$  vanish at the KR point.

We use the vector notation, so that if  $\mathbf{b} = (b_1, \dots, b_i)$  and  $\mathbf{m} = (m_1, \dots, m_i)$  then  $\mathbf{b}^{\mathbf{m}} = \prod_j b_j^{m_j}$ .

**Lemma 4.2.** *Let*

$$y(\mathbf{a}, \mathbf{b}) = \mathbf{b}^{-\mathbf{m}} P(\mathbf{a}, \mathbf{b})$$

where  $P(\mathbf{a}, \mathbf{b})$  is a polynomial in  $\mathbf{b}$  but may have some singularities in  $\{a_i\}$ . That is,

$$P(\mathbf{a}, \mathbf{b}) = \sum_{\mathbf{n} \in \mathbb{Z}^r} C_{\mathbf{n}}(\mathbf{b}) \mathbf{a}^{\mathbf{n}},$$

with only finitely many non-zero terms, and where  $C_{\mathbf{n}}(\mathbf{b})$  are polynomials in the variables  $\mathbf{b}$ . Then  $C_{\mathbf{n}}(\mathbf{b})$  is divisible by

$$\prod_{j: n_j < 0} N_j(\mathbf{b})^{-n_j}.$$

*Proof.* By change of variables to  $\mu_i(a_i) = a'_i$  for all  $i \leq n$ , we have that

$$P(\mathbf{a}, \mathbf{b}) = \sum_{\mathbf{n} \in \mathbb{Z}^r} C_{\mathbf{n}}(\mathbf{b}) \prod_{j=1}^n (a'_j)^{-n_j} N_j(\mathbf{b})^{n_j}.$$

Any terms with  $n_j < 0$  have a positive power of  $N_j(\mathbf{b})$  in the denominator. These must cancel with factors in  $C_{\mathbf{n}}(\mathbf{b})$ , for each  $\mathbf{n}$  which has negative components, because the Laurent property states that a cluster variable cannot be a rational function in the variables  $\mathbf{b}$ . Furthermore, there can be no cancellations between terms with different  $\mathbf{n}$ , since the  $a'_j$  are independent variables.

Thus,  $C_{\mathbf{n}}(\mathbf{b})$  is divisible by  $N_j(\mathbf{b})^{-n_j}$  for each negative  $n_j$ . Therefore, it vanishes at the KR point, where  $N_j(\mathbf{b}) = 0$  for each  $j$ . These are all the terms which have poles at  $a_j = 0$ . This proves the Lemma.  $\square$

Thus we have what we called “strong Laurent phenomenon” in [11] or the polynomiality property:

**Corollary 4.3.** *With the cluster algebra as defined above, at the evaluation of  $\mathbf{b}$  at a point such that  $\mu_i(a_i) = N_i(\mathbf{b}) = 0$ , all cluster variables are polynomials in the variables  $\{a_i\}$ .*

Lemma 4.1 is a special case of this statement (with a rearrangement of the indices). We have  $m = n = r$ ,  $a_i = Q_{i,1}$  and  $b_i = Q_{i,0}$ . The indices of the matrix  $B$  are accordingly rearranged. Note that, in all the algebras introduced so far, we had the property that  $B_{ij} = 0$  if  $r < i, j \leq 2r$ , which is the analog of the condition on  $B$  above.

We conclude that the Laurent phenomenon is responsible for the polynomiality property of the characters  $Q_{\alpha,i}$  in terms of  $Q_{\alpha,1}$  for the KR  $Q$ -system. This property is quite general and is not related to the appearance of a Cartan matrix in the mutation matrix, only on the block structure of the mutation matrix, and the specific nature of the KR point in the parameter space.

Moreover, we can conclude much more.

**Corollary 4.4.** *Any cluster variable at any node of the full cluster graph  $\mathbb{T}_{2r}$  extending  $\mathcal{G}_{\mathfrak{g}}$  as defined in Section 3 is a polynomial in  $R_{\alpha,1}$  after specialization of the parameters to the KR point,  $R_{\alpha,0} = \epsilon_{\alpha}$ .*

We do not, at this point, have a representation-theoretical interpretation for this phenomenon.

## 5. DISCUSSION

**5.1. Deformed Q-systems and the cluster algebra.** In [2], we have defined deformed Q-systems depending on infinite families of parameters  $\mathbf{u} = \{u_\alpha, u_{\alpha,1}, u_{\alpha,2}, \dots\}_{\alpha \in I_r}$ , as well as the extra parameters  $\mathbf{a} = \{a_1, a_2, \dots\}$  in the non-simply-laced case. The deformed Q-system is defined as follows:

$$(5.1) \quad u_{\alpha,n} \mathcal{Q}_{\alpha,n+1} \mathcal{Q}_{\alpha,n-1} = \mathcal{Q}_{\alpha,n}^2 - \prod_{\beta \sim \alpha} \mathcal{U}_n^{(\alpha,\beta)}, \quad \alpha \in I_r, \quad n \geq 1,$$

where

$$\mathcal{U}_n^{(\alpha,\beta)} = \prod_{i=0}^{|C_{\alpha,\beta}|-1} \mathcal{Q}_{\beta, \lfloor \frac{t_\beta n + i}{t_\alpha} \rfloor}, \quad (\alpha, \beta) \neq (s, \ell)$$

and

$$\mathcal{U}_n^{(\alpha,\beta)} = a_n^{t_\alpha(1 + \lfloor \frac{n}{t_\alpha} \rfloor) - n} \prod_{i=0}^{|C_{\alpha,\beta}|-1} \mathcal{Q}_{\beta, \lfloor \frac{n+i}{t_\alpha} \rfloor} \quad (\alpha, \beta) = (s, \ell).$$

The deformed Q-system (5.1) is subject to the initial conditions that  $\mathcal{Q}_{\alpha,0} = 1$  and  $\mathcal{Q}_{\alpha,1} = u_\alpha^{-1}$ , for all  $\alpha \in I_r$ .

The deformed variables  $\mathcal{Q}(\mathbf{u}, \mathbf{a})$  were introduced as a tool for proving the  $M = N$  identity of [7], yielding fermionic expressions for the tensor product multiplicities of KR-modules.

We may view the cluster variables  $R_{\alpha,k}$  of the cluster graph  $\mathcal{G}_{\mathfrak{g}}$  of Section 3 as specializations of the deformed characters  $\mathcal{Q}(\mathbf{u}, \mathbf{a})$  (after normalization). This amounts to relaxing the initial condition on the Q-system, which is the specialization to the KR point.

Let  $\varphi$  denote the evaluation

$$\varphi(a_i) = 1, \quad \varphi(u_{\alpha,i}) = 1, \quad i \geq 2,$$

and

$$\varphi(a_1) = u_{s,1}$$

where  $s$  the unique short root connected to a long root.

The evaluated solutions of the deformed Q-system

$$\overline{\mathcal{Q}}(\{u_\alpha, u_{\alpha,1}\}_{\alpha \in I_r}) := \varphi(\mathcal{Q}(\mathbf{u}, \mathbf{a}))$$

obey a system of equations identical to the Q-system (2.1) for all  $k \geq 1$ , except for the difference that  $\overline{\mathcal{Q}}_{\alpha,0}$  always comes with a prefactor  $u_{\alpha,1}$ .

Therefore, we may absorb the parameter  $u_{\alpha,1}$  into the definition of  $\overline{\mathcal{Q}}_{\alpha,0}$ , namely by introducing

$$Q_{\alpha,k}(\{u_{\beta,1}, u_{\beta}^{-1}\}_{\beta \in I_r}) := u_{\alpha,1}^{\delta_{k,0}} \overline{\mathcal{Q}}_{\alpha,k}(\{u_{\beta}, u_{\beta,1}\}_{\beta \in I_r}).$$

Then  $Q_{\alpha,k}$  obey the Q-system (2.1) for  $k \geq 1$ , with the initial conditions replaced by  $Q_{\alpha,0} = u_{\alpha,1}$  and  $Q_{\alpha,1} = u_\alpha^{-1}$ . This is identical to (2.5).

Upon the renormalization of the variables as in Lemma 2.1, we deduce that the specialized deformed KR characters are related to the variables  $R_{\alpha,k}$  via:

$$(5.2) \quad R_{\alpha,k} = \epsilon_\alpha u_{\alpha,1}^{\delta_{k,0}} \varphi(Q_{\alpha,k}(\mathbf{u}, \mathbf{a})).$$

We note that without the evaluation  $\varphi$ , the variables  $\mathcal{Q}_{\alpha,m}(\mathbf{u}, \mathbf{a})$  are not obviously part of a cluster algebra. For example, they do not have the Laurent property.



In [2], we derived and used a crucial “substitution invariance” property of the deformed KR characters  $\mathcal{Q}$  which, after the specialization  $\varphi$ , reads as follows for the  $Q$ ’s:

$$(5.3) \quad Q_{\alpha, k+t_{\alpha}j}(\{Q_{\beta,0}, Q_{\beta,1}\}) = Q_{\alpha,k}(\{Q_{\beta,t_{\beta}j}, Q_{\beta,t_{\beta}j+1}\}).$$

We showed that this condition is equivalent to the  $Q$ -system, with appropriate initial conditions.

When rephrased in terms of  $R$ ’s, with  $j = 2m$  even, this is nothing but the property of translational invariance of the cluster graph  $\mathcal{G}_{\mathfrak{g}}$ . Namely, the expression of cluster variables at any node  $n$  of  $\mathcal{G}_{\mathfrak{g}}$  in terms of that at cluster variables at the node 0 is the same as that of the cluster variables at the node  $n + m$  in terms of the cluster variables at the node  $m$ . This is obvious from the structure of the graph.

Equation (5.3) is actually slightly more general, as it also includes translations by an odd integer  $j$ . For instance, the case  $j = 1$  corresponds in the cluster formulation to the  $k \rightarrow k'$  (“half-” translation in the simply-laced case, and to  $k \rightarrow k^{(3)}$  in the  $G_2$  case. At the node  $k'$  (or  $k^{(3)}$  for  $G_2$ ), the mutation matrix is indeed  $JBJ$ , where  $J = \begin{pmatrix} 0 & I_{r \times r} \\ I_{r \times r} & 0 \end{pmatrix}$ , hence the nodes  $k$  and  $k'$  (or  $k$  and  $k^{(3)}$  for  $G_2$ ) are equivalent upon interchanging the even and odd cluster variables  $\mathbf{x} \rightarrow J\mathbf{x}$ , and the half-translation invariance property (5.3) for odd  $j$  follows.


**5.2. Extended graphs.** As remarked above, generally there are larger subgraphs of  $\mathbb{T}_{2r}$  which correspond to  $Q$ -system evolutions.

**5.2.1. Graphs corresponding to  $Q$ -system evolutions.** In Example 3.5 for  $A_3$ , we did not describe edges which do not correspond to  $Q$ -system evolutions. But in addition this graph does not describe all edges which are  $Q$ -system evolutions. For example,  $\mu_6$  acting on  $\mu_2 \circ \mu_3(\mathbf{x}[k], B)$  is a  $Q$ -system evolution, which results in the cluster variable

$$\mathbf{x} = (R_{1,2k}, R_{2,2k+2}, R_{3,2k+2}, R_{1,2k+1}, R_{2,2k+1}, R_{3,2k+3}).$$

Similarly,  $\mu_4$  acting on  $\mu_1 \circ \mu_2(\mathbf{x}[k], B)$  is a  $Q$ -system evolution.

We can describe the complete graph corresponding to  $Q$ -system evolutions in the case of  $A_3$ :



**5.3. Generalizations and open questions.** As mentioned in the introduction, there are generalizations of  $Q$ -systems to Cartan matrices corresponding to affine algebras [9]. Moreover there are generalizations of  $Q$ -systems to twisted quantum affine algebras. It is reasonable to expect that these can be recast in the cluster algebra language, although Lemma 2.1 does not, in general, work for affine Cartan matrices.

$Q$ -systems are obtained from  $T$ -systems by taking the combinatorial limit, which amounts to ignoring the dependence on the spectral parameter [12, 13]. The  $T$ -system is a discrete Hirota equation in the case of  $A_n$ , and in general has an integrability property. As a result,  $Q$ -systems inherit some remarkable properties, such as the existence of conserved quantities and associated linear systems. We will address this property in a future publication.

An interesting question arises from the  $M = N$  identity which was proved in [2]. This is a combinatorial identity which relates a restricted sum over products of binomial coefficients to an unrestricted one (see [7] for the original conjecture).

For example, the  $M = N$  identity has the following form for  $A_1$ :

$$\sum_{\mathbf{m} \in \mathbb{Z}_+^k : p_i \geq 0} \prod_i \binom{m_i + p_i}{m_i} = \sum_{\mathbf{m} \in \mathbb{Z}_+^k} \prod_i \binom{m_i + p_i}{m_i}$$

where for a choice of non-negative integers  $\mathbf{n} \in \mathbb{Z}_+^k$ ,  $p_i = \sum_j \min(i, j)(n_j - 2m_j)$  and the sum is taken over  $\mathbf{m} \in \mathbb{Z}_+^k$  such that  $\sum_j j(n_j - 2m_j) = l \in \mathbb{Z}_+$ .

The restricted sum on the left hand side is manifestly positive, whereas the unrestricted sum on the right is an alternating sum. In [2], we have recast this identity in terms of generating functions and their power series expansions in order to prove the original conjecture of [7]. The  $M = N$  identity is a property of the following generating function:

$$Z_{l; \mathbf{n}}^{(k)}(\mathbf{u}) = \frac{Q_1(\mathbf{u})Q_k(\mathbf{u})^{l+1}}{Q_{k+1}(\mathbf{u})^{l+1}} \prod_{i=1}^k \frac{Q_i(\mathbf{u})^{n_i}}{u_i},$$

where  $\mathcal{Q}_m(\mathbf{u})$  are solutions of the deformed  $Q$ -system. The  $M = N$  identity is the following statement: The constant term of  $Z$  in  $u$ , after evaluation at  $u_1 = \cdots = u_k = 1$ , is equal to the constant term in  $u$  of the power series part of  $Z$  in each of the  $u_i$ , after evaluation at the same point.

Our proof relies only on the polynomiality property of the solutions to the KR  $Q$ -system. It would be interesting to understand this identity more fully in the context of the properties of the associated cluster algebras.

## REFERENCES

- [1] Vyjayanthi Chari and Andrew Pressley, *A Guide to Quantum Groups*, Cambridge University Press, Cambridge, 1995.
- [2] Philippe Di Francesco and Rinat Kedem, *Proof of the combinatorial Kirillov-Reshetikhin conjecture* arXiv:0710.4415 [math:QA] (IMRN, to appear).
- [3] Sergey Fomin and Andrei Zelevinsky, *Cluster algebras. I. Foundations*, J. Amer. Math. Soc. **15** (2002), no. 2, 497–529.
- [4] ———, *The Laurent phenomenon*, Adv. in Appl. Math. **28** (2002), no. 2, 119–144.
- [5] ———, *Y-systems and generalized associahedra*, Ann. of Math. (2) **158** (2003), no. 3, 977–1018.
- [6] ———, *Cluster algebras. IV. Coefficients*, Compos. Math. **143** (2007), no. 1, 112–164.
- [7] G. Hatayama, A. Kuniba, M. Okado, T. Takagi, and Y. Yamada, *Remarks on fermionic formula*, Recent developments in quantum affine algebras and related topics (Raleigh, NC, 1998), Contemp. Math., vol. 248, Amer. Math. Soc., Providence, RI, 1999, pp. 243–291.
- [8] Goro Hatayama, Atsuo Kuniba, Masato Okado, Taichiro Takagi, and Zengo Tsuboi, *Paths, crystals and fermionic formulae*, MathPhys Odyssey, 2001, Prog. Math. Phys., vol. 23, Birkhäuser Boston, Boston, MA, 2002, pp. 205–272.
- [9] David Hernandez, *Kirillov-reshetikhin conjecture: The general case*. arXiv:0704.2838 [math:QA].
- [10] ———, *The Kirillov-Reshetikhin conjecture and solutions of T-systems*, J. Reine Angew. Math. **596** (2006), 63–87.
- [11] Rinat Kedem, *Q-systems as cluster algebras*, arXiv:0712.2695 [math:RT].
- [12] A. N. Kirillov and N. Yu. Reshetikhin, *Formulas for the multiplicities of the occurrence of irreducible components in the tensor product of representations of simple Lie algebras*, Zap. Nauchn. Sem. S.-Peterburg. Otdel. Mat. Inst. Steklov. (POMI) **205** (1993), no. Differentsialnaya Geom. Gruppy Li i Mekh. 13, 30–37, 179.
- [13] Atsuo Kuniba, Tomoki Nakanishi, and Junji Suzuki, *Functional relations in solvable lattice models. I. Functional relations and representation theory*, Internat. J. Modern Phys. A **9** (1994), no. 30, 5215–5266.
- [14] Hiraku Nakajima, *t-analogs of q-characters of Kirillov-Reshetikhin modules of quantum affine algebras*, Represent. Theory **7** (2003), 259–274 (electronic).

PDF: INSTITUT DE PHYSIQUE THÉORIQUE DE SACLAY, CEA/DSM/IPhT, URA 2306 DU CNRS, F-91191 GIF SUR YVETTE CEDEX FRANCE. E-MAIL: PHILIPPE.DI-FRANCESCO@CEA.FR

RK: DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ILLINOIS, 1409 WEST GREEN STREET, URBANA, IL 61821. E-MAIL: RINAT@UIUC.EDU