

SMOOTH STRUCTURES AND EINSTEIN METRICS ON $\mathbb{C}\mathbb{P}^2 \# 5, 6, 7 \overline{\mathbb{C}\mathbb{P}^2}$

RAREȘ RĂSDEACONU AND IOANA ȘUVAINA

ABSTRACT. We show that each of the topological 4-manifolds $\mathbb{C}\mathbb{P}^2 \# k \overline{\mathbb{C}\mathbb{P}^2}$, for $k = 6, 7$ admits a smooth structure which has an Einstein metric of scalar curvature $s > 0$, a smooth structure which has an Einstein metric with $s < 0$ and infinitely many non-diffeomorphic smooth structures which do not admit Einstein metrics. We show that there are infinitely many manifolds homeomorphic non-diffeomorphic to $\mathbb{C}\mathbb{P}^2 \# 5 \overline{\mathbb{C}\mathbb{P}^2}$ which don't admit an Einstein metric. We also exhibit new examples of manifolds carrying Einstein metrics of both positive and negative scalar curvature. The main ingredients are recent constructions of exotic symplectic or complex manifolds with small topological numbers.

1. INTRODUCTION

Recently, new methods were developed [Pa05, ABP07] to construct symplectic 4-manifolds with small topology and exotic smooth structures. Moreover, the method proposed by J. Park [Pa05] was later refined [LePa07, PPS07] to produce interesting examples of minimal complex surfaces of general type. In this paper we show how we can use these constructions in regard to the existence or non-existence of Einstein metrics.

LeBrun and Catanese [LeCa98] showed that there exist homeomorphic, but non-diffeomorphic manifolds such that one of them admits an Einstein metric of positive sign while the other admits an Einstein metric of negative sign. The manifold which has positive scalar curvature Einstein metric is $\mathbb{C}\mathbb{P}^2 \# 8 \overline{\mathbb{C}\mathbb{P}^2}$, while the second manifold is a deformation of the Barlow surface. The key ingredient in their proof was to show that the Barlow surface, a simply connected minimal surface of general type with $K^2 = 1$ can be deformed to a surface with ample canonical bundle. In particular, they showed that there exists a simply connected complex surface of general type with $p_g = 0, K^2 = 1$ and having ample canonical bundle. The ampleness of the canonical bundle ensures the existence of a Kähler-Einstein metric of negative scalar curvature.

Based on ideas from [Pa05], Lee and Park [LePa07] and more recently Park, Park, and Shin [PPS07] constructed new examples of simply connected, minimal surfaces of general type with $p_g = 0, K^2 = 1, 2$ or 3 . We show that their examples satisfy the ampleness condition:

Theorem 1.1. *There exist simply connected complex surfaces of general type, with $p_g = 0, K^2 = 2$ or 3 , and ample canonical bundle.*

In his paper, Park [Pa05], used his examples and LeBrun's Obstruction Theorem to exhibit a manifold homeomorphic, non-diffeomorphic to $\mathbb{C}\mathbb{P}^2 \# 8 \overline{\mathbb{C}\mathbb{P}^2}$ which does not admit an Einstein metric. We extend his results to $\mathbb{C}\mathbb{P}^2 \# k \overline{\mathbb{C}\mathbb{P}^2}$, $k = 6, 7$, and we also exhibit an infinite family of homeomorphic manifolds for which no Einstein metric exists.

Date: November 7, 2018.

2000 Mathematics Subject Classification. Primary 70G55; Secondary 32J15.

Key words and phrases. rational blowing down, symplectic 4-manifolds, Einstein metrics.

Theorem 1.2. *Each of the topological 4-manifolds $\mathbb{C}\mathbb{P}^2 \# k\overline{\mathbb{C}\mathbb{P}^2}$, for $k = 6, 7, 8$ admits a smooth structure which has an Einstein metric of scalar curvature $s > 0$, a smooth structure which has an Einstein metric with $s < 0$ and infinitely many non-diffeomorphic smooth structures which do not admit Einstein metrics.*

We can also prove non-existence results for an even smaller topological manifold:

Proposition 1.3. *The canonical smooth structure on $M = \mathbb{C}\mathbb{P}^2 \# 5\overline{\mathbb{C}\mathbb{P}^2}$ admits an Einstein metric of positive scalar curvature, but M also supports infinitely many exotic smooth structures which don't have an Einstein metric.*

It is a well known fact that in dimensions 2 and 3 the sign of the Einstein metric is a topological invariant. This led Besse [Bes87, p.19] to consider the conjecture that no smooth compact n -manifold can admit Einstein metrics with different scalar curvature signs. As we see in the previous theorems, in dimension 4, there are homeomorphic, non-diffeomorphic 4-manifolds which admit Einstein metrics of opposite sign. For higher dimensions, as far as the authors know, LeBrun and Catanese constructed the only known counterexamples to Besse's conjecture. These examples are in dimensions $4k$, where $k \geq 2$. In these dimensions, we are able to provide many new examples with the same property:

Proposition 1.4. *Let $N_1 = \mathbb{C}\mathbb{P}^2 \# 8\overline{\mathbb{C}\mathbb{P}^2}$, $N_2 = \mathbb{C}\mathbb{P}^2 \# 7\overline{\mathbb{C}\mathbb{P}^2}$, $N_3 = \mathbb{C}\mathbb{P}^2 \# 6\overline{\mathbb{C}\mathbb{P}^2}$. Then the manifold N obtained by taking the k -fold products, $k \geq 2$, of arbitrary N_1, N_2 , or N_3 , admits two Einstein metrics g_1, g_2 such that the signs of the scalar curvature are $s_{g_1} = -1, s_{g_2} = +1$. Moreover, these metrics are Kähler-Einstein with respect to two distinct complex structures J_1, J_2 .*

As a consequence of the Kählerian property of the metrics and of our method of construction, we have the following in peculiar fact: their corresponding volumes are equal $Vol_{g_1}(N) = Vol_{g_2}(N)$.

Remark 1.5. *Soon after this paper was finished, Jongil Park and his collaborators found [PPS08] an analogous example of a complex surface of general type homeomorphic to $\mathbb{C}\mathbb{P}^2 \# 5\overline{\mathbb{C}\mathbb{P}^2}$. Using this construction the results of Theorems 1.1, 1.2 and Proposition 1.4 can be extended for $\mathbb{C}\mathbb{P}^2 \# 5\overline{\mathbb{C}\mathbb{P}^2}$.*

The paper is organized as follows: in Section 2 we discuss the ampleness of the canonical line bundles of some interesting examples of complex surfaces and the relation to the existence of Einstein metrics, while in Section 3 we introduce the Seiberg-Witten invariants and use them to get results on non-existence of Einstein metrics. In Section 4 we treat the higher dimensional case.

2. AMPLENESS OF THE (ANTI)CANONICAL BUNDLE: EXISTENCE OF EINSTEIN METRICS

In general the existence or non-existence of Einstein metrics on a given manifold is hard to prove. In the case $c_1(M) > 0$, in the unobstructed situations, i.e. for certain complex surfaces whose underlying differential structure is $\mathbb{C}\mathbb{P}^2 \# k\overline{\mathbb{C}\mathbb{P}^2}$, $3 \leq k \leq 8$, the existence of Kähler-Einstein metrics was proved by Siu [Siu88], or Tian and Yau [TiYa87]. A complete solution to the existence was given by Tian [Ti90]:

Theorem 2.1 (Tian). *A compact complex surface (M^4, J) admits a compatible Kähler-Einstein metric with $s > 0$ if and only if its anti-canonical line bundle K_M^{-1} is ample and its Lie algebra of holomorphic vector fields is reductive.*

In the case when $c_1(M) < 0$, a criterion for the existence of a Kähler-Einstein metric was independently found by Aubin [Aub76] and Yau [Yau77]:

Theorem 2.2 (Aubin-Yau). *A compact complex manifold (M^4, J) admits a compatible Kähler-Einstein metric with $s < 0$ if and only if its canonical line bundle K_M is ample. When such a metric exists, it is unique, up to an overall multiplicative constant.*

In this section, we discuss some examples of smooth structures on $\mathbb{C}P^2 \# k \overline{\mathbb{C}P^2}$, $6 \leq k \leq 8$. For each such k , the natural smooth structure as the complex projective plane blown-up at k generic points, can be endowed [Ti90] by a Kähler-Einstein metric of positive scalar curvature. On the other hand, Lee and Park [LePa07] and more recently Park, Park and Shin [PPS07] constructed new exotic smooth structures on $\mathbb{C}P^2 \# k \overline{\mathbb{C}P^2}$, for any $6 \leq k \leq 8$. Moreover, these admit complex structures yielding interesting examples of minimal surfaces of general type. We are going to prove that the canonical bundle of these surfaces is ample, and so they admit Kähler-Einstein metrics of negative scalar curvature.

The examples constructed in these papers are very similar. In this article we are going to treat just the more complicated example [PPS07] of a simply connected minimal surface of general type with $p_g = 0$ and $K^2 = 3$. The other examples follow similarly. We begin by a brief description of the example of a minimal surface of general type with $p_g = 0$ and $K^2 = 3$, referring frequently to [PPS07].

We start with an appropriate pencil of plane elliptic curves, and we resolve the base locus to obtain a rational elliptic surface Y with the following set of singular fibers: one I_8 -fiber, one reducible I_2 -fiber and two nodal fibers. Moreover, this rational elliptic surface admits 4 sections. Now, Y is blown-up 21 times at well chosen points to obtain a new rational surface Z . These points can be chosen in such a way [PPS07] that Z contains four particular disjoint chains of rational curves consisting of proper transforms of the singular fibers, proper transform of some sections and some of the exceptional divisors introduced in the blowing-up process. According to [PPS07], these chains are $G = \sum_{i=1}^{10} G_i$, $H = \sum_{i=1}^7 H_i$, $I = \sum_{i=1}^6 I_i$ and a chain of length one denoted by A . These chains are represented by continuous lines in Figure 1.




Figure 1: Manifold $Z = \mathbb{C}P^2 \# 30 \overline{\mathbb{C}P^2}$

They all satisfy Artin's contractibility criterion [KoMo98]. By contracting them, we obtain a projective surface X with 4 singularities, each admitting 1-parameter \mathbb{Q} -Gorenstein smoothings. Let $f : Z \rightarrow X$ be the contraction map. Now, it is proved [PPS07] that

the projective surface X admits a \mathbb{Q} –Gorenstein smoothing, which is a smooth, simply-connected, minimal surface of general type with $K_X^2 = 3$. We denote by $F_i, i = 1, \dots, 7$ the seven smooth rational curves of self-intersection -1 , which are marked by discontinuous lines in Figure 1 and by F_8 the remaining exceptional divisor from the last singular fiber, drawn as a dotted line. It is easy to see that the Poincaré duals of the curves $F_i, i = 1, \dots, 7$, together with all of the irreducible components of our 4 chains G, H, I and A form a basis of $H^2(Z, \mathbb{Q})$, which is torsion-free and 31-dimensional.

First, we prove the ampleness of the canonical divisor of the singular surface X . Note that since X has only 1-parameter \mathbb{Q} –Gorenstein smoothing singularities, its canonical divisor is in fact \mathbb{Q} –Cartier.

A common feature of all of the examples appearing in [LePa07, PPS07] is that the pullback of the canonical divisor of the singular variety X to its minimal resolution is effective. For instance, in the example described above, we can write:

$$f^*K_X \equiv_{\mathbb{Q}} \sum_{i=1}^8 a_i F_i + \sum_{i=1}^{10} b_i G_i + \sum_{i=1}^7 c_i H_i + \sum_{i=1}^6 d_i I_i + eA, \quad (2.1)$$

In [PPS07], the coefficients are explicitly given, but for our purpose it is important to know only that a_i, b_i, c_i, d_i and e are strictly positive rational numbers. For convenience, let $Exc(f) = \sum G_i + \sum H_i + \sum I_i + A$ denote the exceptional divisor of f .

By a direct computation, in [PPS07] it is proved that $K_X^2 = 3$ and the nefness of K_X . Suppose now that K_X fails to be ample. Since K_X is already nef, according to the Nakai-Moishezon criterion [KoMo98], there exists an irreducible curve $C \subset X$ such that $(K_X \cdot C) = 0$.

The total transform of C in Z is

$$f^*C \equiv_{\mathbb{Q}} C' + \sum_{i=1}^{10} x_i G_i + \sum_{i=1}^7 y_i H_i + \sum_{i=1}^6 z_i I_i + tA, \quad (2.2)$$

where C' is the strict transform and x_i, y_i, z_i, t are non-negative rational numbers. We should note here that C' is not numerically equivalent to 0. To give an immediate proof for this assertion, note that $(C \cdot H) > 0$ for any ample line bundle H on the projective surface X . But, if $C' \equiv_{\mathbb{Q}} 0$ we would have

$$0 = (C' \cdot f^*H) = (f^*C \cdot f^*H) = (C \cdot H),$$

and this is a contradiction.

Now, a straightforward computation gives:

$$\begin{aligned} (K_X \cdot C) &= (f^*K_X \cdot f^*C) = (f^*K_X \cdot C') \\ &= \sum_{i=1}^8 a_i (F_i \cdot C') + \sum_{i=1}^{10} b_i (G_i \cdot C') + \sum_{i=1}^7 c_i (H_i \cdot C') \\ &\quad + \sum_{i=1}^6 d_i (I_i \cdot C') + e(A \cdot C'). \end{aligned} \quad (2.3)$$

Now, obviously the intersection number of C' with any component of $Exc(f)$ is greater or equal to 0, with equality if and only if C' does not intersect any of the irreducible components of $Exc(f)$, i.e. C does not pass through the singular points of X . Hence, it

follows that

$$\sum_{i=1}^{10} b_i(G_i \cdot C') + \sum_{i=1}^7 c_i(H_i \cdot C') + \sum_{i=1}^6 d_i(I_i \cdot C') + e(A \cdot C') \geq 0.$$

Thus $\sum_{i=1}^8 a_i(F_i \cdot C') \leq 0$. In this case, either there is an $i_0 \in \{1, \dots, 8\}$ such that $(C' \cdot F_{i_0}) < 0$, or $(C' \cdot F_i) = 0$ for all $i = 1, \dots, 8$, and C' does not meet any component of $Exc(f)$. In the first case C' must coincide with F_{i_0} . But the computations in [PPS07, LePa07] show that $(f^*K_X \cdot F_i) > 0$ for all $i = 1, \dots, 8$, which is impossible by our assumption. Therefore C' must have vanishing intersection number with all of the F_i 's and also with the all of the components of $Exc(f)$. But, as the Poincaré duals of the irreducible components of $Exc(f)$ and of the F_i 's generate $H^2(Z, \mathbb{Q})$, C' must be numerically trivial on Z , and this is a contradiction. Hence, X has ample canonical bundle.

As a consequence of Theorem 2.2 we get:

Corollary 2.3. *There exist smooth complex structures on $\mathbb{C}\mathbb{P}^2 \# k \overline{\mathbb{C}\mathbb{P}^2}$, for $k = 6, 7, 8$ which have ample canonical line bundles and hence admitting Kähler-Einstein metrics of negative scalar curvature.*

Proof. As we have already mentioned earlier, we only discuss the case of $\mathbb{C}\mathbb{P}^2 \# 6 \overline{\mathbb{C}\mathbb{P}^2}$. The other cases follow similarly.

We know that a \mathbb{Q} -Gorenstein smoothing of X exists [PPS07] and gives a smooth exotic structure on $\mathbb{C}\mathbb{P}^2 \# 6 \overline{\mathbb{C}\mathbb{P}^2}$. As we showed, the singular projective surface X has ample \mathbb{Q} -Cartier canonical divisor. But, ampleness is an open property [KoMo98], so the canonical bundle of these smoothings is automatically ample. This will allow us now to apply Aubin-Yau's Theorem 2.2 to prove the existence of Kähler-Einstein metrics of negative scalar curvature on the minimal surfaces of general type described above. \square

Remark 2.4. *The examples of complex surfaces with ample canonical bundle we found are new to our knowledge, when $K^2 = 2$ or 3. When $K^2 = 1$, LeBrun and Catanese [LeCa98] were able to find a complex surface with ample canonical bundle homeomorphic to $\mathbb{C}\mathbb{P}^2 \# 8 \overline{\mathbb{C}\mathbb{P}^2}$, as a deformation of the Barlow surface. As was pointed out by [LePa07], it is not known whether the surfaces we found in Corollary 2.3 are deformation equivalent to the Barlow surface, or even diffeomorphic to it.*

3. EXOTIC STRUCTURES: NON-EXISTENCE OF EINSTEIN METRICS

The classical obstruction to the existence of an Einstein metric on an oriented, compact, smooth, Riemannian, four-manifold (M, g) is the Hitchin-Thorpe Inequality:

$$(2\chi \pm 3\tau)(M) \geq 0 \tag{3.1}$$

with equality when the Einstein manifold (M, g) is covered by the hyperkähler $K3$ surface or by the flat four-torus, or by the above with the reversed orientations. Here $\chi(M)$, $\tau(M)$ denote the Euler characteristic and the signature of the manifold M , respectively. Using the Seiberg-Witten theory, LeBrun [LeB96, LeB01] was able to find new obstruction to the existence of Einstein metrics. The novelty of his results is that the existence of Einstein metrics depends on the differential structure considered. The techniques developed by LeBrun rely on the existence of a non-trivial solution of the Seiberg-Witten equations. We briefly introduce the needed notions of the Seiberg-Witten theory and state the main obstruction theorem.

For simplicity, we consider only the case when $H_1(M, \mathbb{Z})$ has no 2-torsion. Then there is a one-to-one correspondence between the set of $Spin^c$ structures and the set $\{c \in$

$H^2(M, \mathbb{Z})|_{c_1 \equiv w_2(M) \pmod{2}}$ of characteristic classes in $H^2(M, \mathbb{Z})$. The Seiberg-Witten invariant is defined [Mo96] as an integer valued function

$$SW_M : \{c \in H^2(M, \mathbb{Z}) \mid c \equiv w_2(M) \pmod{2}\} \longrightarrow \mathbb{Z}$$

We call c a basic class if $SW_M(c) \neq 0$.

The Seiberg-Witten invariant is a diffeomorphism invariant in the case $b_+ > 1$. In the case of $b_+ = 1$, the invariant depends on the choice of an orientation of $H^2(M, \mathbb{Z})$ and $H^1(M, \mathbb{R})$. But, as we are interested in manifolds for which the existence of an Einstein metric is topologically unobstructed, the inequality (3.1) must be strictly satisfied. Hence, in the case when the $SW_M(c) \neq 0$, we have $c^2 \geq (2\chi + 3\tau)(M) > 0$. But for any arbitrary metric g , $c^2 = c_+^2 - c_-^2 > 0$ where we denote by c_{\pm} the (anti)-self-dual part of the harmonic representative 2-form. Thus $c_+ \neq 0$, and in this special situation (see [Mo96, Thm 6.9.2]) the Seiberg-Witten invariant does not depend on the choice of metric.

We are now ready to state LeBrun's obstruction, see [LeB01, Thm. 3.3]:

Theorem 3.1 (LeBrun). *Let X be a compact oriented 4-manifold with a non-trivial Seiberg-Witten invariant and with $(2\chi + 3\tau)(X) > 0$. Then*

$$M = X \#_k \overline{\mathbb{C}\mathbb{P}^2}$$

does not admit an Einstein metric if $k \geq \frac{1}{3}(2\chi + 3\tau)(X)$.

In the last years there was a lot of work done on constructing exotic structures with small topology. In this paper we will use a construction due to Akhmedov, Baykur and Park [ABP07]. Similar results could also be found in papers of above authors or of Baldridge and Kirk.

Here we are interested in some exotic structures on $M = \mathbb{C}\mathbb{P}^2 \# 3\overline{\mathbb{C}\mathbb{P}^2}$. In [ABP07, Section 3.4], the authors construct a minimal symplectic manifold X homeomorphic to M , but not diffeomorphic. To give the reader a better understanding of this new manifold, we sketch its construction. We start with two product manifolds $Y = T^2 \times \Sigma_2$ and T^4 , where T^n is a torus of dimension n and Σ_2 is a Riemann surface of genus 2. If we consider the diagonal class $[T^2 \times \text{point}] + [\text{point} \times T^2] \in H_2(T^4, \mathbb{Z})$, then a smooth representative in this class is a Riemann surface of genus 2 with self-intersection 2. Blowing up two points on the smooth representative gives $\Sigma' \subset (T^4 \# 2\overline{\mathbb{C}\mathbb{P}^2})$, a smooth Riemann surface of genus 2 and 0-self-intersection. If we endow the manifolds Y and $T^4 \# 2\overline{\mathbb{C}\mathbb{P}^2}$ with the canonical symplectic structures, we can choose the representative Σ' to be a symplectic submanifold. We can now take the symplectic sum $X' = Y \#_{\Sigma_2} (T^4 \# 2\overline{\mathbb{C}\mathbb{P}^2})$ along the two genus 2 symplectic submanifolds $\text{point} \times \Sigma_2 \subset Y$ and $\Sigma' \subset (T^4 \# 2\overline{\mathbb{C}\mathbb{P}^2})$. X' is a symplectic manifold with topological invariants: Euler characteristic $\chi(X') = 6$ and signature $\tau(X') = -2$. As X' is not simply connected we need to make six Luttinger surgeries along well chosen tori (see [ABP07]), which yield a new simply connected, minimal, symplectic manifold X , with the same topological invariants as X' . Notice that the definition of the Seiberg-Witten invariant in [ABP07] is for the Poincaré dual of the characteristic class. The results are nevertheless the same. The manifold X has a unique basic class $\beta = c_1(K_X) \in H_2(X, \mathbb{Z})$, such that its Seiberg-Witten invariant $SW_X(\beta) = 1$. Moreover, X contains a null-homologous torus Λ , with a preferred simple loop λ . In Section 4, [ABP07], it is shown that 0-surgery on Λ with respect to λ yields a symplectic manifold X_0 , which also has an unique basic class $\beta_0 = c_1(K_{X_0})$. Then $1/n$ -surgery, $n \geq 1$ on Λ with respect to λ generates a family of manifolds X_n which are homeomorphic to $X_1 = X$. Corresponding to the basic class β of X there is a unique basic class β_n of X_n

for which the Seiberg-Witten invariant is non-zero, and it can be computed as follows:

$$SW_{X_n}(\beta_n) = SW_X(\beta) + (n-1)SW_{X_0}(\beta_0) = 1 + (n-1) = n.$$

Hence, the family $X_n, n \geq 1$ consists of homeomorphic, pairwise non-diffeomorphic manifolds. As β_n is the unique basic class on X_n , and the Seiberg-Witten invariant of $X_n, n > 1$ is not ± 1 , the manifold X_n does not admit a symplectic structure. Of course, its basic class satisfies $\beta_n^2 = \beta^2 = 6 = (2\chi + 3\tau)(X) = (2\chi + 3\tau)(X_n)$.

Let $M_{n,k} = X_n \#^k \overline{\mathbb{C}\mathbb{P}^2}, k = 2, 3, 4, 5$. Then, the conditions of Theorem 3.1 are satisfied and we have the following:

Proposition 3.2. *The manifolds $\mathbb{C}\mathbb{P}^2 \#^l \overline{\mathbb{C}\mathbb{P}^2}$ for $l = 5, 6, 7, 8$ support infinitely many non-diffeomorphic exotic smooth structures none of which admits an Einstein metric.*

Proof. The only thing that needs to be argued is that the manifolds $\{M_{n,k}\}_n$ are homeomorphic but pairwise not diffeomorphic. For this we use again the Seiberg-Witten invariant and the commutativity of our construction. First we blow-up $k = l - 3$ points on $X = X_1$ and denote by E_1, \dots, E_k the exceptional divisors. This gives a new symplectic manifold, which has basic classes $\beta \pm c_1(E_1) \pm \dots \pm c_1(E_k)$. 0-surgery on Λ with respect to λ gives the manifold $X_0 \#^k \overline{\mathbb{C}\mathbb{P}^2}$ with corresponding basic classes $\beta_0 \pm c_1(E_1) \pm \dots \pm c_1(E_k)$. We have a similar relation for the Seiberg-Witten invariant of $M_{n,k}$:

$$\begin{aligned} SW_{M_{n,k}}(\beta_n \pm c_1(E_1) \pm \dots \pm c_1(E_k)) &= SW_X(\beta \pm c_1(E_1) \pm \dots \pm c_1(E_k)) \\ &\quad + (n-1)SW_{X_0}(\beta_0 \pm c_1(E_1) \pm \dots \pm c_1(E_k)) \\ &= 1 + (n-1) = n. \end{aligned}$$

Hence, for any fixed $k \in \{2, 3, 4, 5\}$, the manifolds $M_{n,k}$ are homeomorphic to $\mathbb{C}\mathbb{P}^2 \# (k+3) \overline{\mathbb{C}\mathbb{P}^2}$, but pairwise not diffeomorphic. Moreover, if $n > 1$ they don't support a symplectic structure. \square

Remark 3.3. *We would like to note that since for all of the above smooth structures the Seiberg-Witten invariant is non-trivial the manifolds don't admit Riemannian metrics of positive constant scalar curvature. Moreover, as the $c_1^2(M_{n,k}) > 0$ there is a bound [LeB96] on the scalar curvature:*

$$\int_{M_{n,k}} s_g^2 d\mu \geq 32\pi^2 c_1^2(M_{n,k}) > 0$$

where $d\mu$ is the volume form with respect to g . Hence the manifolds $M_{n,k}$ don't admit non-negative constant scalar curvature metrics, and their Yamabe invariant is negative.

The results of Theorem 1.2 are immediate by putting together the results from 2.2, 2.3, 3.2.

4. HIGHER DIMENSIONAL MANIFOLDS

Let S_1 be a complex surface with ample canonical bundle and $c_1^2 = 1$. As an example of such surface, we can either take the one provided by Corollary 2.3, or the example [LeCa98] constructed by LeBrun and Catanese. Let S_2, S_3 also be complex surfaces of general type, with ample canonical line bundle and $c_1^2 = 2, 3$, respectively, as in Corollary 2. We are now ready to give the proof of Proposition 1.4.

Proof. Let $N = N_{i_1} \times \dots \times N_{i_k}$ and let $S = S_{i_1} \times \dots \times S_{i_k}$, where i_k is either 1, 2 or 3. The manifolds S_i and N_i are homeomorphic, hence by a theorem of Wall [Wa64], they are h-cobordant. These h-cobordisms induce an h-cobordism between N and S . But as N, S

are simply connected manifolds of dimension greater than 5, Smale's h-cobordism theorem tells us that they are diffeomorphic. We know that N_i and S_i admit Kähler-Einstein metrics of positive, negative scalar curvature, respectively. We can rescale these metrics such that the scalar curvatures are ± 1 . On N, S we consider the product metrics associated to the corresponding i_k . As we take products of Kähler-Einstein with the same scalar curvature, the new metrics are going to be Kähler-Einstein. The product complex structures on N, S are of Kodaira dimension $-\infty$ and $4k$, respectively. \square

Remark 4.1. *As the dimension considered increases, the number of manifolds constructed in Proposition 1.4 is $\frac{1}{2}(k+1)(k+2)$.*

ACKNOWLEDGEMENTS

The authors would like to thank Claude LeBrun for suggesting this problem and for his remarks and suggestions on an early version of this paper. This paper was written while the first author was a CNRS post-doc at IRMA, Strasbourg, and the second was visiting IHES as a IPDE fellow. We would like to thank these institutions for their hospitality.

REFERENCES

- [ABP07] A. AKHMEDOV, R. BAYKUR, B. PARK *Constructing infinitely many smooth structures on small 4-manifolds* preprint
- [Aub76] T. AUBIN, *Équations du type Monge-Ampère sur les variétés kähleriennes compactes*, C. R. Acad. Sci. Paris Sr. A-B 283 (1976), no. 3, A119–A121.
- [Bes87] A. BESSE, *Einstein Manifolds*, Ergebnisse der Mathematik und ihrer Grenzgebiete (3), 10, Springer Verlag, Berlin, 1987.
- [LeCa98] F. CATANESE, C. LEBRUN, *On the scalar curvature of Einstein manifolds*, Math. Res. Lett. 4 (1997), no. 6, 843–854.
- [KoMo98] J. KOLLÁR, S. MORI, *Birational geometry of algebraic varieties*, Cambridge Tracts in Mathematics, 134, Cambridge University Press, Cambridge, 1998.
- [LeB96] C. LEBRUN, *Four-manifolds without Einstein metrics*, Math. Res. Lett. 3 (1996), no. 2, 133–147.
- [LeB01] C. LEBRUN, *Ricci curvature, minimal volumes, and Seiberg-Witten theory*, Invent. Math. 145 (2001), no. 2, 279–316.
- [LePa07] Y. LEE, J. PARK *A simply connected surface of general type with $p_g = 0$ and $K^2 = 2$* , Invent. Math. 170 (2007), no. 3, 483–505.
- [Mo96] J. W. MORGAN *The Seiberg-Witten equations and applications to the topology of smooth four-manifolds*, Mathematical Notes, 44, Princeton University Press, Princeton, NJ, 1996
- [Pa05] J. PARK, *Simply connected symplectic 4-manifolds with $b_2^+ = 1$ and $c_1^2 = 2$* , Invent. Math. 159 (2005), no. 3, 657–667
- [PPS07] H. PARK, J. PARK, D. SHIN, *A simply connected surface of general type with $p_g = 0$ and $K^2 = 3$* , arXiv:0708.0273v3
- [PPS08] H. PARK, J. PARK, D. SHIN, *A simply connected surface of general type with $p_g = 0$ and $K^2 = 4$* , arXiv:0803.3667v1 [math.AG]
- [Siu88] Y.T. SIU, *The existence of Kähler-Einstein metrics on manifolds with positive anticanonical line bundle and a suitable finite symmetry group*, Ann. of Math. (2) 127 (1988), no. 3, 585–627.
- [TiYa87] G. TIAN, S.-T. YAU, *Kähler-Einstein metrics on complex surfaces with $c_1 > 0$* , Comm. Math. Phys. 112 (1987), no. 1, 175–203.
- [Ti90] G. TIAN, *On Calabi's conjecture for complex surfaces with positive first chern class*, Invent. Math. 101 (1990), no. 1, 101–172.
- [Yau77] S.-T. YAU, *Calabi's conjecture and some new results in algebraic geometry*. Proc. Nat. Acad. USA 74 (1977), 1789–1799.
- [Wa64] C.T.C. WALL, *On simply-connected 4-manifolds*, J. London Math. Soc. 39 1964 141–149.

RAREȘ RĂSDEACONU, IRMA, 7 RUE RENÉ DESCARTES, 67184 STRASBOURG CEDEX, FRANCE
E-mail address: rasdeaco@math.u-strasbg.fr

IOANA ȘUVAINA, IHES, 35 RUE DE CHARTRES, BURES-SUR-YVETTE, 91440, FRANCE
E-mail address: ioana@ihes.fr