LOCAL INVERSES OF SHIFT MAPS ALONG ORBITS OF FLOWS

SERGIY MAKSYMENKO

ABSTRACT. Let \mathbf{F} be a smooth flow on a smooth manifold Mand $\mathcal{D}(\mathbf{F})$ be the group of diffeomorphisms of M preserving orbits of \mathbf{F} . We study the homotopy type of the identity components $\mathcal{D}_{id}(\mathbf{F})^r$ of $\mathcal{D}(\mathbf{F})$ with respect to distinct Whithey topologies W^r , $(0 \leq r \leq \infty)$. The main result presents a class of flows \mathbf{F} for which $\mathcal{D}_{id}(\mathbf{F})^r$ coincide for all r and are either contractible or homotopy equivalent to the circle. The group $\mathcal{D}_{id}(\mathbf{F})^0$ was studied in the author's paper ["Smooth shifts along trajectories of flows", Topol. Appl. **130** (2003) 183-204]. Unfortunately that article contains a gap in estimations of continuity of local inverses of the so-called shift map. The present paper also repairs these estimations and shows that they hold under additional assumptions on the behavior of regular points of \mathbf{F} .

1. INTRODUCTION

Let M be a smooth (\mathcal{C}^{∞}) , connected, m-dimensional manifold possibly non-compact and with or without boundary. Let also F be a smooth vector field on M tangent to ∂M and generating a global flow $\mathbf{F}: M \times \mathbb{R} \to M$. Denote by Σ_F (or simply by Σ) the set of singular points of F.

Let $\mathcal{E}(\mathbf{F})$ be the subset of $\mathcal{C}^{\infty}(M, M)$ consisting of maps f such that

(1) $f(o) \subset o$ for every orbit o of F;

(2) f is a local diffeomorphism at every singular point $z \in \Sigma_F$.

Let also $\mathcal{D}(M)$ be the group of C^{∞} -diffeomorphisms of M and

$$\mathcal{D}(\mathbf{F}) \stackrel{\text{def}}{=\!\!=} \mathcal{E}(\mathbf{F}) \cap \mathcal{D}(M)$$

be the group of diffeomorphisms preserving orbits of F.

For every $r = 0, 1, ..., \infty$ denote by $\mathcal{E}_{id}(\mathbf{F})^r$ (resp. $\mathcal{D}_{id}(\mathbf{F})^r$) the pathcomponent of the identity map id_M in $\mathcal{E}(\mathbf{F})$ (resp. $\mathcal{D}(\mathbf{F})$) with respect to the weak \mathbf{W}^r Whithey topology, see Definition 1.4. In particular

²⁰⁰⁰ Mathematics Subject Classification. 37C05, 57S05, 57R45.

Key words and phrases. diffeomorphism, flow, homotopy type, shift map, linearizable vector fields, isolating block.

 $\mathcal{E}_{id}(\mathbf{F})^0$ (resp. $\mathcal{D}_{id}(\mathbf{F})^0$) consists of maps f which are homotopic to id_M in $\mathcal{E}(\mathbf{F})$ (resp. $\mathcal{D}(\mathbf{F})$).

Define the following map $\varphi : \mathcal{C}^{\infty}(M, \mathbb{R}) \to \mathcal{C}^{\infty}(M, M)$ by $\varphi(\alpha)(x) = \mathbf{F}(x, \alpha(x))$ for $\alpha \in \mathcal{C}^{\infty}(M, \mathbb{R})$ and $x \in M$. We will call φ the *shift* map along orbits of F and denote by $Sh(\mathbf{F})$ its image $\varphi(\mathcal{C}^{\infty}(V, \mathbb{R}))$ in $\mathcal{C}^{\infty}(M, M)$. Then the following inclusions hold true:

$$Sh(\mathbf{F})\ \subset\ \mathcal{E}_{\mathrm{id}}(\mathbf{F})^\infty\ \subset\ \cdots\ \mathcal{E}_{\mathrm{id}}(\mathbf{F})^1\ \subset\ \mathcal{E}_{\mathrm{id}}(\mathbf{F})^0.$$

The idea of replacing the time in a flow map with a function was extensively used e.g. in [8, 3, 31, 12, 27, 10] for reparametrizations of measure preserving flows and investigations of their mixing properties, see also [19, §1]. Smooth shifts functions were applied in [13, 15, 16, 14, 17] for study of homotopical properties of certain infinite dimensional groups of diffeomorphisms and their actions on spaces of smooth maps. Also in [19, 18] some applications to parameter rigidity of vector fields are given.

Suppose F is inearizable at each $z \in \Sigma_F$, see e.g. [29, 30, 32, 2, 11]. In [13] the author in particular *claimed* that if in addition M is compact, then $\mathcal{D}_{id}(\mathbf{F})^0$ and $\mathcal{E}_{id}(\mathbf{F})^0$ are either contractible or homotopy equivalent to the circle S^1 when endowed with W^{∞} topologies. Unfortunately, it turned out that in such a generality this statement fails and it is necessary to put additional restrictions on the behavior of regular orbits of F. In fact it was shown that $Sh(\mathbf{F}) = \mathcal{E}_{id}(\mathbf{F})^0$ and the mistakes appeared in estimations of continuity of local inverses of φ , see [13, Defn. 15, Th. 17, Lm. 32] and also Remarks 7.4 and 8.1 for detailed discussion.

Furthermore, the above description of $\mathcal{D}_{id}(\mathbf{F})^0$ was essentially used in another author's paper [15] concerning calculations of the homotopy types of stabilizers and orbits of Morse functions on surfaces.

The aim of the present paper is to repair incorrect formulations and proofs of [13] and using [14, 19, 17] extend the classes of vector fields for which the homotopy types of $\mathcal{E}_{id}(\mathbf{F})^r$ and $\mathcal{D}_{id}(\mathbf{F})^r$ can be described (Theorem 3.5). In particular it will be shown that the results of [13] used in [15] remain true (Theorem 3.7).

1.1. Structure of the paper. For the convenience of the reader and due to the length of the paper we will now briefly describe its contents.

In §2 we recall the notion of shift map and review (correct) results obtained in [13]. Also notice that if a flow **F** of a vector field F is not global, then we can find a smooth function $\mu : M \to (0, +\infty)$ such that the flow **F**' generated by vector field $F' = \mu F$ is global, e.g. [6, Cor. 2]. Moreover, F and F' have the same orbit structure, whence $\mathcal{E}(\mathbf{F}) = \mathcal{E}(\mathbf{F}')$ and $\mathcal{D}(\mathbf{F}) = \mathcal{D}(\mathbf{F}')$. Thus it could always be assumed that \mathbf{F} is global. Nevertheless, in §10 and §11 we will consider restriction $F|_W$ of F to open subsets $W \subset M$ and compare shift maps of F and $F|_W$. The latter vector field usually generates non-global flow, therefore in this paper it is chosen to work with local flows from the beginning. In particular, the domain of shift map φ of F changes to certain open and convex subset $\mathsf{func}(\mathbf{F}) \subset \mathcal{C}^{\infty}(M, \mathbb{R})$.

In §3 we introduce a certain class of vector fields $\mathcal{F}(M)$ on M and formulate the principal result of the paper: for every $F \in \mathcal{F}(M)$ its shift map φ is a local homeomorphism of func(**F**) onto its image $Sh(\mathbf{F})$ with respect to S^{∞} topologies (Theorem 3.5). It follows that $Sh(\mathbf{F})$ is either contractible or homotopy equivalent to the circle, and so is the subspace of $Sh(\mathbf{F})$ consisting of diffeomorphisms. On the other hand, in [18] it is considered a class of vector fields $\mathcal{F}'(M)$ which contains $\mathcal{F}(M)$ and such that for every $F \in \mathcal{F}'(M)$ the image of its shift map $Sh(\mathbf{F})$ coincides with either $\mathcal{E}_{id}(\mathbf{F})^1$ or $\mathcal{E}_{id}(\mathbf{F})^0$ for each $F \in \mathcal{F}(M)$. Thus we obtain that $\mathcal{E}_{id}(\mathbf{F})^r$ ($\mathcal{D}_{id}(\mathbf{F})^r$) coincide with each other at least for all $r \geq 1$ and these spaces are either contractible or have homotopy type of S^1 . As an application we prove Theorem 3.7 which extends [15, Th. 1.3]. The rest of the paper is devoted to the proof of Theorem 3.5.

The assumptions of Theorem 3.5 imply that φ is locally injective, therefore in order to prove this theorem it suffices to show that φ is open with respect to S^{∞} topologies. In §4 we present a characterization of $S^{r,s}$ -openness of φ for some $r, s \geq 0$, (Theorem 4.2). First we describe local inverses of φ_V as certain *crossed homomorphisms* and then show that like for homomorphisms of groups $S^{r,s}$ -openness of the whole map φ is equivalent to $S^{s,r}$ -continuity of its local inverse defined *only* on arbitrary small S^s -neighbourhood of the identity map id_M .

Further in §5 we recall notorious examples of irrational flows on the torus and some modifications of them. It is shown that shift maps of these flows are not local homeomorphisms onto their images. This provides counterexamples to [13, Th. 1] and illustrates certain properties prohibited by Theorem 3.5.

In §6 we repair [13, Lm. 28] by giving sufficient conditions for $S^{r,s}$ openness of φ to be inherited by regular extensions of vector fields.

§7 summarizes the formulas for local inverses of shift maps of linear flows obtained in [13] and "reduced" Hamiltonian flows of homogeneous polynomials in two variables obtained in [19, 14]. Estimations of continuity of these formulas for linear flows were based on incorrect "division lemma" [13, Lm. 32], see §7.4. We will show how to avoid referring to this lemma.

SERGIY MAKSYMENKO

§8 provides a correct version of [13, Th. 17], see Theorem 8.2. It reduces verification of openness of φ to openness of a family $\{\varphi_{V_i}\}$ of "local shift maps" corresponding to any locally finite cover $\{V_i\}$ of M e.g. by arbitrary small smooth closed disks. The essentially new additional object here is a finite subset $\Lambda' \subset \Lambda$ which appears due to the construction (8.3) of S^{∞} -open set \mathcal{N} and by existence of singularities for which "local shift map" is $S^{\infty,\infty}$ -open but not $S^{r,d(r)}$ -open for some function $d: \mathbb{N} \to \mathbb{N}$. Without finiteness of Λ' the shift map φ is open if its image is endowed with the so-called *very-strong topology*, [9]. This effect appears of course only on non-compact manifolds.

§9 splits the "global analytical problem" of verification of openness of local shift map φ_V into the following two problems

- (i) openness of local shift map $\varphi_{W,V}$ corresponding to the restriction $F|_W$ to arbitrary open neighbourhood W of V, and
- (ii) openness of the image $Sh(\mathbf{F}_W, V)$ in $Sh(\mathbf{F}, V)$.

Due to Theorem 8.2 the set V can also be taken arbitrary small. Hence to solve (i) it suffices to consider vector fields in \mathbb{R}^n . Thus (i) is a "local analytical" problem. Its solutions for certain vector fields are given in §7.

 $\S10$ and $\S11$ present sufficient conditions for resolving (ii) at regular and singular points respectively (Theorem 10.1 and Lemma 11.2). In both cases the assumptions on F are formulated in the terms of dynamical systems theory, and so the problem (ii) can be regarded as a "global topological" one. In the regular case these conditions are also necessary. Moreover, in the singular case they are relevant with the notion of an isolated block introduced by C. Conley and R. Easton in [4], see Lemma 11.5.

Finally in $\S12$ we prove Theorem 3.5.

1.2. **Preliminaries.** Put $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$ and $\overline{\mathbb{N}}_0 = \mathbb{N} \cup \{0, \infty\}$.

Let M be a smooth manifold of dimension m. Glue to M a collar $\partial M \times [0,1)$ by the identity map $\partial M \times 0 \to \partial M$ and denote the obtained manifold by M'. If $\partial M = \emptyset$, then M' = M. Evidently, M' has smooth structure in which it is diffeomorphic with the interior IntM. Moreover M is a closed subset of M' and $\partial M' = \emptyset$. We will call M' a collaring of M, see Figure 1.

DEFINITION 1.3. A closed subset $V \subset M$ will be called a *D*-submanifold if there exists an m-dimensional submanifold $V' \subset M'$ possibly with boundary and such that $V = M \cap V'$ and the intersection $\partial M \cap \partial V'$ is transversal, see Figure 1.

Denote by IntV the interior of V in M. Then $IntV = V \setminus \partial V'$. We will also say that V is D-neighbourhood of every point $z \in IntV$.

Let $V \subset M$ be a *D*-submanifold. If $V \cap \partial M = \emptyset$, then *V* is a manifold with boundary, otherwise, *V* is a manifold with corners with $\partial M \cap \partial V'$ as the set of corners. Evidently, *M* is a *D*-submanifold of itself, and each $z \in \text{Int}M$ has arbitrary small *D*-neighbourhoods each diffeomorphic to a closed *m*-disk.

Let N be another smooth manifold, N' be its collaring, and $f : V \to N$ be a map. We say that f is of class C^r , $r \in \overline{\mathbb{N}}_0$, (embedding, immersion, etc.) if it extends to a C^r map (embedding, immersion, etc.) $U \to N'$ from some open neighbourhood U of V in M' into the collaring N' of N.

Then it is known well-known that f is C^r if, and only if, the restriction $f|_{\operatorname{Int} V \setminus \partial M}$ is C^r and all its derivatives have continuous limits when x tends to some point $y \in \partial V'$, see e.g. [33, 28, 24, 5] for manifolds with boundary and e.g [22, Pr. 2.1.10] for manifolds with corners.




FIGURE 1. D-submanifolds V and W

Denote by $\mathcal{C}^r(V, N)$, $r \in \overline{\mathbb{N}}_0$, the space of all \mathcal{C}^r maps $V \to N$. Then similarly to [7] this space can be endowed with weak W^r and strong S^r topologies. If V is compact then W^r and S^r coincide for all $r \in \overline{\mathbb{N}}_0$. For a subset $\mathcal{X} \subset \mathcal{C}^r(V, N)$ denote by $(\mathcal{X})^r_W$ (resp. $(\mathcal{X})^r_S$) the set \mathcal{X} with the induced W^r (resp. S^r) topology, $r \in \overline{\mathbb{N}}_0$.

DEFINITION 1.4. A homotopy $H : V \times I \to N$ will be called an *r*-homotopy, $(r \in \overline{\mathbb{N}}_0)$, if for every $t \in I$ the map $H_t = H(\cdot, t) : V \to N$ is \mathcal{C}^r and all its partial derivatives in $x \in V$ up to order r are continuous in (x, t). In other words, H is an *r*-homotopy if, and only if, it yields a continuous path $I \to \mathcal{C}^r(V, N)$ from the standard topology of I to the W^r topology of $\mathcal{C}^r(V, N)$.

Hence for every $f \in \mathcal{X}$ its path connected component in $(\mathcal{X})_W^r$ consists of all $g \in \mathcal{X}$ which are *r*-homotopic to f in \mathcal{X} .

Now let V_1, V_2 be *D*-submanifolds of some smooth manifolds, N_1, N_2 be two smooth manifolds, $\mathcal{X} \subset \mathcal{C}^{\infty}(V_1, N_1)$ and $\mathcal{Y} \subset \mathcal{C}^{\infty}(V_2, N_2)$ be two subsets, and $F : \mathcal{X} \to \mathcal{Y}$ be a map. DEFINITION 1.5. We say that F is $W^{r,s}$ -continuous (-open, etc.) for some $r, s \in \overline{\mathbb{N}}_0$ if it is continuous (open, etc.) as a map $F : (\mathcal{X})_W^r \to (\mathcal{Y})_W^s$.

Similarly we can define $S^{r,s}$ -continuous (-open, etc.) maps with respect to strong topologies.

Notice that the statement that F is $W^{\infty,\infty}$ -continuous at some $x \in \mathcal{X}$ means that for every $s \geq 0$ and a W^s -open neighbourhood $V_{F(x)} \subset \mathcal{Y}$ of F(x) there exist $r \geq 0$ and a W^r -neighbourhood $U_x \subset \mathcal{X}$ of x both depending on s and $V_{F(x)}$ such that $F(U_x) \subset V_{F(x)}$. But in general such a map can be not $W^{r,s}$ -continuous for any r, s, see e.g. [23, p. 93. Eq. (2)], and [25].

2. Shift map

Let F be a vector field on M tangent to ∂M . Then for every $x \in M$ its *orbit* with respect to F is a unique mapping $o_x : \mathbb{R} \supset (a_x, b_x) \to M$ such that $o_x(0) = x$ and $\frac{d}{dt}o_x = F(o_x)$, where $(a_x, b_x) \subset \mathbb{R}$ is the maximal interval on which a map with the previous two properties can be defined. By standard theorems in ODE the following subset of $M \times \mathbb{R}$

$$\mathsf{dom}(\mathbf{F}) = \bigcup_{x \in M} x \times (a_x, b_x),$$

is an open, connected neighbourhood of $M \times 0$ in $M \times \mathbb{R}$. Then the *local flow* of F is the following map

$$\mathbf{F}: M \times \mathbb{R} \supset \operatorname{dom}(\mathbf{F}) \longrightarrow M, \qquad \mathbf{F}(x,t) = o_x(t).$$

If M is compact, then $\mathsf{dom}(\mathbf{F}) = M \times \mathbb{R}$, e.g. [26].

Notice that is $x \in M$ is either fixed or periodic for \mathbf{F} , then $x \times \mathbb{R} \subset \operatorname{\mathsf{dom}}(\mathbf{F})$.

EXAMPLE 2.1. Let $F(x) = x^2 \frac{d}{dx}$ be a vector field on \mathbb{R}^1 . Then it is easy to see that it generates the following local flow $\mathbf{F}(x,t) = \frac{x}{1-xt}$. Hence **F** is defined on the subset $\mathsf{dom}(\mathbf{F}) \subset \mathbb{R}^2$ bounded by the hyperbola xt = 1, see Figure 2.

Let $V \subset M$ be either an open subset or a *D*-submanifold and $i_V : V \subset M$ be the identity inclusion. Denote by $\mathcal{E}(\mathbf{F}, V)$ the subset of $\mathcal{C}^{\infty}(V, M)$ consisting of maps $f : V \to M$ such that (i) $f(o \cap V) \subset o$ for every orbit o of F, and (ii) f is an embedding at every singular point $z \in \Sigma_F$. Let also $\mathcal{D}(\mathbf{F}, V) \subset \mathcal{E}(\mathbf{F}, V)$ be the subset consisting of immersions $V \to M$.




FIGURE 2. Domain dom(F) of the flow F of the vector field $F(x) = x^2 \frac{d}{dx}$

Let $\mathcal{E}_{id}(\mathbf{F}, V)^r$ and $\mathcal{D}_{id}(\mathbf{F}, V)^r$ be the path-components of i_V in the spaces $\mathcal{E}(\mathbf{F}, V)$ and $\mathcal{D}(\mathbf{F}, V)$ respectively endowed with the corresponding topologies W^r .

Denote by $\operatorname{func}(\mathbf{F}, V)$ the subset of $C^{\infty}(V, \mathbb{R})$ consisting of functions α whose graph $\Gamma_{\alpha} = \{(x, \alpha(x)) : x \in V\}$ is contained in dom(**F**). Then we can define the following map

$$\varphi_V: C^{\infty}(V, \mathbb{R}) \ \supset \ \mathsf{func}(\mathbf{F}, V) \ \longrightarrow \ C^{\infty}(V, M)$$

by $\varphi_V(\alpha)(x) = \mathbf{F}(x, \alpha(x))$. We will call φ_V the *(local) shift map* of F on V and denote its image in $\mathcal{C}^{\infty}(M, M)$ by $Sh(\mathbf{F}, V)$. Put

$$\Gamma_V^+ = \{ \alpha \in \mathsf{func}(\mathbf{F}, V) \; : \; d\alpha(F)(x) > -1 \; \forall x \in V \},$$

where $d\alpha(F)$ is a Lie derivative of α along F. Since $\mathsf{func}(\mathbf{F}, V)$ is an S^0 -open and convex subset of $\mathcal{C}^\infty(V, \mathbb{R})$ and the map

$$\mathcal{L}_F : \mathcal{C}^{\infty}(V, \mathbb{R}) \to \mathcal{C}^{\infty}(V, \mathbb{R}), \qquad \mathcal{L}_F(\alpha) = d\alpha(F)$$

is evidently linear and $S^{1,0}$ continuous, we see that Γ_V^+ is convex and S^1 -open in $\mathcal{C}^{\infty}(V,\mathbb{R})$. It also follows from [13, Lm. 20 & Cor. 21] that

(2.1)
$$\Gamma_V^+ = \varphi_V^{-1} (\mathcal{D}_{\mathrm{id}}(\mathbf{F}, V)^1), \qquad Sh(\mathbf{F}, V) \subset \mathcal{E}(\mathbf{F}, V).$$

Lemma 2.2. [17] The following inclusions hold true:

$$\varphi_V(\Gamma_V^+) \subset \mathcal{D}_{\mathrm{id}}(\mathbf{F}, V)^{\infty} \subset \cdots \subset \mathcal{D}_{\mathrm{id}}(\mathbf{F}, V)^r \subset \cdots \subset \mathcal{D}_{\mathrm{id}}(\mathbf{F}, V)^0,$$

$$Sh(\mathbf{F}, V) \subset \mathcal{E}_{\mathrm{id}}(\mathbf{F}, V)^{\infty} \subset \cdots \subset \mathcal{E}_{\mathrm{id}}(\mathbf{F}, V)^r \subset \cdots \subset \mathcal{E}_{\mathrm{id}}(\mathbf{F}, V)^0.$$

$$If Sh(\mathbf{F}, V) \subset \mathcal{E}_{\mathrm{id}}(\mathbf{F}, V)^r, \ then \ \varphi_V(\Gamma_V^+) = \mathcal{D}_{\mathrm{id}}(\mathbf{F}, V)^r.$$

The set $\ker(\varphi_V) \stackrel{\text{def}}{=} \varphi_V^{-1}(i_V)$ will be called the *kernel* of φ_V .

Lemma 2.3. The following properties of shift map hold true:

(1) φ_V is $W^{r,r}$ - and $S^{r,r}$ -continuous for all $r \ge 0$, [13, Lm. 2].

(2) $d\mu(F) = 0$ for every $\mu \in \ker(\varphi_V)$, [13, Lm. 7].

(3) Let $\alpha, \beta \in \text{func}(\mathbf{F}, V)$. Then, [13, Lm. 7], the following conditions are equivalent:

SERGIY MAKSYMENKO

(a) $\varphi_V(\alpha) = \varphi_V(\beta)$ (b) $\alpha - \beta \in \mathsf{func}(\mathbf{F}, V)$ and $\varphi_V(\alpha - \beta) = i_V$, *i.e.* $\alpha - \beta \in \ker(\varphi_V)$.

(4) φ_V is locally injective with respect to each of W^r or S^r topologies of func(\mathbf{F}, V) if and only if $\Sigma_F \cap V$ is nowhere dense in V, [13, Pr. 14].

(5) Suppose that V is connected and that $V \cap \Sigma_F$ is nowhere dense in V. Then, [13, Th. 12(2)],

- (a) either ker(φ_V) = {0} and thus φ_V is injective. This case holds if V contains either a non-periodic point of **F** or a fixed point $z \in V \cap \Sigma_F$ such that the tangent linear flow $T_z \mathbf{F}_t$ on the tangent space $T_z M$ is the identity;
- (b) or $\ker(\varphi_V) = \{n\nu\}_{n\in\mathbb{Z}}$ for some smooth $\nu : V \to (0,\infty)$. In this case $V \times \mathbb{R} \subset \operatorname{dom}(\mathbf{F})$, so $\operatorname{func}(\mathbf{F}, V) = \mathcal{C}^{\infty}(V, \mathbb{R})$.

3. Main result

Let M be a smooth manifold of dimension m. We will introduce a class $\mathcal{F}(M)$ of smooth vector fields F on M satisfying certain *topological* conditions on regular points and certain *analytical* conditions on singular points. The main result describes the homotopy types of the identity path components of $\mathcal{E}(\mathbf{F})$ and $\mathcal{D}(\mathbf{F})$. First we give some definitions.

Recurrent points. Let F be a vector field on M and $z \in M$ be a regular point of F, i.e. $F(z) \neq 0$. Then z is called *recurrent* if there exists a sequence $\{t_i\}_{i\in\mathbb{N}}$ of real numbers such that $\lim_{i\to\infty} |t_i| = \infty$ and $\lim_{i\to\infty} \mathbf{F}(z,t_i) = z$. In particular, every periodic point is recurrent.

First recurrence map. Let z be a periodic point of F, o_z be its orbit, and $B \subset M$ be and open (m-1)-disk passing through z and transversal to o_z . Then we can define a germ at z of the so-called first return (or Poincaré) map $R : (B, z) \to (B, z)$ associating to every $x \in B$ the first point R(x) at which the orbit o_x of x first returns to B. It is well-known that R is a diffeomorphism at z, e.g. [26].

Reduced Hamiltonian vector fields. Let $g : \mathbb{R}^2 \to \mathbb{R}$ be a homogeneous polynomial of degree $p \geq 2$, so we can write

(3.1)
$$g = L_1^{l_1} \cdots L_a^{l_a} \cdot Q_1^{q_1} \cdots Q_b^{q_b},$$

where L_i is a non-zero linear function, Q_j is an irreducible over \mathbb{R} (definite) quadratic form, $l_i, q_j \geq 1$, $L_i/L_{i'} \neq \text{const}$ for $i \neq i'$, and $Q_j/Q_{j'} \neq \text{const}$ for $j \neq j'$. Denote

$$D = L_1^{l_1 - 1} \cdots L_a^{l_a - 1} \cdot Q_1^{q_1 - 1} \cdots Q_b^{q_b - 1}.$$

Then $g = L_1 \cdots L_a \cdot Q_1 \cdots Q_b \cdot D$ and D is the greatest common divisor of the partial derivatives g'_x and g'_y . The following vector field on \mathbb{R}^2 :

$$F(x,y) = -(g'_y/D)\frac{\partial}{\partial x} + (g'_x/D)\frac{\partial}{\partial y}$$

will be called the *reduced Hamiltonian* vector field of g. In particular, if g has no multiple factors, i.e. $l_i = q_j = 1$ for all i, j, then $D \equiv 1$ and F is the usual *Hamiltonian* vector field of g.

Notice that $dg(F) \equiv 0$ and the coordinate functions of F are homogeneous polynomials of degree deg F = a + 2b - 1 being relatively prime in the ring of polynomials $\mathbb{R}[x, y]$.

"Elementary" singularities. Let F be a vector field on \mathbb{R}^k . It can be regarded as a map $F : \mathbb{R}^k \to \mathbb{R}^k$. Define the following types (Z), (L), and (H) of such vector fields.

Type (Z): $F(x) \equiv 0$.

Type (L): $F(x) = \alpha(x) \cdot Ax$, where A is a non-zero $(k \times k)$ -matrix, and $\alpha : \mathbb{R}^n \to (0, +\infty)$ is a \mathcal{C}^{∞} strictly positive function. Let $\lambda_1, \ldots, \lambda_k$ be all the eigen values of A taken with their multiplicities. We can also assume that A has a *real* Jordan normal form. Then we will distinguish the following particular cases.

(L1) $\Re(\lambda_i) \neq 0$ for some $j = 1, \ldots, k$;

(L2) The real Jordan normal form has the block

$$\left\| \begin{array}{ccc} 0 & 1 \\ 0 & 1 \\ & \cdots & \dots \\ & & 0 & 1 \\ & & & 0 \end{array} \right\| \right\} n, \qquad n \ge 2;$$

(L3) The real Jordan normal form has the block

$$\left\| \begin{array}{cccc} 0 & b & 1 & 0 \\ b & 0 & 0 & 1 & \dots & \dots \\ \dots & \dots & \dots & \dots & \dots \\ b & 0 & b & 1 & 0 \\ b & 0 & 0 & 1 & \dots \\ & & 0 & b & 0 \end{array} \right\| \right\} 2n, \qquad (n \ge 2, b \ne 0).$$

In all other cases A is similar to the matrix:

Then we will also distinguish the following two cases:

(L4) there exists $\tau > 0$ such that $b_j \tau \in \mathbb{Z}$ for all j;

(L5) such τ as in (L4) does not exist.

SERGIY MAKSYMENKO

Type (H): $F(x,y) = \alpha(x,y) \cdot (-g'_y/D, g'_x/D)$ for some strictly positive \mathcal{C}^{∞} function $\alpha : \mathbb{R}^2 \to (0, +\infty)$, and a homogeneous polynomial (3.1) such that $a + 2b - 1 \ge 2$, so F is not of type (L). Again we separate the following cases:

- (HE) a = 0 and $b \ge 2$, i.e. g is a product of at least two distinct definite quadratic forms and has no linear multiples. In this case $0 \in \mathbb{R}^2$ is a degenerate global **extreme** of g;
- (HS) $a \ge 1$ and $a + 2b \ge 3$, so g has linear multiples. In this case $0 \in \mathbb{R}^2$ is a degenerate **saddle** critical point for g.

REMARK 3.1. The types (L), (H), (HE), and (HS) coincide with the ones considered in [18].

Also notice that in the case (L4) almost orbits of F are periodic, while in the case (L5) almost orbits are non-periodic and recurrent.

Regular extensions and products of vector fields. Let M, N be manifolds, $G: M \to TM$ be a vector field on M, and F be a vector field on $M \times N$. Then F can be regarded as a map $F: M \times N \longrightarrow TM \times TN$. We say that F is a *regular extension of* G if

$$F(x,y) = (G(x), H(x,y)), \qquad (x,y) \in M \times N,$$

for some smooth map $H: M \times N \to TN$, so G is the "first" coordinate of F and does not depend on y, see e.g. [1, 32]. If H does not depend on x, i.e. is a vector field on N, then

$$F(x,y) = (G(x), H(y))$$

will be called the *product* of G and H. Moreover, if $H \equiv 0$, i.e.

$$F(x,y) = (G(y),0),$$

then F is said to be a *trivial extension* of G.

In the case $N = \mathbb{R}^n$ we will also say that F is a (regular or trivial) *n*-extension. Evidently, a regular (trivial) *m*-extension of a regular (trivial) *n*-extension is a regular (trivial) (m+n)-extension, see e.g. [32, 13]. The following simple statement is left for the reader:

Lemma 3.2. (a) a trivial extension is the same as a product with zero vector field;

(b) a product of vector fields is a regular extension of each of them;

(c) every non-zero linear vector field is a regular extension of a linear vector field defined by one of the following matrices: ||a||, $|| \stackrel{a}{}_{-b} \stackrel{b}{}_{a} ||$, $(b \neq 0)$, or $|| \stackrel{0}{}_{0} \stackrel{1}{}_{0} ||$.

DEFINITION 3.3. Let $z \in F$ and (T) denotes one of the types (Z), (L), (L1), etc., defined above. We will say the germ of F at z is of type

(T)' if the germ of F at z is equivalent to a regular extension of some vector field of type (T). By $\Sigma^{(T)}$ and $\Sigma^{(T)'}$ we will denote the set of singular points of F or types (T) and (T)' respectively.

Evidently, a singular point can belong to distinct types.

DEFINITION 3.4. We will say that a C^{∞} vector field F on a C^{∞} manifold M belongs to class $\mathcal{F}(M)$ if it satisfies the following conditions:

- (a) F is tangent to ∂M and Σ_F is nowhere dense in M;
- (b) every non-periodic regular point of F is non-recurrent;
- (c) for every periodic point z of F the germ at z of its first recurrence map $R: (D, z) \to (D, z)$ is either periodic or the tangent map $T_z R: T_z D \to T_z D$ has eigen value λ such that $|\lambda| \neq 1$;
- (d) for every $z \in \Sigma_F$ the germ of F at z is either
 - (*) a **product** of finitely many vector fields each of which is of type (L) or (H), or
 - (**) belongs to one of the types (L1)', (L2)', (L3)', or (H)', but z is an **isolated** singular point of F.

Thus if the germ of F at z is a regular extension of an (L4)-vector field, then F must in fact be a product of such a vector field with vector fields of types (L) or (H) only. Also notice that singularities of type (L5) are not allowed at all since they have recurrent orbits. Moreover, we will present examples of vector fields with singularities of type (L5) for which the statement of our main result fails, see the end of §5.

The following theorem summarizes the results obtained in [18] and in this paper.

Theorem 3.5. c.f. [13, Th. 1] and [18]. Let $F \in \mathcal{F}(M)$, φ be the shift map of F, and $\Gamma^+ = \{ \alpha \in \mathsf{func}(\mathbf{F}) : d\alpha(F)(x) > -1 \ \forall x \in M \}$. Then

$$Sh(\mathbf{F}) = \mathcal{E}_{id}(\mathbf{F})^1, \qquad \varphi(\Gamma^+) = \mathcal{D}_{id}(\mathbf{F})^1$$

If $\Sigma = \Sigma^{(L)'} \cup \Sigma^{(HS)'}$, then $Sh(\mathbf{F}) = \mathcal{E}_{id}(\mathbf{F})^0$ and $\varphi(\Gamma^+) = \mathcal{D}_{id}(\mathbf{F})^0$. Suppose that $\overline{\Sigma \setminus (\Sigma^{(L1)'} \cup \Sigma^{(L2)'})}$ is compact (this is always true for

Suppose that $\Sigma \setminus (\Sigma^{(L1)'} \cup \Sigma^{(L2)'})$ is compact (this is always true for compact M). Then both maps

$$\varphi: \operatorname{func}(\mathbf{F}) \to \mathcal{E}_{\operatorname{id}}(\mathbf{F})^1, \qquad \qquad \varphi|_{\Gamma}: \Gamma \to \mathcal{D}_{\operatorname{id}}(\mathbf{F})^1$$

are either homeomorphisms or Z-covering maps with respect to S^{∞} topologies. If M is compact, then the inclusion $\mathcal{D}_{id}(\mathbf{F})^1 \subset \mathcal{E}_{id}(\mathbf{F})^1$ is a homotopy equivalence and both spaces are either contractible or homotopy equivalent to the circle. If F has at least one non-closed orbit, or a singular point at which the linear part (i.e. 1-jet) of F vanishes, then $\mathcal{D}_{id}(\mathbf{F})^1$ and $\mathcal{E}_{id}(\mathbf{F})^1$ are contractible.

SERGIY MAKSYMENKO

The first statement about the image of shift maps is established in [18] under more general assumptions on F. Therefore we will be proving the second part of Theorem 3.5, see §12. Notice that in comparison with [13, Th. 1] two additional assumptions are added: "nonergodicity" condition (like absence of recurrent orbits) and compactness of certain subset of singular points. On the other hand the classes of admissible singularities for F are extended due to results of [14, 19, 20].

3.6. Functions on surfaces. As an application of Theorem 3.5 we will now show that [15, Th. 1.3] which used incorrect results of [13] remains true. Moreover, we extend the latter theorem to a large class of functions on surfaces with degenerate homogeneous singularities satisfying certain "secondary" non-degeneracy conditions.

Let M be a compact surface, P be either the real line \mathbb{R} or the circle S^1 . For a smooth map $f: M \to P$ denote by Σ_f the set of its critical points. Let also $\mathcal{S}(f) = \{h \in \mathcal{D}(M) : f \circ h = f\}$ be the stabilizer of f with respect to the action of $\mathcal{D}(M)$ on $\mathcal{C}^{\infty}(M, P)$ and be the identity component of $\mathcal{S}(f)$ with respect to the W^{∞} topology.

Theorem 3.7. c.f. [15, Th. 1.3]. Let $f \in \mathcal{C}^{\infty}(M, P)$. Suppose that

(i) f takes constant values on connected components of ∂M and has no critical points on ∂M ;

(ii) for every $z \in \Sigma_f$ the map f is \mathcal{C}^{∞} equivalent near z to a homogeneous polynomial g_z without multiple factors such that deg $g_z \ge 2$.

If M is orientable and Σ_f consists of non-degenerate local extremes only, i.e. f is a Morse map without critical points of index 1, then $S_{id}(f)$ is homotopy equivalent to the circle. In all other cases $S_{id}(f)$ is contractible as well.

Sketch of proof. It suffices to assume that M is orientable. A nonorientable case will follow from orientable one by arguments of [15, §4.7].

Similarly to [15, Lm. 5.1] using (i) and (ii) it is possible to construct a vector field F on M with the following properties:

- (A) df(F) = 0, in particular F is tangent to ∂M ;
- (B) F(z) = 0 iff $z \in M$ is a critical point of F, i.e. $\Sigma_F = \Sigma_f$;
- (C) for every critical point $z \in \Sigma_f$ there exists a local presentation $f : \mathbb{R}^2 \to \mathbb{R}$ of f in which z = 0, f is a homogeneous polynomial of degree ≥ 2 and without multiple factors, and $F(x, y) = (-f'_y, f'_x)$ is a Hamiltonian vector field of f.

Then it follows from (A), (B), and arguments of [15, Lm. 3.5] that $S_{id}(f) = \mathcal{D}_{id}(\mathbf{F})^{\infty}$.

13

Notice that F belongs to class $\mathcal{F}(M)$. Indeed, conditions (a) and (b) are evident. Moreover, for each periodic point of F its first recurrent map is the identity, which implies (c).

Finally by (C) each non-degenerate saddle of f is of type (L1), each non-degenerate local extreme of f is of type (L4), while all degenerate critical points of f are of types (H). This implies (d).

Hence $S_{id}(f) = \mathcal{D}_{id}(\mathbf{F})^{\infty}$ is either contractible or homotopy equivalent to S^1 . If M is orientable and Σ_f consists of non-degenerate local extremes only, then f is Morse and belongs to one of the types (A)-(D) of [15, Th. 1.9]. In this case the corresponding shift map is not injective, whence $S_{id}(f)$ is homotopy equivalent to S^1 .

In all other cases φ is injective and so $S_{id}(f)$ is contractible. Indeed, if f has a saddle critical point then F has a on-closed orbit, while if f has a degenerate local extreme z then $j^1F(z) = 0$.

4. Openness of φ

Let $V \subset M$ be a *D*-submanifold. Our aim is to find sufficient conditions for the shift map $\varphi_V : \operatorname{func}(\mathbf{F}, V) \to Sh(\mathbf{F}, V)$ to be a local homeomorphism with respect to S^{∞} topologies. Since φ_V is $S^{r,r}$ -continuous for all $r \in \overline{\mathbb{N}}_0$, this is equivalent to $S^{\infty,\infty}$ -openness of φ_V . Moreover, as φ_V is locally injective, we can also require that its local inverses are $S^{\infty,\infty}$ -continuous and defined on S^{∞} -open subsets of $Sh(\mathbf{F}, V)$.

In this section we show that $S^{r,s}$ -openness of φ_V for some $r, s \in \overline{\mathbb{N}}_0$ is equivalent to $S^{s,r}$ -continuity of its local inverse defined *only* on some neighbourhood of the identity inclusion $i_V : V \subset M$ in $Sh(\mathbf{F}, V)$.

DEFINITION 4.1. Let A be a group and S be a semigroup with unit e. Then the **right action** of S on A is a map $*: A \times S \to A$ such that $\alpha * e = \alpha$ and $\alpha * (f g) = (\alpha * f) * g$ for all $\alpha \in A$ and $f, g \in S$. A map $\sigma: S \to A$ is called a **crossed homomorphism** if

$$\sigma(f g) = (\sigma(f) * g) \sigma(g).$$

Suppose for the moment that F generates a global flow \mathbf{F} . Put $S = Sh(\mathbf{F})$ and $A = \mathsf{func}(\mathbf{F}) = \mathcal{C}^{\infty}(M, \mathbb{R})$. Then A is an abelian group and by [13, Eq (8)] S is a subsemigroup of $\mathcal{C}^{\infty}(M, M)$ acting from the right on A as follows:

$$\alpha \circ s : M \xrightarrow{s} M \xrightarrow{\alpha} \mathbb{R}, \qquad \alpha \in A, \, s \in S.$$

Suppose also that the shift map φ of **F** is injective, i.e., $\varphi : \mathcal{C}^{\infty}(M, \mathbb{R}) \to Sh(\mathbf{F})$ is a bijection. Then it follows from [13, Eq (8)] that the inverse map

$$\sigma = \varphi^{-1} : Sh(\mathbf{F}) \to \mathcal{C}^{\infty}(M, \mathbb{R})$$

is a crossed homomorphism, i.e.

$$\sigma(f \circ g) = \sigma(f) \circ g + \sigma(g), \qquad \forall f, g \in S.$$

If φ is not injective, then the local inverse of φ near id_M is a "local crossed homomorphism". Moreover, if \mathbf{F} is not a global flow, then $\mathsf{func}(\mathbf{F})$ is an open subset of $\mathcal{C}^{\infty}(M,\mathbb{R})$ containing the zero function, i.e. a "local group".

Notice that if $Sh(\mathbf{F})$ were a group and φ^{-1} were a homomorphism, then continuity of φ^{-1} would be equivalent to its continuity at the unit element id_M only, which of course is a simpler problem. In our case $Sh(\mathbf{F})$ is just a semigroup, φ^{-1} is a local crossed homomorphism, and $\mathsf{func}(\mathbf{F}, V)$ is in general a local group. Nevertheless we will now show that continuity of local inverses of φ is equivalent to continuity of the local inverse of φ at $i_V : V \subset M$.

Theorem 4.2. Let $V \subset M$ be a D-submanifold and $r, s \in \overline{\mathbb{N}}_0$. Then the following conditions are equivalent:

- (1) The shift map φ_V : func(**F**, V) \rightarrow Sh(**F**, V) is S^{r,s}-open;
- (2) For every $\alpha \in \mathsf{func}(\mathbf{F}, V)$ there exists an S^s -neighbourhood \mathcal{N}_f of $f = \varphi_V(\alpha)$ in $Sh(\mathbf{F}, V)$ and an $S^{s,r}$ -continuous section of φ_V , *i.e.* a map

$$\sigma: \mathcal{N}_f \longrightarrow \mathcal{C}^{\infty}(V, \mathbb{R}),$$

such that $\sigma(f) = \alpha$ and $\varphi_V \circ \sigma = id(\mathcal{N}_f)$. In other words

$$f(x) = \mathbf{F}(x, \sigma(f)(x)),$$

for all $g \in \mathcal{N}_f$ and $x \in V$.

(3) Property (2) holds for the zero function $\alpha = \mathbf{0} : V \to \mathbb{R}$ and the identity inclusion $f = \varphi_V(\mathbf{0}) = i_V : V \subset M$.

Proof. (1) \Rightarrow (2). Suppose that φ_V : func(\mathbf{F}, V) $\rightarrow Sh(\mathbf{F}, V)$ is $S^{r,s}$ open. Since φ_V is locally injective with respect to the S^0 topology, condition (1) means that for every $\alpha \in \mathsf{func}(\mathbf{F}, V)$ there exists an S^r open neighbourhood \mathcal{M}_{α} in func(**F**, V) such that

- a) the restriction $\varphi_V|_{\mathcal{M}_{\alpha}} : \mathcal{M}_{\alpha} \to \varphi_V(\mathcal{M}_{\alpha})$ is a bijection,
- b) $\varphi_V(\mathcal{M}_\alpha)$ is S^s-open in $Sh(\mathbf{F}, V)$, so $\varphi_V(\mathcal{M}_\alpha) = Sh(\mathbf{F}, V) \cap \mathcal{N}_f$, for some S^s -open \mathcal{N}_f neighbourhood of $f = \varphi_V(\alpha)$ in $\mathcal{C}^{\infty}(V, M)$, c) the inverse map $\sigma_V = \varphi_V^{-1} : \varphi_V(\mathcal{M}_\alpha) \to \mathcal{M}_\alpha$ is $S^{s,r}$ -continuous.

Then $\mathcal{N}_f \stackrel{\text{def}}{=} \varphi_V(\mathcal{M}_\alpha) = Sh(\mathbf{F}, V) \cap \mathcal{N}_f$ and σ_V satisfy condition (2). The implication $(2) \Rightarrow (3)$ is evident.

15

 $(3) \Rightarrow (1)$. It suffices to show that for every $\alpha \in \mathsf{func}(\mathbf{F}, V)$ there exists an \mathbf{S}^r -open neighbourhood \mathcal{M}_{α} in $\mathsf{func}(\mathbf{F}, V)$ satisfying conditions a)c) above. Condition (3) means that such a neighbourhood \mathcal{M}_0 exists for the zero function **0**.

For every $\alpha \in \mathcal{C}^{\infty}(V, \mathbb{R})$ define the following subset of $\mathcal{C}^{\infty}(V, M)$

$$\mathcal{U}_{\alpha} = \{ f \in \mathcal{C}^{\infty}(V, M) : (f(x), \alpha(x)) \in \mathsf{dom}(\mathbf{F}), \forall x \in V \}$$

and consider a map $q_{\alpha} : \mathcal{U}_{\alpha} \to \mathcal{C}^{\infty}(V, M)$ defined by

$$q_{\alpha}(f)(x) = \mathbf{F}(f(x), \alpha(x)), \qquad f \in \mathcal{U}_{\alpha}, \ x \in V.$$

It is easy to see that \mathcal{U}_{α} is S^{0} -open in $\mathcal{C}^{\infty}(V, M)$. Also notice that, q_{α} is $S^{r,r}$ -continuous for all $r \in \overline{\mathbb{N}}_{0}$ and that $i_{V} \in \mathcal{U}_{\alpha}$ if and only if $\alpha \in \mathsf{func}(\mathbf{F}, V)$.

Lemma 4.3. The image of q_{α} coincides with $\mathcal{U}_{-\alpha}$ and $q_{-\alpha}$ is its inverse, i.e., $q_{-\alpha} = q_{\alpha}^{-1} : \mathcal{U}_{-\alpha} \to \mathcal{U}_{\alpha}$. Thus q_{α} is an $S^{r,r}$ -homeomorphism ($\forall r \in \overline{\mathbb{N}}_0$) between the S^0 -open sets \mathcal{U}_{α} and $\mathcal{U}_{-\alpha}$.

Proof. Let
$$f \in \mathcal{U}_{\alpha}$$
, i.e. $(f(x), \alpha(x)) \in \mathsf{dom}(\mathbf{F})$ for all $x \in V$. Then
 $(q_{\alpha}(f)(x), -\alpha(x)) = (\mathbf{F}(f(x), \alpha(x)), -\alpha(x)) \in \mathsf{dom}(\mathbf{F}),$

 $q_{-\alpha} \circ q_{\alpha}(f)(x) = \mathbf{F} \big(\mathbf{F} \big(f(x), \alpha(x) \big), -\alpha(x) \big) = \mathbf{F} \big(f(x), \alpha(x) - \alpha(x) \big) = f(x).$ Hence $q_{\alpha}(\mathcal{U}_{\alpha}) \subset \mathcal{U}_{-\alpha}$ and $q_{-\alpha} \circ q_{\alpha} = \mathrm{id}_{\mathcal{U}_{\alpha}}.$ Interchanging α and $-\alpha$ will give the result. \Box

Now we turn to the proof of Theorem 4.2. For every $\alpha \in \mathsf{func}(\mathbf{F}, V)$ set

$$\mathcal{M}_{\alpha} \stackrel{\text{def}}{=} \left(\mathcal{M}_{0} \cap \varphi_{V}^{-1}(\mathcal{U}_{\alpha}) + \alpha \right) \cap \operatorname{func}(\mathbf{F}, V).$$

Then \mathcal{M}_{α} is S^r-open in func(**F**, V).

Moreover $\alpha \in \mathcal{M}_{\alpha}$. Indeed, since $\alpha \in \mathsf{func}(\mathbf{F}, V)$, we have that $i_V \in \mathcal{U}_{\alpha}$, whence $\mathbf{0} \in \varphi_V^{-1}(i_V) \subset \varphi_V^{-1}(\mathcal{U})$ and thus $\alpha \in \mathcal{M}_{\alpha}$.

We will show that \mathcal{M}_{α} satisfies the conditions a)-c) above.

a) The restriction of φ_V to \mathcal{M}_{α} is 1-1, since so is $\varphi_V|_{\mathcal{M}_0}$.

Conditions b) and c) are implied by the following lemma:

Lemma 4.4. Let $\mathcal{M}'_{\alpha} \subset \mathcal{M}_{\alpha}$ be any S^r -open subset. Then $\varphi_V(\mathcal{M}'_{\alpha})$ is S^s -open in $Sh(\mathbf{F}, V)$.

Proof. Denote $\mathcal{M}'_0 = \mathcal{M}'_\alpha - \alpha$. Then $\mathcal{M}'_0 \subset \mathcal{M}_0 \cap \varphi_V^{-1}(\mathcal{U}_\alpha)$ and by the condition b) for \mathcal{M}_0 there exists an S^s-open subset \mathcal{N}' of $\mathcal{C}^{\infty}(V, M)$ such that

 $\varphi_V(\mathcal{M}'_0) = Sh(\mathbf{F}, V) \cap \mathcal{N}'.$

Define the following $S^{r,r}$ -homeomorphism

 $a_{\alpha}: \mathcal{C}^{\infty}(V, \mathbb{R}) \to \mathcal{C}^{\infty}(V, \mathbb{R}), \qquad a_{\alpha}(\beta) = \alpha + \beta.$

Then we obtain the following commutative diagram:

(4.1)
$$\begin{array}{ccc} \mathcal{M}'_{0} & \stackrel{\varphi_{V}}{\longrightarrow} & \varphi_{V}(\mathcal{M}'_{0}) = Sh(\mathbf{F}, V) \cap \mathcal{N}' \cap \mathcal{U}_{\alpha} \\ & & & \downarrow^{q_{\alpha}} \\ & & \mathcal{M}'_{\alpha} & \stackrel{\varphi_{V}}{\longrightarrow} & Sh(\mathbf{F}, V) \cap q_{\alpha}(\mathcal{N}' \cap \mathcal{U}_{\alpha}) \end{array}$$

simply meaning that $\mathbf{F}(x,\beta(x) + \alpha(x)) = \mathbf{F}(\mathbf{F}(x,\beta(x)),\alpha(x))$ for all $\beta \in \mathcal{M}_0$ and $x \in V$. We claim that

$$\varphi_V(\mathcal{M}'_{\alpha}) = Sh(\mathbf{F}, V) \cap q_{\alpha}(\mathcal{N}' \cap \mathcal{U}_{\alpha}).$$

Indeed, by (4.1), $\varphi_V(\mathcal{M}'_{\alpha}) \subset Sh(\mathbf{F}, V) \cap q_{\alpha}(\mathcal{N}' \cap \mathcal{U}_{\alpha}).$

Conversely, let $g \in Sh(\mathbf{F}, V) \cap q_{\alpha}(\mathcal{N}' \cap \mathcal{U}_{\alpha})$, i.e., there exist $\beta \in func(\mathbf{F}, V)$ and $f \in \mathcal{N}' \cap \mathcal{U}_{\alpha}$ such that

(4.2)
$$g(x) = \mathbf{F}(x, \beta(x)) = \mathbf{F}(f(x), \alpha(x)), \quad \forall x \in V.$$

Then $f(x) = \mathbf{F}(x, \beta(x) - \alpha(x))$, i.e. $f \in Sh(\mathbf{F}, V)$ and therefore

$$f = \varphi_V(\beta - \alpha) \in Sh(\mathbf{F}, V) \cap \mathcal{N}' \cap \mathcal{U}_\alpha = \varphi_V(\mathcal{M}'_0).$$

Thus there exists $\gamma \in \mathcal{M}'_0$, possibly distinct from $\beta - \alpha$, such that $f = \varphi_V(\gamma)$. Denote $\beta' = \gamma + \alpha$. Then $\beta' \in \mathcal{M}'_{\alpha}$ and

$$g(x) = \mathbf{F}(f(x), \alpha(x)) = \mathbf{F}\big(\mathbf{F}(x, \gamma(x)), \alpha(x)\big) = \mathbf{F}(x, \beta'(x)).$$

In other words, $g = \varphi_V(\beta') \in \varphi_V(\mathcal{M}'_{\alpha})$. Lemma 4.4 and Theorem 4.2 are completed.

5. Examples when shift map is not open

In this section we discuss four examples of well-known flows whose shift maps turn out not to be homeomorphisms onto their images. They have similar nature, but are given on different types of manifolds. These example provide counterexamples to [13, Th. 1]. All manifolds in this section are compact, therefore we will not distinguish weak and strong topologies. In all the examples below our vector fields will satisfy the assumptions of the following simple lemma:

Lemma 5.1. Let F be a vector field on a compact manifold M tangent to ∂M . Suppose that the shift map φ of F is injective and for every $r \in \mathbb{N}_0$, a \mathbb{W}^r -neighbourhood \mathcal{N} of id_M , and arbitrary large T > 0 there exists $t \in \mathbb{R}$ regarded as a constant function $t : M \to \mathbb{R}$ such that |t| > T and $\mathbf{F}_t = \varphi(t) \in \mathcal{N}$. Then φ is not $\mathbb{W}^{r,s}$ -open for any $r, s \in \overline{\mathbb{N}}_0$. In particular, φ is not a homeomorphism onto its image with respect to \mathbb{W}^{∞} topologies of $\mathcal{C}^{\infty}(M, \mathbb{R})$ and $Sh(\mathbf{F})$. *Proof.* Consider the following W^0 -neighbourhood of the zero function:

$$\mathcal{M}_0 = \{ \alpha \in \mathcal{C}^\infty(M, \mathbb{R}) : |\alpha(x)| < 1 \}.$$

Suppose that there is a W^r-neighbourhood \mathcal{N} of id_M such that $\mathcal{N} \subset \varphi(\mathcal{M}_0)$. By assumption there exists a constant function t > 1 such that $\varphi(t) \in \mathcal{N}$. Since φ is injective, and $t \notin \mathcal{M}_0$, we obtain that $\varphi(t) \notin \varphi(\mathcal{M}_0)$, whence $\mathcal{N} \not\subset \varphi(\mathcal{M}_0)$. This contradiction implies that φ is not W^{r,s}-open for any $r, s \in \overline{\mathbb{N}}_0$.

Irrational flow. For simplicity we will consider the irrational flow on T^2 . Let $\mu \in \mathbb{R}$ be an irrational number and **F** be the *irrational* flow on the 2-torus $T^2 = \mathbb{R}^2/\mathbb{Z}^2$ given by $\mathbf{F}(x, y, t) = (x + t, y + t/\mu)$.

Lemma 5.2. The shift map $\varphi : \mathcal{C}^{\infty}(T^2, \mathbb{R}) \to \mathcal{C}^{\infty}(T^2, T^2)$ of F is not $W^{r,s}$ -open for any $r, s \in \overline{\mathbb{N}}_0$.

Proof. Notice that every orbit of **F** is non-closed and everywhere dense, so φ is injective. First we give convenient formulas for the metrics generating W^r topologies on $\mathcal{C}^{\infty}(T^2, T^2)$. Let $f: T^2 \to T^2$ be a \mathcal{C}^{∞} map. Then f lifts to some \mathbb{Z}^2 -equivariant map $\tilde{f} = (\tilde{f}_1, \tilde{f}_2) : \mathbb{R}^2 \to \mathbb{R}^2$. Let $I^2 = [0, 1] \times [0, 1] \subset \mathbb{R}^2$ be the fundamental domain for the covering map $p: \mathbb{R}^2 \to T^2$. Define the r-th norm of f by

$$||f||^{r} = \sum_{j=1,2} \left(\sup_{(x,y) \in I^{2}} \min(\{|\tilde{f}_{j}|\}, 1-\{|\tilde{f}_{j}|\}) + \sum_{1 \le i_{1}+i_{2} \le r} \sup_{(x,y) \in I^{2}} \left| \frac{\partial^{i_{1}+i_{2}} \tilde{f}_{j}}{\partial x^{i_{1}} \partial y^{i_{2}}} \right| \right),$$

where $\{|t|\}$ is the fractional part of the absolute value of $t \in \mathbb{R}$.

Let $\varepsilon > 0$ and $\mathcal{N}_{\varepsilon}^{r} = \{f \in \mathcal{C}^{\infty}(T^{2}, T^{2}) : \|f - \mathrm{id}_{T^{2}}\|^{r} < \varepsilon\}$ be a base W^r-neighbourhood of $\mathrm{id}_{T^{2}} = \varphi(\mathbf{0}) \in \mathcal{C}^{\infty}(T^{2}, T^{2})$ for some $\varepsilon > 0$. We will show that there exists arbitrary large (by absolute value) $n \in \mathbb{Z}$ such that $\varphi(n\mu) = \mathbf{F}_{n\mu} \in \mathcal{N}_{\varepsilon}^{r}$. Then our lemma will follow from Lemma 5.1.

Notice that $\mathbf{F}_{n\mu}(x, y) = (x + n\mu, y + n) \equiv (x + n\mu, y)$ for all $n \in \mathbb{Z}$, i.e., $\mathbf{F}_{n\mu}$ is just a "rotation along the first coordinate". Since this map is defined by adding constants to coordinates, it follows from formula for $||f||^r$ that for each $r \in \mathbb{N}_0$ the distance between id_{T^2} and $\mathbf{F}_{n\mu}$ with respect to the \mathbf{W}^r topology is equal to

$$\|\mathbf{F}_{n\mu} - \mathrm{id}_{T^2}\|^r = \min(\{|n\mu|\}, 1 - \{|n\mu|\})$$

and therefore does not depend on r.

Since μ is irrational, the set $T_{\mu} = {\min(\{|n\mu|\}, 1 - \{|n\mu|\})}_{n \in \mathbb{Z}}$ is everywhere dense in $S^1 = \mathbb{R}/\mathbb{Z}$. Hence there are arbitrary large (by absolute value) $n \in \mathbb{Z}$ such that $\|\mathbf{F}_{n\mu} - \mathrm{id}_{T^2}\|^r < \varepsilon$, i.e. $\mathbf{F}_{n\mu} \in \mathcal{N}_{\varepsilon}^r$. \Box

Irrational flows on a solid torus. Let $D^2 = \{z \in \mathbb{C} : |z| \le 1\}$ be the unit disk in the complex plane, $T = S^1 \times D^2$ be the solid

SERGIY MAKSYMENKO

torus, and μ be an irrational number. Define the following flow on T by $\mathbf{F}(\phi, z, t) = (\phi + t \mod 1, e^{2\pi t/\mu}z)$. Then by the arguments similar to Lemma 5.2 is can be shown that F satisfies assumptions of Lemma 5.1.

The main feature of this example is that **F** has periodic orbit $\gamma = S^1 \times 0$ and all other ones are recurrent. Let $B = 1 \times D^2$ be and $R: B \to B$ be the first recurrence map of γ defined by $R(z) = e^{2\pi/\mu}z$, i.e. it is the rotation by $\frac{2\pi}{\mu}$. Then R is not periodic, eigen values of its tangent map T_0R at $0 \in B$ have modulus 1, and the iterations of R can be arbitrary close to id_B . Thus **F** satisfies all assumptions but (b) of Definition 3.4 of class $\mathcal{F}(T^2)$.

Periodic linear flows. Given $\lambda_1, \ldots, \lambda_n \in \mathbb{R}$ define the following linear flow on $\mathbb{R}^{2n} = \mathbb{C}^n$ by

(5.1)
$$\mathbf{F}(z_1,\ldots,z_n,t) = (e^{i\lambda_1 t} z_1,\ldots,e^{i\lambda_n t} z_n).$$

Evidently, the closed 2n-disk D_r of radius r and centered at the origin is invariant with respect to **F**.

Lemma 5.3. The following conditions are equivalent:

- (1) $\mathbf{F}_{\tau} = \mathrm{id}_{\mathbb{C}^n}$ for some $\tau > 0$, i.e. $\lambda_j \tau \in \mathbb{Z}$ for all $j = 1, \ldots, n$;
- (2) every $z \in \mathbb{C}^n$ is either fixed or periodic with respect to \mathbf{F} ;
- (3) at least one point $z = (z_1, \ldots, z_n)$ with all non-zero coordinates is periodic.

A flow satisfying one of these conditions will be called **periodic**.

Proof. The implications $(1) \Rightarrow (2) \Rightarrow (3)$ are evident. $(3) \Rightarrow (1)$ Let $z \in \mathbb{C}^n$ be a point with all non-zero coordinates and θ be the period of z with respect to **F**. Then $e^{i\lambda_j\theta}z_j = z_j \neq 0$ for all $j = 1, \ldots, n$, whence $\lambda_j \cdot \theta/2\pi \in \mathbb{Z}$.

Lemma 5.4. If **F** is not periodic then its shift map φ is injective and is not $W^{r,s}$ -open for any $r, s \in \overline{\mathbb{N}}_0$.

Proof. By Lemma 5.3 every point $z \in \mathbb{C}^n$ with all non-zero coordinates is non-periodic and it is easy to see that its orbit is dense on some open subset of the sphere $S_{|x|}^{2n-1} = \partial D_{|x|}$ of radius |x|. In particular, this orbit is non-closed, whence the shift map of \mathbf{F} is injective. Then similarly to Lemma 5.2, we can find arbitrary large $t \in \mathbb{R}$ such that \mathbf{F}_t is arbitrary close to $\mathrm{id}_{\mathbb{R}^{2n}}$ in any of \mathbf{W}^r topologies. \Box

Flow on S^{2n} , $n \ge 2$. We will now extend the last result for the construction of a flow on the sphere S^{2n} with two fixed points at which this flow is linear.

Let \mathbb{R}_1^{2n} and \mathbb{R}_2^{2n} be two copies of $\mathbb{R}^{2n} = \mathbb{C}^n$. Define a diffeomorphism $\eta : \mathbb{R}_1^{2n} \setminus \{0\} \to \mathbb{R}_2^{2n} \setminus \{0\}$ by $\eta(z) = \frac{z}{\|z\|^2}$, where $\|z\|$ is the usual

Euclidean norm in \mathbb{R}^{2n} . Then η maps every sphere of radius r centered at 0 to the sphere of radius 1/r. Gluing \mathbb{R}^{2n}_1 and \mathbb{R}^{2n}_2 via η we obtain a 2*n*-sphere.

Notice that Eq. (5.1) defines the flows on \mathbb{R}_1^{2n} and \mathbb{R}_2^{2n} so that the following diagram is commutative:

Hence these flows determine a unique flow \mathbf{F}' on S^{2n} with two fixed points. Moreover, \mathbf{F}' is *linear* on the charts \mathbb{R}^{2n}_1 and \mathbb{R}^{2n}_2 at these points.

Now suppose that \mathbf{F} is not periodic. Then we can find arbitrary large $t \in \mathbb{R}$ such that \mathbf{F}_t is arbitrary close to $\mathrm{id}_{\mathbb{R}^{2n}}$ in any of W^r topologies. This implies that \mathbf{F}' satisfies assumptions of Lemma 5.1.

6. Regular and trivial extensions

The results of this section will allow to estimate continuity of shift maps for vector fields of types (L) and (H).

Let M, N be two manifolds, G be a vector field on M, F be some regular extension and \overline{G} be the trivial extension of G on $M \times N$. Thus

$$F(x,y) = (G(x), H(x,y)), \qquad \overline{G}(x,y) = (G(x), 0),$$

for some smooth map $H : M \times N \to TN$. By **F**, $\bar{\mathbf{G}}$, and **G** we will denote the corresponding local flows, and by φ , $\bar{\psi}$, and ψ the corresponding shift maps of F, \bar{G} , and G. It is easy to see that

$$(6.1) \quad \mathsf{dom}(\mathbf{F}) \subset \mathsf{dom}(\mathbf{G}) = \mathsf{dom}(\mathbf{G}) \times N.$$

Moreover,

(6.2)
$$\mathbf{F}(x,y,t) = (\mathbf{G}(x,t),\mathbf{H}(x,y,t)), \quad \bar{\mathbf{G}}(x,y,t) = (\mathbf{G}(x,t),y),$$

for some smooth map $\mathbf{H} : \mathsf{dom}(\bar{\mathbf{G}}) \to N$.

Let $V \subset M \times N$ be a connected compact *D*-submanifold. Then it follows from (6.1) that $\operatorname{func}(\mathbf{F}, V)$ is W⁰-open in $\operatorname{func}(\bar{\mathbf{G}}, V)$. Define the map

$$P: Sh(\mathbf{F}, V) \to Sh(\bar{\mathbf{G}}, V)$$

by the rule: if $f \in Sh(\mathbf{F}, V)$, and f(x, y) = (A(x, y), g(x, y)), then P(f)(x, y) = (A(x, y), y). It follows from (6.2) that P is well defined

and is $W^{r,r}$ -continuous for every $r \in \mathbb{N}_0$. Moreover, we have the following commutative diagram:

(6.3)
$$\begin{aligned} & \operatorname{func}(\mathbf{F}, V) \xrightarrow{\varphi_V} Sh(\mathbf{F}, V) \\ & \downarrow & \downarrow_V \\ & \operatorname{func}(\bar{\mathbf{G}}, V) \xrightarrow{\bar{\psi}_V} Sh(\bar{\mathbf{G}}, V) \end{aligned}$$

Theorem 6.1. Suppose that Σ_G is nowhere dense in M and that $\overline{\psi}_V$ is $W^{r,s}$ -open for some $r, s \in \overline{\mathbb{N}}_0$.

- (1) If $\bar{\psi}_V$ is injective, then φ_V is $\mathsf{W}^{r,s}$ -open as well.
- (2) If φ_V is $W^{r,s}$ -open, then P is locally injective with respect to the W^s topology of $Sh(\mathbf{F}, V)$.

Suppose in addition that $\overline{\psi}_V$ is also $\mathsf{W}^{s,t}$ -open for some $t \in \overline{\mathbb{N}}_0$.

- (3) If P is locally injective with respect to the W^s topology of $Sh(\mathbf{F}, V)$, then φ_V is W^{r,t}-open.
- (4) If both ψ_V and φ_V are not injective, then P is locally injective with respect to the W⁰ topology of Sh(**F**, V), whence, by (3), φ_V is W^{r,t}-open.

Proof. (1) If $\bar{\psi}_V$: func(\mathbf{G}, V) \to $Sh(\mathbf{G}, V)$ is a bijection, it follows from (6.3) that so is φ_V . Let $\mathcal{M} \subset \mathsf{func}(\mathbf{F}, V)$ be a W^r-open subset. Then

$$\varphi_V(\mathcal{M}) = P^{-1} \circ \bar{\psi}_V(\mathcal{M})$$

is W^s -open in $Sh(\mathbf{F}, V)$ due to $W^{r,s}$ -openness of $\overline{\psi}_V$ and $W^{s,s}$ -continuity of P.

(2) Let $\alpha \in \operatorname{func}(\mathbf{F}, V)$ and $f = \varphi_V(\alpha)$. We will find a W^s-neighbourhood of f such that $P|_{\mathcal{N}}$ is 1-1.

Since both φ_V and $\bar{\psi}_V$ are $\mathsf{W}^{r,s}$ -open and locally injective with respect to W^0 topologies of $\mathsf{func}(\mathbf{F}, V)$ and $\mathsf{func}(\bar{\mathbf{G}}, V)$ respectively, there exists a W^r -neighbourhood \mathcal{M} of α in $\mathsf{func}(\mathbf{F}, V)$ such that $\mathcal{N} = \varphi_V(\mathcal{M})$ is a W^s -neighbourhood of f in $Sh(\mathbf{F}, V)$ and the restrictions of φ_V and $\bar{\psi}_V$ to \mathcal{M} are 1-1. Then it follows from (6.3) that $P|_{\mathcal{N}}$ is 1-1 as well.

(3) Let $\mathcal{M} \subset \mathsf{func}(\mathbf{F}, V)$ be a W^r -open subset and $\alpha \in \mathcal{M}$. We will show that there exists a W^s -neighbourhood $\mathcal{M}' \subset \mathcal{M}$ of α such that $\varphi_V(\mathcal{M}')$ is W^t -open in $Sh(\mathbf{F}, V)$. Since $\alpha \in \mathcal{M}$ is arbitrary, we will obtain that $\varphi_V(\mathcal{M})$ is W^t -open in $Sh(\mathbf{F}, V)$.

It follows from $W^{r,s}$ -openness of $\bar{\psi}_V|_{\mathcal{M}}$ and $W^{s,s}$ -continuity of P that the set $P^{-1} \circ \bar{\psi}_V(\mathcal{M})$ is a W^s -open neighbourhood of $\varphi_V(\alpha)$ in $Sh(\mathbf{F}, V)$. Moreover, since P is locally injective with respect to the W^s topology of $Sh(\mathbf{F}, V)$, there exists a W^s-neighbourhood $\mathcal{N} \subset P^{-1} \circ \bar{\psi}_V(\mathcal{M})$ of $\varphi_V(\alpha)$ such that the restriction $P|_{\mathcal{N}} : \mathcal{N} \to Sh(\bar{\mathbf{G}}, V)$ is injective.

Denote $\mathcal{M}' = \mathcal{M} \cap \varphi_V^{-1}(\mathcal{N})$. We claim that

$$\varphi_V(\mathcal{M}') = P^{-1} \circ \bar{\psi}_V(\mathcal{M}') \cap \mathcal{N},$$

whence $\varphi_V(\mathcal{M}')$ will be W^t -open in $Sh(\mathbf{F}, V)$. Indeed,

$$P^{-1} \circ \bar{\psi}_V(\mathcal{M}') \cap \mathcal{N} \xrightarrow{(6.3)} P^{-1} \circ P \circ \varphi_V(\mathcal{M}') \cap \mathcal{N} \xrightarrow{\text{injectivity of } P|_{\mathcal{N}}} \varphi_V(\mathcal{M}')$$

(4) Suppose that both $\overline{\psi}_V$ and φ_V are not injective. We will show that for some m > 0 there exists a free *P*-equivariant action of the group \mathbb{Z}_m on $Sh(\mathbf{F}, V)$ such that P(f) = P(g) iff f, g belongs to the same \mathbb{Z}_m -orbit. This will give us a decomposition $P : Sh(\mathbf{F}, V) \xrightarrow{q} Sh(\mathbf{F}, V)/\mathbb{Z}_m \xrightarrow{\sigma} Sh(\mathbf{\bar{G}}, V)$ in which q is a covering map, and σ is a bijection with the image $P(Sh(\mathbf{F}, V)) \subset Sh(\mathbf{\bar{G}}, V)$. Then q will be locally injective with respect to the W^0 topology since \mathbb{Z}_m is a finite group. Hence so will be P.

By Lemma 2.3(5b) func($\mathbf{\bar{G}}, V$) = func(\mathbf{F}, V) = $\mathcal{C}^{\infty}(V, \mathbb{R})$, ker($\bar{\psi}_V$) = $\{n\bar{\eta}\}_{n\in\mathbb{Z}}$, and ker(φ_V) = $\{n\eta\}_{n\in\mathbb{Z}}$ for some smooth functions $\bar{\eta}, \eta$: $M \times N \to (0, \infty)$ such that

$$\overline{\mathbf{G}}(x, y, t + \overline{\eta}(x, y)) \equiv \overline{\mathbf{G}}(x, y, t), \qquad \mathbf{F}(x, y, t + \eta(x, y)) \equiv \mathbf{F}(x, y, t).$$

In other words

(6.4)
$$(\mathbf{G}(x,t+\bar{\eta}),y) = (\mathbf{G}(x,t),y), \\ (\mathbf{G}(x,t+\eta),\mathbf{H}(x,y,t+\eta)) = (\mathbf{G}(x,t),\mathbf{H}(x,y,t)).$$

It follows that $\overline{\mathbf{G}}(x, y, t+\eta) = (\mathbf{G}(x, t+\eta), y) = (\mathbf{G}(x, t), y) = \overline{\mathbf{G}}(x, y, t)$, whence $\eta \in \ker(\overline{\psi}_V)$, i.e. $\eta = m\overline{\eta}$ for some $m \in \mathbb{Z}$.

In particular, by (2) of Lemma 2.3, $\bar{\eta}$ is constant along orbits of F. Since for every $\alpha \in \mathsf{func}(\mathbf{F}, V)$ the map $\varphi_V(\alpha)$ preserves every orbit of F, we have that $\bar{\eta} \circ \varphi_V(\alpha) = \bar{\eta}$, whence

$$\varphi_V(\bar{\eta}) \circ \varphi_V(\alpha) \xrightarrow{[13, \text{ Eq. (8)}]} \varphi_V(\alpha + \bar{\eta} \circ \varphi_V(\alpha)) = \varphi_V(\alpha + \bar{\eta}).$$

Then we can define an action * of the group \mathbb{Z}_m on $Sh(\mathbf{F}, V)$ by

$$k * f = \varphi_V(k\bar{\eta}) \circ f, \qquad k \in \mathbb{Z}_m, \ f \in Sh(\mathbf{F}, V).$$

Evidently this action is free and by (6.4) is equivariant with respect to P.

Moreover, let $f = \varphi_V(\alpha)$, $g = \varphi_V(\beta)$ for some $\alpha, \beta \in \text{func}(\mathbf{F}, V)$ and suppose that P(f) = P(g), i.e. $\bar{\psi}_V(\alpha) = \bar{\psi}_V(\beta)$. Then it follows from (3) of Lemma 2.3 that $\alpha - \beta = k\bar{\eta}$ for some $k \in \mathbb{Z}$, whence $f = \varphi_V(k\bar{\eta}) \circ g$. Moreover, since $\eta = m\bar{\eta}$, we may take k modulo m. In other words, P(f) = P(g) iff f = k * g for some $k \in \mathbb{Z}_m$. \Box

SERGIY MAKSYMENKO

7. VECTOR FIELDS OF TYPES (L) AND (H)

The following lemma shows that property of openness of shift maps near singular points is invariant under reparametrizations.

Lemma 7.1. c.f. [20]. Let F be a \mathcal{C}^{∞} vector field on M, $\nu : M \to (0, +\infty)$ a \mathcal{C}^{∞} strictly positive function, and $G = \nu F$. Let also $V \subset M$ be a D-submanifold and φ_V and ψ_V be shift maps for F and G respectively. Then $Sh(\mathbf{F}, V) = Sh(\mathbf{G}, V)$. Moreover, φ_V is $\mathbf{S}^{r,s}$ -open iff so is ψ_V .

Proof. Define the functions $\alpha : \mathsf{dom}(\mathbf{F}) \to \mathbb{R}$ and $\beta : \mathsf{dom}(\mathbf{G}) \to \mathbb{R}$ by

$$\beta(x,s) = \int_{0}^{s} \nu(\mathbf{G}(x,\tau)) d\tau, \qquad \alpha(x,s) = \int_{0}^{s} \frac{d\tau}{\nu(\mathbf{F}(x,\tau))}$$

Then it is well-known that for each $f \in \mathsf{func}(\mathbf{F}, V)$ and $g \in \mathsf{func}(\mathbf{G}, V)$, see e.g.[20],

$$\mathbf{G}(x,g(x)) = \mathbf{F}(x,\beta(x,g(x))), \qquad \mathbf{F}(x,f(x)) = \mathbf{G}(x,\alpha(x,f(x))),$$

for all $x \in V$. Define the following map

 $\zeta : \operatorname{func}(\mathbf{G}, V) \to \operatorname{func}(\mathbf{F}, V), \qquad \zeta(g)(x) = \beta(x, g(x)).$

Then ζ is a homeomorphism with respect to topologies W^r for all r, and its inverse is given by $\zeta^{-1}(f)(x) = \alpha(x, f(x))$. Moreover $\psi_V = \varphi_V \circ \zeta$. Hence φ_V is $S^{r,s}$ -open iff so is ψ_V .

Thus the study of vector fields of types (L) and (H) is suffices to consider linear and reduced Hamiltonian vector fields only.

Lemma 7.2. [19, 14]. Let $g : \mathbb{R}^2 \to \mathbb{R}$ be a homogeneous polynomial, G be its reduced Hamiltonian vector field, F be a trivial n-extension of G on $\mathbb{R}^n \times \mathbb{R}^2$, and V be a D-neighbourhood of $0 \in \mathbb{R}^{n+2}$. Then the shift map φ_V of F is $\mathbb{W}^{\infty,\infty}$ -open. If deg $G \ge 2$, i.e. G is of type (H), then for every regular n-extension of G its shift map is $\mathbb{W}^{\infty,\infty}$ -open as well.

Proof. It follows from [19, 14] that the shift map $\bar{\psi}_V$ of \bar{G} has a $W^{\infty,\infty}$ continuous local section on some neighbourhood of the identity inclusion $i_V : V \subset \mathbb{R}^{n+2}$, see also [19, Th. 11.1]. By Theorem 4.2 this implies $W^{\infty,\infty}$ -openness of ψ_V .

Moreover, if deg $G \ge 2$, i.e. the linear part of F at 0 vanishes, then by Lemma 2.3 φ_V is injective. Hence by (1) of Theorem 6.1 the shift map of G is $W^{\infty,\infty}$ -open as well. **Lemma** 7.3. Let B be a non-zero $(k \times k)$ -matrix, G(y) = By be the corresponding linear vector field on \mathbb{R}^k , $\mathbf{G}(y,t) = e^{Bt}y$ be its flow, and F be a vector field on \mathbb{R}^{n+k} being a regular (possibly trivial) n-extension of G with respect to the origins of \mathbb{R}^{n+k} and \mathbb{R}^k as indicated in the third column of the table below. Let also $V \subset \mathbb{R}^{n+k}$ be a D-neighbourhood of the origin $0 \in \mathbb{R}^{n+k}$. Then the shift map φ_V of F is $\mathbb{W}^{r,s}$ -open for the values r, s described in the following table.

-		_		<u>^</u>
	type	B	F	openness of φ_V
1	(L1)	$B = \ a\ , \ a \neq 0,$	regular	$W^{r,r+1}, r \ge 0$
2	(L1)	$B = \left\ \begin{smallmatrix} a & -b \\ b & a \end{smallmatrix} \right\ , \ a, b \neq 0$	regular	$W^{r,r+2}, r \ge 0$
3	(L2)	$B = \left\ \begin{smallmatrix} 0 & 1 \\ 0 & 0 \end{smallmatrix} \right\ ,$	regular	$W^{r,r+1}, r \ge 0$
4	(L3)	$B = \begin{vmatrix} 0 & -b & 1 & 0 \\ b & 0 & 0 & 1 \\ 0 & 0 & 0 & -b \\ 0 & 0 & b & 0 \end{vmatrix}, \ b \neq 0$	regular	$W^{\infty,\infty}$
5	(L4)	$B = \begin{bmatrix} 0 & -b \\ b & 0 \end{bmatrix}, \ b \neq 0$	trivial	$W^{\infty,\infty}$
6		$B = \left\ \begin{array}{ccc} 0 & -b_1 \\ b_1 & 0 \\ & \ddots \\ & & 0 & -b_l \\ & & b_l & 0 \end{array} \right\ , \ l \ge 2$		
6a	(L4)	(a) $b_j \tau \in \mathbb{Z}$ for some $\tau > 0$ and	trivial	$W^{\infty,\infty}$
		all $j = 1, \ldots, l$ (periodic case)		
6b	(L5)	(b) otherwise (non-periodic	_	—
		case)		

Proof. In the cases 1-3 we will first suppose that F is a trivial *n*-extension of G. Then F is linear and is generated by the following matrix $\| {}^{0_n}_0 {}^{0}_B \|$, where 0_n is the zero $(n \times n)$ -matrix. Let τ denote the coordinates in \mathbb{R}^n .

Case 1. In this case $F(\tau, x) = ax \frac{\partial}{\partial x}$, $(a \neq 0)$, F generates the following global flow $\mathbf{F}(\tau, x, t) = (\tau, xe^{at})$ on \mathbb{R}^{n+1} , φ_V is injective (because F has non-closed orbits), its image $Sh(\mathbf{F}, V)$ consists of maps $f = (f_1, f_2) \in \mathcal{C}^{\infty}(V, \mathbb{R}^n \times \mathbb{R})$ satisfying the following conditions:

$$f_1 \equiv \tau, \quad f_2(\tau, 0) = 0, \quad \frac{\partial f_2(\tau, 0)}{\partial x} > 0, \quad x f_2(\tau, x) > 0 \ (\forall x \neq 0),$$

and the inverse mapping $\varphi_V^{-1}: Sh(\mathbf{F}, V) \to \mathcal{C}^{\infty}(V, \mathbb{R})$ is given by

(7.1)
$$\varphi_V^{-1}(f)(\tau, x) = \frac{1}{a} \ln \frac{f_2(\tau, x)}{x} = \frac{1}{a} \ln \int_0^1 \frac{\partial f_2(\tau, tx)}{\partial x} dt,$$

see [13, Eq. (23) & (27)]. Hence φ_V^{-1} is $\mathsf{W}^{r+1,r}$ -continuous for all $r \ge 0$, whence φ_V is $\mathsf{W}^{r,r+1}$ -open.

SERGIY MAKSYMENKO

Case 2. We will regard \mathbb{R}^{n+2} as $\mathbb{R}^n \times \mathbb{C}$. Since $a \neq 0$, we have that F has non-closed orbits, whence again its shift map φ_V is injective but now the image $Sh(\mathbf{F}, V)$ can not be described so simply as in the previous cases. Let $\alpha \in \mathcal{C}^{\infty}(V, \mathbb{R})$ and $f = (f_1, f_2) = \varphi_V(\alpha) \in \mathcal{C}^{\infty}(V, \mathbb{R}^n \times \mathbb{C})$. Then $f_1 \equiv \tau$. Notice that we can define complex conjugate \bar{f}_2 and its partial derivatives $\frac{\partial \bar{f}_2}{\partial z}$ and $\frac{\partial \bar{f}_2}{\partial \bar{z}}$ in z and \bar{z} in a usual way. Then it follows from [13, Eq. (29) & Lm. 34] that

$$\alpha(\tau, z) = \frac{1}{2a} \ln \frac{\operatorname{Im}(\omega \ f_2 \ d\bar{f}_2(\bar{\omega}))}{\operatorname{Im}(\omega \ z \ \bar{\omega})} = \frac{1}{2a} \ln \frac{\operatorname{Im}(\omega \cdot f_2 \cdot (\frac{\partial \bar{f}_2}{\partial z} \omega + \frac{\partial \bar{f}_2}{\partial \bar{z}} \bar{\omega}))}{y(a^2 + b^2)}$$

and that the numerator of the last fraction is equal to zero, when y = 0. It follows from this formula and the Hadamard lemma that the expression of α via f contains partial derivatives of f up to order 2. Hence φ_V^{-1} is $W^{r+2,r}$ -continuous for all $r \geq 0$.

Hence φ_V^{-1} is $W^{r+2,r}$ -continuous for all $r \ge 0$. **Case 3.** Now $F(\tau, x, y) = y \frac{\partial}{\partial x}$ for $(\tau, x, y) \in \mathbb{R}^{n+2}$ and it generates the following flow $\mathbf{F}(\tau, x, y, t) = (\tau, x + yt, y)$. Then φ_V is again injective (since F has non-closed orbits), its image $Sh(\mathbf{F}, V)$ consists of mappings $f = (f_1, f_2, f_3) \in \mathcal{C}^{\infty}(V, \mathbb{R}^n \times \mathbb{R} \times \mathbb{R})$ such that

$$f_1 \equiv \tau, \qquad f_2(\tau, x, 0) = 0, \qquad f_3 \equiv y,$$

and the inverse map $\varphi_V^{-1}: Sh(\mathbf{F}, V) \to \mathcal{C}^{\infty}(V, \mathbb{R})$ is given by

(7.2)
$$\varphi_V^{-1}(f) = \frac{f_2(\tau, x, y) - x}{y} = \int_0^1 \frac{\partial f_2(\tau, x, ty)}{\partial y} dt,$$

see [13, Eq. (26)]. Hence φ_V^{-1} is $\mathsf{W}^{r+1,r}$ -continuous for all $r \ge 0$.

Since in the cases 1-3 φ_V is injective, it follows from (1) of Theorem 6.1 that the same estimations of continuity of φ_V^{-1} hold for regular extensions of G.

Notice that in the remaining cases 4-6 F is a regular *n*-extension of the linear vector field $G_5 = -by \frac{\partial}{\partial x} + bx \frac{\partial}{\partial y}$ defined by the matrix $B_5 = \left\| \begin{smallmatrix} 0 & -b \\ b & 0 \end{smallmatrix} \right\|$. In fact, it is easy to see that this vector field is the Hamiltonian vector field of the homogeneous polynomial $g(x,y) = \frac{b}{2}(x^2 + y^2)$. Also notice that the shift map φ_V of any *trivial* extension F of G_5 is not injective and its kernel ker (φ_V) consists of integer multiples of constant function $\frac{2\pi}{b}$.

Case 5. It follows from Lemma 7.2 that local inverses of φ_V are $W^{\infty,\infty}$ -continuous. An independent proof of $W^{\infty,\infty}$ -continuouity of φ_V for this case is also given in [21] under more general settings. Notice that in this case we claim nothing about openness of shift maps of regular extensions F of G_5 , since due to Theorem 6.1 it is necessary to have additional information about F.

25

Case 4. We will regard \mathbb{R}^4 as \mathbb{C}^2 . Suppose at first that F is a trivial *n*-extension of G. Since G is a regular 2-extension of G_5 , we see that F is a regular (n + 2)-extension of G_5 . Let also \overline{G} be a trivial (n + 2)-extension of G_5 . Thus F and \overline{G} are defined on $\mathbb{R}^n \times \mathbb{C} \times \mathbb{C}$ and generate the following global flows:

 $\mathbf{F}(\tau, z_1, z_2, t) = (\tau, e^{ibt}(z_1 + tz_2), e^{ibt}z_2), \quad \bar{\mathbf{G}}(\tau, z_1, z_2, t) = (\tau, z_1, e^{ibt}z_2).$ Denote $\theta = \frac{2\pi}{b}$ and let $V \subset \mathbb{R}^n \times \mathbb{C}^2$ be a *D*-neighbourhood of $0 \in \mathbb{R}^n \times \mathbb{C}^2$. For every $\alpha \in \mathcal{C}^{\infty}(V, \mathbb{R})$ put

$$\mathbf{f}(\alpha)(\tau, z_1, z_2) = e^{ib\,\alpha(\tau, z_1, z_2)}\,z_1, \qquad \mathbf{g}(\alpha)(\tau, z_1, z_2) = e^{ib\,\alpha(\tau, z_1, z_2)}\,z_2,$$

Then the shift maps φ_V and $\bar{\psi}_V$ of F and G respectively are given by (7,2), $(\alpha) = (\pi, f(\alpha)) + \alpha, \pi(\alpha), \pi(\alpha)) = (\pi, \pi, \pi(\alpha))$

(7.3)
$$\varphi_V(\alpha) = (\tau, \mathbf{f}(\alpha) + \alpha \cdot \mathbf{g}(\alpha), \mathbf{g}(\alpha)), \qquad \psi_V(\alpha) = (\tau, z_1, \mathbf{g}(\alpha)).$$

Similarly to (6.3) define $P : Sh(\mathbf{F}, V) \to Sh(\bar{\mathbf{G}}, V)$ by the following rule: if $h = (\tau, f, g) \in Sh(\mathbf{F}, V) \subset \mathcal{C}^{\infty}(V, \mathbb{R}^n \times \mathbb{C} \times \mathbb{C})$, then $P(h)(\tau, z_1, z_2) = (\tau, z_1, g(z_1, z_2))$. Evidently, $\bar{\psi}_V = P \circ \varphi_V$.

By the case 5 the shift map $\bar{\psi}_V : \mathcal{C}^{\infty}(V, \mathbb{R}) \to Sh(\bar{\mathbf{G}}, V)$ is $\mathsf{W}^{\infty,\infty}$ open. We claim that so is $\varphi_V : \mathcal{C}^{\infty}(V, \mathbb{R}) \to Sh(\mathbf{F}, V)$. Since G has non-closed orbits, $\bar{\psi}_V$ is always injective, whence it will follow from (1) of Theorem 6.1 that the shift map of any regular extension of G is $\mathsf{W}^{\infty,\infty}$ -open as well.

By (3) of Theorem 6.1 it suffices to show that P is locally injective with respect to the W¹ topology. Evidently, $\psi_V^{-1} \circ \psi_V(\alpha) = \{\alpha + \theta \, n\}_{n \in \mathbb{Z}}$, whence we obtain from (7.3) that

$$P^{-1} \circ \bar{\psi}_V(\alpha) = \{\varphi_V(\alpha) + (0, \theta \, n \cdot \mathbf{g}(\alpha), 0) \mid n \in \mathbb{Z}\}$$

Now, define the following C_W^1 -neighbourhood of $\varphi_V(\alpha)$ in $Sh(\mathbf{F}, V)$:

$$\mathcal{N} = \{ f \in Sh(\mathbf{F}, V) \mid \| f - \varphi_V(\alpha) \|_V^1 < |\theta|/4 \}.$$

We claim that the restriction $P|_{\mathcal{N}}$ is 1-1.

Since φ_V is injective map, it suffices to establish that whenever both $\varphi_V(\beta)$ and $\varphi_V(\beta + \theta n)$ belong to \mathcal{N} for some $\beta \in \mathcal{C}^{\infty}(V, \mathbb{R})$, then n = 0. Notice that

$$\|\mathbf{g}(\beta)\|_{V}^{1} \ge \left\|\mathbf{g}(\beta)_{z_{2}}^{\prime}\right\|_{(z_{1},z_{2})=0} = \left\|e^{ib\beta}\left(ib\,z_{2}\,\beta_{z_{2}}^{\prime}+1\right)\right\|_{(z_{1},z_{2})=0} = 1.$$

Then

$$\begin{aligned} \|\varphi_V(\beta) - \varphi_V(\beta + \theta n)\|_V^1 &\leq \|\varphi_V(\beta) - \varphi_V(\alpha)\|_V^1 + \\ &+ \|\varphi_V(\beta + \theta n) - \varphi_V(\alpha)\|_V^1 \leq |\theta|/2. \end{aligned}$$

On the other hand,

$$\|\varphi_V(\beta) - \varphi_V(\beta + \theta n)\|_V^1 = \|(0, \theta n \mathbf{g}(\beta), 0)\|_V^1 \ge |\theta n|.$$

Hence n = 0.

Case 6a. In this case F is a regular extension of G_5 . Since the flow **F** is periodic, we have that the shift maps of F and G_5 are not injective and by the case 5 the shift map of G_5 is $W^{\infty,\infty}$ -open. Then by (4) of Theorem 6.1 so is the shift map of F.

Case 6b. In this case φ_V is injective, but as it is shown in Lemma 5.4 its inverse map is not even $W^{\infty,\infty}$ -continuous.

Lemma 7.3 is completed.

REMARK 7.4. Incorrect estimations of continuity of local inverses of φ_V given in [13, pages 199-200] were bases on at the following "division lemma", which was wrongly formulated¹ in [13].

Lemma 7.5. c.f. [13, Lm. 32]. Let \mathbb{F} be either \mathbb{R} or \mathbb{C} , $V \subset \mathbb{F}$ be a D-submanifold, and $Z : \mathcal{C}^{\infty}(V, \mathbb{F}) \to \mathcal{C}^{\infty}(V, \mathbb{F})$ be a map defined by the formula: $Z(\alpha)(x) = x \cdot \alpha(x)$. If $\mathbb{F} = \mathbb{R}$ then the inverse map $Z^{-1} : \operatorname{im} Z \to \mathcal{C}^{\infty}(V, \mathbb{F})$ is $W^{r+1,r}$ -continuous for all $r \geq 0$. If $\mathbb{F} = \mathbb{C}$, then Z^{-1} is only $W^{\infty,\infty}$ -continuous.

The case $\mathbb{F} = \mathbb{R}$ easily follows from the Hadamard lemma, see also [25]. The case $\mathbb{F} = \mathbb{C}$ is more complicated and can be established by the methods of [19, 14] but during the proof of Lemma 7.3 we avoided referring to it.

8. Fragmentation

The aim of this section is to repair [13, Th. 17] by giving a sufficient condition for the shift map $\varphi : \operatorname{func}(\mathbf{F}) \to Sh(\mathbf{F})$ to be either a homeomorphism or an infinite cyclic covering map with respect to S^{∞} topologies, see Theorem 8.2.

REMARK 8.1. The error occurred in the third paragraph of [13, Th. 17], where it was claimed that "... the image $\mathcal{N}_i = \varphi_{U_i}(\mathcal{M}'_i)$ is a C_W^r neighbourhood of $f|_{U_i}$ in $C^{\infty}(U_i, M)$ for all $r \in \mathbb{N}_0$." First of all this phrase contains a misprint: instead of $C^{\infty}(U_i, M)$ the author supposed to be written im φ_{U_i} . But nevertheless the statement that \mathcal{N}_i is C_W^r open in im φ_{U_i} does not follow from the assumptions of [13, Th. 17] and must be included into the formulation of "section" property $(S)^0$ of [13, Defn. 15]. This is the point which was missed.

Theorem 8.2. c.f. [13, Th. 17]. Let F be a vector field on M such that Σ_F is nowhere dense. Suppose that there exists $a \in \mathbb{N}_0$, a function

¹In [13, Lm. 32] it was claimed that Z^{-1} is $W^{r,r}$ -continuous for all $r \in \overline{\mathbb{N}}_0$. But the latter inequality in the proof of [13, Lm. 32] actually shows $W^{r,r}$ -continuity of Z but not of its inverse.

 $d: \mathbb{N}_0 \to \mathbb{N}_0$, a locally finite cover $\{V_i\}_{i \in \Lambda}$ of M by D-submanifolds, and a finite (possibly empty) subset $\Lambda' \subset \Lambda$ such that $\{\operatorname{Int} V_i\}_{i \in \Lambda}$ is also a cover of M, and the local shift map

$$\varphi_{V_i}$$
: func(**F**, V_i) \rightarrow Sh(**F**, V_i)

is $S^{\infty,\infty}$ -open if $i \in \Lambda'$ and $S^{r,d(r)}$ -open for all $r \ge a$ if $i \in \Lambda \setminus \Lambda'$. Then the shift map

$$\varphi : \mathsf{func}(\mathbf{F}) \to Sh(\mathbf{F})$$

is $S^{\infty,\infty}$ -open. Moreover, if $\Lambda' = \emptyset$, then φ is $S^{r,d(r)}$ -open for every $r \ge a$.

Hence if $\varphi : \operatorname{func}(\mathbf{F}) \to Sh(\mathbf{F})$ is injective, then it is a homeomorphism with respect to S^{∞} topologies. Otherwise, φ is a \mathbb{Z} -covering map.

Proof. The proof follows the line of [13, Th. 17]. Since Σ_F is nowhere dense, there exists a continuous function $\delta : M \to (0, \infty)$ such that for every periodic point x of period θ_x we have that $\delta(x) < 3\theta_x$, see [13, Pr. 14].

Let $\alpha \in \mathsf{func}(\mathbf{F}), r \geq a, d^r$ be a metric on the manifold $J^r(M, \mathbb{R})$ of *r*-jets, $\nu : M \to (0, 1)$ be a strictly positive continuous function such that $\nu < \delta$, and

$$\mathcal{M} \stackrel{\text{def}}{=} \{\beta \in \mathsf{func}(\mathbf{F}) \mid d^r(j^r \alpha(x), j^r \beta(x)) < \nu(x), \forall x \in M\}$$

be a base S^r -neighbourhood of α , where $j^r \beta(x)$ denotes the *r*-jet of β at *x*. Then the restriction of φ to \mathcal{M} is injective, [13, Pr. 14]. We will show that $\varphi(\mathcal{M})$ is S^{∞} -open in $Sh(\mathbf{F})$ and if $\Lambda' = \emptyset$, then $\varphi(\mathcal{M})$ is even $S^{d(r)}$ -open in $Sh(\mathbf{F})$. This will complete Theorem 8.2.

For every $i \in \Lambda$ let

(8.1)
$$\mathcal{M}_i \stackrel{\text{def}}{=} \{\beta \in \mathsf{func}(\mathbf{F}, V_i) : d^r(j^r \alpha(x), j^r \beta(x)) < \nu(x), \forall x \in V_i\}$$

be an S^r -neighbourhood of $\alpha|_{V_i}$ in func (\mathbf{F}, V_i) . It follows from assumption about openness of φ_{V_i} that

$$\varphi_{V_i}(\mathcal{M}_i) = Sh(\mathbf{F}, V_i) \cap \mathcal{N}_i \,,$$

where \mathcal{N}_i is S^{∞} -open in $\mathcal{C}^{\infty}(V_i, M)$ for $i \in \Lambda'$ and $S^{d(r)}$ -open for $i \in \Lambda \setminus \Lambda'$. Moreover, the restriction of φ_{V_i} to \mathcal{M}_i is one-to-one.

Let $p_i : \mathcal{C}^{\infty}(M, \mathbb{R}) \to \mathcal{C}^{\infty}(V_i, \mathbb{R})$ and $q_i : \mathcal{C}^{\infty}(M, M) \to \mathcal{C}^{\infty}(V_i, M)$ be the "restriction to V_i " maps. Then we have the following commutative diagram

(8.2)
$$\begin{array}{ccc} \mathsf{func}(\mathbf{F}) & \stackrel{\varphi}{\longrightarrow} & Sh(\mathbf{F}) \\ & & & & \downarrow^{q_i} \\ & & & & \downarrow^{q_i} \\ & & \mathsf{func}(\mathbf{F}, V_i) & \stackrel{\varphi_{V_i}}{\longrightarrow} & Sh(\mathbf{F}, V_i) \end{array}$$

By definition φ and φ_{V_i} as surjective. It also follows from definitions of $\operatorname{func}(\mathbf{F})$ and $\operatorname{func}(\mathbf{F}, V_i)$ and assumption that V_i is a *D*-submanifold, that every $\sigma \in \mathcal{C}^{\infty}(V, \mathbb{R})$ extends to some $\bar{\sigma} \in \mathcal{C}^{\infty}(M, \mathbb{R})$. Moreover, if $\sigma \in \operatorname{func}(\mathbf{F}, V_i)$, we can assume that $\bar{\sigma} \in \operatorname{func}(\mathbf{F})$, whence $p_i(\operatorname{func}(\mathbf{F})) =$ $\operatorname{func}(\mathbf{F}, V_i)$. Therefore q_i is surjective as well. It follows from definition that

$$\mathcal{M} = \bigcap_{i \in \Lambda} p_i^{-1}(\mathcal{M}_i).$$

Put

(8.3)
$$\mathcal{N} \stackrel{\text{def}}{=\!\!=} \mathop{\cap}_{i \in \Lambda} q_i^{-1}(\mathcal{N}_i)$$

Since $\{V_i\}_{i\in\Lambda}$ is a locally finite cover and Λ' is finite², it follows from [13, Lm. 18] that \mathcal{N} is S^{∞} -open in $\mathcal{C}^{\infty}(M, M)$ and even $S^{d(r)}$ -open if $\Lambda' = \emptyset$. We will now show that

(8.4)
$$\varphi(\mathcal{M}) = Sh(\mathbf{F}) \cap \mathcal{N}.$$

This will complete our theorem.

It follows from (8.2) and (8.3) that $\varphi(\mathcal{M}) \subset Sh(\mathbf{F}) \cap \mathcal{N}$. Conversely, let $g \in Sh(\mathbf{F}) \cap \mathcal{N}$. Then $g|_{V_i} = q_i(g) \in Sh(\mathbf{F}, V_i) \cap \mathcal{N}_i = \varphi_{V_i}(\mathcal{M}_i)$ for all $i \in \Lambda$ and $g = \varphi(\beta')$ for some $\beta' \in \mathcal{C}^{\infty}(M, \mathbb{R})$. We have to find (possibly another) function $\beta \in \mathcal{M}$ such that $g = \varphi(\beta)$. Since the restriction of φ_{V_i} to \mathcal{M}_i is injective, $g|_{V_i} = \varphi_{V_i}(\beta_i)$ for a unique $\beta_i \in \mathcal{M}_i$.

It remains to show that $\beta_i = \beta_j$ on $V_i \cap V_j$ for all $i, j \in \Lambda$. Since $\{\operatorname{Int} V_i\}_{i \in \Lambda}$ is a cover of M as well, the family of functions $\{\beta_i\}_{i \in \Lambda}$ will define a unique smooth function $\beta \in \bigcap_{i \in \Lambda} p_i^{-1}(\mathcal{M}_i) = \mathcal{M}$ such that $\beta|_{V_i} = \mathcal{M}$

 β_i and $\varphi(\beta) = g$. This will prove (8.4) and complete our theorem.

Let $x \in \text{Int}V_i \cap \text{Int}V_j$ for some $i, j \in \Lambda$. Then $g(x) = \mathbf{F}(x, \beta_i(x)) = \mathbf{F}(x, \beta_j(x))$.

If x is a non-periodic regular point, then $\beta_i(x) = \beta_j(x)$.

If x is periodic of period θ_x , then $\beta_i(x) - \beta_j(x) = b\theta_x$ for some $b \in \mathbb{Z}$. But $|\beta_i(x) - \beta_j(x)| \leq |\alpha(x) - \beta_i(x)| + |\alpha(x) - \beta_j(x)| < 2\delta(x) < \theta_x$, whence b = 0.

Thus $\beta_i = \beta_j$ on $(\operatorname{Int} V_i \cap \operatorname{Int} V_j) \setminus \Sigma_F$. Since Σ_F is nowhere dense, we see that $\beta_i = \beta_i$ on all of $V_i \cap V_j$ as well.

9. Replacing M with an open subset

Theorem 8.2 reduces the verification of openness of the shift map to openness of a family of shift maps $\{\varphi_{V_i}\}$, where $\{V_i\}$ is any locally

²If Λ' were infinite, then \mathcal{N} would be open in the so-called very-strong topology, see [9]

29

finite cover of M by D-submanifolds. Our next aim is to "localize" the verification an openness of φ_V by replacing all the manifold M with an open neighbourhood W of V.

Let F be a vector field on $M, W \subset M$ be a connected, open subset, and $\mathbf{F}_W : W \times \mathbb{R} \supset \mathsf{dom}(\mathbf{F}_W) \to W$ be the local flow generated by the restriction $F|_W$ of F to W. Then

$$\mathsf{dom}(\mathbf{F}_W) \subset \mathsf{dom}(\mathbf{F}) \cap (W \times \mathbb{R})$$

and $\mathbf{F} = \mathbf{F}_W$ on $\mathsf{dom}(\mathbf{F}_W)$. Let $V \subset W$ be a *D*-submanifold. Then $\mathsf{func}(\mathbf{F}_W, V)$ is an S⁰-open subset of $\mathsf{func}(\mathbf{F}, V)$, and the corresponding shift map $\varphi_{W,V}$ of \mathbf{F}_W coincides with φ_V on $\mathsf{func}(\mathbf{F}_W, V)$, i.e. (9.1)

 $\varphi_{W,V} = \varphi_V|_{\mathsf{func}(\mathbf{F}_W,V)} : \mathsf{func}(\mathbf{F}_W,V) \to Sh(\mathbf{F}_W,V) \subset Sh(\mathbf{F},V).$

The following statement characterizes openness of φ_V via openness of $\varphi_{W,V}$.

Theorem 9.1. Let $V \subset W$ be a *D*-submanifold, and $r, s \in \overline{\mathbb{N}}_0$. Then the following conditions (A)-(C) are equivalent:

- (A) The shift map φ_V is $S^{r,s}$ -open;
- (B) The shift map $\varphi_{W,V}$ is $S^{r,s}$ -open and its image $Sh(\mathbf{F}_W, V)$ is S^s -open in $Sh(\mathbf{F}, V)$, i.e. there exists an S^s -open subset $\mathcal{N} \subset \mathcal{C}^{\infty}(V, W)$ such that

$$Sh(\mathbf{F}, V) \cap \mathcal{N} = Sh(\mathbf{F}_W, V).$$

(C) The shift map $\varphi_{W,V}$ is $S^{r,s}$ -open and there exists an S^{s} -neighbourhood \mathcal{N} of the identity inclusion $i_{V}: V \subset M$ in $\mathcal{C}^{\infty}(V, M)$ such that

$$Sh(\mathbf{F}, V) \cap \mathcal{N} \subset Sh(\mathbf{F}_W, V).$$

Proof. (A) \Rightarrow (B). Since func(\mathbf{F}_W, V) is an S⁰-open subset of func(\mathbf{F}, V) and φ_V is an S^{*r*,*s*}-open map, it follows that the restriction

$$\varphi_V|_{\mathsf{func}(\mathbf{F}_W,V)} = \varphi_{W,V}$$

is an $S^{r,s}$ -open map and its image $Sh(\mathbf{F}_W, V)$ is S^s -open in $Sh(\mathbf{F}, V)$. The implication (B) \Rightarrow (C) is evident.

 $(C) \Rightarrow (A)$. Suppose that $\varphi_{W,V}$ is $S^{r,s}$ -open. Then by (2) of Theorem 4.2 there exists an S^{s} -neighbourhood \mathcal{U} of the identity inclusion $i_{V}: V \subset W$ in $\mathcal{C}^{\infty}(V, W)$ and an $S^{s,r}$ -continuous section of $\varphi_{W,V}$ defined on $\mathcal{U}' = \mathcal{U} \cap Sh(\mathbf{F}_{W}, V)$:

$$\sigma: Sh(\mathbf{F}, V) \supset Sh(\mathbf{F}_W, V) \supset \mathcal{U}' \xrightarrow{\sigma} \mathsf{func}(\mathbf{F}_W, V) \subset \mathsf{func}(\mathbf{F}, V),$$

i.e., $\varphi_{W,V} \circ \sigma = \mathrm{id}_{\mathcal{U}'}$.

Since $Sh(\mathbf{F}, V) \cap \mathcal{N} \subset Sh(\mathbf{F}_W, V)$ and $\mathcal{C}^{\infty}(V, W)$ is an S^0 -open subset of $\mathcal{C}^{\infty}(V, M)$, we obtain that $\mathcal{U}'' = \mathcal{U}' \cap \mathcal{N}$ is an S^s -neighbourhood of $i_V : V \subset M$ in $\mathcal{C}^{\infty}(V, M)$. Moreover, φ_V coincides with $\varphi_{W,V}$ on func (\mathbf{F}_W, V) . Therefore σ is also a section of φ_V defined on \mathcal{U}'' . Then by (2) of Theorem 4.2 φ_V is $S^{r,s}$ -open. \Box

Emphasize that in this theorem W is an *arbitrary* open neighbourhood of V.

10. Openness of $Sh(\mathbf{F}_W, V)$ in $Sh(\mathbf{F})$. Regular case.

Let z be a regular point of F, i.e. $F(z) \neq 0$. In this section we present necessary and sufficient conditions for $W^{r,r}$ -openness of local shift map φ_V , where V is arbitrary small D-neighbourhood of z, see Theorem 10.2. As a consequence we will obtain the following

Theorem 10.1. Suppose that z is a regular point of F having one of the following properties:

- (a) z is non-periodic and non-recurrent;
- (b) z is periodic and the germ at z of its first recurrence map R: (B,z) \rightarrow (B,z) is periodic;
- (c) z is periodic and the tangent map $T_zR : T_zB \to T_zB$ has at least one eigen value λ such that $|\lambda| \neq 1$.

Then for any sufficiently small connected D-neighbourhood V of z the corresponding shift map φ_V of F is $W^{r,r}$ -open for all $r \ge 1$. Moreover, if z satisfies either (a) or (b), then φ_V is $W^{0,0}$ -open as well.

The proof will be given in $\S10.8$.

Since z is a regular point of F, there exist $\varepsilon > 0$, a neighbourhood W' of z, and a diffeomorphism

$$\eta: W' \to \mathbb{R}^{n-1} \times (-4\varepsilon, 4\varepsilon)$$

such that in the coordinates (y, s) on W' induced by η we have that

$$\mathbf{F}((y,s),t) = (y,s+t),$$

whenever $s, s + t \in (-4\varepsilon, 4\varepsilon)$. We will call W' a 4ε -flow-box at z, see Figure 3. Let also $W = \eta^{-1} (\mathbb{R}^{m-1} \times (-\varepsilon, \varepsilon)) \subset W'$ be the "central" ε -flow-box at z. For every D-submanifold $V \subset W$ denote

(10.1)
$$\mathcal{U}_V = \{ \alpha \in \mathsf{func}(\mathbf{F}, V) : \varphi_V(\alpha)(V) \subset W \}.$$

Then $\operatorname{func}(\mathbf{F}_W, V) \subset \mathcal{U}_V$ and

(10.2)
$$\varphi_V(\mathcal{U}_V) = Sh(\mathbf{F}, V) \cap \mathcal{C}^{\infty}(V, W)$$

is a W⁰-open neighbourhood of i_V in $Sh(\mathbf{F}, V)$.




FIGURE 3. Flow box

Let $p_1 : W' \to \mathbb{R}^{m-1}$ and $p_2 : W' \to (-4\varepsilon, 4\varepsilon)$ be the standard projections. Then we can define two maps $P_1 : \mathcal{U}_V \to \mathcal{C}^{\infty}(V, \mathbb{R}^{m-1})$ and $P_2 : \mathcal{U}_V \to \mathcal{C}^{\infty}(V, (-\varepsilon, \varepsilon))$ by

$$P_{1}(\alpha) = p_{1} \circ \varphi_{V}(\alpha) : V \xrightarrow{\varphi_{V}(\alpha)} W \cong \mathbb{R}^{m-1} \times (-\varepsilon, \varepsilon) \xrightarrow{p_{1}} \mathbb{R}^{m-1},$$

$$P_{2}(\alpha) = p_{2} \circ \varphi_{V}(\alpha) : V \xrightarrow{\varphi_{V}(\alpha)} W \cong \mathbb{R}^{m-1} \times (-\varepsilon, \varepsilon) \xrightarrow{p_{2}} (-\varepsilon, \varepsilon),$$

for $\alpha \in \mathcal{U}_{V}$. Thus $\varphi_{V}(\alpha) = (P_{1}(\alpha), P_{2}(\alpha)).$

Theorem 10.2. Let $V \subset W$ be a **connected** compact *D*-submanifold, and $\mathbf{0}: V \to \mathbb{R}$ be the zero function.

- (1) Then the map P_1 is locally constant with respect to the W^0 topology of \mathcal{U}_V and its image $P_1(\mathcal{U}_V) \subset \mathcal{C}^{\infty}(V, \mathbb{R}^{m-1})$ is at most countable.
- (2) The shift map $\varphi_V : \operatorname{func}(\mathbf{F}, V) \to Sh(\mathbf{F}, V)$ is $\mathsf{W}^{r,r}$ -open if and only if $P_1(\mathbf{0})$ is an isolated point of $P_1(\mathcal{U}_V)$ in $\mathcal{C}^{\infty}(V, \mathbb{R}^{m-1})$ with respect to the W^r topology.

Proof. (1) We need the following lemma:

Lemma 10.3. Let $\alpha, \beta \in \mathcal{U}_V$. Then one of the following conditions holds true:

- (i) $|\alpha(x) \beta(x)| < 2\varepsilon$ for all $x \in V$, (ii) $\alpha(x) - \beta(x) > 6\varepsilon$ for all $x \in V$.
- (iii) $\beta(x) \alpha(x) > 6\varepsilon$ for all $x \in V$.

Moreover, condition (i) implies that

(iv) $P_1(\beta) = P_1(\alpha)$ and $P_2(\beta) = P_2(\alpha) + \beta - \alpha$.

Proof. Define the following three open, mutually disjoint subsets of V:

 $K_1 = \{ x \in V : |\alpha(x) - \beta(x)| < 2\varepsilon \},\$ $K_2 = \{ x \in V : \alpha(x) - \beta(x) > 6\varepsilon \},\$ $K_3 = \{ x \in V : \beta(x) - \alpha(x) > 6\varepsilon \}.$

Let $x \in V$ and suppose that $|\beta(x) - \alpha(x)| \ge 2\varepsilon$, i.e. $x \notin K_1$. Since W is the central ε -flow-box of a 4ε -flow-box, and

$$\varphi_V(\alpha)(V) \cup \varphi_V(\beta)(V) \subset W,$$

we see that $|\beta(x) - \alpha(x)| > 6\varepsilon$, i.e. $x \in K_2 \cup K_3$. Thus $V = K_1 \cup K_2 \cup K_3$. Since V is connected and K_i are open and disjoint, we obtain that V coincides with one of them.

(i) \Rightarrow (iv) Denote $f = \varphi_V(\alpha)$, $g = \varphi_V(\beta)$, and $\tau(x) = \beta(x) - \alpha(x)$. Then (10.3)

$$g(x) = \mathbf{F}(x,\beta(x)) = \mathbf{F}\big(\mathbf{F}(x,\alpha(x)),\beta(x) - \alpha(x)\big) = \mathbf{F}\big(f(x),\tau(x)\big)$$

for all $x \in V$. Suppose that $|\tau| < 2\varepsilon$ on all of V. Recall that $\mathbf{F}(y,s;t) = (y,s+t)$ whenever $s,t,s+t \in (-4\varepsilon,4\varepsilon)$ and $y \in \mathbb{R}^{m-1}$. Since $|P_2(\alpha)|, |P_2(\beta)| < \varepsilon$, we obtain that $|P_2(\alpha) + \tau| < 3\varepsilon$ and

$$(P_1(\beta), P_2(\beta)) = g \xrightarrow{(10.3)} \mathbf{F}(P_1(\alpha), P_2(\alpha); \tau) = (P_1(\alpha), P_2(\alpha) + \tau).$$

Hence $P_1(\beta) = P_1(\alpha)$ and $P_2(\beta) = P_2(\alpha) + \beta - \alpha$.

Corollary 10.4. Define the following equivalence relation on \mathcal{U}_V by

$$\alpha \sim \beta$$
 if and only if $|\alpha - \beta| < 2\varepsilon$.

Then every class is W^0 -open and by (iv) of Lemma 10.3 the map P_1 is defined on the equivalence classes, whence P_1 is locally constant.

Moreover, there is a well-defined strict order on the equivalence classes of \sim : if A and B are two distinct classes of \sim then

$$A > B$$
 if and only if $\alpha - \beta > 6\varepsilon$

on V for some $\alpha \in A$ and $\beta \in B$. Hence there are at most countable many classes of \sim , and therefore the image of P_1 is at most countable.

Proof. We have only to show that the definition of ">" does not depend on a particular choice of α and β . Let $\alpha' \in A$ and $\beta' \in B$ be another functions. Then

$$\alpha' - \beta' = (\alpha' - \alpha) + (\alpha - \beta) + (\beta - \beta') > -2\varepsilon + 6\varepsilon - 2\varepsilon = 2\varepsilon.$$

Hence by Lemma 10.3, $\alpha' - \beta' > 6\varepsilon$ as well.

This corollary proves (1).

(2) Let A_0 be the equivalence class of ~ containing the zero function **0**, i.e. $A_0 = \{ \alpha \in \mathcal{U}_V : |\alpha| < 2\varepsilon \}$. Then it is easy to see that

 $A_0 = \mathsf{func}(\mathbf{F}_W, V), \quad \text{whence} \quad Sh(\mathbf{F}_W, V) = \varphi_{W,V}(A_0) = \varphi_V(A_0).$

32

Thus we obtain the following commutative diagram

Lemma 10.5. The image $Sh(\mathbf{F}_W, V)$ is W^r -open in $Sh(\mathbf{F}, V)$ for some $r \in \mathbb{N}_0$ if and only if $P_1(\mathbf{0}) = P_1(A_0)$ is an isolated point of the image im $P_1 = P_1(\mathcal{U}_V) \subset \mathcal{C}^{\infty}(V, \mathbb{R}^{m-1})$ with respect to the W^r topology.

Proof. Sufficiency. Suppose that $P_1(\mathbf{0})$ is an isolated point of im P_1 in the W^r topology, i.e. $P_1(\mathbf{0})$ is a W^r-open subset of im P_1 . Since p_1^* is W^{r,r}-continuous, we obtain that $Sh(\mathbf{F}_W, V) = \varphi_V(A_0)$ is W^r-open in $\varphi_V(\mathcal{U}_V)$ being W^r-open in $Sh(\mathbf{F}, V)$. Hence $Sh(\mathbf{F}_W, V)$ is W^r-open in $Sh(\mathbf{F}, V)$ as well.

Necessity. Conversely, suppose that $P_1(\mathbf{0})$ is not isolated in im P_1 with the W^r topology. We will construct a sequence of functions $\{\beta_i\}$ such that

- (i) $P_1(\beta_i) \neq P_1(\mathbf{0}),$
- (ii) the sequence $\{\varphi_V(\beta_i)\}$ converges to $i_V : V \subset M$ in the W^r topology.

It will follow from (i) that $\varphi_V(\beta_i) \notin Sh(\mathbf{F}_W, V) = \varphi_V(A_0)$, and in particular $\beta_i \notin A_0 = \mathsf{func}(\mathbf{F}_W, V)$ for all $i \in \mathbb{N}$. Then from (ii) we will obtain that $Sh(\mathbf{F}_W, V) = \varphi_{W,V}(A_0)$ is not a W^r-neighbourhood of i_V in $Sh(\mathbf{F}, V)$, i.e. $Sh(\mathbf{F}_W, V)$ is not W^r-open in $Sh(\mathbf{F}, V)$.

Since $P_1(\mathbf{0})$ is not isolated in im P_1 , there exists a sequence of classes $\{A_i\}$ distinct from A_0 such that their images $P_1(A_i)$ converge to $P_1(A_0)$ in the W^r topology.

It is easy to see that for every $i \in \mathbb{N}$ there exists a function $\beta_i \in A_i$ such that $P_2(\beta_i)(x,s) = P_2(\mathbf{0})(x,s) = s$. Indeed, take any $\beta'_i \in A_i$ and set $\beta_i = \beta'_i - P_2(\beta'_i)$.

We claim that the sequence $\{\beta_i\}_{i\in\mathbb{N}}$ satisfies conditions (i) and (ii). Property (i) holds since $P_1(\beta_i) = P_1(A_i) \neq P_1(A_0)$ for all $i \in \mathbb{N}$.

Moreover, we have that $P_2(\beta_i) = P_2(\mathbf{0}) = p_2 \circ i_V$ and $\{P_1(\beta_i)\} \subset \mathcal{C}^{\infty}(V, \mathbb{R}^{m-1})$ converges to $P_1(\mathbf{0}) = p_1 \circ i_V$ in W^r topology. Hence $\varphi_V(\beta_i)$ converges to i_V with respect to the W^r topology. This proves (ii). \Box

Lemma 10.6. The map $\varphi_{W,V}$ is $W^{r,r}$ -open for all $r \geq 0$.

Proof. Notice that

 $\mathsf{func}(\mathbf{F}_W,V) = \{ \alpha \in \mathcal{C}^\infty(V,\mathbb{R}) \ : \ |\alpha(y,s)| < \varepsilon, \ |s+\alpha(y,s)| < \varepsilon \},$

and

$$\varphi_{W,V}(\alpha)(y,s) = (y,\alpha(y,s) + s).$$

Hence

$$Sh(\mathbf{F}_W, V) = \{ f \in \mathcal{C}^{\infty}(V, W) : p_1 \circ f(y, s) = y, |p_2 \circ f| < \varepsilon \}.$$

and the inverse map $\varphi_{W,V}^{-1}: Sh(\mathbf{F}_W, V) \to \mathsf{func}(\mathbf{F}_W, V)$ is given by

$$\varphi_{W,V}^{-1}(f)(y,s) = p_2 \circ f(y,s) - s.$$

Evidently, this map is $W^{r,r}$ -continuous for all $r \in \overline{\mathbb{N}}_0$, whence $\varphi_{W,V}$ is $W^{r,r}$ -open.

Now statement (2) follows from 10.5, 10.6, and statement (B) of Theorem 9.1. Theorem 10.2 is completed. \Box

10.7. **Periodic case.** Let z be a periodic point of F. We will show that for a sufficiently small flow-box at z the images of \sim -classes defined in 10.4 can be described in terms of its first recurrence map only, see (10.5).

Let θ be the period of z, B be a codimension one open disk which transversally intersects the orbit of o_z at z, and $R : (B, z) \to (B, z)$ be the first recurrence map. Let also $\varepsilon < \theta/5$ and $\eta : W' \to \mathbb{R}^{m-1} \times (-4\varepsilon, 4\varepsilon)$ be a 4ε -flow-box at z. Decreasing W' we can assume, in addition, that

(10.4)
$$o_z \cap W' = \mathbf{F}(z \times (-4\varepsilon, 4\varepsilon)),$$

 $B \subset W$, and B is transversal to all orbits of W', so that the restriction $p_1|_B: W' \supset B \to \mathbb{R}^{m-1}$ is a diffeomorphism. Put

$$d = (p_1|_B)^{-1} \circ p_1 : V \xrightarrow{p_1} \mathbb{R}^{m-1} \xrightarrow{(p_1|_B)^{-1}} B.$$

Then d preserves the first coordinate.

Let $W \subset W'$ be the central ε -flow-box, $V \subset W$ be a connected D-neighbourhood of z, and A be the \sim -class of \mathcal{U}_V . Then it follows from (10.4) that there exists a unique $k \in \mathbb{Z}$ such that $|\alpha(z) - k\theta| < \varepsilon$ for every $\alpha \in A$, and

(10.5)
$$P_1(A) = p_1 \circ R^k \circ d : V \xrightarrow{d} B \xrightarrow{R^k} B \xrightarrow{p_1} \mathbb{R}^{m-1}.$$

35

10.8. **Proof of Theorem** 10.1. Due to Theorem 10.2 it suffices to find a 4ε -flow-box neighbourhood W' of z such that for every connected D-neighbourhood V of z, contained in the central ε -flow-box neighbourhood W, the image $P_1(\mathbf{0})$ is an isolated point of $P_1(\mathcal{U}_V)$ in the W^r topology, where $r \geq 0$ in the cases (a) and (b), and $r \geq 1$ in the case (c).

(a) Suppose that z is non-recurrent. Then there exist $\varepsilon > 0$, and a 4ε -flow-box neighbourhood W' of z such that $\mathbf{F}(z,t) \in W'$ iff $|t| < 4\varepsilon$. Let $W \subset W'$ be the central ε -flow-box and $V \subset W$ be a connected *D*-neighbourhood. Then it follows from Lemma 10.3 that there exists only one equivalence class of \sim . Hence im $P_1 = P_1(\mathbf{0})$, and thus $P_1(\mathbf{0})$ is isolated in im P_1 in any of \mathbf{W}^r topologies.

Suppose that z is periodic. Let B be a codimension one open disk which transversally intersects the orbit of o_z at z, $R: (B, z) \to (B, z)$ be the first recurrence map, and W', W, and V be such as in §10.7.

(b) If the germ of R at z is periodic of some period q, then we can assume that there exists an open neighbourhood $E \subset B$ of z such that R(E) = E and $d(V) \subset E$. Then it follows from (10.5) that the image of P_1 is finite and consists of q points. Hence $P_1(\mathbf{0})$ is isolated in im P_1 in any of W^r topologies.

(c) Suppose that the tangent map $T_zR : T_zB \to T_zB$ has at least one non-zero eigen vector $v \in T_zB \subset T_zV$ with eigen value λ such that $|\lambda| \neq 1$. Then $T_zR^k(v) = \lambda^k v$. Let $T_zp_1 : T_zB \to T_0\mathbb{R}^{m-1}$ be the tangent map of p_1 at z and $u = T_zp_1(v)$. Then for every class Acorresponding to some $k \in \mathbb{Z}$ we have that

$$T_z(P_1(A)) \stackrel{(10.5)}{=} T_z(p_1 \circ R^k \circ d)(v) = \lambda^k u.$$

Since $|\lambda| \neq 1$, we see that $P(\mathbf{0})$ is isolated in im P_1 in any of W^r topologies for $r \geq 1$.

11. Openness of $Sh(\mathbf{F}_W, V)$ in $Sh(\mathbf{F})$. Singular case

Let $W \subset M$ be an open subset and $V \subset W$ be a *D*-submanifold. Similarly to (10.1) put

(11.1)
$$\mathcal{U}_V = \{ \alpha \in \mathsf{func}(\mathbf{F}, V) : \varphi_V(\alpha)(V) \subset W \}.$$

Then $\operatorname{func}(\mathbf{F}_W, V) \subset \mathcal{U}_V$ and

(11.2)
$$\varphi_V(\mathcal{U}_V) = Sh(\mathbf{F}, V) \cap \mathcal{C}^{\infty}(V, W)$$

is a W⁰-open neighbourhood of i_V in $Sh(\mathbf{F}, V)$. We will now present a sufficient condition which guarantees that $\mathsf{func}(\mathbf{F}_W, V) = \mathcal{U}_V$. Due to (11.2) this will imply that

(11.3)

 $Sh(\mathbf{F}_W, V) \stackrel{\text{def}}{=} \varphi_V(\mathsf{func}(\mathbf{F}, V)) = \varphi_V(\mathcal{U}_V) = Sh(\mathbf{F}, V) \cap \mathcal{C}^\infty(V, W)$

is W^0 -open in $Sh(\mathbf{F}, V)$. In particular we prove (11.3) for vector fields of types (Z), (L), (H) and their products.

DEFINITION 11.1. Let $V \subset W$ be a subset. We will say that a pair (W, V) has **proper boundary intersection** property (p.b.i.) if for any $x \in V$, $a \in \mathbb{R}$, and T > 1 such that

- $\mathbf{F}(x, \tau a) \in W$ for $\tau \in [0, 1) \cup \{T\}$, but
- $\mathbf{F}(x, a) \in \operatorname{Fr}(W) = \overline{W} \setminus W$,

there exists $\tau' \in (1,T)$ such that $\mathbf{F}(x,\tau'a) \notin \overline{W}$, see Figure 4a).

Roughly speaking, if the orbit o_x of any $x \in V$ leaves W for a certain amount of time and comes back into W, then during this time it must leave the closure \overline{W} . In Figure 4b) the pair (W, V) does not satisfy p.b.i.




FIGURE 4.

The following lemma is the crucial implication of p.b.i.:

Lemma 11.2. Let $z \in W$, F(z) = 0, and $V \subset W$ be a connected *D*neighbourhood of z such that (W, V) has p.b.i. Then $\mathsf{func}(\mathbf{F}_W, V) = \mathcal{U}_V$.

Proof. Let $\alpha \in \mathcal{U}_V$, so $\mathbf{F}(x, \alpha(x)) \in W$ for all $x \in W$. We have to show that $\alpha \in \mathsf{func}(\mathbf{F}_W, V)$. It suffices to verify that the following subset of V:

 $K_{\alpha} = \{ x \in V : \mathbf{F}(x, s\alpha(x)) \in W \text{ for all } s \in I = [0, 1] \}$

coincides with all of V. Notice that $z \in K_{\alpha}$. Moreover, since W is open in M, it follows that K_{α} is open in V. Therefore it suffices to show that $V \setminus K_{\alpha}$ is open in V as well. Since V is connected, we will get that $K_{\alpha} = V$.

Let $x \in V \setminus K_{\alpha}$. Then $x = \mathbf{F}(x, 0 \cdot \alpha(x))$ and $\mathbf{F}(x, 1 \cdot \alpha(x))$ belong to Wand there exists $\tau_0 \in (0, 1)$ such that $\mathbf{F}(x, \tau \alpha(x)) \in W$ for all $\tau \in [0, \tau_0)$

while $\mathbf{F}(x, \tau_0 \alpha(x)) \in \operatorname{Fr}(W)$. Now it follows from p.b.i. for the pair (W, V) that there exists $\tau' \in (\tau_0, 1)$ such that $\mathbf{F}(x, \tau' \alpha(x)) \notin \overline{W}$.

Then there is an open neighbourhood V_x of x in V such that

$$\mathbf{F}(y,\tau'\,\alpha(y))\not\in\overline{W}$$

for all $y \in V_x$. Hence $V_x \subset V \setminus K_\alpha$, and thus $V \setminus K_\alpha$ is open in V. \Box

The following simple lemma is left for the reader.

Lemma 11.3. (1) If (W, V) has p.b.i., then for every subset $V' \subset V$ the pair (W, V') also has this property.

(2) Let W' be an open neighbourhood of \overline{W} . Then (W, V) has property p.b.i. with respect to F iff it has this property with respect to the restriction $F|_{W'}$. In other words, p.b.i. is determined by the behavior of F on arbitrary small neighbourhood of \overline{W} only.

(3) For every $x \in W$ denote by γ_x the connected component of $o_x \cap W$ containing x. Suppose that $\operatorname{Fr}(W)$ is a smooth submanifold of M and for every $x \in V$ its orbit o_x is transversal to $\operatorname{Fr}(W)$ at each $y \in \overline{\gamma_x} \cap$ $\operatorname{Fr}(W)$ whenever such a point exists. Then (W, V) has p.b.i.

11.4. **Isolating blocks.** Statement (3) of Lemma 11.3 is relevant with one of the principal results of [4]. An open subset $U \subset M$ is an *isolating neighbourhood* for the flow **F** if $o_x \not\subset \overline{U}$ for all $x \in \operatorname{Fr} U = \overline{U} \setminus U$. A closed **F**-invariant subset $X \subset M$ is *isolated*, if X is the maximal invariant subset of some isolating neighbourhood U of X.

Let W be a compact D-submanifold of M and $w = \overline{W} \setminus \text{Int}W$ be its boundary. Denote

$$\begin{array}{rcl} w^+ &=& \{x \in w \ : \ \exists \varepsilon > 0 \ \text{with} \ \mathbf{F}(x \times (-\varepsilon, 0)) \cap W = \varnothing\}, \\ w^- &=& \{x \in w \ : \ \exists \varepsilon > 0 \ \text{with} \ \mathbf{F}(x \times (0, \varepsilon)) \cap W = \varnothing\}, \\ \tau &=& \{x \in w \ : \ F \ \text{is tangent to} \ w\}. \end{array}$$

Then W is called an *isolating block* for \mathbf{F} if $w^+ \cap w^- = \tau$ and τ is a smooth submanifold of w with codimension one. Thus w^+ and $w^$ are submanifolds (possibly with corners is so is W) of w with common boundary τ . It follows that the interior of an isolating block is an isolating neighbourhood. Moreover, from (3) of Lemma 11.3 we obtain

Lemma 11.5. If W is an isolating block, then for any subset $V \subset IntW$ the pair (IntW, V) has p.b.i.

Theorem 11.6. [4] Let $X \subset M$ be a closed isolated \mathbf{F} invariant subset and $U \supset X$ be its isolating neighbourhood. Then there exists an isolating block W such that $X \subset \text{Int} W \subset U$.

This result was established for the case when W is a manifold with boundary, but if ∂M is **F**-invariant and $X \cap \partial M \neq \emptyset$, then the proof easily extends to *D*-submanifolds.

Corollary 11.7. Suppose z is an isolated singular point of F. Then there exist a base $\beta = \{W_{\alpha}\}_{\alpha \in A}$ of open neighbourhoods of $0 \in \mathbb{R}^k$ such that for every $W \in \beta$ and any subset $V \subset W$ the pair (W, V) has p.b.i.

11.8. Semi-invariant sets. We say that W is negatively (positively) invariant with respect to F if for every $x \in W$ and $t \leq 0$ ($t \geq 0$) such that $(x,t) \in \mathsf{dom}(\mathbf{F})$ we have $\mathbf{F}(x,t) \in W$. In this case we will also say that W is semi-invariant.

Lemma 11.9. If W is semi-invariant with respect to \mathbf{F} , then for any subset $V \subset W$ the pair (W, V) has p.b.i.

Proof. Notice that if $x, \mathbf{F}(x, Ta) \in W$ for some $a \in \mathbb{R}$ and T > 1, then it follows from semi-invariance of W that $\mathbf{F}(x, \tau a) \in W$ for all $\tau \in [0, T]$. Hence there are no $x \in V$, $a \in \mathbb{R}$ and T > 1 satisfying assumptions of Definition 11.1. Therefore (W, V) has p.b.i. \Box

11.10. **Product of flows.** For i = 1, 2 let M_i be a manifold, F_i be a vector field on M_i , $W_i \subset M_i$ be an open subset, and $V_i \subset W_i$ be a subset. Denote $M = M_1 \times M_2$ and $W = W_1 \times W_2$. Consider the product of these vector fields $F(x, y) = (F_1(x), F_2(y))$ on M. It generates a local flow $\mathbf{F}(x, y, t) = (\mathbf{F}_1(x, t), \mathbf{F}_2(y, t))$.

Lemma 11.11. Suppose that (W_i, V_i) has p.b.i. with respect to F_i , (i = 1, 2), and let $V \subset V_1 \times V_2$ be a subset. Then (W, V) has p.b.i. with respect to F.

Proof. Let $x = (x_1, x_2) \in V$, $a \in \mathbb{R}$, and T > 1 be such that

$$\mathbf{F}(x,\tau a) = (\mathbf{F}_1(x_1,\tau a), \mathbf{F}_2(x_2,\tau a)) \in W, \qquad \tau \in [0,1) \cup \{T\},\$$

 $\mathbf{F}(x,a) = (\mathbf{F}_1(x_1,a), \mathbf{F}_2(x_2,a)) \in \mathrm{Fr}W = (\mathrm{Fr}(W_1) \times W_2) \cup (W_1 \times \mathrm{Fr}(W_2)).$

In particular, $\mathbf{F}_i(x_i, a) \in \operatorname{Fr}(W_i)$ for at least one index i = 1, 2. For definiteness assume that $\mathbf{F}_1(x_1, a) \in \operatorname{Fr}(W_1)$. Since $x_1 \in V_1$, it follows from p.b.i. for (W_1, V_1) that there exists $\tau' \in (1, T)$ such that $\mathbf{F}_1(x_1, \tau'a) \notin \overline{W_1}$. Then $\mathbf{F}(x, \tau'a) \notin \overline{W}$ as well. Hence (W, V) has p.b.i. with respect to F.

Corollary 11.12. Let F be a product of vector fields of types (L) or (H) on \mathbb{R}^m . Then there exist a base $\beta = \{W_\alpha\}_{\alpha \in A}$ of open neighbourhoods of $0 \in \mathbb{R}^k$ such that for every $W \in \beta$ and any subset $V \subset W$ the pair

39

(W, V) has p.b.i. In particular, if V is a connected D-neighbourhood of 0, then func $(\mathbf{F}_W, V) = \mathcal{U}_V$ and thus by (11.2)

$$Sh(\mathbf{F}_W, V) = Sh(\mathbf{F}, V) \cap \mathcal{C}^{\infty}(V, W)$$

is W^0 -open in $Sh(\mathbf{F}, V)$.

Proof. Due to Lemma 11.11 it suffices to assume that F is of type (L) or (H). If F is of type (HE), then 0 has an arbitrary small invariant neighbourhoods W and by Lemma 11.9 (W, V) has p.b.i. for any subset $V \subset W$. If F is of type (HS) then $\{0\} \subset \mathbb{R}^2$ is an isolated invariant subset of any of its small neighbourhood, whence existence of W follows from Theorem 11.6, though it can easily be constructed without referring to this theorem.

It remains to consider the case when F(x) = Bx, where B is a Jordan cell corresponding either to some eigen value $a \in \mathbb{R}$ or to the pair of complex conjugate $a \pm ib$, $(a, b \in \mathbb{R})$.

1) If a > 0 (a < 0), then, e.g.[26], 0 has arbitrary small positively (negatively) **F**-invariant neighbourhoods W. Moreover, if $B = \left\| \begin{smallmatrix} 0 & -b \\ b & 0 \end{smallmatrix} \right\|$, then $0 \in \mathbb{R}^2$ has arbitrary small **F**-invariant neighbourhoods W. Then again by Lemma 11.9 such neighbourhoods have desired properties.

2) Suppose that $B = \left\| \begin{smallmatrix} 0 & 1 \\ 0 & 1 \\ 0 & 1 \\ 0 \end{bmatrix} \right\|$ is nilpotent. Then the coordinate functions of **F** are given by the following formulas:

$$\mathbf{F}_1(x,t) = x_1 + x_2 t + x_3 \frac{t^2}{2!} + \dots + x_m \frac{t^{m-1}}{(m-1)!}, \qquad \mathbf{F}_i = \frac{\partial \mathbf{F}_{i-1}}{\partial t},$$

for i = 2, ..., m. In particular, for each $x \in \mathbb{R}^m$ these functions are polynomials in t of degree $\leq m - 1$. Hence for every $\bar{r} > 0$ there exists T > 0 such that if $|x_0| < \bar{r}$ and x_0 is not a fixed point of **F**, then $\|\mathbf{F}(x,t)\|$ strictly monotone increases when increases t > T (decreases t < -T).

For every $r \ge 0$ let $V_r \subset \mathbb{R}^m$ be the closed *m*-disk of radius *r* centered at the origin and $W_r = \text{Int}(V_r)$ be its interior. Since $\mathbf{F}(0,t) = 0$ for all $t \in \mathbb{R}$, it follows that for every R > 0 there exists $r < \bar{r}$ such that $\mathbf{F}(V_r \times [-T,T]) \subset W_R$, i.e. $\|\mathbf{F}(x,t)\| < R$ for $\|x\| \le r$ and $|t| \le T$. We claim that the pair (W_R, V_r) has p.b.i.

Suppose that for some $x \in V_r$ and $a \in \mathbb{R}$ we have $\mathbf{F}(x, \tau a) \in W_R$ for $\tau \in [0, 1)$ and $\mathbf{F}(x, a) \in \operatorname{Fr}(W_R) = \partial V_R$, i.e. $\|\mathbf{F}(x, \tau a)\| < R$ and $\|\mathbf{F}(x, a)\| = R$. Then |a| > T, whence $\|\mathbf{F}(x, t)\|$ strictly monotone increases when increases |t|. In particular, $\|\mathbf{F}(x, \tau a)\| > R$ for all $\tau > 1$, i.e. $\mathbf{F}(x, \tau a) \notin V_R = \overline{W_R}$.

3) Suppose that $\lambda = ib$, $(b \neq 0)$, is purely imagine but $m = 2k \ge 4$ for some $k \ge 2$. Then regarding \mathbb{R}^m as \mathbb{C}^k we have that in complex

coordinates $A = \left\| \begin{array}{c} \stackrel{ib}{\ldots} \\ \stackrel{1}{\underset{ib}{ib}} \\ \stackrel{1}{\underset{ib}{ib}} \\ \stackrel{1}{\underset{ib}{ib}} \\ \end{array} \right\|$. Hence the coordinate functions of \mathbf{F} are given by formulae similar to the case 2). Denote $z = (z_1, \ldots, z_k)$ and $p(z,t) = z_1 + z_2t + z_3\frac{t^2}{2!} + \cdots + z_k\frac{t^{k-1}}{(k-1)!}$. Then

$$\mathbf{F}_1(z,t) = e^{ib}p(z), \qquad \mathbf{F}_i = e^{ib}\frac{\partial \mathbf{F}_{i-1}}{\partial t}, \qquad i = 2, \dots, k.$$

Since $|e^{ib}| = 1$, it follows that $||\mathbf{F}(x,t)||$ satisfies monotonicity conditions analogous to the case 2). Then by the similar arguments we obtain that for every R > 0 there exists r > 0 such that the pair (W_R, V_r) has property p.b.i.

12. Proof of Theorem 3.5

Let F be a vector field of class $\mathcal{F}(M)$. It follows from results of [18] that $Sh(\mathbf{F}) = \mathcal{E}_{id}(\mathbf{F})^k$, where k = 1 if $\Sigma \neq \Sigma^{(L)'} \cup \Sigma^{(HS)'}$ and k = 0 otherwise.

Therefore we should prove that the map $\varphi : \operatorname{func}(\mathbf{F}) \to Sh(\mathbf{F})$ is a homeomorphism with respect to S^{∞} topologies. Actually we want to apply Theorem 8.2.

Claim 12.1. Suppose that F belongs to class $\mathcal{F}(M)$.

- (1) If $z \in \Sigma$, then for any sufficiently small connected compact *D*neighbourhood *V* of *z* the shift map $\varphi_V : \operatorname{func}(\mathbf{F}, V) \to Sh(\mathbf{F}, V)$ is $W^{\infty,\infty}$ -open. Moreover, if $z \in \Sigma^{(L1)'} \cup \Sigma^{(L2)'}$, then φ_V is even $W^{r,r+2}$ -open for all r > 0.
- (2) If z is a regular point of F, then for any sufficiently small connected compact D-neighbourhood V of z the shift mapping $\varphi_V : \operatorname{func}(\mathbf{F}, V) \to Sh(\mathbf{F}, V)$ is $\mathsf{W}^{r,r}$ -open for all $r \geq 1$.

Proof. By (C) Theorem 9.1 it suffices to find a neighbourhood W of z such that for every connected D-neighbourhood $V \subset W$ of z

- (i) the shift map $\varphi_{W,V}$: func $(\mathbf{F}_W, V) \to Sh(\mathbf{F}_W, V)$ is $\mathsf{W}^{r,s}$ -open for the corresponding values r, s, and
- (ii) its image $Sh(\mathbf{F}_W, V)$ is W^k -open in $Sh(\mathbf{F})$, where k = 0 in the cases (1), and k = 1 in the case (2).

(1) Let $z \in \Sigma_F$. Then by definition of class $\mathcal{F}(M)$ there exists a neighbourhood W of z on which in some local coordinates either

- (a) F is a product of finitely many vector fields G_1, \ldots, G_n each of which is of types (Z), (L), or (H), or
- (b) F belongs to one of the types (L1)', (L2)', (L3)', or (H)' but z is an isolated singular point of F.

41

Decreasing W we can also assume that $Sh(\mathbf{F}_W, V)$ is W⁰-open in $Sh(\mathbf{F})$, i.e. condition (ii) is satisfied. In the case (a) this follows from Corollary 11.12, while in the case (b) from Corollary 11.7.

Then (i) directly follows from Lemmas 7.1-7.3.

(2) Suppose that z is a regular point for F. Since F belongs to $\mathcal{F}(M)$, it follows that the assumptions of Theorem 10.1 are satisfied, whence for any sufficiently small D-neighbourhood of z the shift map φ_V is $\mathsf{W}^{r,r}$ -open for all $r \geq 1$.

For every $z \in M$ let V_z be a neighbourhood guaranteed by Claim 12.1. By assumption of Theorem 3.5 the set $\overline{\Sigma \setminus (\Sigma^{(L1)'} \cup \Sigma^{(L2)'})}$ is compact.

Therefore using paracompactness of M we can find a locally finite cover $\{V_i\}_{i\in\Lambda}$ of M by compact connected D-submanifolds and a finite subset $\Lambda' \subset \Lambda$ such that the map φ_{V_i} is $\mathsf{W}^{\infty,\infty}$ -open for $i \in \Lambda'$ and $\mathsf{W}^{r,r+2}$ -open for all $r \geq 0$ whenever $i \in \Lambda \setminus \Lambda'$. Hence by "fragmentation" Theorem 8.2 φ is a local homeomorphism with respect to S^{∞} topologies. All other statements concerning homotopy types of $\mathcal{D}_{\mathrm{id}}(\mathbf{F})^r$ and $\mathcal{E}_{\mathrm{id}}(\mathbf{F})^r$ follow by the arguments of the proof of [13, §9]. Theorem 3.5 is completed.

13. Acknowledgements

I would like to thank V. V. Sharko, A. Prishlyak, I. Vlasenko, and I. Yurchuk for useful discussions. I am especially thank E. Polulyakh for very helpful conversations and pointing out to the irrational flow on T^2 which lead me to discovering mistakes of [13].

References

- Denis L. Blackmore: On the local normalization of a vector field at a degenerate critical point, J. Differential Equations 14 (1973), 338–359. MR MR0346245 (49 #10970)
- [2] A. D. Bryuno: Analytical form of differential equations, Trudy Moskov. Mat. Obšč. 25 (1971), 119–262.
- [3] R. V. Chacon: Change of velocity in flows, J. Math. Mech. 16 (1966), 417–431.
 MR MR0207955 (34 #7768)
- [4] C. Conley and R. Easton: Isolated invariant sets and isolating blocks, Trans. Amer. Math. Soc. 158 (1971), 35–61. MR MR0279830 (43 #5551)
- [5] Leonhard Frerick: Extension operators for spaces of infinite differentiable Whitney jets, J. Reine Angew. Math. 602 (2007), 123–154. MR MR2300454 (2008h:58015)
- [6] David Hart: On the smoothness of generators, Topology 22 (1983), no. 3, 357–363. MR MR710109 (85b:58105)
- Morris W. Hirsch: *Differential topology*, Graduate Texts in Mathematics, vol. 33, Springer-Verlag, New York, 1994, Corrected reprint of the 1976 original. MR MR1336822 (96c:57001)

SERGIY MAKSYMENKO

- [8] Eberhard Hopf: Ergodentheorie, Chelsea Publishing Co., Berlin, 1937.
- [9] Sören Illman: The very-strong C^{∞} topology on $C^{\infty}(M, N)$ and K-equivariant maps, Osaka J. Math. **40** (2003), no. 2, 409–428. MR MR1988698 (2004d:58052)
- [10] A. V. Kočergin: Change of time in flows, and mixing, Izv. Akad. Nauk SSSR Ser. Mat. 37 (1973), 1275–1298. MR MR0346129 (49 #10855)
- [11] V. A. Kondrat'ev and V. S. Samovol: The linearization of an autonomous system in the neighborhood of a singular point of "knot" type, Mat. Zametki 14 (1973), 833–842. MR MR0338503 (49 #3267)
- [12] Masasi Kowada: The orbit-preserving transformation groups associated with a measurable flow, J. Math. Soc. Japan 24 (1972), 355–373. MR MR0306451 (46 #5577)
- [13] Sergiy Maksymenko: Smooth shifts along trajectories of flows, Topology Appl. 130 (2003), no. 2, 183–204. MR MR1973397 (2005d:37035)
- [14] Sergiy Maksymenko: Hamiltonian vector fields of homogeneous polynomials in two variables, Pr. Inst. Mat. Nats. Akad. Nauk Ukr. Mat. Zastos. 3 (2006), no. 3, 269–308, arXiv:math/0709.2511.
- [15] Sergiy Maksymenko: Homotopy types of stabilizers and orbits of Morse functions on surfaces, Ann. Global Anal. Geom. 29 (2006), no. 3, 241–285. MR MR2248072 (2007k:57067)
- Sergiy Maksymenko: Stabilizers and orbits of smooth functions, Bull. Sci. Math. 130 (2006), no. 4, 279–311. MR MR2237445 (2007d:58066)
- [17] Sergiy Maksymenko: Connected components of partition preserving diffeomorphisms, Methods Funct. Anal. Topology 15 (2009), no. 3, 264–279.
- [18] Sergiy Maksymenko: Image of a shift map along the oribts of a flow, submitted (2009), arXiv:math/0902.2418.
- [19] Sergiy Maksymenko: ∞-jets of diffeomorphisms preserving orbits of vector fields, Cent. Eur. J. Math. 7 (2009), no. 2, 272–298.
- [20] Sergiy Maksymenko: Reparametrization of vector fields and their shift maps, Pr. Inst. Mat. Nats. Akad. Nauk Ukr. Mat. Zastos. 6 (2009), no. 2, 489–498, arXiv:math/0907.0354.
- [21] Sergiy Maksymenko: Symmetries of center singularities of plane vector fields, to appear in Nonlinear Oscilations (2009), arXiv:math/0907.0359.
- [22] Juan Margalef Roig and Enrique Outerelo Domínguez: Differential topology, North-Holland Mathematics Studies, vol. 173, North-Holland Publishing Co., Amsterdam, 1992, With a preface by Peter W. Michor. MR MR1173211 (93g:58005)
- [23] John N. Mather: Stability of C[∞] mappings. I. The division theorem, Ann. of Math. (2) 87 (1968), 89–104. MR MR0232401 (38 #726)
- [24] B. S. Mitjagin: Approximate dimension and bases in nuclear spaces, Uspehi Mat. Nauk 16 (1961), no. 4 (100), 63–132. MR MR0152865 (27 #2837)
- [25] Mark Alan Mostow and Steven Shnider: Joint continuity of division of smooth functions. I. Uniform Lojasiewicz estimates, Trans. Amer. Math. Soc. 292 (1985), no. 2, 573–583. MR MR808738 (87f:58018a)
- [26] Jacob Palis, Jr. and Welington de Melo: Geometric theory of dynamical systems, Springer-Verlag, New York, 1982, An introduction, Translated from the Portuguese by A. K. Manning. MR MR669541 (84a:58004)

- [27] William Parry: Cocycles and velocity changes, J. London Math. Soc. (2) 5 (1972), 511–516. MR MR0320276 (47 #8815)
- [28] R. T. Seeley: Extension of C[∞] functions defined in a half space, Proc. Amer. Math. Soc. 15 (1964), 625–626. MR MR0165392 (29 #2676)
- [29] Carl Ludwig Siegel: Uber die Normalform analytischer Differentialgleichungen in der Nähe einer Gleichgewichtslösung, Nachr. Akad. Wiss. Göttingen. Math.-Phys. Kl. Math.-Phys.-Chem. Abt. **1952** (1952), 21–30. MR MR0057407 (15,222b)
- [30] Shlomo Sternberg: Local contractions and a theorem of Poincaré, Amer. J. Math. 79 (1957), 809–824. MR MR0096853 (20 #3335)
- [31] Haruo Totoki: *Time changes of flows*, Mem. Fac. Sci. Kyushu Univ. Ser. A **20** (1966), 27–55. MR MR0201606 (34 #1488)
- [32] Richard J. Venti: Linear normal forms of differential equations, J. Differential Equations 2 (1966), 182–194. MR MR0188518 (32 #5956)
- [33] Hassler Whitney: Analytic extensions of differentiable functions defined in closed sets, Trans. Amer. Math. Soc. 36 (1934), no. 1, 63–89. MR MR1501735

TOPOLOGY DEPT., INSTITUTE OF MATHEMATICS OF NAS OF UKRAINE, TERESHCHENKIVS'KA ST. 3, KYIV, 01601 UKRAINE

E-mail address: maks@imath.kiev.ua