

EVOLUTION OF THE FIRST STARS WITH DARK MATTER BURNING

SUNG-CHUL YOON¹

UCO/Lick Observatory, Department of Astronomy and Astrophysics, University of California, Santa Cruz, CA 95064, USA,

FABIO IOCCO²

INAF – Osservatorio Astrofisico di Arcetri, Largo Enrico Fermi 5, Firenze 50125, Italy,

AND

SHIZUKA AKIYAMA³

Stanford Linear Accelerator Center and Kavli Institute for Particle Astrophysics and Cosmology, Stanford, CA 94305, USA,

Draft version July 19, 2021

ABSTRACT

Recent theoretical studies have revealed the possibly important role of the capture and annihilation process of weakly interacting massive particles (WIMPs) for the first stars. Using new evolutionary models of metal-free massive stars, we investigate the impact of such “dark matter burning” for the first stars in different environments of dark matter (DM) halos, in terms of the ambient WIMP density (ρ_χ). We find that, in agreement with existing literature, stellar life times can be significantly prolonged for a certain range of ρ_χ (i.e., $10^{10} \lesssim \rho_\chi [\text{GeVcm}^{-3}] \lesssim 10^{11}$ with the current upper limit for the spin-dependent elastic scattering cross section $\sigma_0^{\text{SD}} = 5 \times 10^{-39} \text{cm}^2$). This greatly enhances the role of rotationally induced chemical mixing in rotating stars, in favor of abundant production of primary nitrogen, massive helium stars and long gamma-ray bursts, from the first stars. We also find that stars with $\rho_\chi > 2 \times 10^{11} \text{GeVcm}^{-3}$ may not undergo nuclear burning stages, confirming the previous work, and that ionizing photon fluxes from such DM supported stars are very weak. Delayed metal enrichment and slow reionization in the early universe would have resulted if most of the first stars had been born in DM halos with such high ρ_χ , unless it had been lowered significantly below the threshold for efficient DM burning on a short time scale.

Subject headings: dark matter – early universe – stars: evolution – stars: rotation

1. INTRODUCTION

In the current framework of the Λ CDM cosmological model, the possibility that dark matter (DM) annihilation effects may impact stellar evolution has recently received renewed attention. After the original works of Bouquet, Dearborn, Freese, Gould, Griest, Krauss, Olive, Press, Raffelt, Renzini, Salati, Silk, Spergel, Srednicki and Wilczek in the ‘80s and early ‘90s, several authors have recently re-examined the effects that DM, if made of weakly interacting massive particles (WIMPs), would have on compact objects (Moskalenko & Wai 2007; Bertone & Fairbairn 2008) and on the zero-age main sequence of low-mass stars (Fairbairn et al. 2008). This exciting activity has been motivated by a double scope: any peculiar and distinguishable feature of WIMP annihilation on observable stellar quantities is extremely precious in the “quest” for dark matter evidence; on the other hand, all possible effects impacting the life of celestial objects must be taken into account by astrophysicists in the current precision era.

In particular, the first stellar episode at high redshift occurs under very different conditions from those in the present universe. The higher concentration of dark matter, the short Hubble time which prevents DM self-annihilation from severely affecting the central density, and the characteristic formation of a single PopIII star in the center of the halo are the most favorable conditions for DM annihilation effects to be very efficient in the first stars. In their pioneering work, Spolyar et al.

(2008) first found the possibility of very high DM density in primordial halos at high redshift to such an extent that energy released from DM annihilation at the center may halt the gravitational collapse of the baryonic cloud, calling such a DM powered object a *dark* star. Iocco (2008, hereafter I08) and Freese et al. (2008a) also noticed that WIMP capture is most efficient in Population III stars. More recently, Freese et al. (2008b,c) and Iocco et al. (2008) further investigated the role of annihilation of adiabatically contracted DM in the formation of the first stars. Iocco et al. (2008) have also followed the evolution from the pre-main sequence phase to helium exhaustion in the presence of WIMP capture and annihilation (hereafter, DM burning), showing that this can severely delay the evolution of pre-MS objects in the early universe, as well as extend their MS lifetimes.

All of these studies motivate us to explore possible consequences of DM burning for the final fate of the first stars and their feedback effects on the evolution of the early universe, even if the role of DM annihilation in the formation of the first stars still remains subject to many uncertainties (see First Stars III proceedings 2008, for a recent review). In this Letter, we address the issue by discussing the evolution of the first stars of $20 \leq M/M_\odot \leq 300$ up to the carbon burning stage (Sect. 2). We also investigate the interplay of rotation with DM burning in the evolution of the first stars of $100 M_\odot$, given the particular importance of rotation for the evolution and deaths of metal poor massive stars (e.g. Meynet et al. 2008; Yoon et al. 2008). Implications

of our results for the history of reionization in the early universe are briefly discussed (Sect. 3).

2. PHYSICAL ASSUMPTIONS AND RESULTS

We have implemented the DM capture and annihilation process in a hydrodynamic stellar evolution code, following Gould (1987). The DM capture rate C_* is calculated using Gould's equations as reported in Eqs. (1,2) of I08. Throughout this Letter we assume the DM-baryon scattering cross section σ_0 is 5×10^{-39} cm² for the spin-dependent scattering, to which only hydrogen is sensitive, and 10^{-43} cm² for the spin-independent one. These correspond to the current upper limits of WIMP direct detection search (Desai et al. 2004; Angle et al. 2008). We adopt the same values for other parameters as those used in I08 to calculate C_* , except for the ambient DM density (see below). The WIMPs captured by a star eventually reach thermal equilibrium with the gas, in a configuration dictated by the gravitational potential: the consequent DM density can be given by $n_\chi(r) = n_\chi^c \exp(-r^2/r_\chi^2)$, where $r_\chi = c(3kT_c/2\pi\rho_c m_\chi)^{1/2}$ (Griest & Seckel 1987). Here T_c and ρ_c are temperature and density at the stellar center, c and k are the speed of light and Boltzmann's constant, respectively. The energy generation rate due to DM annihilation is given by

$$\epsilon_\chi(r) = \frac{2}{3} \langle \sigma v \rangle n_\chi^2(r) m_\chi \text{ [erg cm}^{-3} \text{ s}^{-1}] \text{ ,} \quad (1)$$

where m_χ is the mass of the DM particle; we assume it to be $m_\chi=100$ GeV, which is often taken as a fiducial value in astrophysical studies for DM search. We use $\langle \sigma v \rangle = 3 \times 10^{-26}$ cm³ s⁻¹, the value best fitting the relic DM abundance (see, e.g., Bertone et al. 2005, for a recent review). The factor 2/3 is to consider that part of the energy is carried away by neutrinos. We consider the time dependent evolution of the total number of WIMPs (N_{tot}) in the star, and the number of thermally relaxed WIMPs in the core ($N_{\text{th}} := \int n(r)dV$) by the following equations in order to normalize $n(r)$:

$$\frac{dN_{\text{tot}}}{dt} = C_* - \int n(r)^2 \langle \sigma v \rangle dV, \text{ and} \quad (2)$$

$$\frac{dN_{\text{th}}}{dt} = \Gamma_{\text{th}} - \int n(r)^2 \langle \sigma v \rangle dV, \quad (3)$$

where Γ_{th} is the thermalization rate that can be approximated by

$$\Gamma_{\text{th}} = \frac{N_{\text{tot}} - N_{\text{th}}}{\tau_{\text{th}}}, \quad \tau_{\text{th}} = \frac{4\pi}{3\sqrt{2G}} \frac{m_\chi R^{7/2}}{\sigma_0 M^{3/2}}. \quad (4)$$

See Iocco et al. (2008) for detailed discussion on the thermalization time scale τ_{th} . Our models show that equilibrium between C_* and Γ_{th} is well maintained up to the core helium burning phase (see Fig. 2 below).

We consider several different values for the ambient DM energy density ρ_χ ranging from 0 to 2×10^{12} GeV cm⁻³. For the initial composition of the first stars, we assume the mass fractions of ¹H, ⁴He and ³He to be 0.76, 0.23999, and 0.00001, respectively. The mass loss rate from metal free stars is assumed to be zero if the Eddington factor Γ_E is smaller than 0.84 and 10^{-14} M_⊙ yr⁻¹ otherwise, following Krtićka & Kubát (2006). If a star becomes a helium star by rotationally induced mixing (see below), we assume that the




FIG. 1.— *Upper panel:* HR diagram of the non-rotating first star models on the ZAMS of different masses for different adopted values of $\sigma_0^{\text{SD}} \rho_\chi$ as indicated by the labels. Here σ_0^{SD} is the spin dependent WIMP scattering cross section, and ρ_χ is the ambient WIMP density. Stars in the grey shaded region are supposed to be only powered by DM burning, without nuclear reactions (i.e., $T_c < 10^7$ K). We use $\sigma_0^{\text{SD}} = 5 \times 10^{-39}$ cm². *Lower panel:* Life times of the non-rotating first stars as a function of the initial mass for different values of $\sigma_0^{\text{SD}} \rho_\chi$ given in the unit of 10^{-26} GeVcm⁻³. These life times are obtained from stellar models up to carbon burning for $\rho_\chi \leq 4 \times 10^{10}$ GeVcm⁻³ (i.e., $\sigma_0^{\text{SD}} \rho_\chi \leq 0.02 \times 10^{-26}$ GeVcm⁻³), while only approximate estimates are given for $\rho_\chi \geq 10^{11}$ GeVcm⁻³ (i.e., $\sigma_0^{\text{SD}} \rho_\chi \geq 0.05 \times 10^{-26}$ GeVcm⁻³).

mass loss rate from such a metal free Wolf-Rayet (WR) star is the same as that from a corresponding WR star at $Z = 10^{-6}$, as implied by Vink & de Koeter (2005). The code also implements the transport of chemical species and angular momentum due to rotationally induced hydrodynamic instabilities and the Spruit-Taylor dynamo. Other details on the stellar evolution code are described in Yoon et al. (2006).

Fig. 1 shows the HR diagram of the constructed non-rotating first star models on the zero age main sequence (ZAMS), for different values of ρ_χ . Note that the luminosity for a given mass does not significantly change with varying ρ_χ . This represents the well-known nature of the mass-luminosity relation, which is largely independent of any particular mode of energy generation (Kippenhahn & Weigert 1990). The stellar structure is adjusted such that the ratio of the nuclear to the DM luminosity decreases with increasing ρ_χ , while the total luminosity remains nearly constant, leading to prolonged lifetimes of the stars¹ (Fig. 1). If ρ_χ is above a critical value ($\rho_{\chi, \text{crit}} \approx 2 \times 10^{11}$ GeVcm⁻³, see Fig. 1) at a given stel-

¹ For a given ρ_χ , our models give longer life times than those by Iocco et al. (2008). This is because these authors used an approximation for the DM capture rate C_* , while we integrate Eq (2) of I08 over the entire stellar structure.




FIG. 2.— *Upper Panel:* Evolution of the surface luminosity (solid line), neutrino luminosity (three-dotted-dashed line), nuclear luminosities due to hydrogen burning (dot-dashed line), helium burning (dotted line), carbon burning (long dashed line) and DM burning (short dashed line) in the non-rotating $100 M_{\odot}$ model sequence. The contribution of DM annihilation to the neutrino luminosity, which corresponds to $1/3L_{\text{DM}}$, is not included here. The rapid increase in L_{H} and L_{He} after helium exhaustion is due to the hydrogen and helium shell burning. *Lower Panel:* Evolution of the WIMP capture rate (C_* , solid line), the thermalization rate (Γ_{th} , dashed line) and the number of thermalized WIMPs (N_{th} , dotted line) in the corresponding model. See Eqs. (2), (3) and (4).

lar mass the central density and temperature decrease to such an extent that the stars would live forever on the ZAMS, having no nuclear reactions. This is in qualitative agreement with Iocco et al. (2008) who find that pre-MS evolution is “frozen” before reaching the ZAMS at high enough ρ_{χ} ; and with Fairbairn et al. (2008), who find (for $M \leq 4M_{\odot}$) that stars move rightward of the ZAMS locus on the HR, and eventually join the Hayashi track, if “fed” with increasing DM annihilation luminosities.

No meaningful change in the stellar structure according to different values of ρ_{χ} ($< \rho_{\chi, \text{crit}}$) is observed in non-rotating models. Since DM burning only occurs within a very small radius ($r_{\chi} \ll R_{\text{core}}$), stars with different ρ_{χ} at a given mass have similar amounts of energy flux from the core and produce helium cores of a similar size (e.g. $\sim 40 M_{\odot}$ from $100 M_{\odot}$ stars). The luminosity resulting from DM burning gradually increases early on the main sequence as the star expands, but continuously decreases in later stages since the significant reduction of the number of hydrogen atoms lowers the DM capture rate. Rapid increase of the stellar radius after helium exhaustion makes the thermalization time very long, leading to reduction of the number of thermalized WIMPs (see Fig. 2). The DM luminosity accordingly decreases further from $7 \times 10^5 L_{\odot}$ to about $10^5 L_{\odot}$ during the carbon burning phase, in the given example with $100 M_{\odot}$ (Fig. 2). Carbon burning and particularly neutrino cool-




FIG. 3.— Evolution of the internal structure of rotating $100 M_{\odot}$ star models without DM burning (upper panel) and with DM burning (lower panel, $\rho_{\chi} = 4 \times 10^{10} \text{ GeV cm}^{-3}$). Convective layers are hatched, and semi-convective layers are marked by red dots. The color shading indicates nuclear energy generation rates. The adopted initial rotational velocity at the equatorial surface is 132 km s^{-1} , which corresponds to 10 % of the Keplerian value.

ing ($L_{\nu} > 10^{10} L_{\odot}$) dominate the evolution at this stage as shown in Fig 2. As the evolution of the star beyond carbon exhaustion should also be governed by neutrino cooling and other nuclear reactions such as oxygen burning, the effect of DM burning on the pre-supernova structure must be minor. As the situation remains similar in the other models of $20 \leq M/M_{\odot} \leq 300$, we conclude that DM burning may not change the final fate of the non-rotating first stars.

It is noteworthy, however, that rotation can dramatically change the evolution with DM burning. Fluids in rotating massive stars are subject to various rotationally induced hydrodynamic instabilities such as Eddington-Sweet circulations, which can cause mixing of chemical species across the boundary between the hydrogen burning core and the radiative envelope. Such mixing is usually stabilized by the strong buoyancy potential due to the chemical stratification between the hydrogen burning core and the envelope. However, if chemical mixing can occur faster than the building-up of chemical gradients by nuclear burning (i.e., $\tau_{\text{mix}}/\tau_{\text{nuc}} < 1$), then (quasi-) chemical homogeneity can be maintained on the main sequence (so-called chemically homogeneous evolution) (Maeder 1987). The condition, $\tau_{\text{mix}}/\tau_{\text{nuc}} < 1$, can be met either by reducing τ_{mix} or increasing τ_{nuc} . Rapid rotation tends to do the former, and we find that DM burning tends to do the latter. This is because the mixing time scale due to Eddington-Sweet circulations remains

almost unchanged with increasing ρ_χ (for $\rho_\chi < \rho_{\chi,\text{crit}}$), while the nuclear burning time scale increases due to the DM burning. This explains the remarkable impact of DM burning in the evolution of our rotating models as shown in Fig. 3. As shown in the upper panel, rotation in this model ($v_{\text{init}}/v_{\text{Kepler}} = 0.1$) is not rapid enough to cause the strong mixing by itself without DM burning. With $\rho_\chi = 4 \times 10^{10} \text{ GeV cm}^{-3}$ (see lower panel), however, the $100 M_\odot$ star lives about 10 times longer than in the corresponding non-DM burning case, and the star undergoes the quasi-chemically homogeneous evolution even with slow initial rotation because of the DM burning effect on nuclear burning time scale (cf. Yoon et al. 2006). The star is thus gradually transformed into a massive helium star by the end of main sequence as almost all of the hydrogen atoms in the star are fused into heliums due to mixing.

3. DISCUSSION

Our results indicate that DM burning should not significantly alter our view on the final fate of the non-rotating first stars: pair-instability supernovae for $140 \lesssim M/M_\odot \lesssim 260$, and core-collapse events for other masses (Heger & Woosley 2002), although their life times may be significantly prolonged. However, the impact of DM burning appears more important for rotating stars. The quasi-chemically homogeneous evolution can be rather easily realized even with moderate rotation velocities with $10^{10} \lesssim \rho_\chi [\text{GeVcm}^{-3}] \lesssim 10^{11}$. Such evolution can lead to production of massive helium stars that emit large amounts of helium ionizing photons, as well as abundant production of primary nitrogen, as discussed in Yoon et al. (2008). Note also that the quasi-chemically homogeneous evolution scenario (CHES) is one of the favored ones for the production of long GRBs from metal poor stars (Yoon & Langer 2005; Woosley & Heger 2006). Our result therefore indicates that DM burning might promote the production of long gamma-ray bursts from the first stars of $12 \lesssim M/M_\odot \lesssim 60$ via the CHES channel (see Yoon et al. 2006).

DM burning in the first stars must have consequences in the history of reionization in the early universe. Table 1 lists the number of hydrogen and helium ionizing photons emitted from $100 M_\odot$ models. If $\rho_\chi \lesssim 2 \times 10^{11} \text{ GeVcm}^{-3}$, the total number of ionizing photons increases proportionally to the DM density, as a direct consequence of the life-prolonging effect of DM burning. For a given ρ_χ rotation does not significantly alter hydrogen ionizing photon counts nor the lifetime. However, for models that take path of the CHES, the helium ionizing photon counts is increased by more than a factor of 2 compared to the non-rotating case. If ρ_χ is very large, on the other hand, the surface temperature of the star drops significantly enough (Fig. 1) that the total number of ionizing photons is reduced even with much longer lifetimes due to the DM burning. For example, with $\rho_\chi = 2 \times 10^{12} \text{ GeVcm}^{-3}$, it would have to take ~ 10 Gyr to emit as many hydrogen ionizing photons as a non-DM burning counterpart. Therefore, if most of the first stars had been born with such high ρ_χ their contribution to reionization would have been dramatically

reduced. The effect of the temperature drop on the number of helium ionizing photons is even more prominent: with $\rho_\chi = 2 \times 10^{12} \text{ GeVcm}^{-3}$, it decreases by more than

TABLE 1
NUMBER OF HYDROGEN AND HELIUM IONIZING PHOTONS FROM
 $100 M_\odot$ STAR MODELS

$v_{\text{rot}}/v_{\text{K}}$	$\rho_\chi [\text{GeVcm}^{-3}]$	N_{H}	N_{He}	Duration
0.0	0.00	1.2×10^{64}	2.2×10^{62}	3.2 Myr
0.0	2×10^{10}	2.1×10^{64}	3.4×10^{62}	5.5 Myr
0.0	4×10^{10}	8.5×10^{64}	1.5×10^{63}	22.3 Myr
0.0	10^{11}	2.9×10^{65}	6.3×10^{62}	100.0 Myr*
0.0	2×10^{11}	2.0×10^{65}	1.7×10^{61}	100.0 Myr*
0.0	2×10^{12}	1.3×10^{62}	9.4×10^{43}	100.0 Myr*
0.1	0.00	1.5×10^{64}	2.5×10^{62}	3.4 Myr
0.1	2×10^{10}	2.7×10^{64}	5.2×10^{62}	6.0 Myr
0.1	4×10^{10}	8.7×10^{64}	3.8×10^{63}	19.6 Myr

*The numbers are calculated only for the first 100 Myr.

19 orders of magnitude compared to the other cases with lower ρ_χ . Future study on the history of helium ionization at high redshift might therefore be a strong probe of DM burning in the first stars.

In this study we assume that the background DM halo density stays constant throughout the stellar evolution. This assumption may be valid if the stellar life times are shorter than about 100 Myr – which is the expected merger timescale of DM halos at $z \sim 20$ (e.g., Lacey & Cole 1993) – as in our model sequences with $\rho_\chi \lesssim 4 \times 10^{10} \text{ GeVcm}^{-3}$. The evolution of DM burners in halos with higher ρ_χ , however, should be critically determined by the change of DM halo environments. If ρ_χ is sufficiently reduced due to merger events and/or to displacement of the star from the densest region of the DM halo, the DM burners will become “normal” stars, dominated by nuclear burning. The star may then die quickly as we expect for normal stars, or become a “born-again” DM burner if ρ_χ increases again in later stages for some reason. The detailed history of the feedback from the first stars (e.g. metal enrichment and reionization) on the evolution of the early universe may depend on the nature and evolution of DM halos where the first stars are formed. This issue should be addressed in future work.

As a note added in proof, we acknowledge that Taoso et al. (2008) independently report similar results about the effect of DM burning on the MS life time and some stellar properties, using a different numerical code. S. A. appreciates helpful discussion with M. Alvarez and M. Busha. S. A. is supported by a Department of Energy Contract to SLAC DE-AC3-76SF00515. F. I. is grateful to P. Scott and M. Taoso for helpful discussion. F. I. is supported by MIUR through grant PRIN-2006. S.C. Y. is grateful to C. Church for the help with the text, and to W. Hillebrandt and S. Woosley for supporting his visit to MPA, Garching, in June, 2008, where part of the manuscript has been prepared. S.C. Y. is supported by the DOE SciDAC Program (DOE DE-FC02-06ER41438).

REFERENCES

- Bertone, G., & Fairbairn, M., 2008, *Phys. Rev. D*, 77, 043515
- Desai, S., *et al.* [Super-Kamiokande Collaboration], 2004, *Phys. Rev. D*, 70, 109901
- Fairbairn, M., Scott, P., & Edsjo, J., 2008, *Phys. Rev. D*, 77, 047301
- Freese, K., Spolyar, D., & Aguirre, A., 2008a, (astro-ph/0802.1724)
- Freese, K., Gondolo, P., Sellwood, J.A., & Spolyar, D., 2008b, (astro-ph/0805.3540)
- Freese, K., Bodenheimer, P., Spolyar, D., & Gondolo, P., 2008c, (astro-ph/0806.0617)
- "First Stars III", 2008, AIP Conf. Proc., 990, T. Abel, A. Heger, and B. W. 'O. Shea eds
- Gould, A., 1987, *ApJ*, 321, 571
- Griest, K., & Seckel, D., 1987, *Nucl. Phys. B*, 283, 681
- Heger, A., & Woosley, S.E., 2002, *ApJ*, 567, 532
- Iocco, F., 2008, *ApJ*, 677, L1
- Iocco, F., Bressan, A., Ripamonti, E., Schneider, R., Ferrara, A., & Marigo, P., 2008, *MNRAS*, in press
- Kippenhahn, R., & Weigert, A., 1990, *Stellar Structure and Evolution*, Springer-Verlag
- Krtićka, J., & Kubát, J., 2006, *A&A*, 446, 1039
- Lacey, C., & Cole, S., 1993, *MNRAS*, 262, 627
- Maeder, A., 1987, *A&A*, 178, 159
- Meynet, G., Estrom, S., Maeder, A., Hirschi, R., Georgy, C., & Beffa, C., 2008, in *First Stars III*, AIP Conf. Proc., 990, T. Abel, A. Heger, and B. W. 'O. Shea eds
- Moskalenko, I.V., & Wai, L.L., 2007, *ApJ*, 659, L29
- O'Shea, B. W., Abel, T., Whalen, D. & Norman, M., 2005, *ApJ*, 628, 5
- Spolyar, D., Freese, K., & Gondolo, P., 2008, *Phys. Rev. Lett.*, 100, 051101
- Taoso, M., Bertone, G., Meynet, G., & Ekström, S., 2008, (astro-ph/0806.2681)
- Vink, J.S., & de Koter, A., 2005, *A&A*, 442, 587
- Woosley, S.E., & Heger, A., 2006, *ApJ*, 637, 914
- Yoon, S.-C., Cantiello, M., & Langer, N., 2008, in *First Stars III*, AIP Conf. Proc., 990, T. Abel, A. Heger, and B.W.'O. Shea eds.
- Yoon, S.-C. & Langer, N. 2005, *A&A*, 443, 643
- Yoon, S.-C., Langer, N. & Norman, C., 2006, *A&A*, 460, 199